Crouzeix: Folie à Deux

Crouzeix: Folie à Deux

Nic Berkopec

October 2026

bnic101.github.io/crouzeix-folie-a-deux

Two proofs, July 2026. Same method. Both credit ChatGPT.

Crouzeix: Folie à Deux


Numerical range

Numerical range

Def. Let $\mathcal{H}$ be a complex Hilbert space and $T \in \mathcal{B}(\mathcal{H})$ a bounded operator. The numerical range of T is

$$W(T) = \{\langle Tx, x\rangle : \ x \in \mathcal{H}, \ \|x\| = 1\}.$$

In finite dimension closed and convex (Toeplitz–Hausdorff). In infinite dimension convex, $\overline{W(T)} \supseteq \operatorname{conv} \sigma(T)$.



Orange: $\sigma(T)$. Blue: $W(T)$. Normal: $\operatorname{conv} \sigma(T)$. 2×2: ellipse, foci at the eigenvalues.

[3 min] $\langle Nx, x\rangle = x_2 \bar{x}_1$, modulus $\leq \frac{1}{2}$, every value in the disc attained. Minor axis of the ellipse: $|t_{12}|$ in Schur form. Hilbert space case: compress to Krylov subspaces. Website: more examples, $\|T\|/w(T)$.

Crouzeix: Folie à Deux

Spectral sets

[2 min] Spectral set: von Neumann's word.

Spectral sets

Def. A set $S \subseteq \mathbb{C}$ is called a K-spectral set for an operator $T \in \mathcal{B}(\mathcal{H})$ if

$$\|p(T)\| \leq K \sup_{z \in S} |p(z)| \qquad \forall\, p \in \mathbb{C}[z].$$

(S bounded, polynomially convex.)

von Neumann's inequality. The disc is a spectral set for any contraction:

$$\|p(T)\| \leq \sup_{z \in \mathbb{D}} |p(z)|, \qquad \|T\| \leq 1.$$

(Also the closed half plane if $\operatorname{Re} T = \frac{T+T^*}{2} \geq 0$.)

Crouzeix: Folie à Deux

Two proofs of von Neumann

von Neumann	Sz.-Nagy
function theory	dilation the
Blaschke products	unitary dilat
Möbius maps: $I - \varphi_a(T)^*\varphi_a(T) \geq 0$	$p(T) = P_{\mathcal{H}}$

[2 min] von Neumann: reduce to Blaschke products, Möbius maps by hand.

Sz.-Nagy: compression of a unitary. Same split for the two 2026 proofs.

Crouzeix: Folie à Deux

Crouzeix conjecture

[2 min] $\langle Nx, x\rangle = x_2\bar{x}_1$.

Crouzeix conjecture

Conjecture (Crouzeix 2004)

Let T be a bounded operator on a Hilbert space $\mathcal{H}$. The numerical range $\overline{W(T)}$ is a 2-spectral set for T.

Remark. This conjectured bound is sharp:

$$W\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} = D(0, \tfrac{1}{2}), \qquad p(z) = z,$$

$$\left\| \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \right\| = 1 = 2 \cdot \tfrac{1}{2} = 2 \sup_{D(0,1/2)} |p(z)|.$$

Crouzeix: Folie à Deux

Antecedents

Antecedents

- von Neumann: constant 1, only for contractions.
- Okubo–Ando 1975, Berger–Stampfli 1967: the disc is a 2-spectral set.
- Via 2-dilations: $W(T) \subset \overline{\mathbb{D}}$ iff $T^n = 2V^*U^nV$ (Berger).
- Crouzeix 2004: true for 2×2, equality only when $W(T)$ is a disc.

[3 min] Constant 2 for the disc is a theorem, not a triviality: Berger–Stampfli need $p(0) = 0$, Okubo–Ando remove it. 2×2: exact constant as a function of the eccentricity.

Crouzeix: Folie à Deux

Constants

Constants

Ω bounded convex, smooth boundary, $W(T) \subset \Omega$.

$$C_\Omega := \sup\{\|f(T)\| : \ W(T) \subset \Omega, \ \|f\|_\infty \le 1\}.$$

Delyon–Delyon	1999	$C_\Omega < \infty$	depends on Ω
Crouzeix	2004	2	only for 2×2
Crouzeix	2007	11.08	uniform in Ω
Crouzeix–Palencia	2017	$1+\sqrt{2}$	completely bounded
Jin	27 July 2026	2	function theory
Lorist–Schwenninger	Aug 2026	2	dilation theory

Technique (all of the above): the double layer potential.

[2 min] Delyon–Delyon: constant from diameter^2/area. Crouzeix 2007: hard estimates on the Cauchy transform. Crouzeix–Palencia: six pages, survives matrix-valued f. Nine years, then two proofs in a week.

Crouzeix: Folie à Deux

Double layer potential

[2 min] Twice the real part of the Cauchy kernel.

Double layer potential

$\Omega \subset \mathbb{C}$ open bounded convex, smooth boundary, $\overline{W(T)} \subset \Omega$. $A(\Omega)$: holomorphic on Ω, continuous on $\overline{\Omega}$. $C(\partial\Omega)$: continuous on $\partial\Omega$. Sup norms. For $\sigma \in \partial\Omega$ let $\nu = \nu(\sigma)$ be the outward pointing normal; for $z \in \Omega$ the double layer potential is

$$\mu(\sigma, z) = \frac{1}{2\pi}\Big(\underbrace{\frac{\nu}{\sigma - z}}_{\text{first layer}} + \underbrace{\frac{\bar{\nu}}{\bar{\sigma} - \bar{z}}}_{\text{second layer}} \Big)$$


$= \frac{1}{\pi} \operatorname{Re} \frac{\nu}{\sigma - z}$. First layer: the Cauchy kernel.

Crouzeix: Folie à Deux

Double layer potential

[4 min] Twice the real part of the Cauchy kernel. Mass 2 inside, 1 on the boundary: Gauss jump, which gives continuity of the Cauchy transform of $\bar{f}$ up to $\partial\Omega$ (Neumann). Only place $W(T) \subset \Omega$ is used. Website: drag z; push it to the boundary.

Double layer potential



- $\mu\, ds = \frac{1}{\pi}\, d\arg(\sigma - z)$: the angle at which z sees $d\sigma$.
- $\mu \geq 0$: z inside every supporting line.
- $\int_{\partial\Omega} \mu\, ds = 2$ for $z \in \Omega$; 1 for $z \in \partial\Omega$.

Why use? The Cauchy layer leads to a bound on $f(z)$. If we can control the conjugate we win. (Adjoint under the functional calculus.)

Crouzeix: Folie à Deux

Double layer potential, cont.

[2 min] $\mu(\sigma, T) \geq 0$: $\langle Tx, x\rangle \in \Omega$ gives $\operatorname{Re} \bar{\nu}(\sigma - T) \geq 0$; conjugate by the resolvent. Mass $2I$: Cauchy's formula plus its adjoint.

Double layer potential, cont.

Let

$$C(f,T) = \frac{1}{2\pi i} \int_{\partial\Omega} f(\sigma)(\sigma - T)^{-1}\, d\sigma \in \mathcal{B}(\mathcal{H})$$

and

$$S(f,T) = C(f,T) + C(\bar{f},T)^* \in \mathcal{B}(\mathcal{H}).$$

By definition S is the function calculus integration against the double layer potential $\mu(\sigma, z)$, i.e.

$$S(f,T) = \int_{\partial\Omega} f(\sigma)\, \mu(\sigma,T)\, ds,$$

with $\mu(\sigma,T) = \frac{1}{\pi} \operatorname{Re}[\nu(\sigma)(\sigma - T)^{-1}] \geq 0$, $\int \mu\, ds = 2I$.

Crouzeix: Folie à Deux

The question

The question

Lemma (Crouzeix–Palencia 2017)

For $f \in C(\partial\Omega)$,

$$\|S(f,T)\| = \|f(T) + C(\bar{f},T)^*\| \le 2\|f\|_{\partial\Omega}.$$

Proof: $\mu \ge 0$, $\int \mu\, ds = 2I$, Cauchy–Schwarz.

So the question we must answer is: how does $\|S(f,T)\|$ control $\|f(T)\|$? In particular, can we estimate $C(\bar{f},T)^*$ or make it vanish all together?

[3 min] $\mu(\sigma,T) \ge 0$: $\langle Tx, x\rangle \in \Omega$ gives $\operatorname{Re}\bar{\nu}(\sigma - T) \ge 0$; conjugate by the resolvent. Mass $2I$: Cauchy's formula plus its adjoint. First layer of $S(f,T)$ is $f(T)$; second layer is the adjoint of the Cauchy integral of $\bar{f}$, not of f.

Crouzeix: Folie à Deux

Positivity and the double layer formula

[2 min] Same identity, $\Phi = \frac{1}{2}S$.

Positivity and the double layer formula

Consider

$$2\Phi(f) := f(T) + C(\bar{f}, T)^* = f(T) + \alpha(f)(T)^*,$$

where

$$\alpha(f)(z) = \frac{1}{2\pi i} \int_{\partial\Omega} \frac{\overline{f(\sigma)}}{\sigma - z}\, d\sigma.$$

Note that $\alpha : A(\Omega) \to A(\Omega)$ is antilinear and $\alpha(1) = 1$.

Positivity of Φ says $2\|\Phi(f)\| \le 2\|f\|_{\partial\Omega}$: the result from the previous slide, found in Crouzeix–Palencia.

Crouzeix: Folie à Deux

Positivity and the double layer formula

[3 min] Website: $f(z)$ and the companion as vectors; $\Phi(f)(z)$ stays in the convex hull of $f(\partial\Omega)$. Disc: companion constant, so Berger–Stampfli is the identity read off. Extremal f is a Blaschke product (Crouzeix 2004, Nevanlinna–Pick).

Positivity and the double layer formula

Scalar case: $2\Phi(f)(z) = f(z) + \overline{\alpha(f)(z)}$, holomorphic plus anti-holomorphic.

Disc: $\alpha(f) = \overline{f(0)}$, a constant.

Crouzeix–Palencia: estimate the companion, $\|\alpha(f)\| \le \|f\|$. Gives $1 + \sqrt{2}$.

An alternate approach (not taken by either ChatGPT proof) is to "hack" the formula by studying extremal functions: f extremal, $\|f(T)\| \le \|2\Phi(f)\|$, which proves Crouzeix. (Bickel, Gorkin, Greenbaum, Overton.)

Crouzeix: Folie à Deux

Equivalent formulation

Equivalent formulation

Theorem (Twisted real part)

Let $\mathcal{A}$ be a unital commutative operator algebra, let $\theta : \mathcal{A} \to \mathcal{B}(\mathcal{H})$ be a bounded homomorphism and let $\alpha : \mathcal{A} \to \mathcal{A}$ be a unital antilinear map such that the unital linear map

$$\theta_\alpha : \mathcal{A} \to \mathcal{B}(\mathcal{H}), \qquad a \mapsto \tfrac{1}{2}\big(\theta(a) + \theta(\alpha(a))^*\big)$$

is contractive. Then $\|\theta\| \le 2$.

- Ransford–Schwenninger (2018): twisted real part $\Rightarrow$ Crouzeix.
- Michael Hartz (2026) proves the theorem using Jin's ChatGPT solution.

[2 min] $\theta(f) = f(T)$, α the Cauchy transform: $\theta_\alpha = \Phi$. Unital and contractive $\Rightarrow$ positive. Both proofs establish this: Hartz as stated, Lorist–Schwenninger with "completely positive".

Crouzeix: Folie à Deux

Crouzeix: Folie à Deux

Function theory

[3 min] H: Carathéodory function normalised at 0. Class C_ρ: $(\rho-2)I + 2\,\mathrm{Re}(I-zT)^{-1} \geq 0$; $\rho = 2$ is $w(T) \leq 1$.

Function theory

Theorem (Holomorphic → resolvent transfer; Jin, operator form Hartz)

Let $T \in \mathcal{B}(\mathcal{H})$ with $\sigma(T) \subset \overline{\mathbb{D}}$. Let $H : \mathbb{D} \to \mathcal{B}(\mathcal{H})$ be a holomorphic function such that $H(0) = I$ and $\mathrm{Re}\, H(z) \geq 0$ for all $z \in \mathbb{D}$. Assume that

$$H(z) - (I - zT)^{-1}$$

commutes with T^* for all $z \in \mathbb{D}$. Then $\|T\| \leq 2$.

$H(z) - (I-zT)^{-1} = 0$: $\mathrm{Re}(I-zT)^{-1} \geq 0$ iff $W(T) \subset \overline{\mathbb{D}}$ (Berger); conclusion is Okubo–Ando.

Crouzeix: Folie à Deux

Transfer ⇒ twisted real part (Hartz)

[4 min] $\operatorname{Re} c_z = (1-za)^{-*}(I-|z|^2a^*a)(1-za)^{-1} \geq 0$. For $\theta(f) = f(T)$, expanding c_w in powers of f: $H(w) - (I-wT)^{-1} = \sum_{m\geq 1} w^m \alpha(f^m)(T)^*$; the identity for all powers of f. Crouzeix–Palencia used it once.

Transfer ⇒ twisted real part (Hartz)

Let $a \in \mathcal{A}$, $\|a\| \leq 1$, $T = \theta(a)$. For $z \in \mathbb{D}$ the Cayley transform $c_z = (1+za)(1-za)^{-1}$ has $\operatorname{Re} c_z \geq 0$, and θ_α is positive, so

$$H(z) := \theta_\alpha(c_z)$$

is holomorphic with $H(0) = I$ and $\operatorname{Re} H(z) \geq 0$. Since θ is a homomorphism, $\theta(c_z) = 2(I-zT)^{-1} - I$, hence

$$H(z) - (I-zT)^{-1} = \tfrac{1}{2}\big(\theta(\alpha(c_z))^* - I\big),$$

the adjoint of an element of $\theta(\mathcal{A})$, which is commutative and contains T. So $H(z) - (I-zT)^{-1}$ commutes with T^*, and the transfer theorem gives $\|\theta(a)\| \leq 2$.

Crouzeix: Folie à Deux

Proof of the transfer theorem, I

[3 min] Functions vanishing at an operator point form an "ideal" when Δ commutes with W; kernel at 0 kills M_Δ since $\Delta(0) = 0$. Defect removed exactly, not estimated.

Proof of the transfer theorem, I

$R(z) = (I - zT)^{-1}$, $\Delta = H - R$. On $H^2 \otimes \mathcal{H}$ ($\mathcal{H}$-valued power series), $\operatorname{Re} H \geq 0$ gives $\operatorname{Re} M_H \geq 0$, and $M_H = M_R + M_\Delta$.

For W with $\sigma(W) \subset \mathbb{D}$, the functions vanishing at W:

$$N_W = \{F = \textstyle\sum F_n z^n : \ \sum W^n F_n = 0\},$$

and $\Gamma : \mathcal{H} \to N_W$ its reproducing kernel at 0.

- If $\Delta(z)$ commutes with W, then $M_\Delta N_W \subset N_W$.
- Since $\Delta(0) = 0$, $\Gamma^* M_\Delta \Gamma = 0$.

Take $W = \frac{1}{2}T^*$. Then

$$\Gamma^*(M_R + M_R^*)\Gamma = \Gamma^*(M_H + M_H^*)\Gamma \geq 0.$$

Crouzeix: Folie à Deux

Proof of the transfer theorem, II

[4 min] Functions vanishing at an operator point form an "ideal" when Δ commutes with W; kernel at 0 kills M_Δ since $\Delta(0) = 0$. Defect removed exactly, not estimated. Weights $4^{-n}, 2^{-n}$ from sampling radius $\frac{1}{2}$. Jin does this in coordinates: sample the Herglotz kernel at $\lambda_i/2$ and at 0. Appendix: $\Gamma = K_0 - K_W P^{-1}$, eigenvector argument, Stein $P = I + \frac{1}{4} T^* P T$.

Proof of the transfer theorem, II

Explicitly $\Gamma = K_0 - K_W P^{-1}$, with K_W the kernel map at W,

$$P = K_W^* K_W = \sum_{n \geq 0} 4^{-n} T^{*n} T^n,$$

$$Q = K_W^* K_{T^*} = \sum_{n \geq 0} 2^{-n} T^{*n} T^n.$$

The resolvent identity turns the last inequality into

$$4Y - YP - PY \geq 0, \qquad Y = Q - P \geq \tfrac{1}{4} T^* T.$$

If $Px = \alpha x$ with $\alpha > 2$, testing on x gives $(4 - 2\alpha)\langle Yx, x\rangle \geq 0$, so $Yx = 0$, so $Tx = 0$, so $Px = x$. Hence $P \leq 2I$. With $P = I + \frac{1}{4} T^* P T$,

$$T^* T \leq T^* P T = 4(P - I) \leq 4I.$$

Crouzeix: Folie à Deux

Crouzeix: Folie à Deux

Dilation theory

[2 min] Perturbed 2-dilation; error of any size, bounded in n, commuting with T.

Dilation theory

Lemma (Lorist–Schwenninger, ChatGPT)

Let $\mathcal{H}$ be a Hilbert space and $T \in \mathcal{B}(\mathcal{H})$. Suppose there exists a Hilbert space $\mathcal{K}$, a contraction $Q \in \mathcal{B}(\mathcal{K})$ and an isometry $V \in \mathcal{B}(\mathcal{H}, \mathcal{K})$ such that the operators

$$E_n := 2V^* Q^{*n} V - T^{*n}, \qquad n \in \mathbb{N},$$

are uniformly bounded and commute with T. Then $\|T\| \leq 2$.

$E_n = 0$: 2-dilation of T^*, $\|T\| \leq 2$.

Crouzeix: Folie à Deux

Lemma ⇒ Crouzeix

[4 min] A genuine dilation of Φ, a perturbed 2-dilation of $f(A)$, error exactly the companion. Commutes: same functional calculus. Bounded: Crouzeix 2007 had 9.08. Abstract form: θ_α completely positive $\Rightarrow \|\theta\| \le 2$ (Arveson, Stinespring).

Lemma ⇒ Crouzeix

Let $\mathcal{K} = L^2(\partial\Omega, ds; \mathcal{H})$ and $Vx = \frac{1}{\sqrt{2}}\mu(\cdot, A)^{1/2}x$; V is an isometry because $\int \mu\, ds = 2I$. For $\|f\|_\infty = 1$ let Q be multiplication by f, a contraction. Then

$$V^*Q^nV = \tfrac{1}{2}\int f^n \mu\, ds = \Phi(f^n),$$

and the identity for f^n, adjointed, gives with $T = f(A)$

$$E_n = 2\Phi(f^n)^* - f(A)^{*n} = \alpha(f^n)(A).$$

E_n commutes with $f(A)$ (both come from the functional calculus) and $\|E_n\| \le \|\theta\|\|\alpha\|$. The lemma gives $\|f(A)\| \le 2$. The bound $\|\alpha\| \le 1$ is not used.

Crouzeix: Folie à Deux

Proof of the lemma, I

[2 min] $m_1 = \mathrm{Re}\langle (f\alpha(f))(A)x, x\rangle$, central since Crouzeix–Palencia. $m_1 \geq 0$ would finish; false in general.

Proof of the lemma, I

Suppose $\kappa = \|T\| > 1$ and take a unit x with $T^*Tx = \kappa^2 x$ (ultrapower). Put $m_n = \mathrm{Re}\langle E_n T^n x, x\rangle$ and

$$u = Q^*VTx - \kappa Vx,$$

the failure of VTx to be a κ-eigenvector of Q^*.

From the identity for $n = 1$, V isometric and Q contractive:

$$\|u\|^2 \leq 2\kappa^2 - \kappa^3 - \kappa m_1.$$

This alone gives $\kappa \leq 1 + \sqrt{1 - m_1}$; $m_1 \geq -1$ gives $1 + \sqrt{2}$, and $m_1 < 0$ for some 3×3 matrices.

Crouzeix: Folie à Deux

Proof of the lemma, II

[4 min] u: failure of VTx to be a κ-eigenvector of Q^*. Upper bound: $n = 1$ only; $m_1 = \mathrm{Re}\langle (f\alpha(f))(A)x, x\rangle$, central since Crouzeix–Palencia. Lower bound: commutation plus the identity for $n+1$, summed with weights κ^{-n+1}. $m_1 \geq 0$ would finish; false in general. Appendix: both estimates.

Proof of the lemma, II

From the identities for all n and $E_n T = T E_n$, completing a square,

$$\kappa m_n - m_{n+1} \geq -\frac{\|u\|^2}{\kappa(\kappa-1)} \qquad (n \geq 1).$$

Multiply by κ^{-n+1} and sum; the left side telescopes:

$$\kappa m_1 \geq -\frac{\|u\|^2}{(\kappa-1)^2}.$$

With the upper bound,

$$\|u\|^2\big(1-(\kappa-1)^{-2}\big) \leq \kappa^2(2-\kappa),$$

impossible for $\kappa > 2$.

Crouzeix: Folie à Deux

Comparison

[3 min] Same positivity (Carathéodory function / Stinespring), same packaging of all powers, same companion, same commutation hypothesis; compression vs. telescoping.

Comparison

	function theory	dilation theory
positivity	$\operatorname{Re} H \geq 0$	$\Phi = V^* \pi(\cdot) V$
all powers	$H(w) = I + 2\sum_m w^m \Phi(f^m)$	E_n, all n
companion	$\sum_m w^m \alpha(f^m)(T)^*$	$\alpha(f^n)(T)$
used	commutes with T^*	commutes with T
removal	$\Gamma^* M_\Delta \Gamma = 0$	telescoping

Crouzeix–Palencia use $\|\alpha(f)\| \leq \|f\|$; that alone cannot give better than $1+\sqrt{2}$. Both new proofs use instead that $\alpha(f^n)(T)$ commutes with $f(T)$.

[3 min] Both papers say so explicitly. Fill in the reference.

Completely bounded

Conjecture

For every $m \geq 1$ and $F = [f_{ij}] \in M_m(A(\Omega))$, $\|F(T)\| \leq 2 \sup_{z \in \Omega} \|F(z)\|$.

- cb-norm ≤ 2; Paulsen: T similar to an operator with $\overline{W(T)}$ a complete spectral set.
- Known: $1 + \sqrt{2}$ (Crouzeix–Palencia); 2 for discs, weighted shifts (Crouzeix–Greenbaum 2025).
- $M_m(A(\Omega))$ not commutative: both new proofs need the companion to commute with $f(T)$. Crouzeix–Palencia need only $\|\alpha\| \leq 1$.

References

- S. Jin, *The numerical range is a 2-spectral set*, Preprints.org 202607.1919, 27 July 2026.
- M. Hartz, *Notes on the solution of Crouzeix's conjecture*, 31 July 2026.
- E. Lorist, F. L. Schwenninger, *A solution to Crouzeix's conjecture*, arXiv:2608.03841.
- M. Crouzeix, *Bounds for analytical functions of matrices*, IEOT 48 (2004).
- M. Crouzeix, C. Palencia, *The numerical range is a* $(1+\sqrt{2})$*-spectral set*, SIMAX 38 (2017).
- T. Ransford, F. L. Schwenninger, *Remarks on the Crouzeix–Palencia proof*, SIMAX 39 (2018).
- R. Clouâtre, M. Ostermann, T. Ransford, *An abstract approach to the Crouzeix conjecture*, JOT 90 (2023).
- K. Bickel, P. Gorkin, *Blaschke products, level sets, and Crouzeix's conjecture*, J. Anal. Math. 152 (2024).
- B. Delyon, F. Delyon, *Generalization of von Neumann's spectral sets*, Bull. SMF 127 (1999).

`bmic101.github.io/crouzeix-folie-a-deux`