

<https://www.mpa.mpg.de/galform/virgo/millennium/>

Calculation of Large Scale Structure Power Spectra in a Spherical-Fourier Bessel (SFB) Basis

Presented by Brandon Khek, Rice University



Mentors: Olivier Doré and Henry Gebhardt, JPL, Caltech

October 23, 2021



RICE

Matter power spectra are a statistical tool used to constrain cosmological expansion



Physical effects appear in the power spectrum and help us understand structure in the universe




Fig. 1: Depiction of RSD¹

Physical effects considered

- Redshift space distortions (RSD)
- Baryon acoustic oscillations (BAO)
- Non-Gaussianity (NG)







Fig. 2: Cartoon of BAO²

Fig. 3: NG representation




Next-generation galaxy surveys will give us a 3-D map of the universe

E.g. SPHEREx, Nancy Grace Roman Space Telescope




We developed code to calculate the SFB power spectrum to account for wide-angle effects in upcoming surveys







Project

$$P(k) \rightarrow C_\ell(k, k')$$





Clustering amplitude is amplified by RSD and BAO




Depiction of RSD

Non-Gaussianity is evidenced by more clustering on larger scales





Error arises from precision in our numerical integration


ℓ	k Range	Average % Error	Time (s)
2	0.01 to 0.3	$1.7 * 10^{-11}$	1.71
10	0.05 to 0.4	$1.1 * 10^{-4}$	1.27
50	0.15 to 0.5	$1.2 * 10^{-2}$	5.37



Compare to
obtain error



Analysis of modified gravity, performance improvements, and studying off-diagonal terms is underway



Journal of Computational and Applied Mathematics 64 (1995) 217–231

JOURNAL OF
COMPUTATIONAL AND
APPLIED MATHEMATICS

Evaluating infinite integrals involving Bessel functions of arbitrary order

S.K. Lucas*, H.A. Stone

Division of Applied Sciences, Harvard University, Cambridge, MA 02138, USA

Received 6 January 1994; revised 1 August 1994


Abstract

The evaluation of integrals of the form $I_n = \int_0^\infty f(x) J_n(x) dx$ is considered. In the past, the method of dividing an oscillatory integral at its zeros, forming a sequence of partial sums, and using extrapolation to accelerate convergence has been found to be the most efficient technique available where the oscillation is due to a trigonometric function or a Bessel function of order $n = 0, 1$. Here, we compare various extrapolation techniques as well as choices of endpoints in dividing the integral, and establish the most efficient method for evaluating infinite integrals involving Bessel functions of any order n , not just zero or one. We also outline a simple but very effective technique for calculating Bessel function zeros.

Keywords: Quadrature; Infinite integration; Bessel functions; Bessel zeros; ϵ -algorithm; mW transform

1. Introduction

Calculating integrals on $[0, \infty)$ with oscillatory integrands is a more difficult problem than that for the case of an eventually monotonic integrand. Various techniques have been proposed for calculating integrals of the form



Acknowledgements

Henry Gebhardt and Olivier Doré
Jet Propulsion Laboratory
Caltech Student-Faculty Programs



Main Takeaways




The matter power spectrum helps us understand the expansion of space and nature of the universe




We developed code to calculate the matter power spectrum in a spherical Fourier Bessel basis



This code provides a crucial step in the analysis of upcoming galaxy surveys like SPHEREx or NGRST



Spherical harmonics are the Fourier basis on the surface of sphere



Real (Laplace) spherical harmonics $Y_{\ell m}$ for $\ell = 0, \dots, 4$ (top to bottom) and $m = 0, \dots, \ell$ (left to right). Zonal, sectoral, and tesseral harmonics are depicted along the left-most column, the main diagonal, and elsewhere, respectively. (The negative order harmonics $Y_{\ell(-m)}$ would be shown rotated about the z axis by $90^\circ/m$ with respect to the positive order ones.)

https://en.wikipedia.org/wiki/Spherical_harmonics

Formula for the continuous SFB power spectrum

$$C_\ell(k, k') = \int dq \mathcal{W}_\ell(k, q) \mathcal{W}_\ell^*(k', q) P(q) \quad (1)$$

$$\begin{aligned} \mathcal{W}_\ell(k, q) = & \frac{2qk}{\pi} \int dr r^2 \phi(r) D(r) b(r, q) j_\ell(kr) \\ & \times e^{-\frac{1}{2}\sigma_u^2 q^2} \sum_{\Delta\ell} (\delta_{\Delta\ell,0}^K - \beta f_{\Delta\ell}^\ell) j_{\ell+\Delta\ell}(qr) \end{aligned} \quad (2)$$


Formula for the discrete SFB power spectrum

$$\langle \delta_{n\ell m} \delta_{n'\ell m} \rangle = \int \mathcal{W}_{n\ell}(q) \mathcal{W}_{n'\ell}(q) P(q) dq \quad (1)$$

$$\mathcal{W}_{n\ell}(q) = \sqrt{\frac{2q^2}{\pi}} \int_{r_{\min}}^{r_{\max}} g_{n\ell}(r) \Upsilon_{\ell}(r, q) r^2 dr \quad (2)$$

$$\Upsilon_{\ell}(r, q) = \phi(r) D(r) b(r, q) \sum_{\Delta\ell} (\delta_{\Delta\ell, 0}^K - \beta f_{\Delta\ell}^{\ell}) j_{\ell+\Delta\ell}(qr). \quad (3)$$

Why NG results in larger structures at large scales



<https://wwwmpa.mpa-garching.mpg.de/~komatsu/presentation/fnl.pdf>




FIG. 1: Slice through simulation outputs at $z = 0$ generated with the same Fourier phases but with $f_{NL} = -5000, -500, 0, +500, +5000$ respectively from top to bottom. Each slice is $375 h^{-1}$ Mpc wide, and $80 h^{-1}$ Mpc high and deep. We can easily match by eye much of the large scale structure; for example, an overdense region sits on the left, while an underdense region (void) falls on the right, in all panels. Note that for positive f_{NL} , overdense regions are more evolved and produce more clusters than their Gaussian counterparts, while underdense regions are less evolved (*e.g.* grid lines are still visible). For negative f_{NL} , underdense regions are more evolved, producing deeper voids, while overdense regions are less evolved, as illustrated by the grid lines apparent in the left of the top panel.

<https://arxiv.org/pdf/0710.4560.pdf>

Why higher multipole moments give longer runtimes

$$j_n(z) = \frac{z^n}{2^{n+1}n!} \int_0^\pi \cos(z \cos \theta) (\sin \theta)^{2n+1} d\theta.$$

Applications of our code

Theory

$P(\vec{k})$

\longrightarrow

Convert
to SFB

$C_\ell(k, k')$

\longleftarrow

Apply
data

Data

$n(\vec{r})$

Gauss-Legendre quadrature

$$\int_{-1}^1 f(x) dx \approx \sum_{i=1}^n w_i f(x_i)$$

Density contrast

$$\delta = \frac{\rho - \bar{\rho}}{\bar{\rho}}$$