DEPARTMENT OF BIOCHEMISTRY OLABISI ONABANJO UNIVERSITY REMO CAMPUS, IKENNE

the

2009/2010 HARMATTAN SEMESTER EXAMINATION

COURSE CODE : BCH 409 COURSE TITLE: GENETIC ENGINEERING

DATE: 15TH March, 2010 TIME: 2HR ONLY INSTRUCTION: PUT ON YOUR THINKING CAP ANSWER ANY FOUR (4)

OUESTIONS

a. Define a restriction endon clease map

- b. A particular DNA molecule of length 3.0kb is incubated separately with two restriction enzymes A and B. When the cleared DNA is electrophoresed, enzyme A cut the substite DNA into three fragments of lengths 1.7kb, 0.9kb, and 0.4kb respectively. While enzyme B has generated two fragments of 1.6kb and 1.4kb respectively. Double degests produced four fragments of lengths 1.2kb, 0.9kb, 0.5kb and 0.4kb respectively. Construct a restriction map for the DNA molecule showing the relative positions of the cleavage sites of A and B, and the distances in kilo bases between them.
 - 6. What stain has been used to visualize the bands in the gel after lectrophoresis?

How can you determine the sizes of the fragments on the gel? by using polymergianate get

Describe a procedure for cloming a full length cDNA of an eukaryotic mRNA 2, a which you wish to be expressed in E.coli

How would you establish that the cloned gene is expressed in the E. coli

Compare southern blotting technique with that of Northern blotting technique. 3. 11.

Compare sequencing gel with that of Agarose gel

Compare High stringency with that of Low stringency of hybridization.

- There was a case of cholera epidemics in igal a land. The causative agent was found to be present in a plasmid carried in the bacteria. As a scientist working in the medical laboratory, describe vividly how you will isolate and characterize the plasmid gene responsible for the epidemics.
 - Indicate with examples how the following are employed in Gene manipulations. 5
 - S-I Nuclease (a)
 - T4 DNA Ligase (b)
 - Linker Oligonucleotides
 - Terminal Transeferase

Klenow Polymerase