




FIX YOUR GUT

1. ELIMINATE SYMPTOMS
2. LOWER INFLAMMATION
3. REBUILD YOUR GUT

The **first step** has to begin with getting rid of symptoms. IBS has a major impact on a patient's life and they suffer with a lot of different symptoms, including significant pain and cramping.

There are three stages of IBS symptoms that a patient goes through.

Stage 1

In between bouts of symptoms. Your gut is currently 'not bad' and you have no pain. Your bowel movements may not be normal, but they aren't as bad as they've been.

Stage 2

Your gut is just starting to act up. You may be having some digestive discomfort. You can still tolerate some food. Your bowel movements are getting more abnormal.

Stage 3

Your gut is angry. You have pain or cramping. Anything you eat bothers your stomach. Your bowel movements are a mess. You get some pain relief with a bowel movement.

ELIMINATE SYMPTOMS

1. REMOVE ALLERGY

2. REST YOUR GUT

3. THE RIGHT SUPPLEMENTS

1. Remove Allergy

Almost everyone with IBS has food sensitivities or allergies. It can be hard to figure out what food triggers IBS symptoms, especially when you're in stage 3.

Most with IBS have delayed food sensitivities, meaning a food you eat today can bother you tomorrow.

The most common food sensitivities are dairy, eggs, gluten, wheat, soy, and fructose.

If you have IBS that hasn't been resolved yet, it's best to remove all of them, especially if you're in stage 3.

It's important to figure out your IBS triggers.

ELIMINATE SYMPTOMS

1. REMOVE ALLERGY

2. REST YOUR GUT

3. THE RIGHT SUPPLEMENTS

2. Rest Your Gut

This is important. If you have IBS then it's important that you, at the very minimum, fast for more hours than you eat.

If you are in stage 1, then at minimum you should be fasting for 13 hours and eating for 11 hours. This gives more time for digestion.

An example would be eating breakfast at 8 AM and having last food by 7PM. This is an 11 hour eating window. It allows for 13 hours of fasting.

If you are in stage 2, we recommend a longer fast and shorter eating window. Many do well with the 16:8 method, a 16 hour fast and an 8 hour eating window.

An example would be eating breakfast at 11 AM and having last meal around 7PM.

If you are in stage 3, then you have to fast longer out of necessity since all food hurts your gut. To help with a fast in this stage, you can try sipping broth or eating jello until your gut calms down.

ELIMINATE SYMPTOMS

1. REMOVE ALLERGY
2. REST YOUR GUT
3. THE RIGHT SUPPLEMENTS

3. The Right Supplements

Here's the supplements we recommend to eliminate IBS symptoms:



PROBIOTIC
COMPLEX



DIGESTIVE
ENZYME FORMULA



IBS FORMULA

1. **ELIMINATE SYMPTOMS**
2. **LOWER INFLAMMATION**
3. **REBUILD YOUR GUT**

Lower Inflammation

IBS sufferers have higher levels of CORTISOL. High cortisol will hurt your sleep, your energy, your metabolism and it increases inflammation, which leads to more gut symptoms.

We recommend our CORTISOL CONTROL.



1. ELIMINATE SYMPTOMS
2. LOWER INFLAMMATION
3. REBUILD YOUR GUT

Rebuild Your Gut

The same supplements that help eliminate symptoms will also help rebuild your gut. We designed them that way.



PROBIOTIC
COMPLEX



DIGESTIVE
ENZYME
FORMULA



IBS FORMULA