

When does cortical coherence carry shared content?

Compatibility and self-reconstruction as testable conditions

Sebastian Bobadilla-Suarez

Independent Researcher

`sebastian.bobadilla.s@gmail.com`

ORCID: 0000-0002-5951-0772

20 September 2026

Abstract

Synchrony accounts treat collective phase order as the mechanism that integrates distributed cortical activity. We show formally that phase order alone establishes neither shared content nor a self-model, and state what additionally does. We separate three conditions: coherence (collective phase order), compatibility (agreement of independently decoded local descriptions where regions overlap) and self-reconstruction (accurate recovery of the global state from an internal encoding). Compatible local descriptions determine a unique global state. Coherence bounds disagreement between overlapping descriptions only through shared Lipschitz encoders, and the bound grows as the square root of the population read: at the patch order of a simulated cortical sheet it constrains at most 10 sites, so on a larger population the positive link from coherence to content rests entirely on a bridge assumption. A contracting self-map fixes one state but reconstructs accurately only states within a diameter set by its error and contraction factor; a broad family needs input-conditioned maps

and discriminating codes. Lean 4 verifies these implications, and one model satisfies all eight connecting hypotheses together; this shows consistency, not cortical realization. The conditions yield a disconfirmable comparison: the agreement of decoders fitted independently on two regions, in excess of what their separate accuracies imply, should be larger for perceived than for unperceived stimuli at matched phase coherence and single-region accuracy, with perception classified by a no-report marker. An exploratory EEG analysis is a measurement check only; its pooled estimator places a perfectly synchronized signal at zero coherence. Resource and implementation constraints, and a held-out onset comparison during anaesthetic emergence, are stated briefly. No principle yet selects the cover, and the sheaf’s overlaps are spatial where the example’s are overlaps of content, so the proposed correlate of unity has no operational definition. Unity and a minimal self remain interpretations that need an independent experiential endpoint.

1 Introduction

Endogenous extracellular fields can affect spike timing (Fröhlich & McCormick, 2010; Anastassiou & Koch, 2015) and may coordinate activity across cortical regions (McFadden, 2002, 2020; Pockett, 2012). Coordination raises a further question: when do those regions describe one situation, and when does the resulting state represent its own place in it?

Consider reaching for a cup. Vision and movement can describe different features while agreeing on the cup’s location. Matching their timing leaves agreement about that shared quantity untested. An internal description of the perceiver’s relation to the cup asks for still more. We call the three requirements *coherence* (collective phase order), *compatibility* (agreement of local descriptions on overlaps), and *self-representation* (accurate reconstruction

13 from an internal encoding). The mathematical states are spatial measures
14 and fields; identifying them with perceptual content requires independent
15 decoding evidence.

16 Binding-by-synchrony and communication-through-coherence accounts
17 treat phase relations as the medium of integration or of selective routing
18 (Singer & Gray, 1995; Fries, 2015), and field theories place integration in the
19 endogenous electromagnetic field (McFadden, 2020; Pockett, 2012). Both
20 leave open what fact about synchronized activity makes it one content rather
21 than coordinated activity carrying several. The contribution here is to sep-
22 arate the three requirements and prove which follow from which. Phase
23 order constrains shared content only through encoders related on overlaps,
24 with a bound that grows as the square root of the population it is read
25 on. Compatible local descriptions, not synchrony, determine a unique global
26 state. A contracting self-map fixes one state but can reconstruct accurately
27 only a family of bounded diameter. Each result names a measurement that
28 phase data alone cannot supply.

29 The objection that synchrony does not fix content is older than this work.
30 Critiques of the temporal binding hypothesis argued that synchrony, where
31 it occurs, cannot say which features belong together or what they jointly
32 represent: that grouping has to be computed before synchrony could sig-
33 nal it (Shadlen & Movshon, 1999). The formal results here are elementary:
34 uniqueness of gluing, Banach’s fixed-point theorem, the triangle inequality
35 and a pigeonhole count. What formalization adds to the informal argument is

36 quantitative and bookkeeping content. It gives the rate at which a coherence
37 bound on content weakens with population size and the population at which
38 it stops constraining anything, the diameter within which one contracting
39 map can reconstruct, and the number of codes a separated family of states
40 requires. It also gives a complete inventory of the premises a synchrony-to-
41 content argument consumes, each of which names a measurement, together
42 with one model in which all of them hold at once.

43 The familiar noisy mean-field oscillator supplies a stationary coherence
44 threshold (Sakaguchi, 1988; Strogatz & Mirollo, 1991). We add a resource
45 condition for a declared actuator, bounds that connect phase order to con-
46 tent only through specified encoders, and conditions under which compatible
47 descriptions glue and an accurate reconstruction fixes the glued state. A
48 Lean 4 development checks these implications and a joint witness satisfies
49 their eight connecting hypotheses in a specified model. It proves neither that
50 cortex satisfies those hypotheses nor that the glued state is experienced. The
51 latter identification motivates the framework and needs its own endpoint.

52 The empirical question is which measurements could support or discon-
53 firm the proposed connections. A phase resultant can miss a locally or-
54 dered winding field; stationary phase distributions can leave the coupling
55 mechanism unidentified; and coherent adaptation can lose alignment with
56 a designated content template. Independent decoders must identify shared
57 quantities before their agreement can be tested. We therefore distinguish sta-
58 tionary branches from finite observed onset, and compare prespecified field

59 and connectivity onset models on held-out emergence recordings. The com-
60 parison tests a specified spatial prediction, not the presence of experience.
61 Section 10.2 states the comparison that bears on conscious content directly.
62 Sections 2–5 develop the three requirements; Section 6 applies their mea-
63 surement obligations to a digital candidate. Section 7 states what the formal
64 composition establishes, and Sections 8–9 examine observables and the cor-
65 tical test. Table 1 lists the connecting hypotheses the article argues from;
Supplemental Material gives the full chain, derivations and controls.




Figure 1: Three requirements of the proposed account. Each requires the inputs shown below it. Dashed arrows mark proposed connections: phase coherence alone supplies neither compatible local descriptions nor an accurate self-model. Thermodynamics constrains a candidate implementation through separate assumptions. This is the explanatory structure of the article; Table 1 records the connecting hypotheses of the formal composition that the article argues from.

67 **2 Limits of phase order**

68 Several limits hold without the cortical hypothesis. Equal phases can coexist
69 with incompatible local descriptions; phase order constrains shared content
70 only through encoders related on overlaps. A winding field can have zero
71 global resultant while remaining locally ordered. A constant readout can
72 have a unique fixed point without reconstructing its input. A region that
73 does not distinguish relevant states cannot encode them, and a site cannot
74 report a change outside its causal past by a declared deadline. These lim-
75 its reflect familiar phase oscillator and information constraints (Acebrón et
76 al., 2005; Ott & Antonsen, 2008; Dörfler & Bullo, 2014; Breakspear, 2017).
77 The sections below state each one beside the condition or measurement it
78 motivates.

79 **3 Coherence: physical coupling and its cost**

80 **3.1 Dynamics and the stationary threshold**

81 The finite stochastic model is

$$d\theta_i = \left[\omega_i + \sum_j K_{ij}(t) \sin(\theta_j - \theta_i) \right] dt + \sqrt{2D} dW_i, \quad r_N = \left| \frac{1}{N} \sum_i e^{i\theta_i} \right|. \quad (1)$$

82 Here θ_i is the phase of oscillator i , ω_i its natural frequency, $D > 0$ is phase dif-
83 fusion, W_i are independent Wiener processes, and K_{ij} has inverse-time units,

84 with angles dimensionless. Uniform mean-field coupling is $K_{ij} = K/N$; fre-
 85 quency heterogeneity and diffusion are distinct parameters. The stationary
 86 calculation below uses identical frequencies and this scalar coupling. Ap-
 87 plying it to a spatial kernel requires an aggregation rule and a check of the
 88 resulting approximation.

89 For identical frequencies in their rotating frame, the positive normalized
 90 stationary densities of the one-body Fokker–Planck equation are exactly

$$\rho(\theta) = \frac{e^{a \cos(\theta - \psi)}}{2\pi I_0(a)}, \quad a = \frac{Kr}{D}, \quad r = R(a) := \frac{I_1(a)}{I_0(a)}, \quad (2)$$

91 where I_n are modified Bessel functions and ψ is the collective phase. The for-
 92 mal derivation uses the absence of stationary current for a periodic gradient
 93 drift to obtain this density, then verifies normalization and self-consistency.
 94 At $K \leq K_c = 2D$, only $r = 0$ is self-consistent (Sakaguchi, 1988; Strogatz &
 95 Mirollo, 1991). Above threshold, exactly one positive solution coexists with
 96 zero and increases strictly with K . Its onset is

$$\frac{r^2}{K - K_c} \longrightarrow \frac{1}{D}, \quad r \sim \sqrt{\frac{K - K_c}{D}}. \quad (3)$$

97 Figure 2 shows this stationary branch.

98 Linearization about the uniform density gives rate $K/2 - D$ for the first
 99 harmonic and $-Dn^2$ for harmonics $n \geq 2$. A growing mode therefore exists
 100 exactly above threshold. About the coherent density, the quadratic form
 101 is strictly dissipative transverse to collective rotation, whose rate is zero.

102 These linear estimates concern the limiting equation. Relating them to finite
 103 stochastic trajectories requires a dynamical mean-field limit and branch selec-
 104 tion; exchange symmetry alone does not establish the needed independence
 105 (Sznitman, 1991).




Figure 2: Stationary self-consistency for identical frequencies and phase diffusion $D = 1$. The positive coherent branch appears continuously above $K_c = 2D$ and increases with coupling. Zero remains a solution above threshold. The curve is computed from the integral expressions used in the formalization.

106 3.2 A speed limit on changes of coherence

107 One thermodynamic statement bears on coherence directly. For a smooth
 108 positive periodic phase density with probability current J and diffusion D ,

109 the current functional $\sigma_J = \int J^2/(D\rho) d\theta$ satisfies

$$\int_0^\tau \sigma_J dt \geq \frac{[\arcsin r(\tau) - \arcsin r(0)]^2}{D\tau} \quad (4)$$

110 while $0 < r < 1$. A change of coherence completed in time τ therefore has a
111 minimum current cost that grows as $1/\tau$, whereas holding a stationary order
112 has no positive floor. Read the other way, a current budget C sets a minimum
113 duration $\tau \geq [\Delta \arcsin r]^2/(DC)$ for a given change. This bears on ignition-
114 like transitions, in which large-scale coherence rises abruptly (Dehaene et al.,
115 2014), and on recovery latency after anaesthesia: each has a floor set by the
116 current available to drive it. The bound is a heat or power statement only
117 when J is identified with a physical flux, a reservoir at a stated temperature
118 is specified and local detailed balance holds for the steps concerned; only
119 then does σ_J convert to entropy production (Landauer, 1961; Norton, 2005;
120 Shenker, 2000). Installed energy supplies a separate, necessary price: for
121 a declared actuator whose response modes carry installation prices, mean
122 coupling is at most κU_{inst} for a mode-data factor κ , so a coherent stationary
123 branch requires installed energy $U_{\text{inst}} > 2D/\kappa$, and no cortical value of κ
124 is known. Supplemental Material gives both derivations, the register and
125 feedback budgets, and the sensitivity of the metabolic comparison to its
126 assumptions.

127 4 Compatibility: from coordinated activity 128 to a global description

129 4.1 Why common timing is insufficient

130 For overlapping patches U and V , compatibility means that restricting either
131 local description to $U \cap V$ gives the same object. Descriptions can agree on
132 shared quantities while contributing different information elsewhere. Com-
133 patible sections of the specified sheaf have a unique global extension. On
134 a finite cover, local finite spatial measures agreeing as measures on every
135 overlap also have a unique finite-measure extension whose sheaf image is the
136 glued section. No mass-one normalization is imposed.

137 In the cup example, visual and reaching descriptions must express a
138 shared quantity, such as cup location, in common coordinates. Restriction
139 retains that quantity while dropping details particular to each description.
140 Agreement permits surface appearance and hand configuration to contribute
141 to one situation. The mathematical counterexample assigns the same phase
142 to every patch while two local mass profiles disagree on a shared site. Those
143 profiles have no common extension. Because they are chosen independently
144 of phase, synchrony places no constraint on their disagreement.

145 This distinction limits the biological interpretation. Receptive territories
146 and content domains must be related to spatial patches, and the descriptions
147 must support the required extension. Matching overlap marginals of de-

148 coded probability laws need not determine a unique global joint distribution
149 (Abramsky & Brandenburger, 2011).

150 The cover is itself a choice, and the construction constrains it in one way
151 only: synchronized patches agree where they overlap. Where they do not
152 overlap there is nothing to compare, so a cover by pairwise disjoint patches
153 satisfies that obligation for *every* assignment of local descriptions and phases
154 and the gluing theorem still returns a unique global section, which is the
155 tuple it was handed. Membership of the class then reports nothing about the
156 system beyond the patches covering it. What a gluing claim asks of a cover
157 is therefore a non-degeneracy the theorem does not assume: that distinct
158 patches meet, and that what they share is a quantity fixed independently of
159 the agreement it is being used to establish.

160 The cup example exposes a second gap, between two kinds of overlap.
161 The sheaf is built over a spatial substrate, so its patches overlap where they
162 share sites. Visual and motor territories that agree on cup location overlap
163 in what they describe, not in where they lie, and the construction supplies
164 no relation between the two. One route recasts the construction over content
165 variables: each region becomes a partial decoder whose domain is the set of
166 variables it estimates, and two regions overlap exactly where they estimate
167 a common variable. The gluing argument needs only a cover and restriction
168 maps, so it applies formally to such a cover, but no result here is stated
169 for one, and which variables and decoders form it is an empirical question.
170 Until a principle selects the cover, the proposed correlate of unity has no

171 operational definition: gluing on a declared cover is proved, and which cover
 172 cortex realizes is open.

173 4.2 When coherence does constrain content

174 A declared relation between phase and content gives a positive result. Let
 175 contents take values in a real normed vector space $\mathcal{C}$. Patch i reads an
 176 assigned oscillator through $e_{i,x}$ at site x , evaluated on the circle point
 177 $(\cos \theta, \sin \theta)$. Assume a uniform Lipschitz constant L in chord distance and
 178 require $e_{i,x} = e_{j,x}$ wherever patches i and j overlap. Encoders may vary across
 179 sites; their equality on overlaps is a supplied relation between descriptions of
 180 the shared quantity. Write $z_k = e^{i\theta_k}$ and $m = N^{-1} \sum_k z_k$, so $|m| = r$. The
 181 phasors' squared deviations from their mean sum to $\sum_k |z_k - m|^2 = N(1 - r^2)$,
 182 and $|z_i - z_j| \leq |z_i - m| + |z_j - m|$ gives $|z_i - z_j|^2 \leq 2N(1 - r^2)$ for each pair.
 183 Consequently every overlap discrepancy is bounded by

$$\delta_r = L\sqrt{2N(1 - r^2)}. \quad (5)$$

184 Here discrepancy is the norm of the difference, and r and N are read on
 185 whichever population the bound is applied to. That choice is what decides
 186 whether the estimate says anything. A patch carries an order parameter
 187 of its own, and reading the bound there carries that patch's resultant and
 188 that patch's site count, so a large population observed through small patches
 189 is constrained by the small count. Read instead on the whole array, the

190 factor $\sqrt{N}$ makes the estimate weak, because it draws a pointwise guarantee
 191 from an average over N sites. The order cannot be improved in general:
 192 one oscillator in antiphase among N leaves $N(1 - r^2)$ near 4. At $r = 1$ the
 193 observations agree exactly at either scale. A shared scalar encoder is a special
 194 case. Dropping the overlap relation permits disagreement even at perfect
 195 locking. A common nonexpansive linear projection preserves agreement, but
 196 agreement of a projection does not establish agreement of the full vector.

197 What Eq. (5) bounds is a chord, which never exceeds 2, so the estimate
 198 constrains anything only where the right-hand side falls below that. At the
 199 sheet's own patch order, the bound is informative for at most 10 sites; beyond
 200 that, the coherence-to-content link requires the bridge assumption E78 of
 201 the cortical hypothesis (Table 1 gives the full composition and its edges).
 202 Supplemental Material provides the detailed derivation of this saturation,
 203 and states three conditional derivations that replace the uniform factor $\sqrt{N}$
 204 by a count of disagreeing pairs, a chain across a cover's shared sites, or a
 205 spectral gap at a frequency-locked configuration.

206 Mix retained fields with observations at gain $\eta \in [0, 1]$: $s_i \mapsto (1 - \eta)s_i +$
 207 ηo_i . Content and observation discrepancies bounded by ε and δ give a next
 208 discrepancy at most $(1 - \eta)\varepsilon + \eta\delta$. Repeated observations bounded by δ
 209 therefore give

$$\varepsilon_n \leq (1 - \eta)^n \varepsilon_0 + \delta. \quad (6)$$

210 Thus agreeing observations preserve exact compatibility and, for $\eta > 0$, re-

211 duce an initial mismatch. Approximate observations leave a nondecaying
 212 term in the error bound. Taking a uniform bound on Eq. (5) connects this
 213 update to the phase process.

214 **4.3 What approximate agreement determines**

215 Exact gluing requires exact restrictions. For nonnegative profiles on a finite
 216 substrate, suppose every overlap discrepancy is at most ε in uniform distance.
 217 Weights that vanish outside their patches and sum to one at each site select
 218 a global profile by weighted averaging. The selection lies within ε of each
 219 patch there, and any two such selections differ by at most ε , independently
 220 of cover multiplicity. An L -Lipschitz readout consequently varies by at most
 221 $L\varepsilon$ across selections.

222 At zero tolerance, the selection is the unique exact extension. At every
 223 positive tolerance there are families with no exact extension; selection then
 224 gives an approximation whose dependence on weights is bounded. Supple-
 225 mental Material gives a three-site witness attaining the bound. The mass-
 226 profile construction uses the scalar case of the content dynamics and addi-
 227 tionally requires nonnegativity of the initial profiles and observations. The
 228 general vector result supplies no identification with the spatial-measure sheaf.

229 These constructions concern declared objects: a specified cover, contents
 230 valued in a normed space, and encoders related on overlaps. They select
 231 neither the neural cover nor the decoder, and identifying their contents with
 232 perceptual content, or establishing that the identification bears on experi-

233 ence, is a further modelling task; Section 8 sets out what a measurement of
 234 it would have to establish.

235 **5 Self-representation: a state that models it-** 236 **self**

237 **5.1 The reflexive construction**

238 Compatibility gives a global state. Reflexivity relates it to an encoding within
 239 that state. Equip global sections with a complete metric. If the additional
 240 restriction-resonance condition identifies encoding with restriction to a region
 241 $\mathcal{A}$, the reconstruction map is

$$F(s) = \text{readout}(s|_{\mathcal{A}}). \quad (7)$$

242 The region and readout are fixed across the state space. If F contracts and
 243 fixes the section s_* formed from the local descriptions, Banach’s theorem
 244 makes s_* its unique fixed point (Banach, 1922). The formal bridge E89
 245 (Table 1 gives the full composition and its edges) supplies the contraction
 246 and $F(s_*) = s_*$; restriction resonance is separate. Supercritical coupling
 247 makes the stipulated contraction factor $\Lambda = e^{-(K-2D)\tau/2}$ less than one for
 248 a relaxation time $\tau > 0$. The factor is stipulated, not derived. It borrows
 249 the rate $(K - 2D)/2$ at which the first harmonic of the uniform density

grows (Section 3), a rate about a different state from the one F acts on,
and no result derives a Lipschitz law for an encoding or readout from the
phase dynamics. Constructing a map with that factor remains a modelling
obligation. A fixed contraction can reconstruct accurately only a restricted
family of states: if its error is at most ε throughout that family, any two
members are at most $2\varepsilon/(1 - \Lambda)$ apart.

In the cup example, the global state includes the represented scene and the
perceiver’s relation to it. A local encoding must retain enough information
for the readout to recover that modelled state; cup position alone would be
insufficient if hand configuration and viewpoint could not be reconstructed.

For the same map with Lipschitz factor $\Lambda < 1$, approximate reconstruction gives

$$d(s, F(s)) \leq \varepsilon \implies d(s, s_*) \leq \frac{\varepsilon}{1 - \Lambda}. \quad (8)$$

At the stipulated rate this radius grows as $2\varepsilon/((K - 2D)\tau)$ near threshold, for
fixed positive ε and τ . Exact uniqueness survives while the error guarantee
weakens.

5.2 Reconstruction across a family of states

Sensitivity can be required quantitatively. Declare relevant macrostates, an
encoder Φ and readout R , and require $d(s, R(\Phi(s))) \leq \varepsilon$ throughout that

268 family. If two states have the same code, the triangle inequality gives

$$\Phi(s) = \Phi(t) \implies \max\{d(s, R(\Phi(s))), d(t, R(\Phi(t)))\} \geq \frac{1}{2}d(s, t). \quad (9)$$

269 Thus M relevant states pairwise more than 2ε apart require at least M
 270 distinguishable codes; an alphabet of at most 2^b codes requires $M \leq 2^b$.
 271 Applied to the reflexive boundary, E is its local encoding and $R \circ E$ is the
 272 existing self-prediction map.

273 This is an additional accuracy criterion on a family, with no requirement
 274 for multiple exact fixed points. For the same fixed contracting map used
 275 by E89, however, the diameter bound above is an additional restriction: at
 276 zero error the family contains at most one metric-distinct state. A broad
 277 family therefore requires a different, input-conditioned reconstruction map,
 278 one map F_u with its own fixed point for each input u , with its own accuracy
 279 and stability assumptions. No such map is formalized or constructed here.
 280 Under E89 the proposed minimal-self correlate is therefore a condition on
 281 one self-state, not a model of a self that follows a changing scene. A blind
 282 encoder or constant readout fails the stated criterion for separated states
 283 despite having a unique fixed point.

284 Where the encoding is read as the state's own restriction to a region, the
 285 criterion becomes a condition on the region. Restriction resonance requires
 286 the encoder and that restriction to identify the same pairs of states, so a
 287 boundary reconstructing two relevant states more than 2ε apart while its

288 region reads them identically is not resonant. Read forwards, a resonant
 289 boundary that reconstructs a pairwise separated family restricts injectively
 290 on it. Which states a region reads identically is fixed by where the region
 291 sits, before an encoding is chosen, so the resolution the family demands is
 292 the region's and no readout supplies it. The implication runs one way: a
 293 region that does resolve the family is not thereby shown to carry a resonant
 294 encoding, and E89 supplies no such resolution.

295 Resonance and resolution are jointly satisfiable without a local fold. Take
 296 the region to be the whole substrate and use the identity readout: restric-
 297 tion is the identity, so this boundary is resonant and reconstructs every de-
 298 clared family exactly. Its map has Lipschitz factor one and does not satisfy
 299 E89's strict contraction premise on a nontrivial metric space. A properness
 300 condition on the region is therefore a separate requirement, and the fixed-
 301 contraction diameter bound remains in force when E89 is also required. With
 302 an L -Lipschitz readout, a family reconstructed within ε whose members lie
 303 pairwise more than $d_{\min}$ apart has δ -separated codes whenever $\delta L \leq d_{\min} - 2\varepsilon$,
 304 so the family is no larger than the δ -packing number of what the region car-
 305 ries. That is the shape a continuous family of states needs, and at the whole
 306 substrate it says nothing, because the region's range is everything. The gain
 307 L is declared and derived here for no physical readout, and at unbounded
 308 gain the bound says nothing.

309 5.3 Agreement after a change takes communication

310 Snapshot compatibility does not specify how quickly descriptions respond
311 when a shared quantity changes. In a synchronous network whose updates
312 read only the site’s own state and declared neighbours’ messages, two initial
313 configurations agreeing on a site’s causal past through T rounds produce the
314 same state there at round T . The deadline consequence is the one Section 2
315 states, and it follows from the reconstruction bound in Section 5.2: a report
316 from that site is a readout of a code that the two interventions leave identical.

317 A three-site delay line witnesses failure after one round and exact report-
318 ing after two; cutting the path prevents delivery at every round. A report
319 may still happen to be correct about one fixed world. The bound concerns
320 responsiveness across interventions.

321 Throughout this section *prediction* names a reconstruction map, and the
322 relevant family of states, the metric, the tolerance, the communication graph
323 and the deadline are all declared inputs, as are the represented variables and
324 the fidelity counted as accurate. Reading the map as temporal neural pre-
325 diction would require dynamics; reading the resulting code count as physical
326 memory capacity would require a hardware model, which a state-dependent
327 readout would have to extend.

328 6 What the conditions ask of a digital candi- 329 date

330 A claim that a digital system, such as a large language model (LLM) run-
331 ning on graphics processing units (GPUs), has unified content or represents
332 itself commits its maker to the measurements this account asks of cortex.
333 The maker must identify the shared quantities the system’s parts describe,
334 show that decoders fitted independently on those parts agree about them,
335 and show that a declared local code separates the relevant family of states
336 and survives intervention. The indicator-properties approach assesses arti-
337 ficial systems against computational properties drawn from theories of con-
338 sciousness (Butlin et al., 2023); the conditions here are fewer and each is
339 a measurement rather than an architectural property. A digital system is
340 also where they can be measured today, with full access to its state and
341 exact interventions on it: fitting decoders for one shared quantity on two
342 separate layers, heads or token positions, and testing their agreement and
343 whether reconstruction survives intervention, would carry out the proposed
344 measurement directly.

345 One constraint follows from the architecture alone. Within one forward
346 pass of a causally masked model, position t reads only its causal past, so an
347 intervention outside that past cannot be reconstructed at t by that step. This
348 is the deadline result of Section 5 applied to the mask, and it bounds which
349 variables a position can encode about the rest of the sequence within one

350 step. A second constraint needs a premise the architecture does not supply:
 351 a linear encoder and readout factoring through d coordinates pay reconstruction
 352 error at least the discarded spectral tail of the coupling operator, but
 353 query/key dimension does not supply such a bottleneck for softmax attention.
 354 These results rank no architecture family, imply no general inferiority of
 355 GPU hardware and settle no question of machine experience. Supplemental
 356 Material gives the bottleneck derivation, a synthetic two-view task with its
 357 controls, the information ceiling of token outputs and the hardware scope.

358 **7 The formal composition**

359 The Lean development composes the three requirements into one conditional
 360 theorem. Its passive and feedback branches each accept eight connecting hy-
 361 potheses, running from finite capacity through a register, dissipation and a
 362 continuum coupling to the field, coherence, the cover and the self-prediction
 363 fixed point. Each edge is supplied rather than derived from the one above it,
 364 and the inferential content of the composition is its last step, where contrac-
 365 tion and a fixed glued section give uniqueness. Table 1 lists the four edges the
 366 article argues from; Supplemental Material gives the full chain, its thermo-
 367 dynamic edges and the register ledger of the feedback branch. Both branches
 368 also take a cover carrying exact overlap compatibility and a reflexive bound-
 369 ary specifying encoding and readout, and conclude that the glued section is
 370 the unique fixed point of self-prediction. Composing the edges establishes

371 that the premises are consistent, named in one place and machine-checked
372 together; it does not establish that self-representation follows from coher-
373 ence. Joint witnesses satisfy every edge of each branch in a specified model,
374 and establish no common cortical mechanism. The development declares no
375 additional axioms beyond its libraries’ foundational axioms.

Table 1: The four connecting hypotheses the article argues from, shared by both branches of the formal composition. Supplemental Material gives the full eight-edge chain and, in Table S1, maps the node-level results.

Connection	Status	Required assumption
E56 <i>field identification</i>	Physical commitment	A specified physical field realizes the limiting coupling and satisfies the kernel conditions.
E67 <i>above threshold</i>	Physical commitment	That field operates above the synchronization threshold.

Table 1 (continued)

Connection	Status	Required assumption
E78 <i>coherence to cover</i>	Bridge assumption	Coherence relaxes the supplied cover at coupling at least the named mean-field value. Shared phase encoders bound vector-content mismatch; the exact cover and timely responsiveness remain separate requirements. Patches and oscillators carry one index, so one oscillator carries exactly one content patch: a modelling assumption rather than a consequence, and not an innocent one, since field-mediated coupling is spatial while two territories describing one quantity need not be neighbours. Where the cover's nerve is connected the same edge carries through the sites its patches share, at a residual linear in that nerve's diameter; where the configuration is frequency-locked it carries through the coupling graph's algebraic connectivity λ_2 instead of through any patch count, which asks the cover for nothing and is declared hardware data in the standing of κ .
E89 <i>contraction to fixed point</i>	Modelling assumption	A map mediated by local self-encoding fixes the glued section and has the stipulated Lipschitz factor $\Lambda < 1$. Reconstruction over a declared family imposes an additional code-separation criterion.

376 8 What the observations establish

377 The composition holds under its premises. Whether observation can estab-
378 lish those premises is a separate question, and the three requirements it
379 connects concern different objects: phase dynamics, local descriptions and
380 internal encodings. Their observables must retain these distinctions. The

381 following controls show which mechanisms remain indistinguishable, which
382 temporal approximations fail, which states one observable cannot separate,
383 and what independent content measurements must establish. Each is a finite
384 numerical regime on a declared model, with full parameters and controls in
385 Supplemental Material, and none of them calibrates a cortical quantity.

386 **8.1 Stationary distributions leave the mechanism un-** 387 **determined**

388 Equation (2) gives a concentration–coherence relation with no fitted shape
389 parameter. Estimate r from circular resultant length and a from density
390 shape: inverting R from the measured r would make agreement automatic.
391 Even independently estimated agreement identifies a statistical family shared
392 by different mechanisms.

393 Let $q(t)$ cross 2, let $r(q)$ be its stationary branch, and fix $D_0 > 0$. Increas-
394 ing coupling $K(t) = D_0 q(t)$ at diffusion D_0 and decreasing diffusion $D(t) =$
395 $K_0/q(t)$ at coupling K_0 both give concentration $a = q(t)r(q(t))$. Uncoupled
396 oscillators with prescribed drift $-h(t) \sin(\theta - \psi)$, where $h(t) = D_0 q(t)r(q(t))$,
397 and diffusion D_0 have that same density. Their stationary distributions co-
398 incide window by window despite different causes.

399 Resolved dynamics supply additional information. In the ideal model
400 with continuous latent paths, quadratic variation accumulates at rate $2D$
401 and conditional increment means constrain drift; a filtered observation of

402 the phase needs its own observation model before either is read off. Separat-
 403 ing endogenous coupling from a matched common drive additionally requires
 404 a measured drive or controlled perturbation; identifying electromagnetic cou-
 405 pling requires evidence specific to that interaction. Supplemental Material
 406 gives the equivalence construction and its numerical check. Phase statistics
 407 pooled over time and over bipolar pairs cannot supply these estimates: on
 408 public sedation EEG (Bajwa et al., 2025), the pooled pipeline places a per-
 409 fectly phase-locked signal at $r = 0.0001$, so a spatial test needs an estimand
 410 that removes the common rotation and a montage whose effect on spatial
 411 phase is modelled; Supplemental Material reports this estimator exercise and
 412 the dependence-aware design.

413 **8.2 Finite ramps change the observable onset**

414 The square-root law in Eq. (3) describes stationary solutions, and a chang-
 415 ing coupling needs time for the phase distribution to approach them. In a
 416 deterministic mean-field control crossing the threshold, none of the scanned
 417 rates meets the declared tracking tolerance at onset, so a slow control vari-
 418 able supplies no general guarantee of branch tracking. In a finite-population
 419 ramp study ($N = 2000$, $D = 1$, 32 replicas, 8 speeds), faster ramps de-
 420 lay escape. The fitted delay exponent 0.444 has a replica-bootstrap interval
 421 $[0.398, 0.496]$ that excludes the ideal $1/2$, a deterministic reference shows that
 422 the escape criterion changes the fitted exponent, and no common asymp-
 423 totic exponent is established. Delayed passage through a noisy bifurcation

(Berglund & Gentz, 2002) makes the apparent exponent depend on ramp speed and fitting window. Supplemental Material gives the sweep and its figure, the fluctuating-coupling and population-size controls, and the signed-network crossing bracket.

The ideal delayed-onset calculation motivates a square-root scaling hypothesis; its exponent is generic to suitable bifurcations and does not identify the coupling medium. A within-subject emergence design can vary the propofol taper and test how an operational onset changes with that manipulation, but interpreting the result as a coupling-rate law requires a measured or calibrated relation between taper and $\dot{K}$. The escape criterion must be set relative to each recording's pre-threshold fluctuation scale. Section 8.1 describes how D could be estimated from resolved phase traces.

8.3 A winding state leaves the order parameter ambiguous

The global order parameter r discards spatial arrangement. On a periodic sheet, a phase field can wind once around the sheet and have $r = 0$ while neighbouring sites remain locked. At identical frequencies, every such winding is stationary for an isotropic symmetric kernel: contributions at opposite separations cancel. It violates the quarter-turn initial-spread premise of the convergence result, which is why that result can still select the uniform state.

In the spatial control, the winding persists below the interpolated de-

445 cay length 0.1871 mm and unwinds above it (Fig. 3). At 0.1 mm, uniform
 446 and winding starts under the same coupling, diffusion and kernel end at
 447 $r = 0.9657$ and $r = 0.0083$; patch-local order is 0.9015. Under quenched fre-
 448 quency spread 1.5 rad s^{-1} , the measured retention boundary is 0.1421 mm,
 449 and the corresponding order comparison is 0.9418 against 0.0348 over 3 frozen
 450 realizations. These are finite controls, not a cortical wave identification or a
 451 derived transition.




Figure 3: A winding phase field on the spatial sheet, at identical frequencies. Left: global order (solid) against patch-local order (dashed) across a subset of coupling decay lengths. Right: the twisted and uniform initial conditions compared across lengths, with the shaded 0.1–0.3 mm band and the interpolated retention boundary. Below that boundary the two initial conditions reach different stationary values of r at the same coupling, diffusion and kernel, so the resultant does not report the coupling. The sweep is sampled at the lengths marked; it establishes no transition and no biological identification.

452 Linearization about a winding reweights each separation's coupling by
 453 the cosine of its phase advance. The resulting symmetric kernel has explicit
 454 mode rates: nonnegative reweighted coupling gives no growing mode, while

455 a positive rate destabilizes that winding. The criterion concerns linear rates,
 456 not asymptotic stability, and the measured retention boundary need not
 457 equal its limit. An amplitude extension adds an existence condition: $\Gamma + \lambda >$
 458 0 , where Γ is linear gain and $\lambda = \sum_d f_d(\cos \Delta\theta_d - 1)$ comes from the same
 459 kernel. For the nonnegative kernels used here, a broad kernel can remove a
 460 winding from that model even though the phase-only equations retain it as
 461 a stationary state.

462 Neither model turns these states into measured cortical spirals. The pe-
 463 riodic sheet cannot carry a lone phase defect, the relaxational dynamics has
 464 no propagation speed, and the amplitude variable has no calibrated relation
 465 to a recorded envelope. Supplemental Material gives the spectral, amplitude
 466 and defect results. Locally, Eq. (5) can still bound content disagreement on a
 467 winding patch when shared encoders are supplied, although its global $r = 0$
 468 form is uninformative. A nonzero ring winding number q also obstructs a
 469 global real-valued phase lift; it has no identified content meaning here. Thus
 470 an onset criterion based only on r can register a change of basin as well
 471 as a change of coupling. The protocol of Section 9.3 needs a spatial phase
 472 observable.

473 8.4 Coherent adaptation can lose environmental align- 474 ment

475 An adaptive coupling rule might supply representational structure, but its
476 objective must reward the relevant agreement. For symmetric coupling, pair-
477 wise sine terms cancel in the population mean drift. Minimizing squared drift
478 at that fixed mean therefore rewards equalizing drift across nodes. In the
479 frequency-cluster control, this favors coupling between unlike-frequency clus-
480 ters, whereas the diagnostic environmental template rewards coupling within
481 clusters.

482 The formalization proves descent of the squared-drift objective for a spec-
483 ified gradient flow at fixed phases. Its functional differs from thermodynamic
484 entropy production: at a coherent stationary density the probability current
485 vanishes while phase-averaged squared drift is positive. Supplemental Mate-
486 rial gives the operator, its gradient and the conditions of the descent result.

487 The finite control co-evolves noisy phases and symmetric, nonnegative
488 couplings at fixed total resource. Its template is a diagnostic covariance
489 of latent signals, never a training input. Across three illustrative seeds,
490 second-half order remains between 0.9167 and 0.9222, while the within-
491 cluster to between-cluster coupling ratio falls from 0.9824–0.9948 to 0.3507–
492 0.6135 (Fig. 4). Random-update, permuted-gradient and frozen-kernel con-
493 trols are reported in Supplemental Material.

494 The immediate post-update objective reduction disappears when aver-

495 aged over complete update intervals. A prespecified grid of 6 rate–cadence
496 combinations contains no case meeting the 5% sustained-reduction criterion
497 while retaining coherence and losing alignment. The finding is coherent adap-
498 tation away from the specified structure, with no demonstrated sustained cost
499 reduction.

500 Kernel alignment and overlap compatibility are different observables.
501 This control addresses acquisition of its designated template under its par-
502 ticular objective. Learning models can explicitly include data in an objective
503 (Corberi et al., 2026); an account of agreement about the cup must specify
504 the content its learning rule is meant to preserve.

505 8.5 Content agreement needs independent decoders

506 A neural test needs independently identified domains, local descriptions and
507 restriction maps. Two decoders should estimate a shared content on over-
508 lapping domains before their agreement is examined. Discrepancy requires a
509 null under decoding noise and controls using deliberately compatible and in-
510 compatible examples. An implementation on synthetic data separates these
511 cases at $AUC = 0.986$, on a 95% interval of $[0.980, 0.992]$ resampled by trial;
512 the compatible construction scores 0.476 on an interval containing one half.
513 Shared-prior shrinkage can nevertheless manufacture agreement, and regional
514 bias can manufacture disagreement. Per-patch decoding diagnostics detect
515 these failures; the compatibility statistic alone does not.

516 Phase agreement cannot replace that measurement. On constructed con-




Figure 4: Coherence persists while coupling loses alignment with the frequency-cluster template. Gradient updates (solid), random updates (dash-dotted), permuted gradient updates (dash-dot-dotted), and a frozen kernel (dashed) share one phase-noise stream per seed. Panels show the immediate post-update squared-drift objective, phase order, within-cluster to between-cluster coupling, and template distances. The gradient arm preserves order while its coupling ratio falls; growth in both distances mainly reflects kernel norm. The objective panel samples the instant after updates; complete-interval measurements find no sustained reduction in the bounded study.

517 tent, phase carries a share 1.000 of the content entropy when content is built
518 from it and 0.000 when it is not. A phase-derived statistic reports com-
519 patibility on phase-locked patches whose local descriptions disagree. The
520 implemented content statistic detects their incompatibility. These controls
521 validate a measurement on known constructions; a neural content model and
522 decoder remain to be established.

523 Choosing the cover also chooses a candidate subject: trivial descriptions
524 can agree while carrying little content, and a partition can conceal disagree-
525 ment. The useful choice must therefore be constrained by what neural ac-
526 tivity encodes. The exact composition assumes compatibility of its supplied
527 cover. The content mechanisms above identify additional hypotheses under
528 which agreement can arise; they leave the empirical choice of content and
529 the connection to that cover explicit.

530 **9 The cortical hypothesis and an awakening**

531 **test**

532 The cortical proposal identifies a physical source of coupling and asks whether
533 its changes help explain recovery. The tests in Section 8 determine what
534 observations must distinguish before a phase trace can support this identifi-
535 cation.

536 9.1 Physical identification and calibration

537 The cortical hypothesis identifies part of effective coupling with endogenous
538 extracellular electromagnetic interactions. Fields can modulate spike timing
539 (Fröhlich & McCormick, 2010; Anastassiou & Koch, 2015), and field models
540 propose coordination of distributed neural information (Pinotsis & Miller,
541 2022; McFadden, 2020). Phase organization under propofol motivates study-
542 ing this regime (Bastos et al., 2021), but does not measure coupling.

543 The formal field predicate requires a jointly continuous kernel, a normal-
544 ized substrate, coupling equal to the kernel average, and no single site pro-
545 viding more than half that strength. Toy constructions satisfy it. A cortical
546 realization must specify the scale at which the kernel and its reduction ap-
547 ply: calcium imaging reports different state-dependent modularity patterns
548 at single-cell and mesoscale resolution (Kiyooka et al., 2026).

549 Cortical columns are the obvious candidate unit, and they answer one
550 of the two demands the formalism makes. As the oscillator index they are
551 a serious proposal: they are delineated by response properties and anatomy
552 rather than by the agreement they would be used to establish (Mountcastle,
553 1997), and a phase can be estimated from a column-sized territory at the
554 electrode spacings used for cortical wave measurements (Townsend et al.,
555 2015). As the content cover they are a poor one, for three separate reasons.
556 A parcellation into non-overlapping columns is a disjoint cover, which satisfies
557 the compatibility obligation without constraining anything (Section 4). The
558 cover condition additionally requires the patches to exhaust the substrate,

559 while much of association cortex has no accepted columnar parcellation. And
 560 the column's standing as a canonical cortical unit is contested (Horton &
 561 Adams, 2005; Rakic, 2008). Offering columns for both roles at once therefore
 562 supplies the oscillator index; the content cover remains an open empirical
 563 question, which the bridge assumption E78 records.

564 The spatial reduction is also testable. A row-sum aggregation at zero
 565 diagonal tracks the uniform-coupling control over the tested spatial decay
 566 range, while frequency heterogeneity displaces the operational threshold. The
 567 fit window biases both estimates relative to $2D$; agreement with the control
 568 is therefore the supported conclusion. Supplemental Material reports the
 569 reduction, its numerical error and the distinct assumptions needed to identify
 570 the kernel average with the scalar coupling of the installed-energy condition.

571 The field commitment requires physical evidence beyond a choice of math-
 572 ematical representation, and the available support comparisons do not supply
 573 it. A finite matrix spread over positive-mass cells preserves its total cou-
 574 pling and phase correlation exactly, while point-supported vanishing under
 575 an atomless measure and a disadvantage from excluding a better-correlated
 576 pair concern specified support constraints; Supplemental Material reads the
 577 same comparisons against a digital candidate. Neither supplies energy prices.

578 A spike-time shift alone supplies no coupling rate. The dimensionless
 579 product of population size N , timing sensitivity $\mathcal{T}$, field amplitude E and

580 frequency f needs a conversion rate γ :

$$K_{\text{eff}} = \gamma N \mathcal{T} E f. \quad (10)$$

581 This rate describes how induced timing shifts accumulate into drift. Nei-
582 ther it nor the installation factor κ follows from a heat bound. Testing the
583 physical threshold requires independent estimates of the interaction kernel,
584 its aggregation and diffusion in the same cortical regime.

585 **9.2 Extracellular geometry and recovery**

586 The proposed control variable is extracellular geometry. Extracellular vol-
587 ume fraction is approximately $\alpha = 0.2$ at tortuosity $\lambda \approx 1.6$ (Syková &
588 Nicholson, 2008); a sleep study reports a rise from roughly 0.14 to 0.23 (Xie
589 et al., 2013). Below 10 kHz, cortical tissue conducts approximately resistively
590 at $0.3\text{--}0.6\text{ S m}^{-1}$ (Barbour, 2017). At fixed transmembrane current density,
591 an effective-medium description predicts that a larger extracellular fraction
592 raises conductivity and reduces field amplitude.

593 The coupling-gated hypothesis consequently predicts that extracellular
594 contraction on awakening raises E and, holding the other factors in Eq. (10)
595 fixed, raises effective coupling. This argument supplies a direction; the
596 geometry–coupling relation, its magnitude and its role in human recovery
597 require measurement.

598 For a smooth crossing at nonzero speed, evaluating the stationary branch

599 along the changing control gives a deterministic tracking approximation that
600 fails at onset in the tested crossing control:

$$r_{\text{ss}}(K(t)) \sim \sqrt{\frac{\dot{K}(t_0)}{D}} (t - t_0)^{1/2}, \quad K(t_0) = K_c, \quad \dot{K}(t_0) > 0. \quad (11)$$

601 This is a deterministic tracking approximation from Eq. (3) at fixed D ; the
602 crossing controls in Section 8.2 show that it fails as an observed-onset ap-
603 proximation under the tested initialization. A separation of geometry and
604 phase time scales can support tracking away from threshold. The awakening
605 hypothesis needs a calibrated dynamical prediction for an operational onset
606 above the finite-size fluctuation floor. Crossing K_c alone does not supply a
607 recovery latency.

608 **9.3 A protocol for the awakening experiment**

609 Neural inertia describes a behavioural barrier and hysteresis between anaes-
610 thetic induction and emergence (Friedman et al., 2010); recovery can pass
611 through metastable activity states (Hudson et al., 2014; Proekt & Hudson,
612 2018). The $v^{1/2}$ exponent of an ideal delayed bifurcation is generic (Baer
613 et al., 1989; Berglund & Gentz, 2002) and cannot alone discriminate a field
614 mechanism from a synaptic one. A possible discriminator is spatial onset
615 relative to independently estimated local field coupling and network connec-
616 tivity. The present mean-field model has no spatially varying κ and derives
617 no nucleation order, so a field-based onset map needs a separate model spec-

618 ified before the observations are inspected.

619 For that comparison, partition the recording into prespecified regions i
620 and measure pre-onset order $r_i(t_0)$ and diffusion D_i . Define a field score
621 $h_i = N_i \mathcal{T}_i E_i f_i$ from independent pre-onset field and timing measurements,
622 and a connectivity score c_i from a separately estimated structural network.
623 For the same measured emergence drive $g(t)$, compare $K_i^F(t) = a_F g(t) h_i$ with
624 $K_i^C(t) = a_C g(t) c_i$. Each model has one fitted scale. The local first-harmonic
625 approximation predicts

$$\hat{r}_i^M(t) = r_i(t_0) \exp\left(\int_{t_0}^t [K_i^M(s)/2 - D_i] ds\right), \quad M \in \{F, C\}. \quad (12)$$

626 Predicted onset is the first crossing of a region-specific threshold fixed from
627 pre-onset fluctuations. Fit the two scales on training sessions, then com-
628 pare held-out onset times and site order under the same observation model.
629 Larger field scores predict earlier onset only when baseline order and diffu-
630 sion do not reverse the ranking. This declared approximation omits interre-
631 gional coupling and saturation; a successful ranking would support its field
632 covariate, not prove ephaptic causation. Supplemental Material specifies the
633 controls and failure conditions.

634 A within-subject protocol can manipulate emergence rate, with spatially
635 resolved recordings to compare the proposed field and connectivity models:

636 **1. Manipulate emergence rate.** Vary the propofol infusion taper
637 across sessions within the same subject, recording drug concentrations

638 or a validated pharmacokinetic proxy to quantify the manipulation.

639 **2. Measure behavioural recovery.** Record time-to-response to a stan-
640 dard stimulus as the independent behavioural endpoint, rather than
641 estimating ΔK directly.

642 **3. Resolve spatial onset.** Identify anatomical locations where coher-
643 ence first crosses the baseline-scaled threshold and score the two maps
644 of Eq. (12) on held-out sessions. The two covariate maps are fixed
645 before the outcome sessions are inspected.

646 **4. Match physical controls.** Confirm that the observed coherence-to-
647 amplitude relation holds under the specific local diffusion D and ob-
648 servation noise. The model’s quadratic-variation rate $2D$ applies to
649 the latent phase; a band-limited analytic phase is smooth below its
650 correlation time, and on a constructed field its squared-increment rate
651 changes 37-fold between 0.2-ms and 20-ms steps. Estimating D there-
652 fore requires an explicit observation model for filtering and sampling.

653 Predeclare the recording window, regional fluctuation thresholds, and a
654 shared timing-error model. Fit the two scales and that observation model
655 on training sessions only; on held-out sessions, score the observed site-onset
656 times and sites that never cross by the same regional predictive rule. Report
657 site order as a secondary outcome and behavioural time-to-response sepa-
658 rately. Sessions without a crossing are censored observations, not zero delays
659 or exclusions. Compare the two scores only where independently estimated

660 field and connectivity maps predict distinguishable onset patterns at the
661 recording resolution. A held-out field-model advantage would support that
662 specified spatial predictor; it would not identify an ephaptic cause, shared
663 content or experience. Supplemental Material gives the controls and pilot
664 quantities needed to judge whether the comparison is informative.

665 **10 Discussion**

666 **10.1 Unity and a minimal self as interpretive motiva-** 667 **tion**

668 We propose the state glued from compatible local descriptions as a formal
669 correlate of conscious unity, and its accurate internal reconstruction as a cor-
670 relate of a minimal self. In the cup example, this would mean one description
671 that preserves the cup’s relation to the perceiver as well as the scene. Com-
672 patibility permits distinct contents and uncertainty; the shared quantities
673 must be chosen from independent evidence about what each region encodes.
674 The reconstruction need not be a verbal or autobiographical account.

675 The minimal self, in Gallagher’s sense, is the pre-reflective sense of be-
676 ing the subject of an experience, present before any narrative or reflective
677 self-description (Gallagher, 2000). Two of its features motivate the proposed
678 correlate. It is not a separate object of experience but a perspective built
679 into the experienced scene; correspondingly, the reconstruction is not a sec-

680 ond representation beside the glued state but a map whose fixed point is that
 681 state, read through a part of it. It is immediate rather than inferred from
 682 a higher-order state; correspondingly, the fixed point needs no further state
 683 that represents the first. The correspondence is structural and does not show
 684 that a fixed point is experienced as a perspective. Self-model theory locates
 685 the phenomenal self in a transparent model a system has of itself (Met-
 686 zinger, 2003), and interoceptive-inference accounts ground it in prediction of
 687 the body’s own states (Seth, 2013). The present condition is narrower than
 688 either: it asks only that a declared local encoding recover the global state,
 689 including perceiver-relative variables, at a declared fidelity, and it says noth-
 690 ing about transparency or about which bodily variables must be modelled.
 691 Either account could supply the family of states and the fidelity that the con-
 692 dition leaves as inputs. Under E89 that family is one state at zero error and
 693 a family of diameter at most $2\varepsilon/(1 - \Lambda)$ within error ε (Section 5.2), so the
 694 condition as proved constrains a single self-state; a self that persists through
 695 change needs the input-conditioned map that is not constructed here. Its
 696 proposed relation to experienced perspective remains an interpretation. Nei-
 697 ther identification is among the connecting hypotheses of the composition
 698 (Table 1), and satisfying them would not measure experience. The unity
 699 proposal also lacks an operational definition. The gluing theorem holds for
 700 whatever cover it is given, a disjoint cover satisfies it trivially, and the cover of
 701 interest overlaps in content rather than in space (Section 4); until a principle
 702 fixes that cover, the correlate is a schema awaiting one.

703 10.2 A comparison that could disconfirm the link to 704 content

705 The framework predicts that conscious content tracks compatibility, not co-
706 herence. The comparison must also separate compatibility from how well
707 each region encodes the stimulus, because two accurate decoders agree auto-
708 matically. If decoders d_A and d_B estimate a shared stimulus quantity x with
709 errors $e_A = d_A - x$ and $e_B = d_B - x$, then

$$|d_A - d_B| \leq |e_A| + |e_B|, \quad |d_A - d_B|^2 = |e_A|^2 + |e_B|^2 - 2\langle e_A, e_B \rangle. \quad (13)$$

710 Consciously perceived stimuli are usually decoded better, so higher agreement
711 on those trials would follow from single-region fidelity alone. At matched
712 single-region error, agreement differs only through the covariance of the two
713 errors, which is the term compatibility bears on: two regions describing one
714 content should err together.

715 Present matched stimuli that are perceived on some trials and not on
716 others, for example at masking or rivalry threshold, and fit decoders for
717 one shared stimulus quantity independently on two regions that both encode
718 it. Classify trials by perceptual state with a no-report measure, such as
719 reflexive eye movements or pupil responses that track the dominant percept
720 in rivalry (Tsuchiya et al., 2015), so that the classification does not also
721 select for the neural correlates of reporting; report-based classification is a
722 secondary analysis. Match the two classes on phase coherence between the

regions and on each region's own decoding error (or, where the stimulus value is uncertain, on each region's signal strength). The statistic is the excess agreement over a fidelity-matched null: pair region A 's decode on one trial with region B 's decode on another trial of the same stimulus and error stratum, which preserves both marginal accuracies and removes any coupling of the errors.

The proposed link predicts a larger excess on perceived trials at matched coherence and matched single-region accuracy (Fig. 5). Equal excess across classes, or excess that follows coherence and not perceptual state, counts against it. A positive result is not yet specific: noise shared upstream of both regions also correlates their errors, so the excess should be compared against an upstream region's decoding error, matched in the same way. A self-model comparison has the same form: at matched accuracy for the scene variable, the reconstruction of perceiver-relative variables, such as hand position relative to the target, from a declared local encoding should be more accurate on perceived trials. These are correlational tests of proposed correlates, and the no-report classification is itself an indirect marker of experience.

10.3 What a test of the full proposal must establish

A test needs separate estimates of effective coupling and diffusion, shared content on overlapping regions, and reconstruction from a specified encoding at a declared fidelity. The cover, relevant state family and decoder must be fixed without selecting them for agreement. An independent experiential




Figure 5: Design of the disconfirming comparison. Decoders for one shared stimulus quantity are fitted independently on two regions. Trials are classed by a no-report marker of perceptual state and matched on inter-regional phase coherence and on each region’s own decoding error, so that agreement cannot follow from single-region fidelity (Eq. (13)). The statistic is agreement in excess of a null that pairs decodes across trials within the same stimulus and error stratum.

745 endpoint is needed to evaluate the interpretation; absence of a report alone
 746 does not define absence of experience. A unified episode with incompati-
 747 ble decoded descriptions would challenge the unity identification. A unified
 748 episode lacking the proposed self-encoding would challenge the minimal-self
 749 claim. Conversely, satisfying the mathematical conditions without such an
 750 episode would challenge their sufficiency as correlates.

751 For the physical route, cortical actuator modes and prices, the geometry–
 752 coupling relation and a dynamical onset model remain to be measured. The
 753 field/connectivity comparison in Section 9.3 evaluates one spatial onset pre-
 754 diction. Controlled changes in coordination and independently decoded con-

755 tent agreement would address the representational connection. The synthetic
756 decoder exercise calibrates part of these measurements, and the EEG exercise
757 identifies what a pooled bipolar estimand cannot measure; neither identifies
758 a neural content code or tests experience.

759 **10.4 Explanatory scope**

760 Integrated-information, global-workspace and predictive-processing accounts
761 organize unity, access and modelling differently (Tononi et al., 2016; Baars,
762 1997; Dehaene et al., 2014; Friston, 2010). This account specifies conditional
763 relations among coupling, local content and reconstruction. Its formal field
764 condition requires a continuous kernel, a normalized substrate, a specified
765 average coupling and noise, and a site-contribution bound. Those conditions
766 do not identify an electromagnetic substrate; the extracellular interpretation
767 needs the separate calibration of Section 9. The joint witness shows that
768 the formal hypotheses can hold together in a declared model. A cortical
769 realization and an experiential interpretation require additional evidence.

770 Four limitations bound every claim above. Synchrony is not itself an
771 experiential measure, and nothing here establishes a consciousness criterion:
772 the conditions are proposed correlates to be tested against report or another
773 independent endpoint. The present EEG analysis checks a measurement
774 pipeline and does not validate any experiential claim. The formal results are
775 conditional on declared encoders, covers and maps whose cortical counter-
776 parts are unmeasured. No principle selects the cover on which compatibility

777 is assessed, so the unity correlate is not yet operationally defined.

778 The remaining explanatory question is whether these relations occur in
779 neural activity and matter for experience. The framework identifies candidate
780 correlates and tests of their physical and representational premises. It does
781 not explain why satisfying them would be accompanied by experience.

782 **11 Conclusions**

783 Phase coherence, compatibility and self-reconstruction are separable condi-
784 tions. Coherence constrains shared content only through encoders shared on
785 overlaps and with a bound that grows as the square root of the population
786 read. That bound is vacuous beyond 10 sites at the simulated sheet's patch
787 order, so for populations of cortical size the positive link from coherence
788 to content is the bridge assumption E78 and not a derived result. Com-
789 patible local descriptions, not synchrony, determine one global state; and a
790 contracting self-map fixes that state while reconstructing only a family of
791 bounded diameter. Each result replaces an assumption about synchrony by
792 a measurement: independently decoded agreement on overlaps, a declared
793 encoding and its fidelity, and calibrated coupling and diffusion. The compar-
794 ison of Section 10.2 states how such measurements could count against the
795 proposed link to conscious content.

Notation

Table 2: Symbols used throughout, with the section that defines each. Where one letter carries more than one meaning, the meanings are separated by the section they are read in.

Symbol	Defined	Meaning
θ_i, ω_i	Eq. (1)	Phase and natural frequency of oscillator i
N	Eq. (1)	Number of oscillators in the population the quantity is read on
D	Eq. (1), Supp.	Phase diffusion in inverse time; in the Supplemental Material's convergence proof, the phase spread $D \leq \pi/2$
K_{ij}, K	Eq. (1)	Pairwise and scalar mean-field coupling, $K_{ij} = K/N$
K_c	Eq. (2)	Synchronization threshold, $K_c = 2D$
r, r_N	Eq. (1)	Resultant length (order parameter), on the population it is read on
a, ψ	Eq. (2)	Concentration $a = Kr/D$ and collective phase of the stationary density
q	Section 8.3	Winding number of a phase field read around a ring
U_{inst}, κ	Section 3.2	Installed energy of a declared actuator and the mode-data factor bounding its mean coupling by κU_{inst}
L	Section 4.2	Lipschitz constant of an encoder or readout
ε, δ	Section 4.2	Content and observation discrepancy tolerances
M, b	Eq. (9)	Size of a declared family of states and code length in bits
Λ, τ	Eq. (7)	Contraction factor $\Lambda = e^{-(K-2D)\tau/2}$ and its relaxation time; in Eq. (4), τ is the duration of a change
J, σ_J, C	Eq. (4)	Probability current of the phase density, its current functional and a current budget
T	Section 5	Number of communication rounds to a deadline
ν	Supplemental Material	Fan-in of a communication graph: the most sites any one site hears from

Table 2 (continued)

Symbol	Defined	Meaning
$\gamma, \mathcal{T}, f$	Eq. (10)	Field-to-coupling conversion rate, timing sensitivity and frequency
α	Section 9	Extracellular volume fraction
λ	§8.3, §9	Eigenvalue the coupling kernel supplies at a winding; extracellular tortuosity
Ω, λ_2	Section 4.2	Common rotation rate of a frequency-locked configuration, and the algebraic connectivity of the coupling graph
β, k	Section 4.2	Chord diameter of a patch and the diameter of a cover's nerve, in hops
$\mathcal{C}, e_{i,x}$	Section 4.2	Content space and the phase encoder patch i reads at site x
Φ, R	Eq. (9)	Encoder and readout of the reconstruction criterion
$\mathcal{A}, F$	Eq. (7)	Encoding region and the reconstruction map $F(s) = \text{readout}(s _{\mathcal{A}})$
z_i, Γ, A_{ij}	Section 8.3	Complex amplitude, linear gain and coupling matrix of the amplitude layer
$\Delta\theta_d$	Section 8.3	Phase a winding advances across separation d
E	Eq. (10)	Field amplitude, in the cortical identification
s, s_*	§4–5	A global section and the one the reconstruction fixes

797 Data and code availability

798 The Lean development, simulation scripts, tests and saved numerical sum-
799 maries are maintained in the GitHub repository `bobaseb/coupling_gated_`
800 `cortical_coherence`. The formalization uses Lean 4.34.0-rc2 and Math-
801 lib commit `5ce203ecd53f8c3e47ee5530d7c86b94a55f0183`. Publication
802 simulation values are generated from saved summaries without rerunning

803 production sweeps. The exploratory EEG analysis uses OpenNeuro dataset
804 ds005620 (Bajwa et al., 2025) (subjects 1010, 1016, 1022, 1033, 1045,
805 1054, 1060, 1067, resting awake and sedation runs 1–2); preprocessing
806 and selection are described in Supplemental Material and implemented in
807 `simulations/empirical_collapse.py`. The repository contains the repro-
808 ducible analysis and does not redistribute the raw recordings.

809 **Funding and competing interests**

810 This research received no external funding. The author declares no compet-
811 ing interests.

812 **Use of AI tools**

813 AI assistance to this work was substantial and is not separable by task.
814 Several coding assistants and chat interfaces were used in an interleaved way
815 throughout: Google’s Antigravity CLI (Gemini 3.1 Pro, Claude 4.6 Opus,
816 Claude 4.6 Sonnet), OpenAI’s Codex CLI (GPT-6 Astra and Sol; GPT-
817 5.6 Terra, Luna and Sol), Claude Code (Claude Opus 5 and Opus 5.5),
818 Gemini Pro through its web interface, and Anthropic’s Fable and DeepSeek
819 v4 Flash through OpenRouter, accessed with the Hermes CLI. Between them
820 they contributed to the Lean development, the simulation and analysis code,
821 and the drafting and revision of both documents. Because the tools were

822 used in combination and in alternation, no section, proof, figure or result is
823 attributable to any one of them, and none is claimed to be. The author is
824 responsible for all content, including everything a tool produced. Machine
825 checking establishes the stated Lean results under their hypotheses; it does
826 not validate biological interpretations or replace scientific review of generated
827 code and prose.

828 **References**

- 829 Pinotsis, D. A., & Miller, E. K. (2022). Beyond dimension reduction: Stable
830 electric fields emerge from and allow representational drift. *NeuroImage*,
831 253, 119058.
- 832 Tononi, G., Boly, M., Massimini, M., & Koch, C. (2016). Integrated informa-
833 tion theory: from consciousness to its physical substrate. *Nature Reviews*
834 *Neuroscience*, 17(7), 450-461.
- 835 Singer, W., & Gray, C. M. (1995). Visual feature integration and the temporal
836 correlation hypothesis. *Annual Review of Neuroscience*, 18, 555-586.
- 837 Fries, P. (2015). Rhythms for cognition: communication through coherence.
838 *Neuron*, 88(1), 220-235.
- 839 Shadlen, M. N., & Movshon, J. A. (1999). Synchrony unbound: a critical
840 evaluation of the temporal binding hypothesis. *Neuron*, 24(1), 67-77.

- 841 Tsuchiya, N., Wilke, M., Frässle, S., & Lamme, V. A. F. (2015). No-report
842 paradigms: extracting the true neural correlates of consciousness. *Trends*
843 *in Cognitive Sciences*, 19(12), 757-770.
- 844 Metzinger, T. (2003). *Being No One: The Self-Model Theory of Subjectivity*.
845 Cambridge, MA: MIT Press.
- 846 Seth, A. K. (2013). Interoceptive inference, emotion, and the embodied self.
847 *Trends in Cognitive Sciences*, 17(11), 565-573.
- 848 Butlin, P., Long, R., Elmoznino, E., Bengio, Y., Birch, J., et al. (2023).
849 Consciousness in artificial intelligence: insights from the science of con-
850 sciousness. *arXiv preprint* arXiv:2308.08708.
- 851 Gallagher, S. (2000). Philosophical conceptions of the self: implications for
852 cognitive science. *Trends in Cognitive Sciences*, 4(1), 14-21.
- 853 Baars, B. J. (1997). In the Theater of Consciousness: The Workspace of the
854 Mind. *Oxford University Press*.
- 855 Dehaene, S., Charles, L., King, J. R., & Marti, S. (2014). Toward a compu-
856 tational theory of conscious processing. *Current Opinion in Neurobiology*,
857 25, 76-84.
- 858 Friston, K. (2010). The free-energy principle: a unified brain theory?. *Nature*
859 *Reviews Neuroscience*, 11(2), 127-138.

- 860 Milinkovic, B., & Aru, J. (2026). On biological and artificial consciousness:
861 A case for biological computationalism. *Neuroscience & Biobehavioral Re-*
862 *views*, 181, 106524. doi:10.1016/j.neubiorev.2025.106524.
- 863 Aru, J., Larkum, M. E., & Shine, J. M. (2023). The feasibility of artificial
864 consciousness through the lens of neuroscience. *Trends in Neurosciences*,
865 46(12), 1008–1017. doi:10.1016/j.tins.2023.09.009.
- 866 Seth, A. K. (2026). Conscious artificial intelligence and biological naturalism.
867 *Behavioral and Brain Sciences*, 49, e315. doi:10.1017/S0140525X25000032.
- 868 Seth, A. K. (2026). The stuff matters: Consciousness, compu-
869 tation, and biology. *Behavioral and Brain Sciences*, 49, e366.
870 doi:10.1017/S0140525X26106906.
- 871 Gurnee, W., et al. (2026). Verbalizable Representations Form a
872 Global Workspace in Language Models. *Transformer Circuits Thread*.
873 arXiv:2607.15495.
- 874 Aguilera, M., Millidge, B., Tschantz, A., & Buckley, C. L. (2022). How par-
875 ticular is the physics of the free energy principle?. *Physics of Life Reviews*,
876 40, 24-50.
- 877 Biehl, M., Pollock, F. A., & Kanai, R. (2021). A technical critique of some
878 parts of the free energy principle. *Entropy*, 23(3), 293.
- 879 Bruineberg, J., Dolega, K., Dewhurst, J., & Baltieri, M. (2022). The Em-
880 peror’s New Markov Blankets. *Behavioral and Brain Sciences*, 45, e183.

881 Pinotsis, D. A., & Miller, E. K. (2023). In vivo ephaptic coupling allows
882 memory network formation. *Cerebral Cortex*, 33(17), 9877-9895.

883 Pinotsis, D. A., Fridman, G., & Miller, E. K. (2023). Cytoelectric coupling:
884 Electric fields sculpt neural activity and "tune" the brain's infrastructure.
885 *Progress in Neurobiology*, 226, 102465.

886 McFadden, J. (2002). The conscious electromagnetic information (cemi) field
887 theory: the hard problem made easy?. *Journal of Consciousness Studies*,
888 9(8), 45-60.

889 Pockett, S. (2012). The electromagnetic field theory of consciousness: a
890 testable hypothesis about the characteristics of conscious as opposed to
891 non-conscious fields. *Journal of Consciousness Studies*, 19(11-12), 191-223.

892 McFadden, J. (2020). Integrating information in the brain's EM field: the
893 cemi field theory of consciousness. *Neuroscience of Consciousness*, 2020(1),
894 niaa016.

895 Fröhlich, F., & McCormick, D. A. (2010). Endogenous electric fields may
896 guide neocortical network activity. *Neuron*, 67(1), 129-143.

897 Anastassiou, C. A., & Koch, C. (2015). Ephaptic coupling to endogenous
898 electric field activity: why bother?. *Current Opinion in Neurobiology*, 31,
899 95-103.

900 Pockett, S. (2002). Difficulties with the electromagnetic field theory of con-
901 sciousness. *Journal of Consciousness Studies*, 9(4), 51-56.

902 Vöröslakos, M., et al. (2018). Direct effects of transcranial electric stimulation
903 on brain circuits in rats and humans. *Nature Communications*, 9, 483.

904 Verkhratsky, A., & Nedergaard, M. (2018). Physiology of Astroglia. *Physio-*
905 *logical Reviews*, 98(1), 239-389.

906 Mountcastle, V. B. (1997). The columnar organization of the neocortex.
907 *Brain*, 120(4), 701-722. doi:10.1093/brain/120.4.701.

908 Horton, J. C., & Adams, D. L. (2005). The cortical column: a structure
909 without a function. *Philosophical Transactions of the Royal Society B*,
910 360(1456), 837-862. doi:10.1098/rstb.2005.1623.

911 Rakic, P. (2008). Confusing cortical columns. *Proceedings of the National*
912 *Academy of Sciences*, 105(34), 12099-12100. doi:10.1073/pnas.0807271105.

913 Bastos, A. M., Donoghue, J. A., Brincat, S. L., Mahnke, M., Yanar, J., Cor-
914 rea, J., ... & Miller, E. K. (2021). Neural effects of propofol-induced uncon-
915 sciousness and its reversal using thalamic stimulation. *eLife*, 10, e60824.

916 Bajwa, I. J., Nilsen, A. S., Skukies, R., Aamodt, A., Ernst, G., Storm, J.
917 F., & Juel, B. E. (2025). A repeated awakening study exploring the capac-
918 ity of complexity measures to capture dreaming during propofol sedation.
919 *Scientific Reports*, 15, 32746.

920 Xie, L., Kang, H., Xu, Q., Chen, M. J., Liao, Y., Thiyagarajan, M., ... &
921 Nedergaard, M. (2013). Sleep drives metabolite clearance from the adult
922 brain. *Science*, 342(6156), 373-377.

- 923 Syková, E., & Nicholson, C. (2008). Diffusion in brain extracellular space.
924 *Physiological Reviews*, 88(4), 1277-1340.
- 925 Barbour, B. (2017). Analysis of claims that the brain extracellular impedance
926 is high and non-resistive. *Biophysical Journal*, 113(7), 1636-1638.
- 927 Logothetis, N. K., Kayser, C., & Oeltermann, A. (2007). In vivo measure-
928 ment of cortical impedance spectrum in monkeys: implications for signal
929 propagation. *Neuron*, 55(5), 809–823. doi:10.1016/j.neuron.2007.07.027.
- 930 Gabriel, S., Lau, R. W., & Gabriel, C. (1996). The dielectric properties of bio-
931 logical tissues: II. Measurements in the frequency range 10 Hz to 20 GHz.
932 *Physics in Medicine and Biology*, 41(11), 2251–2269. doi:10.1088/0031-
933 9155/41/11/002.
- 934 Norton, J. D. (2005). Eaters of the lotus: Landauer’s principle and the return
935 of Maxwell’s demon. *Studies in History and Philosophy of Science Part B:*
936 *Studies in History and Philosophy of Modern Physics*, 36(2), 375-411.
- 937 Landauer, R. (1961). Irreversibility and heat generation in the computing
938 process. *IBM Journal of Research and Development*, 5(3), 183-191.
- 939 Kibble, T. W. B. (1976). Topology of cosmic domains and strings. *Journal*
940 *of Physics A: Mathematical and General*, 9(8), 1387-1398.
- 941 Sakaguchi, H. (1988). Cooperative phenomena in coupled oscillator systems
942 under external fields. *Progress of Theoretical Physics*, 79(1), 39-46.

943 Strogatz, S. H., & Mirollo, R. E. (1991). Stability of incoherence in a popu-
944 lation of coupled oscillators. *Journal of Statistical Physics*, 63, 613–635.

945 Acebrón, J. A., Bonilla, L. L., Pérez Vicente, C. J., Ritort, F., &
946 Spigler, R. (2005). The Kuramoto model: A simple paradigm for
947 synchronization phenomena. *Reviews of Modern Physics*, 77, 137–185.
948 doi:10.1103/RevModPhys.77.137.

949 Ott, E., & Antonsen, T. M. (2008). Low dimensional behavior of
950 large systems of globally coupled oscillators. *Chaos*, 18, 037113.
951 doi:10.1063/1.2930766.

952 Dörfler, F., & Bullo, F. (2014). Synchronization in complex net-
953 works of phase oscillators: A survey. *Automatica*, 50, 1539–1564.
954 doi:10.1016/j.automatica.2014.04.012.

955 Breakspear, M. (2017). Dynamic models of large-scale brain activity. *Nature*
956 *Neuroscience*, 20, 340–352. doi:10.1038/nn.4497.

957 Still, S., Sivak, D. A., Bell, A. J., & Crooks, G. E. (2012). Thermodynamics
958 of prediction. *Physical Review Letters*, 109(12), 120604.

959 Kawai, R., Parrondo, J. M. R., & Van den Broeck, C. (2007). Dissipation:
960 The phase-space perspective. *Physical Review Letters*, 98(8), 080602.

961 Horowitz, J. M., & Esposito, M. (2014). Thermodynamics with continuous
962 information flow. *Physical Review X*, 4, 031015. doi:10.1103/PhysRevX.4.
963 031015.

- 964 Ito, S., & Sagawa, T. (2013). Information thermodynamics on
 965 causal networks. *Physical Review Letters*, 111, 180603. doi:10.1103/
 966 PhysRevLett.111.180603.
- 967 Corberi, F., dello Russo, S., Messuti, G., Scarpetta, S., & Smaldone, L.
 968 (2026). Thermodynamic learning. *arXiv preprint*. arXiv:2609.04732.
- 969 Banach, S. (1922). Sur les opérations dans les ensembles abstraits et leur ap-
 970 plication aux équations intégrales. *Fundamenta Mathematicae*, 3(1), 133-
 971 181.
- 972 Shenker, O. R. (2000). Logic and entropy. *PhilSci-Archive preprint*. philsci-
 973 archive.pitt.edu/115.
- 974 Rosas, F. E., Mediano, P. A. M., Luppi, A. I., Boyd, A. B., Berta, M., Jensen,
 975 H. J., Gastpar, M., & Carroll, S. M. (2026). Four faces of information in
 976 natural systems. *Nature Reviews Physics*. doi:10.1038/s42254-026-00971-4.
- 977 Sznitman, A.-S. (1991). Topics in propagation of chaos. In *École d'Été de*
 978 *Probabilités de Saint-Flour XIX—1989*, Lecture Notes in Mathematics,
 979 vol. 1464, pp. 165–251. Springer. doi:10.1007/BFb0085169.
- 980 Dai Pra, P., & den Hollander, F. (1996). McKean–Vlasov limit for interacting
 981 random processes in random media. *Journal of Statistical Physics*, 84, 735–
 982 772. doi:10.1007/BF02179656.
- 983 Bertini, L., Giacomin, G., & Pakdaman, K. (2010). Dynamical aspects of

- mean field plane rotators and the Kuramoto model. *Journal of Statistical Physics*, 138, 270–290. doi:10.1007/s10955-009-9908-9.
- Berglund, N., & Gentz, B. (2002). Pathwise description of dynamic pitchfork bifurcations with additive noise. *Probability Theory and Related Fields*, 122, 341–388. doi:10.1007/s004400100174.
- Abramsky, S., & Brandenburger, A. (2011). The sheaf-theoretic structure of non-locality and contextuality. *New Journal of Physics*, 13, 113036. doi:10.1088/1367-2630/13/11/113036.
- Townsend, R. G., Solomon, S. S., Chen, S. C., Pietersen, A. N. J., Martin, P. R., Solomon, S. G., & Gong, P. (2015). Emergence of complex wave patterns in primate cerebral cortex. *The Journal of Neuroscience*, 35(11), 4657–4662. doi:10.1523/JNEUROSCI.4509-14.2015.
- Xu, Y., Long, X., Feng, J., & Gong, P. (2023). Interacting spiral wave patterns underlie complex brain dynamics and are related to cognitive processing. *Nature Human Behaviour*, 7(7), 1196–1215. doi:10.1038/s41562-023-01626-5.
- Nicoletti, G., Di Terlizzi, I., & Busiello, D. M. (2026). Balancing information and dissipation with partially observed fluctuating signals. *Physical Review Letters*, 137(12), 127101. doi:10.1103/8gx2-rbns.
- Kiyooka, D., Oomoto, I., Kitazono, J., Saito, Y., Kobayashi, M., Matsubara, C., Kobayashi, K., Murayama, M., & Oizumi, M. (2026). Single-

1005 cell resolution functional networks during unconsciousness are segre-
 1006 gated into spatially intermixed modules. *Cell Reports*, 45(2), 116902.
 1007 doi:10.1016/j.celrep.2025.116902.

1008 Oomoto, I., Kiyooka, D., Oizumi, M., & Murayama, M. (2026). Multi-area
 1009 single-cell calcium imaging dataset of the mouse cortex across wakefulness,
 1010 sleep, and anesthesia. *Scientific Data*. doi:10.1038/s41597-026-08202-2.

1011 Attwell, D., & Laughlin, S. B. (2001). An energy budget for signaling in the
 1012 grey matter of the brain. *Journal of Cerebral Blood Flow & Metabolism*,
 1013 21(10), 1133-1145. doi:10.1097/00004647-200110000-00001.

1014 Harris, J. J., Jolivet, R., & Attwell, D. (2012). Synaptic energy use and
 1015 supply. *Neuron*, 75(5), 762-777. doi:10.1016/j.neuron.2012.08.019.

1016 Nicholls, D. G., & Ferguson, S. J. (2013). *Bioenergetics 4*. Academic Press.

1017 Baer, S. M., Erneux, T., & Rinzel, J. (1989). The slow passage through a
 1018 Hopf bifurcation: delay, memory effects, and resonance. *SIAM Journal on*
 1019 *Applied Mathematics*, 49(1), 55–71. doi:10.1137/0149003.

1020 Friedman, E. B., Sun, Y., Moore, J. T., Hung, H.-T., Meng, Q. C., Perera, P.,
 1021 Joiner, W. J., Thomas, S. A., Eckenhoff, R. G., Sehgal, A., & Kelz, M. B.
 1022 (2010). A conserved behavioral state barrier impedes transitions between
 1023 anesthetic-induced unconsciousness and wakefulness: evidence for neural
 1024 inertia. *PLoS ONE*, 5(7), e11903. doi:10.1371/journal.pone.0011903.

- 1025 Hudson, A. E., Calderon, D. P., Pfaff, D. W., & Proekt, A. (2014). Recovery
1026 of consciousness is mediated by a network of discrete metastable activity
1027 states. *Proceedings of the National Academy of Sciences*, 111(25), 9283–
1028 9288. doi:10.1073/pnas.1408296111.
- 1029 Proekt, A., & Hudson, A. E. (2018). A stochastic basis for neural inertia
1030 in emergence from general anaesthesia. *British Journal of Anaesthesia*,
1031 121(1), 86–94. doi:10.1016/j.bja.2018.02.035.