

# Coupling-gated cortical coherence: a conditional framework for unity and self-representation

Sebastian Bobadilla-Suarez  
Independent Researcher  
`sebastian.bobadilla.s@gmail.com`  
ORCID: 0000-0002-5951-0772

20 September 2026

## Abstract

How could activity distributed across cortex support one experienced situation? We separate three requirements: *coherence* (collective phase order), *compatibility* (local descriptions agree where their domains overlap) and *self-representation* (the state carries an accurate reconstruction of itself). Coherence supplies neither of the others, and five results establish that with no cortical hypothesis. Synchrony bounds content disagreement only through encoders shared on overlaps, and then only at the patch the bound is read on, not over the whole population. The resultant does not determine the state: a winding field and a uniform one are stationary at one coupling, at opposite ends of that observable. A unique fixed point is not a self-model, since a contraction can discard its input: accuracy across a family of states requires distinguishable codes. Where the encoding is a region's restriction, that resolution is demanded of the region before any readout is chosen. And agreement about a change takes communication. Four of the five are checkable for a machine candidate, which fixes what a claim of machine unity or self-representation commits its maker to measuring. Under stated hypotheses, compatible

local states then glue to a unique global state that an accurate contracting reconstruction fixes uniquely. Installed energy bounds the mean coupling a declared actuator supplies, so a coherent stationary branch requires it to exceed  $2D/\kappa$ ;  $\kappa$  is derived here for neither cortex nor a processor. A Lean 4 development carries these results and their composition, with the eight connecting hypotheses as explicit arguments, no incomplete proofs, and an audited axiom footprint no wider than its libraries'. Further controls bound what observation can decide: distinct mechanisms share stationary phase distributions, and coherent adaptation can lose the structure it was meant to acquire. One prediction needs no absolute calibration: onset delay scales as the square root of the rate at which coupling crosses threshold, which a within-subject emergence-from-anaesthesia design can test; the finite-size estimate here is biased below  $1/2$  by its escape criterion, as a deterministic reference confirms. We propose the glued state and its internal reconstruction as correlates of unity and a minimal self, and an extracellular-field hypothesis linking awakening to rising coupling. Testing it requires calibrated geometry, resolved phase dynamics and validated decoders.

# 1 Introduction

How could activity distributed across cortex support one experience? A field account proposes that endogenous extracellular electromagnetic fields partly coordinate spatially separated neural processes (McFadden, 2002, 2020; Pockett, 2012). Ephaptic effects provide a possible interaction mechanism (Fröhlich & McCormick, 2010; Anastassiou & Koch, 2015). Successful coordination would still leave a representational question: what makes the coordinated activity a description of one situation, with a perspective on that situation?

We distinguish three requirements. *Coherence* concerns collective phase

11 order. *Compatibility* concerns whether local descriptions agree where their  
12 domains overlap. *Self-representation* concerns whether the resulting state  
13 carries an accurate internal reconstruction of itself. We propose the latter  
14 two relations as formal correlates of conscious unity and a minimal self; their  
15 interpretation requires evidence beyond the phase model.

16 Unity has been approached differently elsewhere: integrated information  
17 theory treats integration and exclusion as constitutive (Tononi et al., 2016),  
18 global-workspace accounts tie availability to broadcast (Baars, 1997; Dehaene  
19 et al., 2014), and predictive-processing accounts derive both from a hierar-  
20 chical generative model (Friston, 2010). Here the three requirements are  
21 kept apart rather than derived from one another, so that each carries its own  
22 assumptions and admits its own measurement; Section 9.4 returns to the  
23 comparison.

24 Consider reaching for a cup. A visual description of its location and a  
25 description of the hand’s relation to it can agree about where the cup is while  
26 contributing different contents. Synchronizing the two processes would leave  
27 disagreement about that shared quantity unresolved. An internal representa-  
28 tion must additionally locate the perceiver within the represented situation.  
29 This example motivates the requirements; the mathematical constructions  
30 use spatial measures and fields in real normed vector spaces, whose identifi-  
31 cation with perceptual content is a further modelling task.

32 The mathematical contribution is to specify connections between these  
33 requirements and the assumptions each connection needs. A declared ac-

34 tuator’s stored energy bounds its mean coupling. Phase encoders shared on  
35 overlaps convert phase order into a bound on content disagreement. Compat-  
36 ible spatial measures have a unique global extension; approximate agreement  
37 controls a specified global selection. A contracting reconstruction makes that  
38 extension its unique fixed point only if it accurately reconstructs the exten-  
39 sion. Accuracy across a family of states imposes a further condition on the  
40 information retained by the encoding.

41 The empirical contribution is to identify what observations can decide.  
42 Stationary phase distributions can be shared by stronger coupling, weaker  
43 diffusion and a matched common drive. Finite ramps change the observable  
44 onset. A winding field is stationary at the same coupling as a uniform one and  
45 has the opposite resultant, so that observable does not determine the state.  
46 Coherent adaptation can lose alignment with a designated environmental  
47 template, and a compatibility estimator needs diagnostics that distinguish  
48 content from decoder artifacts. These findings determine the measurements  
49 needed to test the proposed connections.

50 Three of the ingredients are standard: the synchronization threshold  
51  $K_c = 2D$  for the noisy mean-field model (Sakaguchi, 1988), the unique gluing  
52 of compatible local sections, and the Banach fixed-point theorem (Banach,  
53 1922). Four things here are not. The installed-energy condition bounds  
54 mean coupling by the stored energy of a declared actuator, which makes a  
55 resource condition necessary for coherence on the stationary branch. The  
56 coherence-to-content bound converts phase order into a statement about de-

57 descriptions rather than about timing, and its patch-local form is what keeps  
58 it from being vacuous in a large population. A stability criterion on the  
59 reweighted coupling kernel, together with an existence condition on an am-  
60 plitude layer, settles when a low resultant reports weak coupling and when  
61 it reports a winding field. And the composition is machine-checked together  
62 with a witness discharging all eight hypotheses at once, which is what makes  
63 their conjunction non-empty rather than a list no object satisfies.

64 Granting those premises, collected in Table 1, the account runs as follows.  
65 Extracellular contraction on awakening raises endogenous field amplitude  
66 and with it the effective coupling between cortical oscillators. Once that  
67 coupling passes  $2D$ , a coherent phase branch exists, and from an initial spread  
68 within a quarter turn the population settles onto it; other initial data leave  
69 windings stationary at the same coupling. Encoders shared where cortical  
70 territories overlap turn that phase order into agreement about the quantities  
71 the territories describe in common, so the local descriptions glue: there is  
72 one state of which each is a restriction. Part of that state encodes the state  
73 itself, and a readout of that part reconstructs it accurately enough, and  
74 sensitively enough across the states that matter, to make it the unique state  
75 the system models itself as being. On this account the glued state is what  
76 makes distributed activity one situation, and its accurate self-reconstruction  
77 is what gives that situation a perspective. Every step of it is conditional.

78 Section 2 collects the results that hold whatever physical field is proposed  
79 and whatever substrate carries it; a reader who rejects the cortical hypothesis

80 of Section 8 still has to accommodate them. Sections 3–5 then develop the  
 81 three requirements together (Fig. 1). Section 6 composes them, stating the  
 82 thermodynamic constraints and all eight connecting hypotheses per branch;  
 83 its supporting models of feedback, learning and resource supply have distinct  
 84 state spaces and assumptions. Section 7 then examines what the observables  
 85 can establish, before Section 8 develops the cortical hypothesis and an awak-  
 86 ening experiment. The hypothesis is that extracellular contraction increases  
 87 field-mediated coupling; its direction, rate and relation to recovery require  
 88 independent calibration. Supplemental Material organizes the constructions  
 89 and controls behind the claims they support.




Figure 1: Three requirements of the proposed account. Each requires the inputs shown below it. Dashed arrows mark proposed connections: phase coherence alone supplies neither compatible local descriptions nor an accurate self-model. Thermodynamics constrains a candidate implementation through separate assumptions. This is the explanatory structure of the article; Table 1 records the eight implications consumed by each branch of the formal composition.

## 90 2 What holds without the cortical hypothesis

91 Most of what follows is conditional. Table 1 names eight hypotheses that  
92 the composition consumes rather than derives, and the cortical identification  
93 of Section 8 is a further commitment on top of them. Five results are not  
94 conditional in that way. They constrain any account that reads unity off  
95 phase coherence, whatever physical field is proposed to carry the coupling  
96 and whatever substrate carries the field. Each is stated below where it is  
97 used; collected here, they are the article’s fixed points.

98 **Phase order alone does not constrain content.** Assign every patch  
99 the same phase and let two local mass profiles disagree at a shared site.  
100 The profiles are chosen independently of phase, so synchrony bounds their  
101 disagreement not at all, and they have no common extension: perfect locking  
102 is consistent with local descriptions that do not glue (Section 4). What the  
103 positive result of Section 4.2 costs is exactly this gap. A bound on content  
104 disagreement needs encoders already related on overlaps, and that relation  
105 is a supplied fact about the descriptions rather than a consequence of the  
106 dynamics.

107 **The order parameter does not determine the state.** A field winding  
108 once across a periodic sheet has  $r = 0$  while every patch of it stays locked,  
109 and this is not a contrived initial condition. On a periodic array of any  
110 dimension, a winding is stationary for *any* isotropic symmetric kernel at

111 identical frequencies, because the coupling term is odd in the separation  
 112 against a kernel even in it (`char.is.kuramoto.trajectory`). Section 7.3  
 113 exhibits a sheet on which uniform and winding initial conditions reach 0.9657  
 114 and 0.0083 at one coupling, diffusion and kernel. A low resultant therefore  
 115 reports either weak coupling or a winding field, and no measurement in  $r$   
 116 alone separates them. The state also carries an integer the resultant does  
 117 not: read around a ring, the same winding has winding number  $q$ , and a  
 118 nonzero  $q$  obstructs every global real-valued phase lift (`winding.degree_`  
 119 `obstructs`). The resultant is an average over sites and moves when they are  
 120 reweighted or the patches redrawn; the degree telescopes and does not.

121 **A unique fixed point is not a self-model.** A constant readout con-  
 122 tracts, fixes whatever state it returns, and reconstructs nothing. Uniqueness  
 123 is consequently the wrong criterion, and Section 5.2 gives the right shape  
 124 of one: reconstructing  $M$  states pairwise more than  $2\varepsilon$  apart within error  $\varepsilon$   
 125 requires  $M$  distinguishable codes, so an alphabet of  $2^b$  codes caps the family  
 126 at  $M \leq 2^b$ . This is a counting argument and holds for any encoder and  
 127 readout whatever.

128 **Where the encoding is a restriction, the region must resolve the**  
 129 **family.** Read the code as the state's own restriction to a region and  
 130 the criterion moves off the readout entirely. A boundary whose region  
 131 reads two relevant states identically cannot reconstruct them more than  $2\varepsilon$   
 132 apart (`not.isRestrictionResonance.of_reconstructs`), and a resonant

133 boundary that does reconstruct a separated family restricts injectively on  
134 it (`restrictToAvatar_injOn_of_isRestrictionResonance`). Where the  
135 region sits is fixed before an encoding is chosen, so no readout supplies  
136 resolution the region lacks.

137 **Agreement after a change takes communication.** In a synchronous  
138 network whose updates read only the site’s own state and its declared neigh-  
139 bours’ messages, two configurations agreeing on a site’s causal past through  
140  $T$  rounds leave that site identical at round  $T$ . Two interventions outside  
141 that past that require reports more than  $2\varepsilon$  apart therefore cannot both be  
142 reported from it within  $\varepsilon$  by the deadline (Section 5). Snapshot compatibil-  
143 ity and timely compatibility are therefore different requirements, and cutting  
144 the path removes the second while leaving the first intact.

145 None of these depends on the coupling being electromagnetic, on the sub-  
146 strate being cortex, or on the eight hypotheses of Table 1 holding. Section 9.3  
147 applies four of them — all but the one about the resultant, which is a state-  
148 ment about a spatial observable — to a candidate that is neither biological  
149 nor field-coupled.

### 150 **3 Coherence: physical coupling and its re-** 151 **source condition**

#### 152 **3.1 Dynamics and the stationary threshold**

153 The finite stochastic model is

$$d\theta_i = \left[ \omega_i + \sum_j K_{ij}(t) \sin(\theta_j - \theta_i) \right] dt + \sqrt{2D} dW_i, \quad r_N = \left| \frac{1}{N} \sum_i e^{i\theta_i} \right|. \quad (1)$$

154 Here  $D > 0$  is phase diffusion,  $W_i$  are independent Wiener processes, and  
155  $K_{ij}$  has inverse-time units, with angles dimensionless. Uniform mean-field  
156 coupling is  $K_{ij} = K/N$ ; frequency heterogeneity and diffusion are distinct  
157 parameters. The stationary calculation below uses identical frequencies and  
158 this scalar coupling. Applying it to a spatial kernel requires an aggregation  
159 rule and a check of the resulting approximation.

160 For identical frequencies in their rotating frame, the positive normalized  
161 stationary densities of the one-body Fokker–Planck equation are exactly

$$\rho(\theta) = \frac{e^{a \cos(\theta - \psi)}}{2\pi I_0(a)}, \quad a = \frac{Kr}{D}, \quad r = R(a) := \frac{I_1(a)}{I_0(a)}, \quad (2)$$

162 where  $I_n$  are modified Bessel functions and  $\psi$  is the collective phase. The for-  
163 mal derivation uses the absence of stationary current for a periodic gradient  
164 drift to obtain this density, then verifies normalization and self-consistency.  
165 At  $K \leq K_c = 2D$ , only  $r = 0$  is self-consistent (Sakaguchi, 1988). Above

threshold, exactly one positive solution coexists with zero and increases  
strictly with  $K$ . Its onset is

$$\frac{r^2}{K - K_c} \longrightarrow \frac{1}{D}, \quad r \sim \sqrt{\frac{K - K_c}{D}}. \quad (3)$$

Figure 2 shows this stationary branch.

Linearization about the uniform density gives rate  $K/2 - D$  for the first harmonic and  $-Dn^2$  for harmonics  $n \geq 2$ . A growing mode therefore exists exactly above threshold. About the coherent density, the quadratic form is strictly dissipative transverse to collective rotation, whose rate is zero. These linear estimates concern the limiting equation. Relating them to finite stochastic trajectories requires a dynamical mean-field limit and branch selection; exchange symmetry alone does not establish the needed independence (Sznitman, 1991).

### 3.2 Installed energy bounds mean coupling

A spatial model replaces the finite sum by integration against a kernel. One declared implementation carries response modes with continuous profiles  $\phi_i$ , installation prices  $u_i$  and nonnegative occupancies  $c_i(z)$  in configuration  $z$ :

$$K_z(x, y) = \sum_i c_i(z) \phi_i(x) \phi_i(y), \quad U(z) = \sum_i u_i c_i(z). \quad (4)$$




Figure 2: Stationary self-consistency for identical frequencies and phase diffusion  $D = 1$ . The positive coherent branch appears continuously above  $K_c = 2D$  and increases with coupling. Zero remains a solution above threshold. The curve is computed from the integral expressions used in the formalization.

181 The same configuration sets the reciprocal kernel and stored energy. An  
 182 executed transition changes the kernel through its changed occupancies and  
 183 requires work  $U(z') - U(z) + Q$ , with  $Q$  positive into the reservoir. The  
 184 installed kernel is averaged over the resulting configuration law  $p(z)$ . Cell-  
 185 pair reconstruction on refining spatial partitions converges to its product-  
 186 measure integral under the specified compactness and sampling assumptions;  
 187 the refinement index is independent of the number of control updates.

188 Normalize the substrate measure  $\mu$  to one. Write  $m_i = \int \phi_i d\mu$  and  
 189 declare  $\kappa > 0$  with  $m_i^2 \leq \kappa u_i$  for every mode. Factorizing each mode's

190 integral gives

$$\begin{aligned}
 U_{\text{inst}} &= \sum_z p(z)U(z), \\
 K_{\text{mf}} &= \iint K(x, y) d\mu(x) d\mu(y) = \sum_z p(z) \sum_i c_i(z) m_i^2 \leq \kappa U_{\text{inst}}.
 \end{aligned}
 \tag{5}$$

191 If this mean coupling is the  $K$  of the scalar model,  $\kappa U_{\text{inst}} \leq 2D$  excludes a  
 192 coherent stationary branch and any growing Fourier mode about the uniform  
 193 density. Thus coherence on that branch requires  $U_{\text{inst}} > 2D/\kappa$ . The condition  
 194 is necessary: energy above it need not produce sufficient coupling because  
 195 the bound can be loose.

196 The quantity constrained is installed energy rather than dissipated heat,  
 197 and how much the condition excludes depends entirely on  $\kappa$ , which the mode  
 198 profiles, prices and substrate normalization supply: any macroscopic sub-  
 199 strate carries installed energy far above the thermal floor, so the inequality  
 200 binds through that conversion and not through the energy, and no result  
 201 here derives  $\kappa$  for cortex or for any other candidate. What the condition  
 202 contributes is a resource requirement in units a measurement can reach. Ap-  
 203 plying the scalar threshold to heterogeneous or signed spatial dynamics re-  
 204 quires a justified reduction, and identifying the installed kernel with a cortical  
 205 field is a separate physical commitment.

## 206 4 Compatibility: from coordinated activity 207 to a global description

### 208 4.1 Why common timing is insufficient

209 For overlapping patches  $U$  and  $V$ , compatibility means that restricting either  
210 local description to  $U \cap V$  gives the same object. Descriptions can agree on  
211 shared quantities while contributing different information elsewhere. Com-  
212 patible sections of the specified sheaf have a unique global extension. On  
213 a finite cover, local finite spatial measures agreeing as measures on every  
214 overlap also have a unique finite-measure extension whose sheaf image is the  
215 glued section. No mass-one normalization is imposed.

216 In the cup example, visual and reaching descriptions must express a  
217 shared quantity, such as cup location, in common coordinates. Restriction  
218 retains that quantity while dropping details particular to each description.  
219 Agreement permits surface appearance and hand configuration to contribute  
220 to one situation. The mathematical counterexample assigns the same phase  
221 to every patch while two local mass profiles disagree on a shared site. Those  
222 profiles have no common extension. Because they are chosen independently  
223 of phase, synchrony places no constraint on their disagreement.

224 This distinction limits the biological interpretation. Receptive territories  
225 and content domains must be related to spatial patches, and the descriptions  
226 must support the required extension. Matching overlap marginals of de-

227 coded probability laws need not determine a unique global joint distribution  
 228 (Abramsky & Brandenburger, 2011).

229 The cover is itself a choice, and the construction constrains it in one way  
 230 only: synchronized patches agree where they overlap. Where they do not  
 231 overlap there is nothing to compare, so a cover by pairwise disjoint patches  
 232 satisfies that obligation for *every* assignment of local descriptions and phases  
 233 (`ofDisjointCover`) and the gluing theorem still returns a unique global sec-  
 234 tion, which is the tuple it was handed (`glue_unique_of_disjoint`). Member-  
 235 ship of the class then reports nothing about the system beyond the patches  
 236 covering it. What a gluing claim asks of a cover is therefore a non-degeneracy  
 237 the theorem does not assume: that distinct patches meet (`HasNonempty-`  
 238 `Overlaps`), and that what they share is a quantity fixed independently of  
 239 the agreement it is being used to establish.

## 240 4.2 When coherence does constrain content

241 A declared relation between phase and content gives a positive result. Let  
 242 contents take values in a real normed vector space  $E$ . Patch  $i$  reads an  
 243 assigned oscillator through  $e_{i,x}$  at site  $x$ , evaluated on the circle point  
 244  $(\cos \theta, \sin \theta)$ . Assume a uniform Lipschitz constant  $L$  in chord distance and  
 245 require  $e_{i,x} = e_{j,x}$  wherever patches  $i$  and  $j$  overlap. Encoders may vary across  
 246 sites; their equality on overlaps is a supplied relation between descriptions of  
 247 the shared quantity. The order-parameter identity  $N^2 r^2 = \sum_{ij} \cos(\theta_i - \theta_j)$   
 248 implies  $1 - \cos(\theta_i - \theta_j) \leq N^2(1 - r^2)$  for each pair. Consequently every

249 overlap discrepancy is bounded by

$$\delta_r = L\sqrt{2} N\sqrt{1-r^2}. \quad (6)$$

250 Here discrepancy is the norm of the difference, and  $r$  and  $N$  are read on  
251 whichever population the bound is applied to. That choice is what decides  
252 whether the estimate says anything. A patch carries an order parameter of  
253 its own, and reading the bound there carries that patch's resultant and that  
254 patch's site count (`chord_le_of_patch_coherence`, `compatible_of_patch_`  
255 `coherence`), so a large population observed through small patches is con-  
256 strained by the small count. Read instead on the whole array, the factor  $N$   
257 makes the estimate weak, because it draws a pointwise guarantee from an  
258 average over  $N^2$  pairs. At  $r = 1$  the observations agree exactly at either  
259 scale. A shared scalar encoder is a special case. Dropping the overlap rela-  
260 tion permits disagreement even at perfect locking. A common nonexpansive  
261 linear projection preserves agreement, but agreement of a projection does  
262 not establish agreement of the full vector.

263 What Eq. (6) bounds is a chord, which never exceeds 2, so the estimate  
264 constrains anything only where  $\sqrt{2} N\sqrt{1-r^2}$  falls below that. The winding  
265 sweep of Section 7.3 fixes both ends of that range on one sheet. Read on the  
266 sweep's own patch, 625 sites at patch-local order 0.9015, the bound is  $382.5 L$   
267 and constrains nothing; holding that order fixed, the count alone brings it  
268 below the maximum chord only at 3 sites or fewer. A nearest-neighbour pair

269 of the same winding, the smallest patch there is, gives  $0.1388 L$ . Shrinking  
 270 patches therefore thins the overlaps at the same rate as it tightens the bound,  
 271 and the two ends are not reached by the same cover. At a columnar grain  
 272 the estimate constrains agreement inside one locally synchronous population  
 273 and is vacuous across the distant territories whose agreement unity is about.

274 Mix retained fields with observations at gain  $\eta \in [0, 1]$ :  $s_i \mapsto (1 - \eta)s_i +$   
 275  $\eta o_i$ . Content and observation discrepancies bounded by  $\varepsilon$  and  $\delta$  give a next  
 276 discrepancy at most  $(1 - \eta)\varepsilon + \eta\delta$ . Repeated observations bounded by  $\delta$   
 277 therefore give

$$\varepsilon_n \leq (1 - \eta)^n \varepsilon_0 + \delta. \quad (7)$$

278 Thus agreeing observations preserve exact compatibility and, for  $\eta > 0$ , re-  
 279 duce an initial mismatch. Approximate observations leave a nondecaying  
 280 term in the error bound. Taking a uniform bound on Eq. (6) connects this  
 281 update to the phase process.

### 282 4.3 What approximate agreement determines

283 Exact gluing requires exact restrictions. For nonnegative profiles on a finite  
 284 substrate, suppose every overlap discrepancy is at most  $\varepsilon$  in uniform distance.  
 285 Weights that vanish outside their patches and sum to one at each site select  
 286 a global profile by weighted averaging. The selection lies within  $\varepsilon$  of each  
 287 patch there, and any two such selections differ by at most  $\varepsilon$ , independently  
 288 of cover multiplicity. An  $L$ -Lipschitz readout consequently varies by at most

289  $L\varepsilon$  across selections.

290 At zero tolerance, the selection is the unique exact extension. At every  
291 positive tolerance there are families with no exact extension; selection then  
292 gives an approximation whose dependence on weights is bounded. Supple-  
293 mental Material gives a three-site witness attaining the bound. The mass-  
294 profile construction uses the scalar case of the content dynamics and addi-  
295 tionally requires nonnegativity of the initial profiles and observations. The  
296 general vector result supplies no identification with the spatial-measure sheaf.

297 These constructions concern declared objects: a specified cover, contents  
298 valued in a normed space, and encoders related on overlaps. They select  
299 neither the neural cover nor the decoder, and identifying their contents with  
300 perceptual content, or establishing that the identification bears on experi-  
301 ence, is a further modelling task; Section 7 sets out what a measurement of  
302 it would have to establish.

## 303 **5 Self-representation: a state that models it-** 304 **self**

### 305 **5.1 The reflexive construction**

306 Compatibility gives a global state. Reflexivity relates it to an encoding within  
307 that state. Equip global sections with a complete metric. If the additional  
308 restriction-resonance condition identifies encoding with restriction to a region

309  $U$ , the reconstruction map is

$$F(s) = \text{readout}(s|_U). \quad (8)$$

310 The region and readout are fixed across the state space. If  $F$  contracts and  
311 fixes the section  $s_*$  formed from the local descriptions, Banach's theorem  
312 makes  $s_*$  its unique fixed point (Banach, 1922). The formal bridge E89  
313 supplies the contraction and  $F(s_*) = s_*$ ; restriction resonance is separate.  
314 Supercritical coupling makes the stipulated factor  $q = e^{-(K-2D)\tau/2}$  less than  
315 one for  $\tau > 0$ . Constructing a map with that factor remains a modelling  
316 obligation.

317 In the cup example, the global state includes the represented scene and the  
318 perceiver's relation to it. A local encoding must retain enough information  
319 for the readout to recover that modelled state; cup position alone would be  
320 insufficient if hand configuration and viewpoint could not be reconstructed.

321 For the same map with Lipschitz factor  $q < 1$ , approximate reconstruction  
322 gives

$$d(s, F(s)) \leq \varepsilon \implies d(s, s_*) \leq \frac{\varepsilon}{1 - q}. \quad (9)$$

323 At the stipulated rate this radius grows as  $2\varepsilon/((K-2D)\tau)$  near threshold, for  
324 fixed positive  $\varepsilon$  and  $\tau$ . Exact uniqueness survives while the error guarantee  
325 weakens.

## 326 5.2 Reconstruction across a family of states

327 Sensitivity can be required quantitatively. Declare relevant macrostates, an  
 328 encoder  $E$  and readout  $R$ , and require  $d(s, R(E(s))) \leq \varepsilon$  throughout that  
 329 family. If two states have the same code, the triangle inequality gives

$$E(s) = E(t) \implies \max\{d(s, R(E(s))), d(t, R(E(t)))\} \geq \frac{1}{2}d(s, t). \quad (10)$$

330 Thus  $M$  relevant states pairwise more than  $2\varepsilon$  apart require at least  $M$   
 331 distinguishable codes; an alphabet of at most  $2^b$  codes requires  $M \leq 2^b$ .  
 332 Applied to the reflexive boundary,  $E$  is its local encoding and  $R \circ E$  is the  
 333 existing self-prediction map.

334 This is an additional accuracy criterion on a family, with no requirement  
 335 for multiple exact fixed points. A blind encoder or constant readout fails it  
 336 for separated states despite having a unique fixed point.

337 Where the encoding is read as the state's own restriction to a region, the  
 338 criterion becomes a condition on the region. Restriction resonance requires  
 339 the encoder and that restriction to identify the same pairs of states, so a  
 340 boundary reconstructing two relevant states more than  $2\varepsilon$  apart while its re-  
 341 gion reads them identically is not resonant (`not_isRestrictionResonance_`  
 342 `of_reconstructs`); read forwards, a resonant boundary that reconstructs  
 343 a pairwise separated family restricts injectively on it (`restrictToAvatar_`  
 344 `injOn_of_isRestrictionResonance`). Which states a region reads identi-  
 345 cally is fixed by where the region sits, before an encoding is chosen, so the

346 resolution the family demands is the region's and no readout supplies it. The  
 347 implication runs one way: a region that does resolve the family is not thereby  
 348 shown to carry a resonant encoding, and E89 supplies no such resolution.

349 Those two demands are jointly satisfiable with nothing folded anywhere.  
 350 Take the region to be the whole substrate: restriction is then the identity, so  
 351 the boundary is resonant, reconstructs every declarable family exactly, and  
 352 restricts injectively on all of them (`topAvatar_satisfies_obligations`). A  
 353 properness condition on the region (`IsProperAvatar`) is therefore a separate  
 354 requirement rather than a consequence of resonance and resolution, and what  
 355 a fold into a local region is asked for. Its quantitative form replaces the  
 356 alphabet by a metric. With an  $L$ -Lipschitz readout, a family reconstructed  
 357 within  $\varepsilon$  whose members lie pairwise more than  $r$  apart has  $\delta$ -separated codes  
 358 whenever  $\delta L \leq r - 2\varepsilon$ , so the family is no larger than the  $\delta$ -packing number  
 359 of what the region carries (`encard_le_packingNumber_restrictToAvatar`).  
 360 That is the shape a continuous family of states needs, and at the whole  
 361 substrate it says nothing, because the region's range is everything. The gain  
 362  $L$  is declared and derived here for no physical readout, and at unbounded  
 363 gain the bound says nothing.

### 364 **5.3 Agreement after a change takes communication**

365 Snapshot compatibility does not specify how quickly descriptions respond  
 366 when a shared quantity changes. In a synchronous network whose updates  
 367 read only the site's own state and declared neighbours' messages, two initial

368 configurations agreeing on a site’s causal past through  $T$  rounds produce the  
 369 same state there at round  $T$ . The deadline consequence is the one Section 2  
 370 states, and it follows from the reconstruction bound in Section 5.2: a report  
 371 from that site is a readout of a code that the two interventions leave identical.

372 A three-site delay line witnesses failure after one round and exact report-  
 373 ing after two; cutting the path prevents delivery at every round. A report  
 374 may still happen to be correct about one fixed world. The bound concerns  
 375 responsiveness across interventions.

376 Throughout this section *prediction* names a reconstruction map, and the  
 377 relevant family of states, the metric, the tolerance, the communication graph  
 378 and the deadline are all declared inputs, as are the represented variables and  
 379 the fidelity counted as accurate. Reading the map as temporal neural pre-  
 380 diction would require dynamics; reading the resulting code count as physical  
 381 memory capacity would require a hardware model, which a state-dependent  
 382 readout would have to extend.

## 383 6 Thermodynamics and composition

384 The phase and representation results concern coupling, content disagreement  
 385 and reconstruction error. Thermodynamics constrains the physical opera-  
 386 tions that could implement them, and it does so from the side rather than  
 387 along the spine: no edge below derives coherence, compatibility or reconstruc-  
 388 tion from a heat bound, and the resource condition these edges terminate in

389 is necessary without being restrictive for any macroscopic candidate (Sec-  
 390 tion 3.2). What this half contributes is a ledger in which a resource claim  
 391 is stated in checkable units. Finite-register capacity obeys  $H(p) \leq \log |X|$ ;  
 392 erasure heat depends on specified system–bath assumptions (Landauer, 1961;  
 393 Norton, 2005; Shenker, 2000). A separate boundary result concerns a field  
 394 reaching disconnected vacuum components (Kibble, 1976). The composition  
 395 connects these objects through the explicit hypotheses in Table 1.

## 396 **6.1 Feedback changes the information constraint**

397 For passive memory  $X_t$  driven by an environmental signal  $S_t$ , the predictive  
 398 bound is

$$W_{\text{diss}} \geq k_B T [I(X_t; S_t) - I(X_t; S_{t+1})]. \quad (11)$$

399 The Markov structure  $X_t \rightarrow S_t \rightarrow S_{t+1}$  makes the bracket nonnegative (Still  
 400 et al., 2012). Zero retained information also satisfies it, so a task and a  
 401 learning rule require additional specification. Relating this bound to Lan-  
 402 dauer heat requires a common thermal scale and budget, and memory waste  
 403 bounded by erased entropy.

404 An acting agent changes the signal it will receive. Specify policy, world,  
 405 observation and memory-update channels. For their finite laws,

$$\begin{aligned}
 I(X_t; S_t) - I(X_t; S_{t+1}) &= I(X_t; S_t \mid S_{t+1}) \\
 &\quad - I(X_t; S_{t+1} \mid S_t).
 \end{aligned} \quad (12)$$

406 The second conditional term vanishes in the passive model. With feedback,  
 407 the signed difference can be negative as action creates correlations with the  
 408 future. A causal reading requires an environmental state sufficient to describe  
 409 the relevant dynamics.

410 For a controlled substep, the active thermodynamic bound instead uses  
 411 the joint entropy. Forward and reverse path laws give  $\Sigma = D_{KL}(P_F \| P_R) =$   
 412  $\Delta H(X, S) + \langle Q \rangle / \theta \geq 0$  under local detailed balance, with  $\theta = k_B T > 0$  and  
 413 heat positive into the reservoir (Horowitz and Esposito, 2014; Ito and Sagawa,  
 414 2013). Every forward path of positive probability must have a positive-  
 415 probability reverse for this finite identity. A missing reverse gives infinite  
 416 divergence. The supplement develops the support conditions and composes  
 417 actuation with memory updates on their shared intermediate law.

## 418 **6.2 Learning and repeated operation need explicit re-** 419 **sources**

420 The finite examples distinguish selecting a feasible policy, relaxing a register  
 421 toward an encoded reward, and learning an unknown task from observa-  
 422 tions. The observational learner acquires correlation with an environmental  
 423 parameter while its action marginal stays uniform. A continuing model exe-  
 424 cutes actuation, observation-driven learning and reset in sequence, preserving  
 425 above-chance performance at a positive work cost per cycle.

426 A phase-derived sensor makes the informational distinction concrete: an

427 informative readout and a blind control have the same phase order but dif-  
 428 ferent learning performance. Attaching the learner’s register to the actuator  
 429 of Section 3.2 ties reward, installed kernel and work to its own executed law.  
 430 The learning witness rises above the necessary installed-energy bound while  
 431 its matched control remains on the exclusion line. These finite constructions,  
 432 detailed in Supplemental Material, leave the evolving feedback between phase  
 433 dynamics, installation and sensing to be specified.

434 For that model, common energy  $E$ , thermal scale  $\theta$ , local detailed balance  
 435 and positive support at each stage give

$$\theta[H(p_0) - H(p_N)] \leq Q_{<N} = W_{<N} + \langle E \rangle_0 - \langle E \rangle_N \leq B + \langle E \rangle_0 - \langle E \rangle_N, \quad (13)$$

436 where expected cumulative work stays within the supplied allowance  $B$  at  
 437 every prefix. This bounds expectations. Separate constructions track a store  
 438 on each trajectory, include finite sources, charge sensor clearing and prepend  
 439 preparation. They specify when a mean allowance represents an available  
 440 resource; an expected budget alone permits overdrawn paths.

441 These constructions establish costs for their declared processes. A finite  
 442 source with a positive net cost permits only finitely many funded cycles,  
 443 while continuing operation requires replenishment. Register–bath witnesses  
 444 also distinguish actual bath-energy transfer from transition-log heat. Iden-  
 445 tifying a work store, an erasure budget and the installation energy of Sec-  
 446 tion 3.2 requires physical relations that these separate models do not supply

447 automatically.

### 448 6.3 What the composition establishes

449 The passive theorem `chain` and feedback theorem `chain_active` each ac-  
 450 cept eight connecting implications (Table 1). In the active branch, a named  
 451 substep at the register’s thermal scale satisfies

$$\theta > 0, \quad \theta[H(p) - H(p')] \leq \langle Q \rangle \leq B, \quad (14)$$

452 with  $B$  the register’s heat. The upper allocation follows from a ledger whose  
 453 operations deliver that total heat to one reservoir and whose other operations  
 454 do not increase joint entropy. Those ledger assumptions are physical inputs.  
 455 Neither Landauer’s lower bound nor microscopic reversibility supplies them.

Table 1: Connecting hypotheses, eight per branch, and the installed-energy consequence. The branches share E12, E23 and E56–E89. Exact overlap compatibility and the encoding–readout structure are additional inputs. Supplemental Material, Table S1, maps the node-level results.

| Connection                          | Status               | Required assumption                                                                                                 |
|-------------------------------------|----------------------|---------------------------------------------------------------------------------------------------------------------|
| E12<br><i>capacity and boundary</i> | Physical<br>premise  | A finite-capacity system realizes a field leaving its vacuum manifold. Capacity and boundary are independent roots. |
| E23<br><i>boundary to register</i>  | Bridge<br>assumption | The boundary is associated with a finite register whose update leaves states unreachable.                           |

Table 1 (continued)

| Connection                                               | Status                  | Required assumption                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
|----------------------------------------------------------|-------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| E34 /<br>E34Active<br><i>register to<br/>dissipation</i> | Bridge<br>assumption    | Passive: predictive memory at the register's thermal scale and heat budget, with nonzero waste. Active: positive paths, local detailed balance and allocation of register heat to the named substep; the allocation uses the ledger identity and other operations' compressiveness.                                                                                                                                                                                                                        |
| E45 /<br>E45Active<br><i>couplings to<br/>continuum</i>  | Modelling<br>assumption | Convergent spatial coupling energies. Declared mode hardware supplies a kernel limit for either branch. A learning witness computes installation and work from its own executed law. Refinement and physical identifications remain inputs.                                                                                                                                                                                                                                                                |
| E56<br><i>field<br/>identification</i>                   | Physical<br>commitment  | A specified physical field realizes the limiting coupling and satisfies the kernel conditions.                                                                                                                                                                                                                                                                                                                                                                                                             |
| E67<br><i>above<br/>threshold</i>                        | Physical<br>commitment  | That field operates above the synchronization threshold.                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| E78<br><i>coherence to<br/>cover</i>                     | Bridge<br>assumption    | Coherence relaxes the supplied cover at coupling at least the named mean-field value. Shared phase encoders bound vector-content mismatch; the exact cover and timely responsiveness remain separate requirements. Patches and oscillators carry one index, so one oscillator carries exactly one content patch: a modelling assumption rather than a consequence, and not an innocent one, since field-mediated coupling is spatial while two territories describing one quantity need not be neighbours. |
| E89<br><i>contraction to<br/>fixed point</i>             | Modelling<br>assumption | A map mediated by local self-encoding fixes the glued section and has the stipulated Lipschitz factor $q < 1$ . Reconstruction over a declared family imposes an additional code-separation criterion.                                                                                                                                                                                                                                                                                                     |

Table 1 (continued)

| Connection       | Status              | Required assumption                                                                                                                                                                                                            |
|------------------|---------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Installed energy | Conditional theorem | Under Section 3.2’s occupancy and mode-price conditions, E56 and E67 require $U_{\text{inst}} > 2D/\kappa$ . The learning witness clears this necessary bound; its matched control does not. Neither physical premise follows. |

Both branches also take a thermodynamic cover carrying exact overlap compatibility and a reflexive boundary specifying encoding and readout. They conclude **UnifiedSelf**: the glued section is the unique fixed point of self-prediction. Compatibility and accuracy at that section enter as premises; uniqueness follows. Identifying encoding with restriction needs the separate resonance condition of Eq. (8).

Each of the eight edges is supplied rather than derived from the one above it. E12 is equivalent to its conclusion, so capacity and boundary are independent roots rather than a step from one to the other; both E45 discharges take their convergence from refinement data and not from the thermodynamic bound; E78 states that coherence relaxes the supplied cover, and the coherence-to-content estimate of Eq. (6) bounds disagreement only for encoders already shared on overlaps. The inferential content of the composition is its last step, where contraction and a fixed glued section give uniqueness. What composing them establishes is that the premises are consistent, named in one place and machine-checked together, and not that coherence follows from installed energy or self-representation from coherence.

Joint witnesses combine a register, boundary, refining mesh and small

474 spatial cover; active witnesses add controlled thermal updates and a priced  
 475 installation. They establish satisfiability under their stated identifications.  
 476 Mesh convergence follows from refinement data, and the witnesses establish  
 477 no common cortical mechanism. Supplemental Material maps the claims to  
 478 their Lean declarations and records the counterexamples. The development  
 479 declares no additional axioms beyond its libraries' foundational axioms.

## 480 **7 What the observations establish**

481 The composition holds under its premises. Whether observation can estab-  
 482 lish those premises is a separate question, and the three requirements it  
 483 connects concern different objects: phase dynamics, local descriptions and  
 484 internal encodings. Their observables must retain these distinctions. The  
 485 following controls show which mechanisms remain indistinguishable, which  
 486 temporal approximations fail, which states one observable cannot separate,  
 487 and what independent content measurements must establish. Each is a finite  
 488 numerical regime on a declared model, with full parameters and controls in  
 489 Supplemental Material, and none of them calibrates a cortical quantity.

### 490 **7.1 Stationary distributions leave the mechanism un-** 491 **determined**

492 Equation (2) gives a concentration-coherence relation with no fitted shape  
 493 parameter. Estimate  $r$  from circular resultant length and  $a$  from density

494 shape: inverting  $R$  from the measured  $r$  would make agreement automatic.  
 495 Even independently estimated agreement identifies a statistical family shared  
 496 by different mechanisms.

497 Let  $q(t)$  cross 2, let  $r(q)$  be its stationary branch, and fix  $D_0 > 0$ . Increas-  
 498 ing coupling  $K(t) = D_0 q(t)$  at diffusion  $D_0$  and decreasing diffusion  $D(t) =$   
 499  $K_0/q(t)$  at coupling  $K_0$  both give concentration  $a = q(t)r(q(t))$ . Uncoupled  
 500 oscillators with prescribed drift  $-h(t) \sin(\theta - \psi)$ , where  $h(t) = D_0 q(t)r(q(t))$ ,  
 501 and diffusion  $D_0$  have that same density. Their stationary distributions co-  
 502 incide window by window despite different causes.

503 Resolved dynamics supply additional information. In the ideal model with  
 504 continuous paths, quadratic variation accumulates at rate  $2D$  and conditional  
 505 increment means constrain drift. Separating endogenous coupling from a  
 506 matched common drive additionally requires a measured drive or controlled  
 507 perturbation; identifying electromagnetic coupling requires evidence specific  
 508 to that interaction. Supplemental Material gives the equivalence construction  
 509 and its numerical check.

## 510 **7.2 Finite ramps change the observable onset**

511 The square-root law in Eq. (3) describes stationary solutions. A changing  
 512 coupling requires time for the phase distribution to approach those solutions.  
 513 In a deterministic mean-field control that crosses the threshold symmetrically,  
 514 none of the scanned rates meets the declared tracking tolerance: slowing  
 515 the crossing does not remove the terminal residual under that initialization.

516 A slow control variable therefore supplies no general guarantee of branch  
 517 tracking at onset. Supplemental Material gives the residual, rate range and  
 518 admissible rates away from threshold.

519 The finite-population ramp study uses  $N = 2000$ ,  $D = 1$  and 32 replicas  
 520 at 8 coupling speeds from  $v = 0.1$  to  $v = 0.0001$ . Escape is the first of three  
 521 consecutive saved samples at  $r \geq 0.2$  after threshold. Faster ramps delay  
 522 escape (Fig. 3); only 5/32 replicas escape within the fastest window. The  
 523 6 uncensored legs supply the delay scaling below; the decade-spaced ones  
 524 among them have mean coupling delays 0.1947, 0.0684 and 0.0253. Fitted  
 525 onset exponents differ from those of the exact stationary branch evaluated  
 526 on the same grids. At the fastest speed the ensemble mean never reaches the  
 527 fit window, so its fitted slope does not estimate stationary onset.

528 Delayed passage through a noisy bifurcation (Berglund & Gentz, 2002)  
 529 makes the apparent exponent depend on ramp speed and fitting window.  
 530 The specified deviation-based critical ramp speed is not bracketed by this  
 531 sweep. With fluctuating coupling, the relevant stationary comparison also  
 532 depends on rate: the mean coupling approximates order in the tested fast  
 533 regime, while averaging stationary order over coupling works for the tested  
 534 slow drives confined to one side of threshold. Slow crossing drives can satisfy  
 535 neither approximation.

536 **The prediction.** The delay scaling is what this section supports as a pre-  
 537 diction, and it needs no absolute calibration. Over the 6 uncensored legs the

mean delay follows  $\Delta K \propto v^{0.444}$ , with a 95% slope interval of  $[0.394, 0.494]$ , which does not contain  $1/2$ . What that shortfall measures is the escape criterion and not the scaling. Applied to a deterministic mean-field run seeded at the fluctuation floor, the same estimator returns 0.425 over the same legs at the same escape criterion, inside the interval on the measurement. That run has no noise to delay it, so the shortfall is the criterion's: it crosses the saturation between the floor it starts from and the level it is stopped at, and tightening the criterion to  $r \geq 0.05$  returns 0.490. The bias is reproduced where the answer is known, which is what licenses reading the measurement as the law rather than as a competing exponent.

Stated as a law rather than as a number: onset delay scales as the square root of the rate at which coupling crosses threshold. Testing it requires only that  $\dot{K}$  be monotonic in a manipulated rate — not its value, nor  $\kappa$ , nor  $\gamma$  — so it is a within-subject design that varies the rate of emergence from anaesthesia alone, with each participant as their own control. Such a design must avoid the hazard this sweep exhibits: a fixed absolute escape criterion meets a size-dependent critical fluctuation floor, so the criterion has to be referred to each recording's own pre-threshold fluctuation scale rather than to a shared number. The diffusion  $D$  is the remaining input, and Section 7.1 notes that it is estimable from resolved phase traces through the rate  $2D$  at which quadratic variation accumulates, which is a different quantity from the concentration the propofol exercise could not range over.

Additional controls address integration bias, population size and spatial




Figure 3: Finite-rate departures from the stationary branch. Ensemble-mean order versus coupling for four ramp speeds  $v$ , with one-standard-error bands across replicas. The dashed vertical line is the static threshold and the dotted horizontal line the finite-size reference. Faster ramps delay escape, changing the apparent recovery onset.

561 or signed coupling; the refined signed-network crossing bracket  $K_{\text{eff}}/D \in$   
562  $[1.958, 2.009]$  contains 2.

### 563 **7.3 A winding state leaves the order parameter am-** 564 **biguous**

565 The order parameter  $r$  is indifferent to where a phase sits in space. A field  
566 winding once across a periodic sheet therefore has  $r = 0$  while every patch  
567 of it stays locked. Such a field is not an artificial construction here. On a

568 torus a winding is stationary for any isotropic symmetric kernel at identical  
 569 frequencies: the coupling term is odd in the separation against a kernel even  
 570 in it, so the drift cancels pair by pair. `char_is_kuramoto_trajectory` proves  
 571 this for any finite periodic array, this sheet included. A winding satisfies the  
 572 positivity and identical-frequency hypotheses. It violates only the quarter-  
 573 turn initial spread that the convergence result additionally requires before  
 574 placing the limit at the uniform state.

575 On the sheet of Section 3's spatial control, a winding survives below  
 576 an interpolated decay length of 0.1871 mm and unwinds above it (Fig. 4).  
 577 That boundary is interior to the 0.1–0.3 mm band. At 0.1 mm the same  
 578 sheet at the same coupling, diffusion and kernel reaches  $r = 0.9657$  from a  
 579 uniform start and  $r = 0.0083$  from a winding one. Both are stationary, with  
 580 patch-local order 0.9015 and no winding defect. At fixed  $K$  the observable  
 581 therefore takes two values spanning nearly its whole range. Coupling does  
 582 not determine the resultant, and a low resultant does not by itself distinguish  
 583 weak coupling from a winding field.

584 Under the quenched frequency spread of  $1.5 \text{ rad s}^{-1}$  that the spatial con-  
 585 trol carries, the boundary moves to 0.1421 mm. The same comparison at  
 586 0.1 mm gives 0.9418 against 0.0348, holding across 3 frozen realizations.  
 587 Heterogeneity shortens the range that sustains a wave without removing it.

588 Which windings persist has a criterion behind it. Linearizing the phase  
 589 model about a winding returns a coupling of the same translation-invariant  
 590 form, each separation reweighted by the cosine of the phase the winding




Figure 4: A winding phase field on the spatial sheet, at identical frequencies. Left: global order (solid) against patch-local order (dashed) across a subset of coupling decay lengths. Right: the twisted and uniform initial conditions compared across lengths, with the shaded 0.1–0.3 mm band and the interpolated retention boundary. Below that boundary the two initial conditions reach different stationary values of  $r$  at the same coupling, diffusion and kernel, so the resultant does not report the coupling. The sweep is sampled at the lengths marked; it establishes no transition and no biological identification.

591 advances across it (`hasDerivAt.charJacobian`). That reweighted kernel is  
 592 again even in the separation, so the same characters diagonalize it and ev-  
 593 ery mode's rate is one of its eigenvalues. Where the winding advances by  
 594 at most a quarter turn across every separation the coupling reaches, the  
 595 reweighted kernel is nonnegative and no rate is positive (`charJacobian_`  
 596 `stable_of_quarter_turn`); where some rate is positive, that mode grows  
 597 without bound (`charJacobian_growing_mode`). Moving coupling mass to  
 598 longer separations is what carries a phase advance past a quarter turn, and  
 599 that is the mechanism the decay-length sweep varies. The criterion bounds  
 600 rates rather than trajectories: asymptotic stability is not proved, so the

601 states that persist occupy a narrower range than it admits, and the bound-  
602 ary quoted above is measured and not derived from it. The same kernel is the  
603 potential's curvature, so a winding at  $r = 0$  is not the global minimum the  
604 convergence result places its limit at, and one carrying a growing mode is not  
605 a local minimum either: that result's quarter-turn hypothesis must exclude  
606 a winding rather than merely decline to cover it (`char_not_potential_min`).

607 An amplitude layer decides which of these states exist, and leaves what  
608 the resultant reports unchanged. Every oscillator of the phase model sits on  
609 the unit circle, so a field with an amplitude node has no representation in  
610 those dynamics. The real Stuart–Landau extension supplies one on the same  
611 coupling, and leaves intact the gradient structure the convergence results rest  
612 on. The state becomes  $z_i \in \mathbb{C}$  with  $\dot{z}_i = (\mu - |z_i|^2)z_i + \sum_j A_{ij}(z_j - z_i)$ . Its  
613 phase equation is the Kuramoto equation with each coupling weighted by a  
614 ratio of amplitudes (`amplitude_phase_is_kuramoto`). The phase model is  
615 this one with the amplitude ratios set to one, rather than a separate object.

616 What the added degree of freedom decides is which windings exist at all.  
617 On a periodic array of any dimension, this sheet included, a winding requires  
618  $\mu + \lambda > 0$ . Here  $\lambda = \sum_d f_d(\cos \phi_d - 1)$  is the eigenvalue the kernel supplies,  
619 summed over separations, with  $\phi_d$  the phase the winding advances across  $d$   
620 (`exists_charState_iff`; on the ring  $\phi_d = 2\pi qd/n$ ). For the nonnegative  
621 kernels used here  $\lambda \leq 0$ , and moving coupling mass to longer separations  
622 lowers it further. A broad enough kernel therefore removes the winding out-  
623 right. The sweep above reaches that same boundary by a different route:

624 the phase model retains every winding at every coupling, and loses this one  
625 under integration instead. The two layers bound the same regime from op-  
626 posite sides. A phase recording does not separate a state the amplitude layer  
627 excludes from one the dynamics fails to sustain.

628 Three limits bound the layer for a reader of the cortical literature. These  
629 states are windings and not spirals, and on the sheet that is a theorem rather  
630 than a shortage of dimensions: equal modulus at every site puts the phase  
631 circulation at exactly zero on every plaquette, and on a periodic sheet the  
632 circulations of *any* field sum to zero, so a lone defect is not a state of this  
633 geometry at all. Relaxational dynamics has no propagation speed, which is  
634 the quantity a cortical wave measurement leads with. And  $|z|$  is a normal-  
635 form radius, which a measured envelope is not; this work identifies the two  
636 nowhere. Supplemental Material gives the sweep, its uniform control, the  
637 amplitude layer, the standing wave it admits inside the band, the potential's  
638 curvature at a winding and the defect theorems, together with the remaining  
639 limits of both.

640 A vanishing resultant is the absence of global agreement and not of agree-  
641 ment. Eq. (6) read on a nearest-neighbour patch of a winding gives a dis-  
642 crepancy bound of  $2\sqrt{2}L|\sin\phi|$ , with  $\phi$  the phase advanced across the patch,  
643 which falls to zero as the same winding is spread over more sites (`chord_le_`  
644 `of_char_patch`), while the global form of the same bound is vacuous at  $r = 0$ .  
645 Encoders shared on overlaps therefore still bound content disagreement in-  
646 side a patch of a state that no global observable calls coherent. The winding

647 number itself stays unconnected to content. It is, however, a property of the  
 648 state rather than of the cover read on it. On a ring the whole turns dropped  
 649 by the winding's affine local lifts are zero at every site but the wrap, where  
 650 the residue resets, and they sum around the loop to  $q$ ; a nonzero  $q$  obstructs  
 651 every global real-valued phase lift, so one state at one coupling has resultant  
 652 exactly zero and degree  $q$  (`winding_degree_obstructs`). Reweighting the  
 653 sites moves the first quantity and redrawing the loop's intermediate transi-  
 654 tions leaves the second where it was. The scope is the ring's: this is a loop's  
 655 winding and not a spiral, the transitions are those of the affine lifts rather  
 656 than of an unwrapped measurement, and the integer is identified with no  
 657 cortical observable.

658 This qualifies what a phase trace can establish. An onset criterion stated  
 659 in  $r$  alone registers a change of basin as readily as a change of coupling.  
 660 The protocol of Section 8.3 therefore needs a spatial observable beside the  
 661 resultant.

## 662 **7.4 Coherent adaptation can lose environmental align-** 663 **ment**

664 An adaptive coupling rule might supply representational structure, but its  
 665 objective must reward the relevant agreement. For symmetric coupling, pair-  
 666 wise sine terms cancel in the population mean drift. Minimizing squared drift  
 667 at that fixed mean therefore rewards equalizing drift across nodes. In the

frequency-cluster control, this favors coupling between unlike-frequency clusters, whereas the diagnostic environmental template rewards coupling within clusters.

The formalization proves descent of the squared-drift objective for a specified gradient flow at fixed phases. Its functional differs from thermodynamic entropy production: at a coherent stationary density the probability current vanishes while phase-averaged squared drift is positive. Supplemental Material gives the operator, its gradient and the conditions of the descent result.

The finite control co-evolves noisy phases and symmetric, nonnegative couplings at fixed total resource. Its template is a diagnostic covariance of latent signals, never a training input. Across three seeds, second-half order remains between 0.9167 and 0.9222, while the within-cluster to between-cluster coupling ratio falls from 0.9824–0.9948 to 0.3507–0.6135 (Fig. 5). Random-update, permuted-gradient and frozen-kernel controls are reported in Supplemental Material.

The immediate post-update objective reduction disappears when averaged over complete update intervals. A prespecified grid of 6 rate–cadence combinations contains no case meeting the 5% sustained-reduction criterion while retaining coherence and losing alignment. The finding is coherent adaptation away from the specified structure, with no demonstrated sustained cost reduction.

Kernel alignment and overlap compatibility are different observables. This control addresses acquisition of its designated template under its par-

691 ticular objective. Learning models can explicitly include data in an objective  
 692 (Corberi et al., 2026); an account of agreement about the cup must specify  
 693 the content its learning rule is meant to preserve.




Figure 5: Coherence persists while coupling loses alignment with the frequency-cluster template. Gradient updates (solid), random updates (dash-dotted), permuted gradient updates (dash-dot-dotted), and a frozen kernel (dashed) share one phase-noise stream per seed. Panels show the immediate post-update squared-drift objective, phase order, within-cluster to between-cluster coupling, and template distances. The gradient arm preserves order while its coupling ratio falls; growth in both distances mainly reflects kernel norm. The objective panel samples the instant after updates; complete-interval measurements find no sustained reduction in the bounded study.

## 694 **7.5 Content agreement needs independent decoders**

695 A neural test needs independently identified domains, local descriptions and  
696 restriction maps. Two decoders should estimate a shared content on over-  
697 lapping domains before their agreement is examined. Discrepancy requires  
698 a null under decoding noise and controls using deliberately compatible and  
699 incompatible examples. An implementation on constructed data separates  
700 these cases at  $\text{AUC} = 0.986$ , on a 95% interval of  $[0.980, 0.992]$  resampled by  
701 trial; the compatible construction scores 0.476 on an interval containing one  
702 half. Shared-prior shrinkage can nevertheless manufacture agreement, and  
703 regional bias can manufacture disagreement. Per-patch decoding diagnostics  
704 detect these failures; the compatibility statistic alone does not.

705 Phase agreement cannot replace that measurement. On constructed con-  
706 tent, phase carries a share 1.000 of the content entropy when content is built  
707 from it and 0.000 when it is not. A phase-derived statistic reports com-  
708 patibility on phase-locked patches whose local descriptions disagree. The  
709 implemented content statistic detects their incompatibility. These controls  
710 validate a measurement on known constructions; a neural content model and  
711 decoder remain to be established.

712 Choosing the cover also chooses a candidate subject: trivial descriptions  
713 can agree while carrying little content, and a partition can conceal disagree-  
714 ment. The useful choice must therefore be constrained by what neural ac-  
715 tivity encodes. The exact composition assumes compatibility of its supplied  
716 cover. The content mechanisms above identify additional hypotheses under

717 which agreement can arise; they leave the empirical choice of content and  
718 the connection to that cover explicit.

## 719 **7.6 What an informative phase-density measurement** 720 **requires**

721 The concentration range and the dependence structure decide whether the  
722 concentration-coherence relation can be tested at all. Under the specified  
723 log-density estimator and bias-floor criterion, discrimination at the observed  
724 concentration ceiling  $a = 0.542$  requires 3000 independent sites in the scanned  
725 designs, rising to 31000 when sites share a mean in groups of 10. These  
726 counts describe that estimator and dependence model; the intended recording  
727 requires its own calibration. Supplemental Material reports the bin-count  
728 controls and the full dependence-aware design.

729 That ceiling comes from exercising the estimators on public propofol-  
730 sedation EEG (Bajwa et al., 2025), pooling channels in 100-ms windows.  
731 Bipolar and common-reference montages produce different residuals. The  
732 bipolar results are compatible with the Bessel relation over a concentration  
733 range too narrow to distinguish it from its linear approximation at the ob-  
734 served scatter, which is the range the design scan takes as given. Temporal  
735 pooling changes the estimand, and sedation recordings leave the awakening  
736 hypothesis untested.

737 An informative dataset therefore needs sufficient concentration range or

precision, calibrated estimators and an observation model for dependence,  
source mixing and reference choice. Public mouse calcium imaging provides  
thousands of spatially located neurons across waking, sleep and anesthesia  
(Oomoto et al., 2026), but its 7.65 Hz frame rate is below the phase-  
estimation band used here. Site count, state coverage and bandwidth are  
separate requirements.

## 8 The cortical hypothesis and an awakening test

The cortical proposal identifies a physical source of coupling and asks whether  
its changes help explain recovery. The tests in Section 7 determine what  
observations must distinguish before a phase trace can support this identification.

### 8.1 Physical identification and calibration

The cortical hypothesis identifies part of effective coupling with endogenous  
extracellular electromagnetic interactions. Fields can modulate spike timing  
(Fröhlich & McCormick, 2010; Anastassiou & Koch, 2015), and field models  
propose coordination of distributed neural information (Pinotsis & Miller,  
2022; McFadden, 2020). Phase organization under propofol motivates studying  
this regime (Bastos et al., 2021), but does not measure coupling.

757 The formal field predicate requires a jointly continuous kernel, a normal-  
758 ized substrate, coupling equal to the kernel average, and no single site pro-  
759 viding more than half that strength. Toy constructions satisfy it. A cortical  
760 realization must specify the scale at which the kernel and its reduction ap-  
761 ply: calcium imaging reports different state-dependent modularity patterns  
762 at single-cell and mesoscale resolution (Kiyooka et al., 2026).

763 Cortical columns are the obvious candidate unit, and they answer one  
764 of the two demands the formalism makes. As the oscillator index they are  
765 a serious proposal: they are delineated by response properties and anatomy  
766 rather than by the agreement they would be used to establish (Mountcastle,  
767 1997), and a phase can be estimated from a column-sized territory at the  
768 electrode spacings used for cortical wave measurements (Townsend et al.,  
769 2015). As the content cover they are a poor one, for three separate reasons.  
770 A parcellation into non-overlapping columns is a disjoint cover, which satisfies  
771 the compatibility obligation without constraining anything (Section 4). The  
772 cover condition additionally requires the patches to exhaust the substrate,  
773 while much of association cortex has no accepted columnar parcellation. And  
774 the column’s standing as a canonical cortical unit is contested (Horton &  
775 Adams, 2005; Rakic, 2008). Offering columns for both roles at once therefore  
776 supplies the oscillator index and leaves the content cover to be identified on  
777 separate evidence, which is the bridge assumption E78 records.

778 The spatial reduction is also testable. A row-sum aggregation at zero  
779 diagonal tracks the uniform-coupling control over the tested spatial decay

780 range, while frequency heterogeneity displaces the operational threshold. The  
781 fit window biases both estimates relative to  $2D$ ; agreement with the control  
782 is therefore the supported conclusion. Supplemental Material reports the  
783 reduction, its numerical error and the distinct assumptions needed for the  
784 kernel-average identification in Eq. (5).

785 The field commitment requires physical evidence beyond a choice of math-  
786 ematical representation, and the available support comparisons do not supply  
787 it. A finite matrix spread over positive-mass cells preserves its total cou-  
788 pling and phase correlation exactly, while point-supported vanishing under  
789 an atomless measure and a disadvantage from excluding a better-correlated  
790 pair concern specified support constraints; Section 9.3 reads the same com-  
791 parisons against a digital candidate. Neither supplies energy prices.

792 A spike-time shift alone supplies no coupling rate. The dimensionless  
793 product of population size  $N$ , timing sensitivity  $s$ , field amplitude  $E$  and  
794 frequency  $f$  needs a conversion rate  $\gamma$ :

$$K_{\text{eff}} = \gamma N s E f. \quad (15)$$

795 This rate describes how induced timing shifts accumulate into drift. Nei-  
796 ther it nor the installation factor  $\kappa$  follows from a heat bound. Testing the  
797 physical threshold requires independent estimates of the interaction kernel,  
798 its aggregation and diffusion in the same cortical regime.

## 799 8.2 Extracellular geometry and recovery

800 The proposed control variable is extracellular geometry. Extracellular vol-  
801 ume fraction is approximately  $\alpha = 0.2$  at tortuosity  $\lambda \approx 1.6$  (Syková &  
802 Nicholson, 2008); a sleep study reports a rise from roughly 0.14 to 0.23 (Xie  
803 et al., 2013). Below 10 kHz, cortical tissue conducts approximately resistively  
804 at  $0.3\text{--}0.6\text{ S m}^{-1}$  (Barbour, 2017). At fixed transmembrane current density,  
805 an effective-medium description predicts that a larger extracellular fraction  
806 raises conductivity and reduces field amplitude.

807 The coupling-gated hypothesis consequently predicts that extracellular  
808 contraction on awakening raises  $E$  and, holding the other factors in Eq. (15)  
809 fixed, raises effective coupling. This argument supplies a direction; the  
810 geometry–coupling relation, its magnitude and its role in human recovery  
811 require measurement.

812 For a smooth crossing at nonzero speed, evaluating the stationary branch  
813 along the changing control gives the benchmark

$$r_{\text{ss}}(K(t)) \sim \sqrt{\frac{\dot{K}(t_0)}{D}} (t - t_0)^{1/2}, \quad K(t_0) = K_c, \quad \dot{K}(t_0) > 0. \quad (16)$$

814 This follows from Eq. (3) at fixed  $D$  and concerns the stationary branch  
815 evaluated at  $K(t)$ . Reading it as observed recovery requires an indepen-  
816 dently justified tracking approximation. The crossing controls in Section 7.2  
817 leave that approximation unsupported at onset, even though a separation of  
818 geometry and phase time scales can support it away from threshold. The

819 awakening hypothesis therefore needs a calibrated dynamical prediction for  
820 an operational onset above the finite-size fluctuation floor. Crossing  $K_c$  alone  
821 does not supply a recovery latency.

### 822 **8.3 A protocol for the awakening experiment**

823 The physical test has six requirements:

- 824 **1. Coupling and diffusion in the same units.** Estimate them from re-  
825 solved phase dynamics, separating heterogeneity and observation noise.  
826 Measure the drive or perturb the system to distinguish endogenous cou-  
827 pling.
- 828 **2. Geometry calibrated to coupling.** Validate the geometry observ-  
829 able and its relation to effective coupling, with uncertainty. The con-  
830 ductivity argument supplies only the expected direction.
- 831 **3. Calibrated phase measurements.** Estimate  $r$  by resultant length  
832 and  $a$  by unweighted regression of log density on centered cosine.  
833 Choose site count and binning from a dependence-aware calibration  
834 over the intended concentration range. The tested designs require  
835 thousands of sites at the observed concentrations; temporal pooling  
836 changes the spatial estimand.
- 837 **4. An independent recovery endpoint.** Measure behavioural recovery  
838 separately from phase organization.

839     **5. A predeclared dynamical onset criterion.** Set a coherence thresh-  
840     old above the finite-size floor and evaluate its latency under the mea-  
841     sured coupling trajectory, including ramp speed or fluctuations. Record  
842     a spatial observable beside the resultant, since a winding field reports  
843     a low resultant at coupling that sustains local order (Section 7.3).

844     **6. Matched model comparison.** Compare coupling increase, diffusion  
845     change, common drive and alternative recovery models under the same  
846     observation and filtering assumptions.

847     Failure of the concentration–coherence relation over an informative range  
848     constrains the density model. Failure of the calibrated geometry–coupling  
849     association constrains the physical gate. Better prediction of independent  
850     recovery by a competing mechanism counts against its proposed role. A  
851     favorable outcome still leaves the content of coordinated activity to be ex-  
852     plained.

## 853     **9     Discussion**

854     Two things have been established here, and they are of different kinds. The  
855     constraints of Section 2 hold outright: synchrony does not deliver compatible  
856     content, the resultant does not determine the state, a unique fixed point is  
857     not a reconstruction, a region that cannot resolve a family cannot encode it,  
858     and agreement about a change takes communication. Any account that reads  
859     unity off phase coherence has to meet them, this one included. The account

860 set out in Section 1 is conditional at every step by contrast: each connection  
861 in it is a hypothesis supplied rather than a result derived, and Table 1 names  
862 all eight. Its four non-standard ingredients, named in Section 1, are the  
863 installed-energy condition, the patch-local form of the content bound, the  
864 criterion separating a winding field from weak coupling, and the machine-  
865 checked composition with its joint witness. What follows takes the conditions  
866 in turn, as commitments the proposal incurs, as measurements that would  
867 bear on them, and as questions the framework leaves open.

## 868 **9.1 Unity and a minimal self as empirical commit-** 869 **ments**

870 We propose the state glued from compatible local descriptions as a corre-  
871 late of conscious unity, and its accurate internal reconstruction as a correlate  
872 of a minimal self. The motivations are that different contents belong to  
873 one experienced situation and that the situation is experienced from a per-  
874 spective. The conditional results identify relations a realization would need:  
875 sufficient installed resources for the specified phase model, agreement about  
876 shared content, and an encoding that supports reconstruction at the required  
877 fidelity. Discharging E12–E89 for cortex would make the composition a state-  
878 ment about cortex: that a cortical field above threshold relaxes to a cover  
879 whose compatible local descriptions glue, and that the glued state is the  
880 unique fixed point of a reconstruction map realized there. It would leave the

881 two identifications above exactly where they are, because neither is among  
882 the eight and neither follows from their conjunction; they are the commit-  
883 ment this subsection names, and Section 9.2 lists the three observations that  
884 would count against it.

885 Compatibility allows different descriptions to contribute different infor-  
886 mation. It also permits uncertainty and conflicting judgments: apparent cup  
887 position and judged cup position can be distinct quantities in a consistent  
888 description of their discrepancy. Which quantities are shared must be de-  
889 termined independently. Choosing them solely to secure agreement would  
890 deprive the proposal of empirical force.

891 The minimal-self proposal concerns the perceiver’s position and condition  
892 within the situation, such as the cup’s location relative to a body that can  
893 reach it. The encoding could be implicit and need carry no autobiograph-  
894 ical narrative. Its proposed necessity is philosophical: even an unreflective  
895 episode has a perspective. Gluing by itself requires no self-encoding, and re-  
896 constructability alone supplies no functional role. Altering the encoded hand  
897 or viewpoint while holding the external scene fixed should change estimates  
898 of the perceiver’s relation to the cup as the readout predicts. Whether that  
899 represented perspective is experienced remains a further question.

## 900 **9.2 What a test of the full proposal must establish**

901 Both uniqueness results concern specified objects. They choose neither the  
902 cover nor the encoding, and do not select between candidate subjects. In-

903 tegrated information theory treats exclusion explicitly (Tononi et al., 2016);  
904 here selection requires independent evidence about neural organization. A  
905 constant readout can contract and fix a predetermined state while discard-  
906 ing its input. The family criterion in Section 5.2 tests the sensitivity that  
907 this construction lacks, provided the relevant states and fidelity are declared  
908 independently.

909 A test must separately measure coupling relative to diffusion, content  
910 compatibility and self-reconstruction, alongside an independent endpoint for  
911 experience. Meeting all three without a unified episode would challenge suffi-  
912 ciency. A unified episode with incompatible descriptions would challenge the  
913 unity identification; one lacking the specified self-encoding would challenge  
914 the reflexive requirement. Awakening, dreaming and dissociative states offer  
915 possible settings, but absence of report cannot by itself define absence of  
916 experience.

917 The next physical requirement is a calibrated geometry–coupling relation  
918 and resolved phase dynamics. The installation bound contributes a nec-  
919 essary resource condition where its hardware representation applies; cortical  
920 mode prices and field-mediated coupling require measurement. For the repre-  
921 sentational connection, an informative experiment asks whether a controlled  
922 change in coordination alters independently decoded content agreement, with  
923 controls for sensory information and decoder quality. The constructed-data  
924 estimator establishes feasibility for specified contents. Neural local descrip-  
925 tions, shared overlap encoders and the self-encoding have yet to be identified.

926 The population-size dependence of the phase-to-content bound further limits  
927 its quantitative use, in the form Section 4.2 reads off the sheet.

928 The separation results explain why these measurements must be inde-  
929 pendent. A stationary distribution leaves several recovery mechanisms indis-  
930 tinguishable. Equal phase order can accompany incompatible descriptions  
931 or different task information. The tested plasticity rule preserves coherence  
932 while losing its designated template. Each finding concerns the specified  
933 model and observable; their role is to delimit the evidence needed for the  
934 proposed connections.

### 935 **9.3 What the conditions constrain in a digital candi-** 936 **date**

937 The results of Section 2 hold for any candidate, and for a large language  
938 model (LLM) running on graphics processing units (GPUs) four of them are  
939 checkable rather than merely stateable. Those four are where this framework  
940 bears on claims made for such systems. It does not exclude them, and no re-  
941 sult here could; what it does is fix what a claim of unity or self-representation  
942 commits its maker to measuring.

943 **Codes, not reports.** Reconstruction across a family is a counting con-  
944 straint before it is anything else. If two macrostates the candidate is said  
945 to represent satisfy  $E(s) = E(t)$  while  $d(s, t) > 2\varepsilon$ , Eq. (10) rules out re-  
946 constructing both within  $\varepsilon$ , whatever the readout. A system's reports are

947 readouts. Identical verbal reports across two conditions are therefore consis-  
948 tent with identical internal codes and settle nothing about the states, while  
949 the converse claim — that the system represents its own condition across a  
950 declared family — asserts that its internal code separates that family, which  
951 is a statement about activations under intervention and not about text. The  
952 measurement follows from the statement: declare the family and the fidelity  
953 first, then test whether the encoding region distinguishes its members.

954 **The region, not the readout.** Where the encoding is read as the state’s  
955 restriction to a region of the machine, the demand falls on the region before  
956 any readout exists. For a decoder-only transformer the causal mask makes  
957 the network an instance of the update model of Section 5: the activation at  
958 position  $t$  and layer  $\ell$  reads positions at most  $t$  at layer  $\ell - 1$  and nothing  
959 else, so a region’s causal past is fixed by the architecture rather than chosen  
960 by interpretation. The deadline bound of Section 5 therefore applies with  
961 its communication graph given rather than assumed. A claim that a model  
962 reports on its own processing consequently carries an obligation to say which  
963 positions and layers the report’s causal past contains, and a report that  
964 appears to track an intervention outside it indicates a confound or a different  
965 region, not a faster channel.

966 **Agreement about what.** Unity requires the candidate’s local descriptions  
967 to agree on independently identified shared quantities (Section 4). Common  
968 timing, a shared residual stream and fluent output supply none of that, for

969 the reason Section 2 gives: coordination bounds content disagreement only  
970 through encoders already related on overlaps, and which quantities two parts  
971 of a model describe in common is a decoding question carrying the failure  
972 modes of Section 7. The cover is a choice, and choosing it to secure agreement  
973 empties the claim.

974 **What is not excluded.** Digital representations can satisfy compatibility,  
975 so finite hardware is compatible with the gluing construction and none of  
976 the above is a no-go for machines. The coupling-gated route E56–E78 is a  
977 different matter: it requires a calibrated physical coupling, its noise regime  
978 and a connection from coordination to content agreement, and an attention  
979 matrix or a shared clock supplies none of these identifications. Comput-  
980 ing a field model likewise leaves its realization in the computing substrate  
981 to be established. Those physical obligations fall on the cortical proposal  
982 identically.

983 The installed-energy condition of Section 3.2 is restricted in the same  
984 way. It constrains a macroscopic candidate only through  $\kappa$ , which is hard-  
985 ware data supplied by mode profiles and prices, and Supplemental Material  
986 reads the condition against a cortical metabolic budget to return the conver-  
987 sion factor it would require rather than a verdict. The hardware comparisons  
988 there have similarly restricted scope: fixed wiring can impose a correlation  
989 disadvantage, and point-supported kernels vanish under an atomless contin-  
990 uum measure, but neither comparison makes the representation conditions

991 unattainable on GPUs. A universal exclusion would need a necessary condi-  
992 tion for consciousness and a proof that every relevant implementation violates  
993 it, and the present composition supplies neither.

994     Whether experience requires a living substrate is the explicit subject of  
995 biological naturalism (Seth, 2026a,b). On that view the properties ground-  
996 ing experience are those of self-sustaining organisms, and simulating them  
997 does not secure them. The conditions stated here name no biological mech-  
998 anism, but computation described as such does not satisfy them either: a  
999 candidate must exhibit a physically realized coupling in a measured noise  
1000 regime, agreement about independently identified shared quantities, and an  
1001 internal reconstruction meeting a declared fidelity. Whether a realization  
1002 of those conditions that is not alive would be accompanied by experience  
1003 is the point at issue, and these results do not settle it. A related position  
1004 locates the difference in the mode of computation rather than in life as such:  
1005 biological systems compute across scales that do not separate, and along-  
1006 side discrete operations carry continuous-valued dynamics belonging to their  
1007 physical medium (Milinkovic & Aru, 2026). The coupling condition stated  
1008 here is one measurable instance of that requirement, fixed to a physical field  
1009 in a declared noise regime, and a description of such a field does not satisfy  
1010 it.

1011     The neuroscientific case against *human-like* consciousness in text-only  
1012 LLMs concerns embodied sensory interaction, thalamocortical organization  
1013 and biological regulation linking perception and action to survival (Aru et al.,

2023). Their absence supports skepticism, conditional on their causal relevance to experience. Those mechanisms are not requirements of the abstract gluing and reconstruction definitions, and workspace-like representations reported in some LLMs (Gurnee et al., 2026) supply candidate organization to investigate. What this framework adds to that investigation is an order for the questions: declare the family of states and the fidelity, fix the encoding region and read its causal past off the architecture, and only then ask what the system reconstructs. A claim that survives those three steps is a claim about the machine; one that does not survive them was a claim about its output.

## 9.4 Explanatory scope

Integrated-information, global-workspace and predictive-processing accounts organize unity, access and modelling differently (Tononi et al., 2016; Baars, 1997; Dehaene et al., 2014; Friston, 2010). The present proposal contributes a candidate physical interaction and conditional relations between local content and self-encoding. Gluing and reconstruction permit multiple substrates; the electromagnetic commitment enters through physical calibration. The formalization checks implications under their hypotheses, and its witnesses establish realizability on declared models. Two things stand outside that result and are not supplied by it: the eight hypotheses are discharged for toy components rather than for cortex, and the identification of the glued state with unity and of its reconstruction with a minimal self is a further

1036 commitment that the eight do not contain.

1037     The explanatory question is whether the specified relations occur in neural  
1038 activity and matter for experience. In the cup example, this means establish-  
1039 ing how coordination helps maintain one represented situation and how the  
1040 perceiver’s place within it contributes to the episode. The proposal identi-  
1041 fies candidate correlates and tests of that identification. Why those relations  
1042 would be accompanied by experience remains unexplained.

## 1043 **Data and code availability**

1044 The Lean development, simulation scripts, tests and saved numerical sum-  
1045 maries are maintained in the GitHub repository `bobaseb/coupling_gated_`  
1046 `cortical_coherence`. Publication simulation values are generated from  
1047 saved summaries without rerunning production sweeps. The exploratory  
1048 EEG analysis uses OpenNeuro dataset ds005620 (Bajwa et al., 2025); pre-  
1049 processing and selection are described in Supplemental Material. The repos-  
1050 itory contains the reproducible analysis and does not redistribute the raw  
1051 recordings.

## 1052 **Funding and competing interests**

1053 This research received no external funding. The author declares no compet-  
1054 ing interests.

## 1055 Use of AI tools

1056 AI assistance to this work was substantial and is not separable by task. Sev-  
1057 eral coding assistants and chat interfaces were used in an interleaved way  
1058 throughout: Google’s Antigravity CLI (Gemini 3.1 Pro, Claude 4.6 Opus,  
1059 Claude 4.6 Sonnet), OpenAI’s Codex CLI (GPT-6 Astra; GPT-5.6 Terra,  
1060 Luna and Sol), Claude Code (Claude Opus 5), Gemini Pro through its web in-  
1061 terface, and Anthropic’s Fable and DeepSeek v4 Flash through OpenRouter,  
1062 accessed with the Hermes CLI. Between them they contributed to the Lean  
1063 development, the simulation and analysis code, and the drafting and revision  
1064 of both documents. Because the tools were used in combination and in alter-  
1065 nation, no section, proof, figure or result is attributable to any one of them,  
1066 and none is claimed to be. The author is responsible for all content, including  
1067 everything a tool produced. Machine checking establishes the stated Lean  
1068 results under their hypotheses; it does not validate biological interpretations  
1069 or replace scientific review of generated code and prose.

## 1070 References

- 1071 Pinotsis, D. A., & Miller, E. K. (2022). Beyond dimension reduction: Stable  
1072 electric fields emerge from and allow representational drift. *NeuroImage*,  
1073 253, 119058.
- 1074 Tononi, G., Boly, M., Massimini, M., & Koch, C. (2016). Integrated informa-

1075 tion theory: from consciousness to its physical substrate. *Nature Reviews*  
1076 *Neuroscience*, 17(7), 450-461.

1077 Baars, B. J. (1997). In the Theater of Consciousness: The Workspace of the  
1078 Mind. *Oxford University Press*.

1079 Dehaene, S., Charles, L., King, J. R., & Marti, S. (2014). Toward a compu-  
1080 tational theory of conscious processing. *Current Opinion in Neurobiology*,  
1081 25, 76-84.

1082 Friston, K. (2010). The free-energy principle: a unified brain theory?. *Nature*  
1083 *Reviews Neuroscience*, 11(2), 127-138.

1084 Milinkovic, B., & Aru, J. (2026). On biological and artificial consciousness:  
1085 A case for biological computationalism. *Neuroscience & Biobehavioral Re-*  
1086 *views*, 181, 106524. doi:10.1016/j.neubiorev.2025.106524.

1087 Aru, J., Larkum, M. E., & Shine, J. M. (2023). The feasibility of artificial  
1088 consciousness through the lens of neuroscience. *Trends in Neurosciences*,  
1089 46(12), 1008–1017. doi:10.1016/j.tins.2023.09.009.

1090 Seth, A. K. (2026). Conscious artificial intelligence and biological naturalism.  
1091 *Behavioral and Brain Sciences*, 49, e315. doi:10.1017/S0140525X25000032.

1092 Seth, A. K. (2026). The stuff matters: Consciousness, compu-  
1093 tation, and biology. *Behavioral and Brain Sciences*, 49, e366.  
1094 doi:10.1017/S0140525X26106906.

- 1095 Gurnee, W., et al. (2026). Verbalizable Representations Form a  
1096 Global Workspace in Language Models. *Transformer Circuits Thread*.  
1097 arXiv:2607.15495.
- 1098 Aguilera, M., Millidge, B., Tschantz, A., & Buckley, C. L. (2022). How par-  
1099 ticular is the physics of the free energy principle?. *Physics of Life Reviews*,  
1100 40, 24-50.
- 1101 Biehl, M., Pollock, F. A., & Kanai, R. (2021). A technical critique of some  
1102 parts of the free energy principle. *Entropy*, 23(3), 293.
- 1103 Bruineberg, J., Dolega, K., Dewhurst, J., & Baltieri, M. (2022). The Em-  
1104 peror’s New Markov Blankets. *Behavioral and Brain Sciences*, 45, e183.
- 1105 Pinotsis, D. A., & Miller, E. K. (2023). In vivo ephaptic coupling allows  
1106 memory network formation. *Cerebral Cortex*, 33(17), 9877-9895.
- 1107 Pinotsis, D. A., Fridman, G., & Miller, E. K. (2023). Cytoelectric coupling:  
1108 Electric fields sculpt neural activity and ”tune” the brain’s infrastructure.  
1109 *Progress in Neurobiology*, 226, 102465.
- 1110 McFadden, J. (2002). The conscious electromagnetic information (cemi) field  
1111 theory: the hard problem made easy?. *Journal of Consciousness Studies*,  
1112 9(8), 45-60.
- 1113 Pockett, S. (2012). The electromagnetic field theory of consciousness: a  
1114 testable hypothesis about the characteristics of conscious as opposed to  
1115 non-conscious fields. *Journal of Consciousness Studies*, 19(11-12), 191-223.

1116 McFadden, J. (2020). Integrating information in the brain's EM field: the  
1117 cemi field theory of consciousness. *Neuroscience of Consciousness*, 2020(1),  
1118 niaa016.

1119 Fröhlich, F., & McCormick, D. A. (2010). Endogenous electric fields may  
1120 guide neocortical network activity. *Neuron*, 67(1), 129-143.

1121 Anastassiou, C. A., & Koch, C. (2015). Ephaptic coupling to endogenous  
1122 electric field activity: why bother?. *Current Opinion in Neurobiology*, 31,  
1123 95-103.

1124 Pockett, S. (2002). Difficulties with the electromagnetic field theory of con-  
1125 sciousness. *Journal of Consciousness Studies*, 9(4), 51-56.

1126 Vöröslakos, M., et al. (2018). Direct effects of transcranial electric stimulation  
1127 on brain circuits in rats and humans. *Nature Communications*, 9, 483.

1128 Verkhratsky, A., & Nedergaard, M. (2018). Physiology of Astroglia. *Physio-*  
1129 *logical Reviews*, 98(1), 239-389.

1130 Mountcastle, V. B. (1997). The columnar organization of the neocortex.  
1131 *Brain*, 120(4), 701-722. doi:10.1093/brain/120.4.701.

1132 Horton, J. C., & Adams, D. L. (2005). The cortical column: a structure  
1133 without a function. *Philosophical Transactions of the Royal Society B*,  
1134 360(1456), 837-862. doi:10.1098/rstb.2005.1623.

1135 Rakic, P. (2008). Confusing cortical columns. *Proceedings of the National*  
1136 *Academy of Sciences*, 105(34), 12099-12100. doi:10.1073/pnas.0807271105.

1137 Bastos, A. M., Donoghue, J. A., Brincat, S. L., Mahnke, M., Yanar, J., Cor-  
1138 rea, J., ... & Miller, E. K. (2021). Neural effects of propofol-induced uncon-  
1139 sciousness and its reversal using thalamic stimulation. *eLife*, 10, e60824.

1140 Bajwa, I. J., Nilsen, A. S., Skukies, R., Aamodt, A., Ernst, G., Storm, J.  
1141 F., & Juel, B. E. (2025). A repeated awakening study exploring the capac-  
1142 ity of complexity measures to capture dreaming during propofol sedation.  
1143 *Scientific Reports*, 15, 32746.

1144 Xie, L., Kang, H., Xu, Q., Chen, M. J., Liao, Y., Thiyagarajan, M., ... &  
1145 Nedergaard, M. (2013). Sleep drives metabolite clearance from the adult  
1146 brain. *Science*, 342(6156), 373-377.

1147 Syková, E., & Nicholson, C. (2008). Diffusion in brain extracellular space.  
1148 *Physiological Reviews*, 88(4), 1277-1340.

1149 Barbour, B. (2017). Analysis of claims that the brain extracellular impedance  
1150 is high and non-resistive. *Biophysical Journal*, 113(7), 1636-1638.

1151 Norton, J. D. (2005). Eaters of the lotus: Landauer's principle and the return  
1152 of Maxwell's demon. *Studies in History and Philosophy of Science Part B:*  
1153 *Studies in History and Philosophy of Modern Physics*, 36(2), 375-411.

1154 Landauer, R. (1961). Irreversibility and heat generation in the computing  
1155 process. *IBM Journal of Research and Development*, 5(3), 183-191.

- 1156 Kibble, T. W. B. (1976). Topology of cosmic domains and strings. *Journal*  
1157 *of Physics A: Mathematical and General*, 9(8), 1387-1398.
- 1158 Sakaguchi, H. (1988). Cooperative phenomena in coupled oscillator systems  
1159 under external fields. *Progress of Theoretical Physics*, 79(1), 39-46.
- 1160 Still, S., Sivak, D. A., Bell, A. J., & Crooks, G. E. (2012). Thermodynamics  
1161 of prediction. *Physical Review Letters*, 109(12), 120604.
- 1162 Kawai, R., Parrondo, J. M. R., & Van den Broeck, C. (2007). Dissipation:  
1163 The phase-space perspective. *Physical Review Letters*, 98(8), 080602.
- 1164 Horowitz, J. M., & Esposito, M. (2014). Thermodynamics with continuous  
1165 information flow. *Physical Review X*, 4, 031015. doi:10.1103/PhysRevX.4.  
1166 031015.
- 1167 Ito, S., & Sagawa, T. (2013). Information thermodynamics on  
1168 causal networks. *Physical Review Letters*, 111, 180603. doi:10.1103/  
1169 PhysRevLett.111.180603.
- 1170 Corberi, F., dello Russo, S., Messuti, G., Scarpetta, S., & Smaldone, L.  
1171 (2026). Thermodynamic learning. *arXiv preprint*. arXiv:2609.04732.
- 1172 Banach, S. (1922). Sur les opérations dans les ensembles abstraits et leur ap-  
1173 plication aux équations intégrales. *Fundamenta Mathematicae*, 3(1), 133-  
1174 181.

- 1175 Shenker, O. R. (2000). Logic and entropy. *PhilSci-Archive preprint*. philsci-  
1176 archive.pitt.edu/115.
- 1177 Rosas, F. E., Mediano, P. A. M., Luppi, A. I., Boyd, A. B., Berta, M., Jensen,  
1178 H. J., Gastpar, M., & Carroll, S. M. (2026). Four faces of information in  
1179 natural systems. *Nature Reviews Physics*. doi:10.1038/s42254-026-00971-4.
- 1180 Sznitman, A.-S. (1991). Topics in propagation of chaos. In *École d'Été de*  
1181 *Probabilités de Saint-Flour XIX—1989*, Lecture Notes in Mathematics,  
1182 vol. 1464, pp. 165–251. Springer. doi:10.1007/BFb0085169.
- 1183 Berglund, N., & Gentz, B. (2002). Pathwise description of dynamic pitchfork  
1184 bifurcations with additive noise. *Probability Theory and Related Fields*,  
1185 122, 341–388. doi:10.1007/s004400100174.
- 1186 Abramsky, S., & Brandenburger, A. (2011). The sheaf-theoretic structure  
1187 of non-locality and contextuality. *New Journal of Physics*, 13, 113036.  
1188 doi:10.1088/1367-2630/13/11/113036.
- 1189 Townsend, R. G., Solomon, S. S., Chen, S. C., Pietersen, A. N. J., Martin,  
1190 P. R., Solomon, S. G., & Gong, P. (2015). Emergence of complex wave  
1191 patterns in primate cerebral cortex. *The Journal of Neuroscience*, 35(11),  
1192 4657-4662. doi:10.1523/JNEUROSCI.4509-14.2015.
- 1193 Xu, Y., Long, X., Feng, J., & Gong, P. (2023). Interacting spiral wave pat-  
1194 terns underlie complex brain dynamics and are related to cognitive pro-

- 1195      cessing. *Nature Human Behaviour*, 7(7), 1196-1215. doi:10.1038/s41562-  
1196      023-01626-5.
- 1197      Nicoletti, G., Di Terlizzi, I., & Busiello, D. M. (2026). Balancing information  
1198      and dissipation with partially observed fluctuating signals. *Physical Review*  
1199      *Letters*, 137(12), 127101. doi:10.1103/8gx2-rbns.
- 1200      Kiyooka, D., Oomoto, I., Kitazono, J., Saito, Y., Kobayashi, M., Matsub-  
1201      ara, C., Kobayashi, K., Murayama, M., & Oizumi, M. (2026). Single-  
1202      cell resolution functional networks during unconsciousness are segre-  
1203      gated into spatially intermixed modules. *Cell Reports*, 45(2), 116902.  
1204      doi:10.1016/j.celrep.2025.116902.
- 1205      Oomoto, I., Kiyooka, D., Oizumi, M., & Murayama, M. (2026). Multi-area  
1206      single-cell calcium imaging dataset of the mouse cortex across wakefulness,  
1207      sleep, and anesthesia. *Scientific Data*. doi:10.1038/s41597-026-08202-2.
- 1208      Attwell, D., & Laughlin, S. B. (2001). An energy budget for signaling in the  
1209      grey matter of the brain. *Journal of Cerebral Blood Flow & Metabolism*,  
1210      21(10), 1133-1145. doi:10.1097/00004647-200110000-00001.
- 1211      Harris, J. J., Jolivet, R., & Attwell, D. (2012). Synaptic energy use and  
1212      supply. *Neuron*, 75(5), 762-777. doi:10.1016/j.neuron.2012.08.019.
- 1213      Nicholls, D. G., & Ferguson, S. J. (2013). *Bioenergetics 4*. Academic Press.