EARLY TOPICS AND EXAMPLES IN OPTIMAL TRANSPORT

ASHER CHRISTIAN

Abstract. The goal of this text is to develop the foundations of optimal transport using only the tools of measure theory taught in Math 245A and some basic probability theory. I establish necessary results from measure theory, including Prokhorov's theorem and properties of weak convergence, to provide a self-contained proof of the existence of optimal transport plans.

1. Introduction

Optimal transport seeks to provide the most efficient way to move mass from one distribution to another while minimizing a sepcific cost. The theory has applications in numerous fields from machine learning to economics. The concept of optimal transport was first proposed by French mathematician Gaspard Monge in 1781 [5]. Later, during World War II, the Soviet mathematician Leonid Kantorovich greatly advanced the topic by proving that solving the problem of finding an optimal transport plan is equivalent under certain conditions to maximizing a dual variable [4]. I will mainly be following the works of Cédric Villani [9] in establishing the existence of solutions to the problem as well as providing examples.

2. Notation

$(\mathcal{X}, \mathcal{B})$	-	A measurable space $\mathcal{X}$ and the set of measurable sets $\mathcal{B} \subset 2^{\mathcal{X}}$.
$(\mathcal{X}, \mathcal{B}, \mu)$	-	A measure space with measure μ.
$(\mathcal{X}, \mu)$	-	$(\mathcal{X}, \mathcal{B}, \mu)$ where $\mathcal{B}$ are the Borel measurable sets under the condition that $\mathcal{X}$ has a topology.
$P(\mathcal{X})$	-	The set of all Borel probability measures on $\mathcal{X}$.
$X_*(\mathbb{P})$	-	The Push forward measure induced by $X : (\Omega, \mathcal{F}, \mathbb{P}) \to (\mathcal{X})$ measurable.
	-	$X_*(\mathbb{P})(A) = \mathbb{P}(X^{-1}(A))$
$\sigma(\mathcal{P})$	-	The sigma algebra generated by $\mathcal{P}$ whenever $\mathcal{P} \subset 2^{\Omega}$ is a collection of sets.
$\mathbb{R}^+$	-	$\mathbb{R} \cap [0, \infty)$ the nonnegative real numbers.
$\langle A, B \rangle_F$	-	$\sum_{i=1}^{n} \sum_{j=1}^{m} A_{i,j} B_{i,j}$ the Frobenius inner product of two $n \times m$ real matrices A, B.

3. Introducing the Problem

Before we state the problem, we must first introduce a basic notion of couplings at the level of measures.

Definition 3.1 (Couplings)**.** Let $(\mathcal{X}, \mu), (\mathcal{Y}, \nu)$ be two probability spaces. The set $\Pi(\mu, \nu)$ is defined to be the set of all Borel probability measures π on $\mathcal{X} \times \mathcal{Y}$ satisfying $\pi(A \times \mathcal{Y}) = \mu(A), \pi(\mathcal{X} \times B) = \nu(B)$ for all $A \subset \mathcal{X}, B \subset \mathcal{Y}$ measurable sets. That is π admits as marginals the measures μ and ν. These measures are referred to as **couplings**.

Remark 3.2. The set of couplings is non-empty since one can always construct the independent coupling $\mu \times \nu$ referred to as the product measure as shown in Proposition 1.7.11 of Tao [8]. The existence of this measure was proven under the condition that $(\mathcal{X}, \mu)$ and $(\mathcal{Y}, \nu)$ are σ-finite measure spaces which they trivially are since $\mu(\mathcal{X}) = \nu(\mathcal{Y}) = 1$.

Remark 3.3. The set of couplings is also closed under convex combinations. Formally if $\pi_1, \ldots, \pi_k$ are couplings of μ and ν and $\lambda_1, \ldots, \lambda_k \geq 0$ satisfy $\sum_{i=1}^k \lambda_i = 1$ then

$$\pi := \sum_{i=1}^k \lambda_i \pi_i$$

is a coupling. Indeed for any $A \in \mathcal{X}$

$$\pi(A \times \mathcal{Y}) = \sum_{i=1}^k \lambda_i \pi(A \times \mathcal{Y}) = \sum_{i=1}^k \lambda_i \mu(A) = \mu(A)$$

with a similar result for any $B \in \mathcal{Y}$. Once we have multiple couplings we can construct infinitely many more by creating convex combinations of our existing couplings.

Definition 3.4 (MKp)**.** Fix $(\mathcal{X}, \mu), (\mathcal{Y}, \nu)$ two probability spaces and let $c : \mathcal{X} \times \mathcal{Y} \to \mathbb{R} \cup \{+\infty\}$ be a cost function. We say a coupling $\pi_{ot} \in \Pi(\mu, \nu)$ is a solution to the **Monge-Kantorovich minimization problem** (MKp) if

$$\int_{\mathcal{X} \times \mathcal{Y}} c(x, y) d\pi_{ot}(x, y) = \inf_{\pi \in \Pi(\mu, \nu)} \int_{\mathcal{X} \times \mathcal{Y}} c(x, y) d\pi(x, y) \tag{1}$$

4. Examples of Couplings

Clearly many couplings are not optimal. To keep the first example simple, consider a village living on the real line $\mathcal{X} = \mathcal{Y} = \mathbb{R}$. The village has three olive trees at positions 3 and 10 and 15 which produce a third of a ton of olives each. The village also has two olive presses at positions 3 and 12 which each require half a ton of olives to efficiently produce olive oil. Mathematically we let $\mu = \frac{1}{3}(\delta_3 + \delta_{10} + \delta_{15})$ $\nu = \frac{1}{2}(\delta_3 + \delta_{12})$ where δ_x is the Dirac measure centered at point x. Then in the absence of a cost function we can construct the independent coupling $\pi = \mu \times \nu$ as in figure 1.



Figure 1. Independent coupling between two discrete measures.

Additionally, using the fact that every coupling of sums of Dirac measures is a sum of Dirac measures, we can say that every coupling is of the form:

$$\pi_{\vec{c}} = c_1\delta_{3,3} + c_2\delta_{10,3} + c_3\delta_{15,3} + c_4\delta_{3,12} + c_5\delta_{10,12} + c_6\delta_{15,12} \tag{2}$$

such that

$$c_1 + c_4 = \tfrac{1}{3} \qquad c_1 + c_2 + c_3 = \tfrac{1}{2}$$
$$c_2 + c_5 = \tfrac{1}{3} \qquad c_4 + c_5 + c_6 = \tfrac{1}{2}$$
$$c_3 + c_6 = \tfrac{1}{3}$$

Now suppose we introduce a natural cost function

$$c(x, y) = (x - y)^2$$

then we want to minimize

$$\int_{\mathbb{R}^2} (x - y)^2 d\pi(x, y)$$

this integral is equal to

$$c_2 7^2 + c_3 12^2 + c_4 9^2 + c_5 2^2 + c_6 3^2$$

Thus our total cost is linear with respect to the measure. this is a linear optimization problem with solution

$$(c_1, c_2, c_3, c_4, c_5, c_6) = (\frac{1}{3}, \frac{1}{6}, 0, 0, \frac{1}{6}, \frac{1}{3})$$



FIGURE 2. Optimal coupling between two discrete measures.

Rather than proving the optimality of this linear problem, in Section 6, we will establish the condition of c-cyclic monotonicity which generalizes the method to solve this discrete probelm.

5. Existence of MKp solution

The example given above was very simple, but the problem is stated very generally, and it will be important to consider continuous cases, even those where μ and ν are neither discrete nor continuous. To get closer to achieving this goal we must first prove that the goal is feasible. In other words we must prove that there is a coupling that satisfies (1).

Theorem 5.1 (Existence of Optimal Coupling)**.** *Let* $(\mathbb{R}, \mu), (\mathbb{R}, \nu)$ *be two probability spaces on* $\mathbb{R}$*. Let* $c : \mathbb{R}^2 \to \mathbb{R}^+ \cup \{+\infty\}$ *be an unsigned continuous cost function. Then there exists* $\pi \in \Pi(\mu, \nu)$ *satisfying* (1)

Remark 5.2. In reality this theorem is true even when $(\mathbb{R}, \mathbb{R})$ are replaced by $(\mathcal{X}, \mathcal{Y})$ any separable complete metrizable topological spaces known as Polish spaces and c is only upper semicontinuous and bounded below by two lower semicontinuous functions. However, in the interest of time we will focus on this simpler case.

To prove this theorem in full generality we must use the power of weak convergence covered in Tao's *An Epsilon of Room* [7], but luckily with our reduced assumptions we can prove it directly. We will still need some building blocks.

Definition 5.3 (Weak Convergence)**.** A sequence of probability measures $(\pi_k)_{k\in\mathbb{N}}$ on $\mathbb{R}^d$ converges weakly to a measure π on $\mathbb{R}^d$ if for every continuous bounded function $f : \mathbb{R}^d \to \mathbb{R}$

$$\int_{\mathbb{R}^d} f d\pi_k \to \int_{\mathbb{R}^d} f d\pi$$

as $k \to \infty$.

This definition opens up a way for us to guarantee an optimal coupling via a sequential compactness argument.

Lemma 5.4. *Let $c : \mathbb{R}^2 \to \mathbb{R}^+ \cup \{+\infty\}$. Let $(\pi_k)_{k\in\mathbb{N}}$ be a sequence of probability measures on $\mathbb{R}^2$ converging weakly to some π probability measure on $\mathbb{R}^2$. Then*

$$\int_{\mathbb{R}^2} c d\pi \leq \liminf_{k\to\infty} \int_{\mathbb{R}^2} c d\pi_k$$

Proof. Let

$$c_l(x, y) = \begin{cases} c(x, y) & c(x, y) \leq l \\ l & c(x, y) > l \end{cases}$$

then $(c_l)_{l\in\mathbb{N}}$ is a monotone nondecreasing sequence of continuous bounded functions and by the monotone convergence theorem

$$\int_{\mathbb{R}^2} c d\pi = \lim_{l\to\infty} \int_{\mathbb{R}^2} c_l d\pi$$

By the definition of weak convergence on our continuous bounded functions we have

$$\int_{\mathbb{R}^2} c_l d\pi = \lim_{k\to\infty} \int_{\mathbb{R}^2} c_l d\pi_k$$

Additionally for all l and monotonicity of $(c_l)_l$

$$\int_{\mathbb{R}^2} c_l d\pi_k \leq \int_{\mathbb{R}^2} c d\pi_k$$

putting this together we have

$$\int_{\mathbb{R}^2} c d\pi = \lim_{l\to\infty} \lim_{k\to\infty} \int_{\mathbb{R}^2} c_l d\pi_k \leq \liminf_{k\to\infty} \int_{\mathbb{R}^2} c d\pi_k$$

□

This result is important because if we can construct a sequence of couplings converging weakly to another coupling and the infimum of the MKp we can conclude that the limiting coupling is optimal.

Lemma 5.5. *Let μ, ν be two probability measures on $\mathbb{R}$ then for all $\epsilon > 0$ there exists $K \subset \mathbb{R}^2$ compact, such that for all $\pi \in \Pi(\mu, \nu)$, $\pi(\mathbb{R}^2 \setminus K) < \epsilon$.*

Proof. Fix $\epsilon > 0$. By continuity from below of measure we have

$$\lim_{R\to\infty} \mu([-R, R]) = \mu(\mathbb{R}) = 1$$

There exists R such that

$$\mu(\mathbb{R} \setminus [-R, R]) < \frac{\epsilon}{2} \qquad \nu(\mathbb{R} \setminus [-R, R]) < \frac{\epsilon}{2}$$

$K := [-R, R] \times [-R, R]$ is a compact set in $\mathbb{R}^2$. Note that the complement of this rectangle is contained in the union of the complements of its sides

$$\mathbb{R}^2 \setminus K \subset ((\mathbb{R} \setminus [-R, R]) \times \mathbb{R}) \cup (\mathbb{R} \times (\mathbb{R} \setminus [-R, R]))$$

Let $\pi \in \Pi(\mu, \nu)$ be arbitrary, using marginal properties and subadditivity of measures:

$$\begin{aligned}\pi(\mathbb{R}^2 \setminus K) &\leq \pi(\mathbb{R} \setminus [-R, R]) \times \mathbb{R}) + \pi(\mathbb{R} \times (\mathbb{R} \setminus [-R, R])) \\ &= \mu(\mathbb{R} \setminus [-R, R]) + \nu(\mathbb{R} \setminus [-R, R]) \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon\end{aligned}$$

Since ϵ was arbitrary we have proved this lemma. □

Remark 5.6. This proof is also valid when the sets are Polish spaces.

Now that we have established tightness of the couplings we can say more about the existence of a limit. This fundamental result shows that under this assumption, a space of measures is relatively compact, originally formulated by Yuri Prokhorov in 1956 [6].

Theorem 5.7 (Prokhorov's Theorem)**.** *Let $(\pi_k)_{k\in\mathbb{N}}$ be a sequence of probability measures on $\mathbb{R}$. If the sequence is tight (satisfies the condition in Lemma 5.5), then there exists a subsequence $(\pi_{k_j})_{j\in\mathbb{N}}$ that converges weakly to some probability measure π.*

To prove the existence of a weak limit, we utilize the correspondence between probability measures on $\mathbb{R}^d$ and their cumulative distribution functions (CDFs).

Lemma 5.8 (Helly's Selection Principle)**.** *Let $(F_k)_{k\in\mathbb{N}}$ be a sequence of uniformly bounded, non-decreasing functions on $\mathbb{R}$. Then there exists a subsequence (F_{k_j}) and a non-decreasing, right-continuous function F such that $F_{k_j}(x) \to F(x)$ pointwise at all continuity points of F.*

Proof. Let $S = \{q_1, q_2 \ldots, \}$ be an enumeration of $\mathbb{Q}$. Consider the sequence at q_1 $(F_k(q_1))_k$. Since the functions are uniformly bounded this is a bounded sequence of real numbers and so there exists a convergent subsequence. Call it $F_{1,k}$. Proceeding in this way inductively create a subsequence $F_{n,k}$ of the previous sequence $F_{n-1,k}$ such that it converges for q_n. Then the sequence $F_{k,k}$ converges for every $q \in \mathbb{Q}$. Let $G(q) = \lim_{k\to\infty} F_{k,k}(q)$ for all $q \in \mathbb{Q}$. Extend G to $F : \mathbb{R} \to \mathbb{R}$ by letting

$$F(x) = \inf\{G(q) : q \in \mathbb{Q}, q > x\}$$

First if $y > x$ then it is easy to see that $F(y) \geq F(x)$ since the infimum is being taken over a smaller set. Additionally F is right continuous. Let $(x_n)_n \subset \mathbb{R}^d$ be a sequence such that $x_1 \geq x_2 \geq \ldots$ and $\lim_{i\to\infty} x_i = x$. For any $\epsilon > 0$ there exists a rational $q > x$ such that $G(q) < F(x) + \epsilon$. For sufficiently large n, we have $x \leq x_n < q$ and by monotonicity of F we have

$$F(x) \leq F(x_n) \leq G(q) < F(x) + \epsilon$$

thus

$$\lim_{n\to\infty} F(x_n) = F(x)$$

proving right continuity. Finally we must show that $F_{k,k}$ converges to F at every point x where F is continuous. Let x be a continuity point of F. For any rational $q > x$ monotonicity of the F_k implies

$$F_{k,k}(x) \leq F_{k,k}(q)$$

so

$$\limsup_{k\to\infty} F_{k,k}(x) \leq \lim_{k\to\infty} F_{k,k}(q) = G(q)$$

taking the infimum over all rational $q > x$ we get

$$\limsup_{k\to\infty} F_{k,k}(x) \leq F(x)$$

similarly for all rational $r < x$ we have $F_{k,k}(r) \leq F_{k,k}(x)$ and so we have

$$G(r) \leq \liminf_{k\to\infty} F_{k,k}(x)$$

but since x is a continuity point $\sup\{G(r) : r \in \mathbb{Q}, r \leq x\} = F(x)$ by monotonicity of G and that $G = F$ on $\mathbb{Q}$ so we can conclude

$$F(x) \leq \liminf_{k\to\infty} F_{k,k}(x)$$

taking these two facts together we get

$$\lim_{k\to\infty} F_{k,k}(x) = F(x)$$

for all x that are continuity points of F. □

Remark 5.9 (Extension to $\mathbb{R}^d$). The proof of Lemma 5.8 relies on the countability of the dense subset $\mathbb{Q} \subset \mathbb{R}$. Since $\mathbb{Q}^d$ is also a countable dense subset of $\mathbb{R}^d$, the same diagonalization argument allows us to extract a subsequence of CDFs $F_{k_j}(x_1, \ldots, x_d)$ that converges at every continuity point. Consequently, Helly's Selection Principle holds for $\mathbb{R}^d$. A rigorous proof can be found in Theorem 11.1.1 of Dudley [2].

With this tool, we can prove Prokhorov's Theorem.

Proof of Theorem 5.7. Let F_k be the CDF corresponding to μ_k, that is $F_k(x) = \mu_k(\{y \in \mathbb{R}, y \leq x\})$. Since μ_k are probability measures $0 \leq F_k(x) \leq 1$ so the sequence is uniformly bounded. Note that F_k satisfies the conditions of Helly's Selection Principle 5.8.

By Helly's Selection Principle (Lemma 5.8), there exists a subsequence (which we still denote as F_k) and a non-decreasing right-continuous function F such that $F_k \to F$ at continuity points. We must show that F is the CDF of a valid probability measure. Recall the moving bump example provided in lecture, we do not want our sequence to lose mass in the limit. Fix $\epsilon > 0$. Since the sequence is tight, there exists a compact interval $K = [-M, M]$ such that for all k, $\mu_k(K^c) < \epsilon$. In terms of our CDFs, this says that for all k $F_k(M) - F_k(-M) \geq 1 - \epsilon$. In particular for all continuity points $|x| > M$ $F(x) - F(-x) \geq 1 - \epsilon$ Sending ϵ to 0 and picking larger M's we see that

$$\lim_{x\to\infty} F(x) - F(-x) = 1 \quad \lim_{x\to\infty} F(x) = 1 \quad \lim_{x\to-\infty} F(x) = 0$$

Thus F is a CDF. Additionally to this CDF there exists a unique Borel probability measure μ satisfying $\mu([a, b]) = F_+(b) - F_-(a)$ as outlined in Theorem 1.7.9 of Tao [8], the Lebesgue-Stieltjes measure. The fact that μ_k converges weakly to μ is unfortunately involved, but can be found in theorems 3.2.8 and 3.2.9 of Durett's Probability [3]. □

Remark 5.10. As with 5.8, this Theorem is true in the general setting of Polish spaces, see Theorem 5.1 Billingsley [1] for exact details.

Despite my efforts to build this theory using only the tools taught in [8] I will introduce one theorem that will be useful in the next lemma. The Pi-Lambda theorem, attributed to Eugene Dynkin allows statements that are true on a "basic set" to be expanded to the sigma algebra completion of the set.

Theorem 5.11 (Pi-Lambda Theorem). *A set $\mathcal{P} \subset 2^\Omega$ is a π-system if*

(1) *$\mathcal{P}$ is non-empty*
(2) *$A, B \in \mathcal{P} \implies A \cap B \in \mathcal{P}$*

A set $\mathcal{L} \subset 2^\Omega$ is a λ-system if

(1) *$\emptyset \in \mathcal{L}$*
(2) *$A \in \mathcal{L} \implies \Omega \setminus A \in \mathcal{L}$*
(3) *if $A_1, A_2, \ldots$ are a sequence of pairwise disjoint sets in $\mathcal{L}$ then $\bigcup_{i=1}^\infty A_i \in \mathcal{L}$*

If $\mathcal{P}$ is a π-system and $\mathcal{L}$ is a λ-system and $\mathcal{P} \subset \mathcal{L}$ then $\sigma(\mathcal{P}) \subset \mathcal{L}$

A proof can be found in Theorem A.1.4 of Durrett [3].

Lemma 5.12. *Let $(\pi_k)_k$ be a sequence of couplings in $\Pi(\mu,\nu)$ for some probability measures μ,ν on $\mathbb{R}$. Suppose further that π_k converges weakly to π, a measure on $\mathbb{R}^2$ then $\pi \in \Pi(\mu,\nu)$.*

Proof. We wish to show that the marginals of π are exactly μ and ν. Let $\phi \in C_b(\mathbb{R})$ be a continuous bounded function on $\mathbb{R}$. By the definition of the marginal

$$\int_{\mathbb{R}^2} \phi(x) d\pi(x,y) = \int_{\mathbb{R}} \phi(x) d\pi_1(x)$$

where π_1 is the first marginal of π but we also have

$$\int_{\mathbb{R}^2} \phi(x) d\pi(x,y) = \lim_{k\to\infty} \int_{\mathbb{R}^2} \phi(x) d\pi_k(x,y) = \int_{\mathbb{R}} \phi(x) d\mu(x)$$

by definition of weak convergence so we can conclude that for all continuous bounded functions ϕ

$$\int_{\mathbb{R}} \phi(x) d\pi_1(x) = \int_{\mathbb{R}} \phi(x) d\mu(x).$$

Let F be any closed set and define

$$g_l(x) = \max(0, 1 - l d(x,F))$$

where $d(x,F) = \inf\{d(x,y) : y \in F\}$.
Then $g_l(x) \geq 1_F(x)$ and is continuous, with $\lim_{l\to\infty} g_l(x) = 1_F(x)$ and so by the dominated convergence theorem since $|g_l| \leq 1$ which is absolutely integrable with respect to every probability measure

$$\pi_1(F) = \int_{\mathbb{R}} 1_F(x) d\pi_1(x) = \lim_{l\to\infty} \int_{\mathbb{R}} g_l(x) d\pi_1(x) = \lim_{l\to\infty} \int_{\mathbb{R}} g_l(x) d\mu(x) = \int_{\mathbb{R}} 1_F(x) d\mu(x) = \mu(F)$$

so π_1 and μ agree on all closed sets. Let $\mathcal{P}$ be the set of all closed sets. Let $\mathcal{S} \subset \mathcal{B}(\mathbb{R})$ be the collection of Borel sets on $\mathbb{R}$ such that $A \in \mathcal{S} \implies \mu(A) = \pi_1(A)$. Then $\mathcal{S}$ contains all closed sets on $\mathbb{R}$ and

$$\begin{aligned} \mathbb{R} \in S \qquad & \text{since } \mathbb{R} \text{ is closed} \\ A \in S \implies & \pi_1(A^c) = 1 - \pi_1(A) = 1 - \mu(A) = \mu(A^c) \\ A_1, A_2, \ldots, \in S, A_i \cap A_j = \emptyset, \forall i \neq j \implies & \pi_1\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} \pi_1(A_i) = \sum_{i=1}^{\infty} \mu(A_i) = \mu\left(\bigcup_{i=1}^{\infty} A_i\right) \end{aligned}$$

thus $\mathcal{S}$ is a λ-system and since the set of closed sets is closed under finite intersections it is a π-system and so by the $\pi - \lambda$ theorem 5.11, $\sigma(\mathcal{P}) \subset \mathcal{S}$ but $\sigma(\mathcal{P}) = \mathcal{B}(\mathbb{R})$ so we can conclude that $\pi_1(A) = \mu(A)$ for all Borel sets A on $\mathbb{R}$. Thus $\pi_1 = \mu$ and π admits μ as its first marginal.
The argument for the other direction is equivalent and we conclude that $\pi \in \Pi(\mu,\nu)$ whenever it is the weak limit of couplings in $\Pi(\mu,\nu)$. □

We have now shown that our space of couplings $\Pi(\mu,\nu)$ is sequentially compact with respect to weak convergence. We have also shown that when a sequence of couplings $(\pi_k)_k$ converges weakly to π our transport cost cannot increase in the limit. Putting these together we can finally argue the existence of an optimal transport plan:

Proof of Theorem 5.1. Fix μ,ν probability spaces on $\mathbb{R}$ and $c : \mathbb{R}^2 \to \mathbb{R}^+ \cup \{+\infty\}$ unsigned continuous cost function. Then by Lemma 5.5, $\Pi(\mu,\nu)$ is tight. Let $(\pi_k)_{k\in\mathbb{N}}$ be a sequence of couplings such that

$$\lim_{k\to\infty} \int_{\mathbb{R}^2} c(x,y) d\pi_k(x,y) = \inf_{\pi\in\Pi(\mu,\nu)} \int_{\mathbb{R}^2} c(x,y) d\pi(x,y)$$

then by Prokhorov's Theorem 5.7 there exists a subsequence $(\pi_{k_j})_{j\in\mathbb{N}}$ converging to some probability measure π weakly. By Lemma 5.12 we know $\pi \in \Pi(\mu,\nu)$ and by Lemma 5.4

$$\int_{\mathbb{R}^2} c(x,y)d\pi(x,y) \leq \liminf_{j\to\infty} \int_{\mathbb{R}^2} c(x,y)d\pi_{k_j}(x,y) = \inf_{\eta\in\Pi(\mu,\nu)} \int_{\mathbb{R}^2} c(x,y)d\eta(x,y)$$

and so π is an optimal transport plan. □

5.1. **Counterexample.** To illustrate why we need the above conditions to guarantee the existence of an optimal coupling, consider the following counterexample. Consider the space $\mathcal{X} = \mathcal{Y} = [0,1]$ equipped with the Lebesgue measure $\mu = \nu = \lambda|_{[0,1]}$. Define the cost function

$$c(x,y) := \begin{cases} 2 & x = y \\ |x-y| & x \neq y. \end{cases}$$

Then clearly c is continuous everywhere except at all points where $x = y$. Consider the sequence of transport maps $T_n(x) = (x + \frac{1}{n}) \pmod 1$ where $\pi_n = (\mathrm{Id}, T_n)_*(\mu)$ the push forward measure. Then the couplings π_n are supported on the lines $y = (x + \frac{1}{n}) \pmod 1$ and as $n \to \infty$ the π_n converge to $\pi_{id} = (\mathrm{Id}, \mathrm{Id})_*(\mu)$. Additionally

$$\lim_{n\to\infty} \int c(x,y)d\pi_n(x,y) = \lim_{n\to\infty} \int c(x,T_n(x))d\mu(x) = \lim_{n\to\infty} \frac{1}{n} = 0.$$

However

$$\int c(x,y)d\pi_{id}(x,y) = \int_0^1 c(x,x)dx = \int_0^1 2dx = 2.$$

We can see that the infimum cost must be 0 since our cost function is strictly positive and there exists a sequence of couplings approaching this value. The infimum is unable to be attained since any coupling π with

$$\int c(x,y)d\pi(x,y) = 0$$

would mean that $c(x,y) = 0$ π-almost everywhere, however there is no point (x,y) on $[0,1]\times[0,1]$ where the cost is zero, so this can never be the case. Thus the optimal coupling is not achievable. From this we can see why one of our condition of continuity (more exactly, lower semicontinuity) is necessary to guarantee an optimal coupling.

6. Characterizing Optimality: c-Cyclic Monotonicity

Now that we are guaranteed an optimal solution, let's see what we can say about the nature of optimal couplings.

Definition 6.1 (c-Cyclic Monotonicity [9])**.** Let $\mathcal{X}, \mathcal{Y}$ be sets and $c : \mathcal{X}\times\mathcal{Y} \to (-\infty,\infty]$ be a function. A subset $A \subset \mathcal{X}\times\mathcal{Y}$ is c-cyclically monotone if, for any $N \in \mathbb{N}$, and any collection $(x_1,y_1),\ldots,(x_N,y_N)$ of points in A

$$\sum_{i=1}^N c(x_i,y_i) \leq \sum_{i=1}^N c(x_i,y_{i+1})$$

with $y_{N+1} = y_1$. A coupling is said to be c-cyclically monotone if it is supported on a c-cyclically monotone set.

Intuitively, our optimal coupling should be c-cyclically monotone since if it were not the case we can "shuffle mass around" to reduce our total cost.

Proposition 6.2 (c-Cyclic Monotonicity if and only if Optimality)**.** *Let $\mu,\nu \in P(\mathbb{R})$ be discrete measures (i.e. $\mu = \sum_{i=1}^n p_i\delta_{x_i}, \nu = \sum_{i=1}^m q_i\delta_{y_i}$) and let $c : \mathbb{R}^2 \to \mathbb{R}\cup\{+\infty\}$ be a continuous cost function. A coupling $\pi \in \Pi(\mu,\nu)$ is optimal if and only if it is c-cyclically monotone.*

I will split this proof into its two parts.

Proof of $\Rightarrow$ direction. Let $\pi \in \Pi(\mu, \nu)$ be optimal. Assume for contradiction that it is not supported on a c-cyclically monotone set, then there exists a set of points $A = \{(x_{j_1}, y_{j_1}), \dots, (x_{j_k}, y_{j_k})\}$ such that $\pi(x_{j_l}, y_{j_l}) > \epsilon > 0$ for each l, and

$$\sum_{l=1}^{k} c(x_{j_l}, y_{j_{l+1}}) < \sum_{l=1}^{k} c(x_{j_l}, y_{j_l})$$

Then construct $\tilde{\pi}$ by adjusting π along this cycle.

(1) First we remove the extra mass from each pair.

$$\tilde{\pi}(x_{j_l}, y_{j_l}) = \pi(x_{j_l}, y_{j_l}) - \epsilon$$

(2) We then add that mass to the new pair

$$\tilde{\pi}(x_{j_l}, y_{j_{l+1}}) = \pi(x_{j_l}, y_{j_{l+1}}) + \epsilon$$

(3) Keep $\tilde{\pi} = \pi$ for all other pairs.

We must check that $\tilde{\pi} \in \Pi(\mu, \nu)$. First all entries remain nonnegative by our choice of ϵ. Additionally $\tilde{\pi}$ is a probability measure since for each point that is weighted ϵ less, there is another point weighted ϵ ensuring the total mass is constant. Fix x_t on the support of μ then

$$\tilde{\pi}(\{x_t\} \times \mathbb{R}) = \sum_{i=1}^{m} \tilde{\pi}(x_t, y_i) = \sum_{i=1}^{m} \pi(x_t, y_i) = \pi(\{x_t\} \times \mathbb{R}) = \mu(\{x_t\})$$

since in the sums, every term where an epsilon is subtracted, an epsilon is added to a different term. The same result holds for the second projections so we can conclude that $\tilde{\pi} \in \Pi(\mu, \nu)$. Finally we compute

$$\int_{\mathbb{R}^2} c(x, y) d\tilde{\pi}(x, y) - \int_{\mathbb{R}^2} c(x, y) d\pi(x, y) = \epsilon \sum_{l=1}^{k} c(x_{j_l}, y_{j_{l+1}}) - \epsilon \sum_{l=1}^{k} c(x_{j_l}, y_{j_l}) < 0$$

by our assumption, thus the cost of coupling $\tilde{\pi}$ is less than that of π, a contradiction. Therefore every optimal coupling is c-cyclically monotone. □

Proof of $\Leftarrow$. Let $\pi \in \Pi(\mu, \nu)$ be c-cyclically monotone. I wish to show that for any other $\tilde{\pi} \in \Pi(\mu, \nu)$ π admits a lower cost. First let's write our couplings in matrix form

$$M(\pi) = \begin{pmatrix} s_{1,1} & \dots & s_{1,m} \\ \vdots & \ddots & \vdots \\ s_{n,1} & \dots & s_{n,m} \end{pmatrix} \qquad M(\tilde{\pi}) = \begin{pmatrix} \tilde{s}_{1,1} & \dots & \tilde{s}_{1,m} \\ \vdots & \ddots & \vdots \\ \tilde{s}_{n,1} & \dots & \tilde{s}_{n,m} \end{pmatrix}$$

where $\pi(x_i, y_j) = s_{i,j}, \tilde{\pi}(x_i, y_j) = \tilde{s}_{i,j}$ and $\sum_{j=1}^{m} s_{i,j} = p_i = \sum_{j=1}^{m} \tilde{s}_{i,j}$ and $\sum_{i=1}^{n} s_{i,j} = q_j = \sum_{i=1}^{n} \tilde{s}_{i,j}$. Then consider the matrix $M(\tilde{\pi}) - M(\pi)$, an $n \times m$ matrix whose rows and columns sum to 0. I claim that $M(\tilde{\pi}) - M(\pi) = \sum_{l=1}^{L} \alpha_l D_l$ where each D_l is a matrix supported on an even number indices $(i_1, j_1), (i_1, j_2), (i_2, j_2), \dots, (i_r, j_1)$, such that the first term is negative, then positive, and so on alternating. To show that this is possible consider a bipartite directed graph with left nodes $(x_1, x_2, \dots, x_n)$ and right nodes $(y_1, y_2, \dots, y_m)$. Let the edge between any two nodes x_i, y_j have weight $\tilde{s}_{i,j} - s_{i,j}$. Then every node has zero net flow. If every edge has zero weight we are done, if not choose an entry (i_1, j_1) such that $\tilde{s}_{i_1,j_1} - s_{i_1,j_1} < 0$ (a point x_{i_1}, y_{j_1} that π weights more than $\tilde{\pi}$). Since the rows sum to zero there exists an index j_2 such that $s_{i_1,j_2} - \tilde{s}_{i_1,j_2} > 0$, continuing inductively we can construct an alternating sequence $(i_1, j_1), (i_1, j_2), (i_2, j_2), \dots$ since there are only finitely many indices we must eventually visit one index twice producing a closed alternating cycle. In this cycle one term $|\tilde{s}_{i,j} - s_{i,j}|$ must be the smallest but nonzero. Let ϵ be this value and construct

the matrix $D^1_{i_1,j_1} = -\epsilon, D^1_{i_1,j_2} = \epsilon, \dots$ and zero everywhere else. Then $\sum_{i=1}^n D^1_{i,j} = 0$ since every index j comes in pairs of opposite signed points which cancel out the sum. The same holds for sums over columns and we see that the matrix D^1 obeys the same property as $M(\tilde{\pi}) - M(\pi)$. Continuing inductively in this way we generate a sequence of $D^1, D^2, \dots$ of finitely many cycle matrices who sum to $M(\tilde{\pi}) - M(\pi)$. To see that this process ends in finite time note that in each stage at least one index turns to zero in the remaining matrix. If we let the cost matrix $C \in \mathbb{R}^{n\times m}$ be such that $C_{i,j} = c(x_i, y_j)$ we can see that

$$\int c(x,y)d\tilde{\pi}(x,y) - \int c(x,y)d\pi(x,y) = \langle M(\tilde{\pi}) - M(\pi), C\rangle_F = \sum_{l=1}^{L} \langle D_l, C\rangle_F$$

If I can show that $\langle D_l, C\rangle_F \geq 0$ for each l I am done. But

$$\langle D_1, C\rangle_F = \epsilon_l \left(\sum (\text{cost of positive entries of } D^1) - \sum (\text{cost of negative entries in } D^1\ ) \right)$$

However the negative points are all areas of the support of π of the form $(i_1, j_1), (i_2, j_2), \dots$ and each positive term is $(i_1, j_{1+1}), (i_2, j_{2+1}), \dots$ and we have

$$\sum_{p=1}^{N} c(x_{i_p}, y_{i_p}) \leq \sum_{p=1}^{N} c(x_i, y_{i_{p+1}})$$

so we conclude

$$\langle D_1, C\rangle_F \geq 0$$

which further implies that $\tilde{\pi}$ is not more optimal than π and since $\tilde{\pi}$ was arbitrary we have that π is the optimal solution. □

Remark 6.3. A natural expansion to this paper would be to expand further and delve into other ways of finding optimal solutions including solving the dual problem as proposed by *Kantorovich* [4]. We now have a strong understanding of why optimal couplings exist and their behavior under certain assumptions. I am encouraged to learn more and keep exploring this fascinating subject in the future.

Acknowledgments

I would like to acknowledge Elias Manuelides for helping me with this project. He inspired me to choose the topic of optimal transport and gave helpful feedback on an initial draft of this paper.

References

[1] Patrick Billingsley. *Convergence of Probability Measures.* John Wiley & Sons, New York, 2nd edition, 1999. ISBN 978-0471197454.

[2] Richard M. Dudley. *Real Analysis and Probability.* Cambridge University Press, Cambridge, 2nd edition, 2002. ISBN 978-0521007542.

[3] Rick Durrett. *Probability: Theory and Examples.* Cambridge University Press, Cambridge, UK, 5 edition, 2019. ISBN 9781108473682. URL `https://sites.math.duke.edu/~rtd/PTE/PTE5_011119.pdf`. Version 5 (PDF available from Duke University Mathematics Dept.).

[4] L. V. Kantorovich. On the translocation of masses. *Management Science*, 5(1):1–4, 1958. doi: 10.1287/mnsc.5.1.1. English translation of the 1942 Russian paper in Doklady Akademii Nauk SSSR.

[5] Gaspard Monge. Mémoire sur la théorie des déblais et des remblais, 1781. Histoire de l'Académie Royale des Sciences de Paris, avec les Mémoires de Mathématique et de Physique pour la même année pages 666–704.

[6] Yuri V Prokhorov. Convergence of random processes and limit theorems in probability theory. *Theory of Probability & Its Applications*, 1(2):157–214, 1956. doi: 10.1137/1101016.

[7] Terence Tao. An epsilon of room: Pages from year three of a mathematical blog. Online PDF, 2010. URL `https://terrytao.wordpress.com/wp-content/uploads/2010/02/epsilon.pdf`. Accessed: 2025-12-06.

[8] Terence Tao. *An Introduction to Measure Theory*, volume 126 of *Graduate Studies in Mathematics*. American Mathematical Society, 2011. ISBN 978-1-4704-6640-4.

[9] Cédric Villani. *Optimal Transport: Old and New*, volume 338 of *Grundlehren der Mathematischen Wissenschaften*. Springer, Berlin, 2008. doi: 10.1007/978-3-540-71050-9.