

Light element partitioning between silicates, oxides and metallic melts

Insights into the formation and composition of Earth's core

R. Myhill*, D. Rubie, D. J. Frost

* Corresponding author: bob.myhill@uni-bayreuth.de

MOTIVATION

- ▶ The Earth's core is an iron-nickel alloy which contains several weight percent of one or more unknown light elements. Likely candidates include Si, O, S, C (and perhaps H and Mg).
- ▶ The formation of Earth's core involved iron segregation through a silicate magma ocean and underlying solids after impacts between planetesimals and planetary embryos.
- ▶ Light elements were partitioned between the metallic alloy and the surrounding silicates during segregation.
- ▶ These early processes had a dominant effect on the oxygen and carbon budget of the shallow Earth.




Figure 1: Schematic diagram of the proto-Earth after the impact of a large differentiated planetesimal.

The aims of this study:

- ▶ Create a self-consistent thermodynamic model of iron-rich solids and liquids within the software *burnman* [1].
- ▶ Predict the chemical evolution of the Earth and other planetary bodies during planetary growth, core-mantle differentiation and magma ocean crystallisation.
- ▶ Provide estimates of seismic velocities and densities in planetary cores.

SOLID ENDMEMBERS

- ▶ Popular thermodynamic models for minerals use a Debye-type lattice model [2] or polynomial parameterisation [3] for thermal properties.
- ▶ We adopt the Debye model approach, with additional terms to account for magnetic, order-disorder, electronic and anharmonic contributions to the thermal pressure.





Figure 2: The phase diagram of iron. The experimental data are from [4].

Figure 3: The volume of HCP iron along the Hugoniot. Red dots are experimental data [5].

MELT ENDMEMBERS

- ▶ There is significant disagreement on the properties of melts reported in the literature.
- ▶ We use the constraints provided by the volume, heat capacity and thermal expansivity at 1 bar, and use the piston-cylinder/multi-anvil-constrained melting curve.
- ▶ To extrapolate to high pressures, we attempt to satisfy the following equations [6]: $\Delta S_{\text{melt}} = R \ln 2 / (1 - \alpha K_T dT/dP)$; $\Delta V_{\text{melt}} = \Delta S_{\text{melt}} dT/dP$.





Figure 4: The melting curve of FeO (0.94 ≤ x < 1). Model curve for FeO in red. Constraints are from PC [7], MA [8, 9] and LH-DAC experiments [10, 11, 12].

Figure 5: Liquid properties along the FeO melting curve. The dots correspond to the predicted properties based on a model for simple melting [6].

SOLUTION MODELS

- ▶ We adapt the subregular solution model to incorporate excess bulk moduli and heat capacities, enabling accurate modelling over the P-T range of the mantle.
- ▶ Metallic melt solutions are fit to immiscibility gaps and liquidus curves.





Figure 6: Interaction parameters of Fe-FeO and FeO-Fe. Also shown are extrapolations of published low pressure [13] and high pressure [14] parameterisations.

Figure 7: The Fe-FeO solvus as a function of pressure and temperature. Data points are from the published literature [13, 15].

EQUILIBRIUM CALCULATIONS





Figure 8: Model Mg_{0.8}Fe_{0.2}O-melt equilibria, with published experimental data [13, 16] for comparison.

Figure 9: Mg_{0.9}Fe_{0.1}SiO₃ bridgmanite-melt equilibria

ACKNOWLEDGEMENTS

The authors would like to thank Rebecca Fischer for data and useful discussions. This project is funded by the European Research Council (ERC) Advanced Grant "Accretion and Early Differentiation of the Earth and Terrestrial Planets" (ACCRETE).

REFERENCES

[1] Cottar, S., Heister, T., Rose, I. & Unterborn, C. BurnMan: A lower mantle mineral physics toolkit. *Geochimica, Geophysica, Geosystems* 15, 1164–1179 (2014).

[2] Stixrude, L. & Lithgow-Bertelloni, C. Thermodynamics of mantle minerals - II. Phase equilibria. *Geophysical Journal International* 184, 1180–1213 (2011).

[3] Holland, T. & Powell, R. An improved and extended internally consistent thermodynamic dataset for phases of petrological interest, involving a new equation of state for solids. *Journal of Metamorphic Geology* 29, 333–383 (2011).

[4] Anzellini, S., Dewaele, A., Mezouar, M., Loubeyre, P. & Morard, G. Melting of iron at Earth's inner core boundary based on fast X-ray diffraction. *Science* 340, 464–466 (2013).

[5] Brown, J. M., Fritz, J. N. & Hixson, R. S. Hugoniot data for iron. *Journal of Applied Physics* 88, 5496–5498 (2000).

[6] Tallon, J. L. The entropy change on melting of simple substances. *Physics Letters A* 76, 139–142 (1980).

[7] Lindsley, D. H. Pressure-temperature relations in the system FeO-SiO₂. *Carnegie Institute Washington Yearbook* 65, 226–230 (1966).

[8] Ringwood, A. E. & Hibberson, W. The system Fe-FeO revisited. *Physics and Chemistry of Minerals* 17, 313–319 (1990).

[9] Zhang, L. & Fei, Y. Melting behavior of (Mg,Fe)O solid solutions at high pressure. *Geophysical Research Letters* 35, 13302 (2008).

[10] Knittle, E. & Jeanloz, R. The high-pressure phase diagram of Fe(0.94)O - A possible constituent of the Earth's core. *Journal of Geophysical Research (Solid Earth)* 96, 16169 (1991).

[11] Seagle, C. T., Heinz, D. L., Campbell, A. J., Prakapenka, V. B. & Wanless, S. T. Melting and thermal expansion in the Fe-FeO system at high pressure. *Earth and Planetary Science Letters* 265, 655–665 (2008).

[12] Fischer, R. A. & Campbell, A. J. High-pressure melting of wüstite. *American Mineralogist* 95, 1473–1477 (2010).

[13] Frost, D. J. et al. Partitioning of oxygen between the Earth's mantle and core. *Journal of Geophysical Research (Solid Earth)* 115, 2202 (2010).

[14] Komabayashi, T. Thermodynamics of melting relations in the system Fe-FeO at high pressure: Implications for oxygen in the Earth's core. *Journal of Geophysical Research (Solid Earth)* 119, 4164–4177 (2014).

[15] Tsuno, K., Ohtani, E. & Terasaki, H. Immiscible two-liquid regions in the Fe-O-S system at high pressure: Implications for planetary cores. *Physics of the Earth and Planetary Interiors* 160, 75–85 (2007).

[16] Ozawa, H. et al. Chemical equilibrium between ferropericlase and molten iron to 134 GPa and implications for iron content at the bottom of the mantle. *Geophysical Research Letters* 35, 5308 (2008).