

Quenchable water-rich aluminous post-stishovite, the SiO₂ - phase H - δ AlOOH solid solution, and seismic scatterers in the lower mantle

R. Myhill, D. J. Frost, T. Boffa-Ballaran, H. Bureau, C. Raepsaet

Bayerisches Geoinstitut, Universität Bayreuth, Universitätsstrasse 30, 95447 Bayreuth, Germany

Abstract

Keywords: high pressure, post-stishovite, water, slab, scatterers

1. Introduction

The stishovite structure (tetragonal, P4₂/mm, no. 136) undergoes a weak first-order transition (Andrault et al., 1998; Hemley et al., 2000) with second-order characteristics of Landau/ferroelastic type (Tsuchida and Yagi, 1989; Carpenter et al., 2000) to the CaCl₂-type structure (orthorhombic, Pnnm, no. 58). This transition occurs at pressures of \sim 50 GPa at room temperature (Kingma et al., 1995; Andrault et al., 1998), increasing with temperature to \sim 70 GPa at 2200 K (Hirose et al., 2005; Nomura et al., 2010). A somewhat higher dT/dP was observed by Ono et al. (2002). The nature of this transition means that the shear modulus $C_{11} - C_{12}$ decreases with increasing pressure and vanishes at the transition pressure, as observed in spectroscopic and high pressure diffraction studies (Kingma et al., 1995; Shieh et al., 2002). The stishovite-post-stishovite transition is therefore of interest to seismologists, as it should produce zones of low shear wave velocity which act as scatterers in the deep mantle.


Deviatoric stresses in the diamond anvil cell can cause large variations in the

*Corresponding author: R. Myhill


Email address: myhill.bob@gmail.com (R. Myhill, D. J. Frost, T. Boffa-Ballaran, H. Bureau, C. Raepsaet)

pressure of the stishovite to post-stishovite transition. Nonhydrostaticity greatly decreases the transition pressure from 60 GPa (Andrault et al., 2003; Hemley et al., 2000) to 40 GPa (Kingma et al., 1996; Singh et al., 2012). Molecular and lattice dynamic studies suggest that only 1.5–2.5 GPa differential stress is
20 required to cause this decrease in pressure (Dubrovinsky and Belonoshko, 1996).

Aluminium has a profound effect on the stability of the post-stishovite phase. Al_2O_3 can be incorporated into stishovite by the substitution of Si by Al with the formation of oxygen vacancies (Smyth et al., 1995; Hirose et al., 2005; Bromiley et al., 2006):




25 An alternative mechanism, where charge balance is accomplished by Al occupying the large interstitial sites is probably minor Smyth et al. (1995):



It has been argued that the addition of 4 wt% Al_2O_3 in the absence of any other components causes the transition to shift from 50 to 23 GPa at room temperature by inducing a ‘chemical pressure’ (Bolfan-Casanova et al., 2009). In
30 stishovite, there is one octahedral cation position (Wyckoff notation 2a) located at the origin and the body center of the tetragonal cell. One oxygen at position 4f creates a moderately distorted octahedron with point symmetry mmm. The addition of Al *decreases* distortion of this octahedron (Smyth et al., 1995). The addition of Al should therefore have little effect on the stishovite-post-stishovite
35 transition (Panero, 2006). The results of Bolfan-Casanova et al. (2009) are therefore unexpected; it seems possible that another component (such as H_2O) or deviatoric stress acted to stabilise the CaCl_2 structure.

It has also been suggested that the coupled substitution of Al and H may have a significant effect on the transition pressure. Lakshtanov et al. (2007) has shown
40 that the addition of 6 wt% Al_2O_3 and 0.24 wt% H_2O reduces the transition pressure to 24 GPa at room temperature. The substitution mechanism in this case is probably



Given that high pressure δ -AlOOH is isostructural with the post-stishovite phase above ca. 19 GPa (Sano-Furukawa et al., 2008; Kuribayashi et al., 2014),
 45 after transforming from the P2₁nm structure (no. 31) (Suzuki et al., 2000; Komatsu et al., 2006; Vanpeteghem et al., 2007), it seems sensible to suggest that addition of an AlOOH component lowers the transition pressure. Ab initio calculations support this suggestion (Umemoto et al., 2015). The equilibrium transition for 6.25 mol% AlOOH is at ca. 15 GPa at room temperature, im-
 50 plying an even more marked reduction than observed in the experimental data (Lakshtanov et al., 2007). This could be due to inaccuracies in the ab-initio data, or metastable preservation of the tetragonal phase in the experiments. Umemoto et al. (2015) also suggest that the larger number of hydrogen sites stabilises the tetragonal structure at high temperatures, changing the slope of
 55 the transition. The tetragonal \rightarrow orthorhombic transition in their simulations is associated with splitting of the hydrogen sites on the equatorial oxygens into two groups, one occupied and the other unoccupied. Additionally, the redistribution of hydrogens among equatorial oxygens in the tetragonal \rightarrow orthorhombic transition implies that it is now first order and no longer ferroelastic. Anelasticity
 60 by hydrogen hopping seems a plausible alternative to reduce seismic body wave speeds.

Despite this, H₂O contents in tetragonal stishovite measured using the FTIR calibration of Pawley et al. (1993) are less than 20% of that expected from an AlOOH component (Panero et al., 2003; Bromiley et al., 2006; Litasov et al.,
 65 2007) and in MORB compositions (Chung and Kagi, 2002).

This study was designed to investigate water solubility in Al-rich stishovite by synthesising crystals at relatively high temperature (Ono, 1999). In doing so, we created and quenched an Al-H rich post-stishovite phase.

1.1. Hysteresis vs. stabilisation

70 (Umemoto et al., 2015)

1.2. Mechanism of shear wave velocity reduction

Ferroelasticity (Carpenter et al., 2000) Snoek relaxation (anelasticity via H mobility) (Snoek, 1941; Nowick and Berry, 1972; McKnight et al., 2007).

1.3. Mechanisms of Al-, H incorporation

75 We found that hydrogen was most stable when bonded to the apical oxygen of the Al-octahedron, with the hydroxyl bond along $\langle 110 \rangle$ and co-planar with Al (Panero and Stixrude, 2004).

The relatively short O-O distance and correspondingly long O-H distance in the stishovite-AlOOH solid solution would lead one to expect a low OH stretch-
80 ing frequency compared to other nominally anhydrous minerals, consistent with the value of 3111 cm^{-1} observed by Pawley et al. (1993).

Symmetric bonding (Panero and Stixrude, 2004)

1.4. Water concentrations in stishovite

Pawley et al. (1993) calibration Panero et al. (2003) Bromiley et al. (2006),
85 Litasov et al. (2007), MORB Chung and Kagi (2002),

1.5. Elastic properties

Lakshtanov et al. (2005) Ono et al. (2002) Sano-Furukawa et al. (2009)

1.6. Phase relations in the lower mantle

Irifune and Ringwood (1993) Wood (2000) Hirose (2002) Walter et al. (2015)
90 Litasov and Ohtani (2005)

1.7. Datasets and modelling

Holland and Powell (2011) Stixrude and Lithgow-Bertelloni (2011) Cottaar et al. (2014)

1.7.1. Elastic properties AlOOH

95 Suzuki (2009) (Sano-Furukawa et al., 2008) Vanpeteghem et al. (2002) Tsuchiya and Tsuchiya (2009) Li et al. (2006)

2. The solid solution (MgSi , Fe^{2+}Si , Fe_2^{3+} , Al_2) O_4H_2

2.1. Endmembers

2.1.1. SiO_2

100 2.1.2. $\delta\text{-AlOOH}$

2.1.3. $\delta\text{-FeOOH}$

Pnma (no.62 α) to $\text{P2}_1\text{nm}$ (no. 31, ε) (Gleason et al., 2008) to Pnnm (no. 58, high spin) to Pnnm (no. 58, low spin) (Gleason et al., 2013)

2.1.4. “Phase H”

105 The addition of an MgSiO_4H_2 component also stabilises the Pnnm structure in stishovite (Komatsu et al., 2011). A previous candidate space group (Pnn2; no. 34 Kudoh et al., 2004) is now thought to be unlikely.

Stability (Ohtani et al., 2014) Ab initio $\text{P2}/\text{m}$ (no. 10) (Tsuchiya, 2013) Pnnm (no. 58) (Bindi et al., 2014) $\text{P2}_1\text{nm}$ (no. 31) (Nishi et al., 2014)

110 Kudoh et al. (2004) suggested a hypothetical high pressure form of $\text{Mg}(\text{OH})_2$ also in the space group Pnnm from Mg incorporation. Importance of the AlOOH - phase H solid solution Ohira et al. (2014)

3. Other phases

3.1. Water in seifertite

115 A few studies have suggested that the CaCl_2 type may undergo a phase transition to a mineral named seifertite with the scrutinyite ($\alpha\text{-PbO}_2$) structure (orthorhombic, Pbcn, no. 60 or Pb2n) within mantle pressures.

AlOOH to $\text{Pa}\bar{3}$ (Tsuchiya and Tsuchiya, 2011)

3.2. $\delta\text{-Al}(\text{OH})_3$

120 $\text{P2}_12_12_1$ (no. 19) to Pnma (no. 62) at ~ 67 GPa Matsui et al. (2011). Xue and Kanzaki (2007)

3.3. *Ca-perovskite*

Tetragonal (probably P4/mmm, no. 123) (Shim et al., 2002) → Cubic (Pm3m) for low Al contents. For pure CaSiO₃, transition occurs at ca. 580 K at 50 GPa (Kurashina et al., 2004; Komabayashi et al., 2007)

Orthorhombic (Pbnm) → Cubic (Pm3m) for Al-rich CaSiO₃ (Kurashina et al., 2004). At 5.9 wt%, transition occurs at ca. 1840 K at 50 GPa (Kurashina et al., 2004).

In a pyrolitic mantle composition, the CaSiO₃ perovskite contains 1.0–2.3 wt.% Al₂O₃ at upper mantle pressures and somewhat less (0.7–1.6 wt.%) at lower mantle pressures (Kesson et al., 1998; Wood, 2000; Hirose, 2002)

In a MORB composition, 2.0–4.8 wt.% and 1.2–4.5 wt.% Al₂O₃ are included in CaSiO₃-rich perovskite in garnetite and perovskitite lithologies, respectively (Kesson et al., 1994; Irifune and Ringwood, 1993; Hirose and Fei, 2002).

3.4. *Reaction with metal*

(Terasaki et al., 2012)

4. Lower mantle scatterers

Deuss et al. (2013) Kaneshima and Helffrich (1998, 2003) Niu et al. (2003) Krüger et al. (2001) Kaneshima (2009) Bina et al. (2010) Kaneshima et al. (2010) Bentham and Rost (2014) Asahara et al. (2013) Mookherjee (2011)

5. Experimental and analytical techniques

6. Chemical composition

Figure 1: Single crystal XRD spectra of post-stishovite

7. Conclusions

References

- 145 Andrault, D., Angel, R.J., Mosenfelder, J.L., Le Bihan, T., 2003. Equation of state of stishovite to lower mantle pressures. *American Mineralogist* 88, 301–307.
- Andrault, D., Fiquet, G., Guyot, F., Hanfland, M., 1998. Pressure-Induced Landau-Type Transition in Stishovite. *Science* 282, 720.
- 150 Asahara, Y., Hirose, K., Ohishi, Y., Hirao, N., Ozawa, H., Murakami, M., 2013. Acoustic velocity measurements for stishovite across the post-stishovite phase transition under deviatoric stress: Implications for the seismic features of subducting slabs in the mid-mantle. *American Mineralogist* 98, 2053–2062.
- Benthams, H.L.M., Rost, S., 2014. Scattering beneath Western Pacific subduc-
155 tion zones: evidence for oceanic crust in the mid-mantle. *Geophysical Journal International* 197, 1627–1641.
- Bina, C.R., Suetsugu, D., Bina, C., Inoue, T., Wiens, D., Jellinek, M., 2010. Scale limits of free-silica seismic scatterers in the lower mantle. *Physics of the Earth and Planetary Interiors* 183, 110–114.
- 160 Bindi, L., Nishi, M., Tsuchiya, J., Irifune, T., 2014. Crystal chemistry of dense hydrous magnesium silicates: The structure of phase H, MgSiH_2O_4 , synthesized at 45 GPa and 1000 °C. *American Mineralogist* 99, 1802–1805.
- Bolfan-Casanova, N., Andrault, D., Amiguet, E., Guignot, N., 2009. Equation of state and post-stishovite transformation of Al-bearing silica up to 100 GPa
165 and 3000 K. *Physics of the Earth and Planetary Interiors* 174, 70–77.
- Bromiley, G.D., Bromiley, F.A., Bromiley, D.W., 2006. On the mechanisms for H and Al incorporation in stishovite. *Physics and Chemistry of Minerals* 33, 613–621.
- Carpenter, M.A., Hemley, R.J., Mao, H.K., 2000. High-pressure elasticity of
170 stishovite and the $\text{P4}_2/\text{mmn} \rightarrow \text{Pnnm}$ phase transition. *Journal of Geophysical Research* 105, 10807.

- Chung, J.I., Kagi, H., 2002. High concentration of water in stishovite in the MORB system. *Geophysical Research Letters* 29, 16–1–16–4. 2020.
- Cottaar, S., Heister, T., Rose, I., Unterborn, C., 2014. BurnMan: A lower
175 mantle mineral physics toolkit. *Geochemistry, Geophysics, Geosystems* 15, 1164–1179.
- Deuss, A., Andrews, J., Day, E., 2013. Seismic Observations of Mantle Discontinuities and Their Mineralogical and Dynamical Interpretation. pp. 295–323.
- Dubrovinsky, L.S., Belonoshko, A.B., 1996. Pressure-induced phase transi-
180 tion and structural changes under deviatoric stress of stishovite to CaCl_2 -like structure. *Geochimica et Cosmochimica Acta* 60, 3657–3663.
- Gleason, A., Jeanloz, R., Kunz, M., 2008. Pressure-temperature stability studies of feooh using x-ray diffraction. *American Mineralogist* 93, 1882–1885.
- Gleason, A., Quiroga, C., Suzuki, A., Pentcheva, R., Mao, W., 2013. Sym-
185 metrization driven spin transition in ε -FeOOH at high pressure. *Earth and Planetary Science Letters* 379, 49 – 55.
- Hemley, R.J., Shu, J., Carpenter, M.A., Hu, J., Mao, H.K., Kingma, K.J., 2000. Strain/order parameter coupling in the ferroelastic transition in dense SiO_2 . *Solid State Communications* 114, 527–532.
- Hirose, K., 2002. Phase transitions in pyrolitic mantle around 670-km depth:
190 Implications for upwelling of plumes from the lower mantle. *Journal of Geophysical Research (Solid Earth)* 107, 2078.
- Hirose, K., Fei, Y., 2002. Subsolidus and melting phase relations of basaltic composition in the uppermost lower mantle. *Geochimica et Cosmochimica*
195 *Acta* 66, 2099–2108.
- Hirose, K., Takafuji, N., Sata, N., Ohishi, Y., 2005. Phase transition and density of subducted MORB crust in the lower mantle. *Earth and Planetary Science Letters* 237, 239–251.

- Holland, T.J.B., Powell, R., 2011. An improved and extended internally consistent thermodynamic dataset for phases of petrological interest, involving a
200 new equation of state for solids. *Journal of Metamorphic Geology* 29, 333–383.
- Irifune, T., Ringwood, A.E., 1993. Phase transformations in subducted oceanic crust and buoyancy relationships at depths of 600–800 km in the mantle. *Earth and Planetary Science Letters* 117, 101–110.
- 205 Kaneshima, S., 2009. Seismic scatterers at the shallowest lower mantle beneath subducted slabs. *Earth and Planetary Science Letters* 286, 304–315.
- Kaneshima, S., Helffrich, G., 1998. Detection of lower mantle scatterers north-east of the Marianna subduction zone using short-period array data. *Journal of Geophysical Research* 103, 4825–4838.
- 210 Kaneshima, S., Helffrich, G., 2003. Subparallel dipping heterogeneities in the mid-lower mantle. *Journal of Geophysical Research (Solid Earth)* 108, 2272.
- Kaneshima, S., Helffrich, G., Suetsugu, D., Bina, C., Inoue, T., Wiens, D., Jellinek, M., 2010. Small scale heterogeneity in the mid-lower mantle beneath the circum-Pacific area. *Physics of the Earth and Planetary Interiors* 183, 91–
215 103.
- Kesson, S.E., Fitz Gerald, J.D., Shelley, J.M., 1998. Mineralogy and dynamics of a pyrolite lower mantle. *Nature* 393, 252–255.
- Kesson, S.E., Fitz Gerald, J.D., Shelley, J.M.G., 1994. Mineral chemistry and density of subducted basaltic crust at lower-mantle pressures. *Nature* 372,
220 767–769.
- Kingma, K., Mao, H.K., Hemley, R., 1996. Synchrotron X-ray diffraction of SiO_2 to multimegabar pressures. *High Pressure Research* 14, 363–374.
- Kingma, K.J., Cohen, R.E., Hemley, R.J., Mao, H.K., 1995. Transformation of stishovite to a denser phase at lower-mantle pressures. *Nature* 374, 243–245.

- 225 Komabayashi, T., Hirose, K., Sata, N., Ohishi, Y., Dubrovinsky, L.S., 2007. Phase transition in CaSiO₃ perovskite. *Earth and Planetary Science Letters* 260, 564 – 569.
- Komatsu, K., Kuribayashi, T., Sano, A., Ohtani, E., Kudoh, Y., 2006. Redetermination of the high-pressure modification of AlOOH from single-crystal
230 synchrotron data. *Acta Crystallographica Section E* 62, 216–218.
- Komatsu, K., Sano-Furukawa, A., Kagi, H., 2011. Effects of Mg and Si ions on the symmetry of δ -AlOOH. *Physics and Chemistry of Minerals* 38, 727–733.
- Krüger, F., Baumann, M., Scherbaum, F., Weber, M., 2001. Mid mantle scatterers near the Mariana Slab detected with a double array method. *Geophysical
235 Research Letters* 28, 667–670.
- Kudoh, Y., Kuribayashi, T., Suzuki, A., Ohtani, E., Kamada, T., 2004. Space group and hydrogen sites of δ -AlOOH and implications for a hypothetical high-pressure form of Mg(OH)₂. *Physics and Chemistry of Minerals* 31, 360–364.
- 240 Kurashina, T., Hirose, K., Ono, S., Sata, N., Ohishi, Y., 2004. Phase transition in Al-bearing CaSiO₃ perovskite: implications for seismic discontinuities in the lower mantle. *Physics of the Earth and Planetary Interiors* 145, 67–74.
- Kuribayashi, T., Sano-Furukawa, A., Nagase, T., 2014. Observation of pressure-induced phase transition of δ -AlOOH by using single-crystal synchrotron X-ray diffraction method. *Physics and Chemistry of Minerals* 41, 303–312.
245
- Lakshtanov, D.L., Sinogeikin, S.V., Litasov, K.D., Prakapenka, V.B., Hellwig, H., Wang, J., Sanches-Valle, C., Perrillat, J.P., Chen, B., Somayazulu, M., Li, J., Ohtani, E., Bass, J.D., 2007. The post-stishovite phase transition in hydrous alumina-bearing SiO₂ in the lower mantle of the Earth. *Proceedings of the National Academy of Sciences* 104, 13588–13590.
250
- Lakshtanov, D.L., Vanpeteghem, C.B., Jackson, J.M., Bass, J.D., Shen, G., Prakapenka, V.B., Litasov, K., Ohtani, E., 2005. The equation of state of

Al,H-bearing SiO₂ stishovite to 58 GPa. *Physics and Chemistry of Minerals* 32, 466–470.

255 Li, S., Ahuja, R., Johansson, B., 2006. The elastic and optical properties of the high-pressure hydrous phase δ -AlOOH. *Solid State Communications* 137, 101–106.

Litasov, K.D., Kagi, H., Shatskiy, A., Ohtani, E., Lakshtanov, D.L., Bass, J.D., Ito, E., 2007. High hydrogen solubility in Al-rich stishovite and water transport in the lower mantle. *Earth and Planetary Science Letters* 262, 620 –
260 634.

Litasov, K.D., Ohtani, E., 2005. Phase relations in hydrous MORB at 18-28 GPa: implications for heterogeneity of the lower mantle. *Physics of the Earth and Planetary Interiors* 150, 239–263.

265 Matsui, M., Komatsu, K., Ikeda, E., Sano-Furukawa, A., Gotou, H., Yagi, T., 2011. The crystal structure of δ -Al(OH)₃: Neutron diffraction measurements and ab initio calculations. *American Mineralogist* 96, 854–859.

McKnight, R.E., Carpenter, M.A., Darling, T.W., Buckley, A., Taylor, P.A., 2007. Acoustic dissipation associated with phase transitions in lawsonite,
270 caal₂si₂o₇(oh)₂h₂o. *American Mineralogist* 92, 1665–1672.

Mookherjee, M., 2011. Mid-mantle anisotropy: Elasticity of aluminous phases in subducted MORB. *Geophysical Research Letters* 38, n/a–n/a. L14302.

Nishi, M., Irifune, T., Tsuchiya, J., Tange, Y., Nishihara, Y., Fujino, K., Higo, Y., 2014. Stability of hydrous silicate at high pressures and water transport
275 to the deep lower mantle. *Nature Geoscience* 7, 224–227.

Niu, F., Kawakatsu, H., Fukao, Y., 2003. Seismic evidence for a chemical heterogeneity in the midmantle: A strong and slightly dipping seismic reflector beneath the Mariana subduction zone. *Journal of Geophysical Research (Solid Earth)* 108, 2419.

- 280 Nomura, R., Hirose, K., Sata, N., Ohishi, Y., Suetsugu, D., Bina, C., Inoue, T., Wiens, D., Jellinek, M., 2010. Precise determination of post-stishovite phase transition boundary and implications for seismic heterogeneities in the mid-lower mantle. *Physics of the Earth and Planetary Interiors* 183, 104–109.
- Nowick, A., Berry, D., 1972. Anelastic relaxation in crystalline solids, acad. Press, New York .
- 285 Ohira, I., Ohtani, E., Sakai, T., Miyahara, M., Hirao, N., Ohishi, Y., Nishijima, M., 2014. Stability of a hydrous δ -phase, $\text{AlOOH-MgSiO}_2(\text{OH})_2$, and a mechanism for water transport into the base of lower mantle. *Earth and Planetary Science Letters* 401, 12–17.
- 290 Ohtani, E., Amaiike, Y., Kamada, S., Sakamaki, T., Hirao, N., 2014. Stability of hydrous phase H MgSiO_4H_2 under lower mantle conditions. *Geophysical Research Letters* 41, 8283–8287. 2014GL061690.
- Ono, S., 1999. High temperature stability limit of phase egg, $\text{AlSiO}_3(\text{OH})$. *Contributions to Mineralogy and Petrology* 137, 83–89.
- 295 Ono, S., Hirose, K., Murakami, M., Isshiki, M., 2002. Post-stishovite phase boundary in SiO_2 determined by in situ X-ray observations. *Earth and Planetary Science Letters* 197, 187–192.
- Ono, S., Suto, T., Hirose, K., Kuwayama, Y., Komabayashi, T., Kikegawa, T., 2002. Equation of state of Al-bearing stishovite to 40 GPa at 300 K. *American Mineralogist* 87, 1486–1489.
- 300 Panero, W.R., 2006. Aluminum incorporation in stishovite. *Geophysical Research Letters* 33, n/a–n/a. L20317.
- Panero, W.R., Benedetti, L.R., Jeanloz, R., 2003. Transport of water into the lower mantle: Role of stishovite. *Journal of Geophysical Research (Solid Earth)* 108, 2039.
- 305

- Panero, W.R., Stixrude, L.P., 2004. Hydrogen incorporation in stishovite at high pressure and symmetric hydrogen bonding in δ -AlOOH. *Earth and Planetary Science Letters* 221, 421 – 431.
- Pawley, A.R., McMillan, P.F., Holloway, J.R., 1993. Hydrogen in Stishovite, with Implications for Mantle Water Content. *Science* 261, 1024–1026.
- Sano-Furukawa, A., Kagi, H., Nagai, T., Nakano, S., Fukura, S., Ushijima, D., Iizuka, R., Ohtani, E., Yagi, T., 2009. Change in compressibility of δ -AlOOH and δ -AlOOD at high pressure: A study of isotope effect and hydrogen-bond symmetrization. *American Mineralogist* 94, 1255–1261.
- Sano-Furukawa, A., Komatsu, K., Vanpeteghem, C.B., Ohtani, E., 2008. Neutron diffraction study of δ -AlOOD at high pressure and its implication for symmetrization of the hydrogen bond. *American Mineralogist* 93, 1558–1567.
- Shieh, S.R., Duffy, T.S., Li, B., 2002. Strength and Elasticity of SiO_2 across the Stishovite- CaCl_2 -type Structural Phase Boundary. *Physical Review Letters* 89, 255507.
- Shim, S.H., Jeanloz, R., Duffy, T.S., 2002. Tetragonal structure of CaSiO_3 perovskite above 20 GPa. *Geophysical Research Letters* 29, 2166.
- Singh, A.K., Andrault, D., Bouvier, P., 2012. X-ray diffraction from stishovite under nonhydrostatic compression to 70 GPa: Strength and elasticity across the tetragonal \rightarrow orthorhombic transition. *Physics of the Earth and Planetary Interiors* 208209, 1 – 10.
- Smyth, J.R., Swope, R.J., Pawley, A.R., 1995. H in rutile-type compounds; II, Crystal chemistry of Al substitution in H-bearing stishovite. *American Mineralogist* 80, 454–456.
- Snoek, J.L., 1941. Effect of small quantities of carbon and nitrogen on the elastic and plastic properties of iron. *Physica* 8, 711–733.

- Stixrude, L., Lithgow-Bertelloni, C., 2011. Thermodynamics of mantle minerals
- II. Phase equilibria. *Geophysical Journal International* 184, 1180–1213.
- Suzuki, A., 2009. Compressibility of the high-pressure polymorph of AlOOH to
335 17 GPa. *Mineralogical Magazine* 73, 479–485.
- Suzuki, A., Ohtani, E., Kamada, T., 2000. A new hydrous phase δ -AlOOH
synthesized at 21 GPa and 1000 °C. *Physics and Chemistry of Minerals* 27,
689–693.
- Terasaki, H., Ohtani, E., Sakai, T., Kamada, S., Asanuma, H., Shibazaki, Y.,
340 Hirao, N., Sata, N., Ohishi, Y., Sakamaki, T., Suzuki, A., Ichi Funakoshi, K.,
2012. Stability of FeNi hydride after the reaction between Fe-Ni alloy and
hydrous phase (δ -AlOOH) up to 1.2 Mbar: Possibility of H contribution to
the core density deficit. *Physics of the Earth and Planetary Interiors* 194–195,
18–24.
- 345 Tsuchida, Y., Yagi, T., 1989. A new, post-stishovite high-pressure polymorph
of silica. *Nature* 340, 217–220.
- Tsuchiya, J., 2013. First principles prediction of a new high-pressure phase of
dense hydrous magnesium silicates in the lower mantle. *Geophysical Research
Letters* 40, 4570–4573.
- 350 Tsuchiya, J., Tsuchiya, T., 2009. Elastic properties of δ -AlOOH under pressure:
First principles investigation. *Physics of the Earth and Planetary Interiors*
174, 122 – 127. *Advances in High Pressure Mineral Physics: from Deep Mantle
to the Core*.
- Tsuchiya, J., Tsuchiya, T., 2011. First-principles prediction of a high-pressure
355 hydrous phase of AlOOH. *Phys. Rev. B* 83, 054115.
- Umemoto, K., Kawamura, K., Hirose, K., , Revenaugh, J., Wentzcovitch, R.,
2015. Post-stishovite transition in hydrous aluminous SiO_2 . *PNAS* -.

- Vanpeteghem, C., Sano, A., Komatsu, K., Ohtani, E., Suzuki, A., 2007. Neutron
diffraction study of aluminous hydroxide δ -AlOOD. *Physics and Chemistry*
360 *of Minerals* 34, 657–661.
- Vanpeteghem, C.B., Ohtani, E., Kondo, T., 2002. Equation of state of the
hydrous phase δ -AlOOH at room temperature up to 22.5 GPa. *Geophysical*
Research Letters 29, 1119.
- Walter, M., Thomson, A., Wang, W., Lord, O., Ross, J., McMahon, S., Baron,
365 M., Melekhova, E., Kleppe, A., Kohn, S., 2015. The stability of hydrous
silicates in Earth’s lower mantle: Experimental constraints from the systems
MgO–SiO₂–H₂O and MgO–Al₂O₃–SiO₂–H₂O. *Chemical Geology*, –.
- Wood, B.J., 2000. Phase transformations and partitioning relations in peridotite
under lower mantle conditions. *Earth and Planetary Science Letters* 174, 341–
370 354.
- Xue, X., Kanzaki, M., 2007. High-Pressure δ -Al(OH)₃ and δ -AlOOH Phases
and Isostructural Hydroxides/Oxyhydroxides: New Structural Insights from
High-Resolution ¹H and ²⁷Al NMR. *The Journal of Physical Chemistry B*
111, 13156–13166.