

# Quenchable water-rich aluminous post-stishovite: implications for water and seismic scatterers in the lower mantle

R. Myhill, D. J. Frost, T. Boffa-Ballaran, H. Bureau, C. Raepsaet

*Bayerisches Geoinstitut, Universität Bayreuth, Universitätsstrasse 30, 95447 Bayreuth,  
Germany*

---

## Abstract

*Keywords:* high pressure, post-stishovite, water, slab, scatterers

---

## 1. Introduction

The stishovite structure (tetragonal,  $P4_2/mnm$ , no. 136) undergoes a weak first-order transition (Andrault et al., 1998; Hemley et al., 2000) with second-order characteristics of Landau/ferroelastic type (Tsuchida and Yagi, 1989; Carpenter et al., 2000) to the  $\text{CaCl}_2$ -type structure (orthorhombic,  $Pnmm$ , no. 58). This transition occurs at pressures of  $\sim 50$  GPa at room temperature (Kingma et al., 1995; Andrault et al., 1998), increasing with temperature to  $\sim 70$  GPa at 2200 K (Hirose et al., 2005; Nomura et al., 2010). A somewhat higher  $dT/dP$  was observed by Ono et al. (2002). The nature of this transition means that the shear modulus  $C_{11} - C_{12}$  decreases with increasing pressure and vanishes at the transition pressure, as observed in spectroscopic and high pressure diffraction studies (Kingma et al., 1995; Shieh et al., 2002). The stishovite-post-stishovite transition is therefore of interest to seismologists, as it should produce zones of low shear wave velocity which act as scatterers in the deep mantle.

Deviatoric stresses in the diamond anvil cell can cause large variations in the


---

\*Corresponding author: R. Myhill


*Email address:* myhill.bob@gmail.com (R. Myhill, D. J. Frost, T. Boffa-Ballaran, H. Bureau, C. Raepsaet)

pressure of the stishovite to post-stishovite transition. Nonhydrostaticity greatly decreases the transition pressure from 60 GPa (Andrault et al., 2003; Hemley et al., 2000) to 40 GPa (Kingma et al., 1996; Singh et al., 2012). Molecular and lattice dynamic studies suggest that only 1.5–2.5 GPa differential stress is  
20 required to cause this decrease in pressure (Dubrovinsky and Belonoshko, 1996).

Aluminium has a profound effect on the stability of the post-stishovite phase.  $\text{Al}_2\text{O}_3$  can be incorporated into stishovite by the substitution of Si by Al with the formation of oxygen vacancies (Smyth et al., 1995; Hirose et al., 2005; Bromiley et al., 2006; Lakshtanov et al., 2007a):




25 An alternative mechanism, where charge balance is accomplished by Al occupying the large interstitial sites is probably minor Smyth et al. (1995):



It has been argued that the addition of 4 wt%  $\text{Al}_2\text{O}_3$  in the absence of any other components causes the transition to shift from 50 to 23 GPa at room temperature by inducing a ‘chemical pressure’ (Bolfan-Casanova et al., 2009). In  
30 stishovite, there is one octahedral cation position (Wyckoff notation 2a) located at the origin and the body center of the tetragonal cell. One oxygen at position 4f creates a moderately distorted octahedron with point symmetry mmm. The addition of Al *decreases* distortion of this octahedron (Smyth et al., 1995). The addition of Al should therefore have little effect on the stishovite-post-stishovite  
35 transition (Panero, 2006). The results of Bolfan-Casanova et al. (2009) are therefore unexpected; it seems possible that another component (such as  $\text{H}_2\text{O}$ ) or deviatoric stress acted to stabilise the  $\text{CaCl}_2$  structure.

It has also been suggested that the coupled substitution of Al and H may have a significant effect on the transition pressure. Lakshtanov et al. (2007b)  
40 have shown that the addition of 6 wt%  $\text{Al}_2\text{O}_3$  and 0.24 wt%  $\text{H}_2\text{O}$  reduces the transition pressure to 24 GPa at room temperature. The substitution mechanism in this case is probably



Given that high pressure  $\delta$ -AlOOH is isostructural with the post-stishovite phase above ca. 19 GPa (Sano-Furukawa et al., 2008; Kuribayashi et al., 2014),  
 45 after transforming from the P2<sub>1</sub>nm structure (no. 31) (Suzuki et al., 2000; Komatsu et al., 2006; Vanpeteghem et al., 2007), it seems sensible to suggest that addition of an AlOOH component lowers the transition pressure. Ab initio calculations support this suggestion (Umemoto et al., 2015). The equilibrium transition for 6.25 mol% AlOOH is at ca. 15 GPa at room temperature, im-  
 50 plying an even more marked reduction than observed in the experimental data (Lakshtanov et al., 2007b). This could be due to inaccuracies in the ab-initio data, or metastable preservation of the tetragonal phase in the experiments. Umemoto et al. (2015) also suggest that the larger number of hydrogen sites stabilises the tetragonal structure at high temperatures, changing the slope of  
 55 the transition. The tetragonal  $\rightarrow$  orthorhombic transition in their simulations is associated with splitting of the hydrogen sites on the equatorial oxygens into two groups, one occupied and the other unoccupied. Additionally, the redistribution of hydrogens among equatorial oxygens in the tetragonal  $\rightarrow$  orthorhombic transition implies that it is now first order and no longer ferroelastic. Anelasticity  
 60 by hydrogen hopping seems a plausible alternative to reduce seismic body wave speeds.

Despite this, H<sub>2</sub>O contents in tetragonal stishovite measured using the FTIR calibration of Pawley et al. (1993) are less than 20% of that expected from an AlOOH component (Panero et al., 2003; Bromiley et al., 2006; Litasov et al.,  
 65 2007) and in MORB compositions (Chung and Kagi, 2002).

This study was designed to investigate water solubility in Al-rich stishovite by synthesising crystals at relatively high temperature (Ono, 1999). In doing so, we created and quenched an Al-H rich post-stishovite phase.

### 1.1. Hysteresis vs. stabilisation

70 (Umemoto et al., 2015)

### 1.2. Mechanism of shear wave velocity reduction

Ferroelasticity (Carpenter et al., 2000) Snoek relaxation (anelasticity via H mobility) (Snoek, 1941; Nowick and Berry, 1972; McKnight et al., 2007).

### 1.3. Mechanisms of Al-, H incorporation

75 We found that hydrogen was most stable when bonded to the apical oxygen of the Al-octahedron, with the hydroxyl bond along  $\langle 110 \rangle$  and co-planar with Al (Panero and Stixrude, 2004).

The relatively short O-O distance and correspondingly long O-H distance in the stishovite-AlOOH solid solution would lead one to expect a low OH stretch-  
80 ing frequency compared to other nominally anhydrous minerals, consistent with the value of  $3111 \text{ cm}^{-1}$  observed by Pawley et al. (1993).

Symmetric bonding (Panero and Stixrude, 2004)

### 1.4. Water concentrations in stishovite

Pawley et al. (1993) calibration Panero et al. (2003) Bromiley et al. (2006),  
85 Litasov et al. (2007), MORB Chung and Kagi (2002),

### 1.5. Elastic properties

Andrault et al. (2003) Lakshtanov et al. (2005) Ono et al. (2002) Sano-Furukawa et al. (2009)

### 1.6. Phase relations in the lower mantle

90 Irifune and Ringwood (1993) Wood (2000) Hirose (2002) Walter et al. (2015) Litasov and Ohtani (2005)

### 1.7. Datasets and modelling

Holland and Powell (2011) Stixrude and Lithgow-Bertelloni (2011) Cottaar et al. (2014)

#### 95 1.7.1. Elastic properties AlOOH

Suzuki (2009) (Sano-Furukawa et al., 2008) Vanpeteghem et al. (2002) Tsuchiya and Tsuchiya (2009) Li et al. (2006)

## 2. The solid solution ( $\text{MgSi}$ , $\text{Fe}^{2+}\text{Si}$ , $\text{Fe}_2^{3+}$ , $\text{Al}_2$ ) $\text{O}_4\text{H}_2$

### 2.1. Endmembers

100 2.1.1.  $\text{SiO}_2$

2.1.2.  $\delta\text{-AlOOH}$

2.1.3.  $\delta\text{-FeOOH}$

Pnma (no.62  $\alpha$ ) to  $\text{P2}_1\text{nm}$  (no. 31,  $\varepsilon$ ) (Gleason et al., 2008) to Pnnm (no. 58, high spin) to Pnnm (no. 58, low spin) (Gleason et al., 2013)

105 2.1.4. “Phase H”

The addition of an  $\text{MgSiO}_4\text{H}_2$  component also stabilises the Pnnm structure in stishovite (Komatsu et al., 2011). A previous candidate space group (Pnn2; no. 34 Kudoh et al., 2004) is now thought to be unlikely.

Stability (Ohtani et al., 2014) Ab initio  $\text{P2}/\text{m}$  (no. 10) (Tsuchiya, 2013)  
110 Pnnm (no. 58) (Bindi et al., 2014)  $\text{P2}_1\text{nm}$  (no. 31) (Nishi et al., 2014)

Kudoh et al. (2004) suggested a hypothetical high pressure form of  $\text{Mg}(\text{OH})_2$  also in the space group Pnnm from Mg incorporation. Importance of the AlOOH - phase H solid solution Ohira et al. (2014)

## 3. Other phases

115 3.1. *Water in seifertite*

A few studies have suggested that the  $\text{CaCl}_2$  type may undergo a phase transition to a mineral named seifertite with the scrutinyite ( $\alpha\text{-PbO}_2$ ) structure (orthorhombic, Pbcn, no. 60 or Pb2n) within mantle pressures.

AlOOH to  $\text{Pa}\bar{3}$  (Tsuchiya and Tsuchiya, 2011)

120 3.2.  $\delta\text{-Al}(\text{OH})_3$

$\text{P2}_1\text{2}_1\text{2}_1$  (no. 19) to Pnma (no. 62) at  $\sim 67$  GPa Matsui et al. (2011). Xue and Kanzaki (2007)

### 3.3. *Ca-perovskite*

Tetragonal (probably  $P4/mmm$ , no. 123) (Shim et al., 2002)  $\rightarrow$  Cubic  
125 (Pm3m) for low Al contents. For pure  $\text{CaSiO}_3$ , transition occurs at ca. 580  
K at 50 GPa (Kurashina et al., 2004; Komabayashi et al., 2007)

Orthorhombic (Pbnm)  $\rightarrow$  Cubic (Pm3m) for Al-rich  $\text{CaSiO}_3$  (Kurashina  
et al., 2004). At 5.9 wt%, transition occurs at ca. 1840 K at 50 GPa (Kurashina  
et al., 2004).

130 In a pyrolitic mantle composition, the  $\text{CaSiO}_3$  perovskite contains 1.0–2.3  
wt.%  $\text{Al}_2\text{O}_3$  at upper mantle pressures and somewhat less (0.7–1.6 wt.%) at  
lower mantle pressures (Kesson et al., 1998; Wood, 2000; Hirose, 2002)

In a MORB composition, 2.0–4.8 wt.% and 1.2–4.5 wt.%  $\text{Al}_2\text{O}_3$  are included  
in  $\text{CaSiO}_3$ -rich perovskite in garnetite and perovskitite lithologies, respectively  
135 (Kesson et al., 1994; Irifune and Ringwood, 1993; Hirose and Fei, 2002).

### 3.4. *Reaction with metal*

(Terasaki et al., 2012)

## 4. Lower mantle scatterers

Deuss et al. (2013) Kaneshima and Helffrich (1998, 2003) Niu et al. (2003)  
140 Krüger et al. (2001) Kaneshima (2009) Bina et al. (2010) Kaneshima et al.  
(2010) Bentham and Rost (2014) Asahara et al. (2013) Mookherjee (2011)

## 5. Experimental and analytical techniques

## 6. Chemical composition

Figure 1: Single crystal XRD spectra of post-stishovite

## 7. Conclusions

145 **References**

- Andrault, D., Angel, R.J., Mosenfelder, J.L., Le Bihan, T., 2003. Equation of state of stishovite to lower mantle pressures. *American Mineralogist* 88, 301–307.
- Andrault, D., Angel, R.J., Mosenfelder, J.L., Le Bihan, T., 2003. Equation  
150 of state of stishovite to lower mantle pressures. *American Mineralogist* 88, 301–307.
- Andrault, D., Fiquet, G., Guyot, F., Hanfland, M., 1998. Pressure-Induced Landau-Type Transition in Stishovite. *Science* 282, 720.
- Asahara, Y., Hirose, K., Ohishi, Y., Hirao, N., Ozawa, H., Murakami, M.,  
155 2013. Acoustic velocity measurements for stishovite across the post-stishovite phase transition under deviatoric stress: Implications for the seismic features of subducting slabs in the mid-mantle. *American Mineralogist* 98, 2053–2062.
- Benthams, H.L.M., Rost, S., 2014. Scattering beneath Western Pacific subduction zones: evidence for oceanic crust in the mid-mantle. *Geophysical Journal  
160 International* 197, 1627–1641.
- Bina, C.R., Suetsugu, D., Bina, C., Inoue, T., Wiens, D., Jellinek, M., 2010. Scale limits of free-silica seismic scatterers in the lower mantle. *Physics of the Earth and Planetary Interiors* 183, 110–114.
- Bindi, L., Nishi, M., Tsuchiya, J., Irifune, T., 2014. Crystal chemistry of dense  
165 hydrous magnesium silicates: The structure of phase H,  $\text{MgSiH}_2\text{O}_4$ , synthesized at 45 GPa and 1000 °C. *American Mineralogist* 99, 1802–1805.
- Bolfan-Casanova, N., Andrault, D., Amiguet, E., Guignot, N., 2009. Equation of state and post-stishovite transformation of Al-bearing silica up to 100 GPa and 3000 K. *Physics of the Earth and Planetary Interiors* 174, 70–77.
- 170 Bromiley, G.D., Bromiley, F.A., Bromiley, D.W., 2006. On the mechanisms for H and Al incorporation in stishovite. *Physics and Chemistry of Minerals* 33, 613–621.

- Carpenter, M.A., Hemley, R.J., Mao, H.K., 2000. High-pressure elasticity of stishovite and the  $P4_2/mnm \rightarrow Pnnm$  phase transition. *Journal of Geophysical Research* 105, 10807.
- 175
- Chung, J.I., Kagi, H., 2002. High concentration of water in stishovite in the MORB system. *Geophysical Research Letters* 29, 16–1–16–4. 2020.
- Cottaar, S., Heister, T., Rose, I., Unterborn, C., 2014. BurnMan: A lower mantle mineral physics toolkit. *Geochemistry, Geophysics, Geosystems* 15, 1164–1179.
- 180
- Deuss, A., Andrews, J., Day, E., 2013. Seismic Observations of Mantle Discontinuities and Their Mineralogical and Dynamical Interpretation. pp. 295–323.
- Dubrovinsky, L.S., Belonoshko, A.B., 1996. Pressure-induced phase transition and structural changes under deviatoric stress of stishovite to  $\text{CaCl}_2$ -like structure. *Geochimica et Cosmochimica Acta* 60, 3657–3663.
- 185
- Gleason, A., Jeanloz, R., Kunz, M., 2008. Pressure-temperature stability studies of feooh using x-ray diffraction. *American Mineralogist* 93, 1882–1885.
- Gleason, A., Quiroga, C., Suzuki, A., Pentcheva, R., Mao, W., 2013. Symmetrization driven spin transition in  $\varepsilon\text{-FeOOH}$  at high pressure. *Earth and Planetary Science Letters* 379, 49 – 55.
- 190
- Hemley, R.J., Shu, J., Carpenter, M.A., Hu, J., Mao, H.K., Kingma, K.J., 2000. Strain/order parameter coupling in the ferroelastic transition in dense  $\text{SiO}_2$ . *Solid State Communications* 114, 527–532.
- Hirose, K., 2002. Phase transitions in pyrolitic mantle around 670-km depth: Implications for upwelling of plumes from the lower mantle. *Journal of Geophysical Research (Solid Earth)* 107, 2078.
- 195
- Hirose, K., Fei, Y., 2002. Subsolvus and melting phase relations of basaltic composition in the uppermost lower mantle. *Geochimica et Cosmochimica Acta* 66, 2099–2108.



- 200 Hirose, K., Takafuji, N., Sata, N., Ohishi, Y., 2005. Phase transition and density  
of subducted MORB crust in the lower mantle. *Earth and Planetary Science  
Letters* 237, 239–251.
- Holland, T.J.B., Powell, R., 2011. An improved and extended internally con-  
sistent thermodynamic dataset for phases of petrological interest, involving a  
205 new equation of state for solids. *Journal of Metamorphic Geology* 29, 333–383.
- Irifune, T., Ringwood, A.E., 1993. Phase transformations in subducted oceanic  
crust and buoyancy relationships at depths of 600–800 km in the mantle.  
*Earth and Planetary Science Letters* 117, 101–110.
- Kaneshima, S., 2009. Seismic scatterers at the shallowest lower mantle beneath  
210 subducted slabs. *Earth and Planetary Science Letters* 286, 304–315.
- Kaneshima, S., Helffrich, G., 1998. Detection of lower mantle scatterers north-  
east of the Marianna subduction zone using short-period array data. *Journal  
of Geophysical Research* 103, 4825–4838.
- Kaneshima, S., Helffrich, G., 2003. Subparallel dipping heterogeneities in the  
215 mid-lower mantle. *Journal of Geophysical Research (Solid Earth)* 108, 2272.
- Kaneshima, S., Helffrich, G., Suetsugu, D., Bina, C., Inoue, T., Wiens, D.,  
Jellinek, M., 2010. Small scale heterogeneity in the mid-lower mantle beneath  
the circum-Pacific area. *Physics of the Earth and Planetary Interiors* 183, 91–  
103.
- 220 Kesson, S.E., Fitz Gerald, J.D., Shelley, J.M., 1998. Mineralogy and dynamics  
of a pyrolite lower mantle. *Nature* 393, 252–255.
- Kesson, S.E., Fitz Gerald, J.D., Shelley, J.M.G., 1994. Mineral chemistry and  
density of subducted basaltic crust at lower-mantle pressures. *Nature* 372,  
767–769.
- 225 Kingma, K., Mao, H.K., Hemley, R., 1996. Synchrotron X-ray diffraction of  
SiO<sub>2</sub> to multimegabar pressures. *High Pressure Research* 14, 363–374.

- Kingma, K.J., Cohen, R.E., Hemley, R.J., Mao, H.K., 1995. Transformation of stishovite to a denser phase at lower-mantle pressures. *Nature* 374, 243–245.
- Komabayashi, T., Hirose, K., Sata, N., Ohishi, Y., Dubrovinsky, L.S., 2007. Phase transition in CaSiO<sub>3</sub> perovskite. *Earth and Planetary Science Letters* 260, 564 – 569.
- Komatsu, K., Kuribayashi, T., Sano, A., Ohtani, E., Kudoh, Y., 2006. Redetermination of the high-pressure modification of AlOOH from single-crystal synchrotron data. *Acta Crystallographica Section E* 62, 216–218.
- Komatsu, K., Sano-Furukawa, A., Kagi, H., 2011. Effects of Mg and Si ions on the symmetry of  $\delta$ -AlOOH. *Physics and Chemistry of Minerals* 38, 727–733.
- Krüger, F., Baumann, M., Scherbaum, F., Weber, M., 2001. Mid mantle scatterers near the Mariana Slab detected with a double array method. *Geophysical Research Letters* 28, 667–670.
- Kudoh, Y., Kuribayashi, T., Suzuki, A., Ohtani, E., Kamada, T., 2004. Space group and hydrogen sites of  $\delta$ -AlOOH and implications for a hypothetical high-pressure form of Mg(OH)<sub>2</sub>. *Physics and Chemistry of Minerals* 31, 360–364.
- Kurashina, T., Hirose, K., Ono, S., Sata, N., Ohishi, Y., 2004. Phase transition in Al-bearing CaSiO<sub>3</sub> perovskite: implications for seismic discontinuities in the lower mantle. *Physics of the Earth and Planetary Interiors* 145, 67–74.
- Kuribayashi, T., Sano-Furukawa, A., Nagase, T., 2014. Observation of pressure-induced phase transition of  $\delta$ -AlOOH by using single-crystal synchrotron X-ray diffraction method. *Physics and Chemistry of Minerals* 41, 303–312.
- Lakshtanov, D.L., Litasov, K.D., Sinogeikin, S.V., Hellwig, H., Li, J., Ohtani, E., Bass, J.D., 2007a. Effect of al<sup>3+</sup> and h<sup>+</sup> on the elastic properties of stishovite. *American Mineralogist* 92, 1026–1030.

- Lakshtanov, D.L., Sinogeikin, S.V., Litasov, K.D., Prakapenka, V.B., Hellwig, H., Wang, J., Sanches-Valle, C., Perrillat, J.P., Chen, B., Somayazulu, M.,  
 255 Li, J., Ohtani, E., Bass, J.D., 2007b. The post-stishovite phase transition in hydrous alumina-bearing SiO<sub>2</sub> in the lower mantle of the Earth. *Proceedings of the National Academy of Sciences* 104, 13588–13590.
- Lakshtanov, D.L., Vanpeteghem, C.B., Jackson, J.M., Bass, J.D., Shen, G., Prakapenka, V.B., Litasov, K., Ohtani, E., 2005. The equation of state of  
 260 Al,H-bearing SiO<sub>2</sub> stishovite to 58 GPa. *Physics and Chemistry of Minerals* 32, 466–470.
- Li, S., Ahuja, R., Johansson, B., 2006. The elastic and optical properties of the high-pressure hydrous phase  $\delta$ -AlOOH. *Solid State Communications* 137, 101–106.
- 265 Litasov, K.D., Kagi, H., Shatskiy, A., Ohtani, E., Lakshtanov, D.L., Bass, J.D., Ito, E., 2007. High hydrogen solubility in Al-rich stishovite and water transport in the lower mantle. *Earth and Planetary Science Letters* 262, 620 – 634.
- Litasov, K.D., Ohtani, E., 2005. Phase relations in hydrous MORB at 18-28  
 270 GPa: implications for heterogeneity of the lower mantle. *Physics of the Earth and Planetary Interiors* 150, 239–263.
- Matsui, M., Komatsu, K., Ikeda, E., Sano-Furukawa, A., Gotou, H., Yagi, T., 2011. The crystal structure of  $\delta$ -Al(OH)<sub>3</sub>: Neutron diffraction measurements and ab initio calculations. *American Mineralogist* 96, 854–859.
- 275 McKnight, R.E., Carpenter, M.A., Darling, T.W., Buckley, A., Taylor, P.A., 2007. Acoustic dissipation associated with phase transitions in lawsonite, CaAl<sub>2</sub>Si<sub>2</sub>O<sub>7</sub>(OH)<sub>2</sub>H<sub>2</sub>O. *American Mineralogist* 92, 1665–1672.
- Mookherjee, M., 2011. Mid-mantle anisotropy: Elasticity of aluminous phases in subducted MORB. *Geophysical Research Letters* 38, n/a–n/a. L14302.

- 280 Nishi, M., Irifune, T., Tsuchiya, J., Tange, Y., Nishihara, Y., Fujino, K., Higo, Y., 2014. Stability of hydrous silicate at high pressures and water transport to the deep lower mantle. *Nature Geoscience* 7, 224–227.
- Niu, F., Kawakatsu, H., Fukao, Y., 2003. Seismic evidence for a chemical heterogeneity in the midmantle: A strong and slightly dipping seismic reflector  
285 beneath the Mariana subduction zone. *Journal of Geophysical Research (Solid Earth)* 108, 2419.
- Nomura, R., Hirose, K., Sata, N., Ohishi, Y., Suetsugu, D., Bina, C., Inoue, T., Wiens, D., Jellinek, M., 2010. Precise determination of post-stishovite phase transition boundary and implications for seismic heterogeneities in the  
290 mid-lower mantle. *Physics of the Earth and Planetary Interiors* 183, 104–109.
- Nowick, A., Berry, D., 1972. *Anelastic relaxation in crystalline solids*, acad. Press, New York .
- Ohira, I., Ohtani, E., Sakai, T., Miyahara, M., Hirao, N., Ohishi, Y., Nishijima, M., 2014. Stability of a hydrous  $\delta$ -phase,  $\text{AlOOH-MgSiO}_2(\text{OH})_2$ , and a mechanism for water transport into the base of lower mantle. *Earth and Planetary  
295 Science Letters* 401, 12–17.
- Ohtani, E., Amaike, Y., Kamada, S., Sakamaki, T., Hirao, N., 2014. Stability of hydrous phase H  $\text{MgSiO}_4\text{H}_2$  under lower mantle conditions. *Geophysical Research Letters* 41, 8283–8287. 2014GL061690.
- 300 Ono, S., 1999. High temperature stability limit of phase egg,  $\text{AlSiO}_3(\text{OH})$ . *Contributions to Mineralogy and Petrology* 137, 83–89.
- Ono, S., Hirose, K., Murakami, M., Isshiki, M., 2002. Post-stishovite phase boundary in  $\text{SiO}_2$  determined by in situ X-ray observations. *Earth and Planetary Science Letters* 197, 187–192.
- 305 Ono, S., Suto, T., Hirose, K., Kuwayama, Y., Komabayashi, T., Kikegawa, T., 2002. Equation of state of Al-bearing stishovite to 40 GPa at 300 K. *American Mineralogist* 87, 1486–1489.

- Panero, W.R., 2006. Aluminum incorporation in stishovite. *Geophysical Research Letters* 33, n/a–n/a. L20317.
- 310 Panero, W.R., Benedetti, L.R., Jeanloz, R., 2003. Transport of water into the lower mantle: Role of stishovite. *Journal of Geophysical Research (Solid Earth)* 108, 2039.
- Panero, W.R., Stixrude, L.P., 2004. Hydrogen incorporation in stishovite at high pressure and symmetric hydrogen bonding in  $\delta$ -AlOOH. *Earth and Planetary Science Letters* 221, 421 – 431.
- 315 Pawley, A.R., McMillan, P.F., Holloway, J.R., 1993. Hydrogen in Stishovite, with Implications for Mantle Water Content. *Science* 261, 1024–1026.
- Sano-Furukawa, A., Kagi, H., Nagai, T., Nakano, S., Fukura, S., Ushijima, D., Iizuka, R., Ohtani, E., Yagi, T., 2009. Change in compressibility of  $\delta$ -AlOOH and  $\delta$ -AlOOD at high pressure: A study of isotope effect and hydrogen-bond symmetrization. *American Mineralogist* 94, 1255–1261.
- 320 Sano-Furukawa, A., Komatsu, K., Vanpeteghem, C.B., Ohtani, E., 2008. Neutron diffraction study of  $\delta$ -AlOOD at high pressure and its implication for symmetrization of the hydrogen bond. *American Mineralogist* 93, 1558–1567.
- 325 Shieh, S.R., Duffy, T.S., Li, B., 2002. Strength and Elasticity of  $\text{SiO}_2$  across the Stishovite- $\text{CaCl}_2$ -type Structural Phase Boundary. *Physical Review Letters* 89, 255507.
- Shim, S.H., Jeanloz, R., Duffy, T.S., 2002. Tetragonal structure of  $\text{CaSiO}_3$  perovskite above 20 GPa. *Geophysical Research Letters* 29, 2166.
- 330 Singh, A.K., Andrault, D., Bouvier, P., 2012. X-ray diffraction from stishovite under nonhydrostatic compression to 70 GPa: Strength and elasticity across the tetragonal  $\rightarrow$  orthorhombic transition. *Physics of the Earth and Planetary Interiors* 208209, 1 – 10.

- Smyth, J.R., Swope, R.J., Pawley, A.R., 1995. H in rutile-type compounds;  
 335 II, Crystal chemistry of Al substitution in H-bearing stishovite. *American Mineralogist* 80, 454–456.
- Snoek, J.L., 1941. Effect of small quantities of carbon and nitrogen on the elastic and plastic properties of iron. *Physica* 8, 711–733.
- Stixrude, L., Lithgow-Bertelloni, C., 2011. Thermodynamics of mantle minerals  
 340 - II. Phase equilibria. *Geophysical Journal International* 184, 1180–1213.
- Suzuki, A., 2009. Compressibility of the high-pressure polymorph of AlOOH to 17 GPa. *Mineralogical Magazine* 73, 479–485.
- Suzuki, A., Ohtani, E., Kamada, T., 2000. A new hydrous phase  $\delta$ -AlOOH synthesized at 21 GPa and 1000 °C. *Physics and Chemistry of Minerals* 27,  
 345 689–693.
- Terasaki, H., Ohtani, E., Sakai, T., Kamada, S., Asanuma, H., Shibazaki, Y., Hirao, N., Sata, N., Ohishi, Y., Sakamaki, T., Suzuki, A., ichi Funakoshi, K., 2012. Stability of FeNi hydride after the reaction between Fe-Ni alloy and hydrous phase ( $\delta$ -AlOOH) up to 1.2 Mbar: Possibility of H contribution to  
 350 the core density deficit. *Physics of the Earth and Planetary Interiors* 194–195, 18–24.
- Tsuchida, Y., Yagi, T., 1989. A new, post-stishovite high-pressure polymorph of silica. *Nature* 340, 217–220.
- Tsuchiya, J., 2013. First principles prediction of a new high-pressure phase of  
 355 dense hydrous magnesium silicates in the lower mantle. *Geophysical Research Letters* 40, 4570–4573.
- Tsuchiya, J., Tsuchiya, T., 2009. Elastic properties of  $\delta$ -AlOOH under pressure: First principles investigation. *Physics of the Earth and Planetary Interiors* 174, 122 – 127. *Advances in High Pressure Mineral Physics: from Deep Mantle*  
 360 *to the Core*.

- Tsuchiya, J., Tsuchiya, T., 2011. First-principles prediction of a high-pressure hydrous phase of AlOOH. *Phys. Rev. B* 83, 054115.
- Umemoto, K., Kawamura, K., Hirose, K., , Revenaugh, J., Wentzcovitch, R., 2015. Post-stishovite transition in hydrous aluminous  $\text{SiO}_2$ . *PNAS* -.
- 365 Vanpeteghem, C., Sano, A., Komatsu, K., Ohtani, E., Suzuki, A., 2007. Neutron diffraction study of aluminous hydroxide  $\delta$ -AlOOD. *Physics and Chemistry of Minerals* 34, 657–661.
- Vanpeteghem, C.B., Ohtani, E., Kondo, T., 2002. Equation of state of the hydrous phase  $\delta$ -AlOOH at room temperature up to 22.5 GPa. *Geophysical*  
370 *Research Letters* 29, 1119.
- Walter, M., Thomson, A., Wang, W., Lord, O., Ross, J., McMahon, S., Baron, M., Melekhova, E., Kleppe, A., Kohn, S., 2015. The stability of hydrous silicates in Earth’s lower mantle: Experimental constraints from the systems  $\text{MgO-SiO}_2\text{-H}_2\text{O}$  and  $\text{MgO-Al}_2\text{O}_3\text{-SiO}_2\text{-H}_2\text{O}$ . *Chemical Geology* , -.
- 375 Wood, B.J., 2000. Phase transformations and partitioning relations in peridotite under lower mantle conditions. *Earth and Planetary Science Letters* 174, 341–354.
- Xue, X., Kanzaki, M., 2007. High-Pressure  $\delta$ -Al(OH)<sub>3</sub> and  $\delta$ -AlOOH Phases and Isostructural Hydroxides/Oxyhydroxides: New Structural Insights from  
380 High-Resolution  $^1\text{H}$  and  $^{27}\text{Al}$  NMR. *The Journal of Physical Chemistry B* 111, 13156–13166.