

Quenchable water-rich aluminous post-stishovite: implications for water and seismic scatterers in the lower mantle

R. Myhill, T. Boffa-Ballaran, N. Miyajima, D. J. Frost, H. Bureau, C.
Raepsaet

*Bayerisches Geoinstitut, Universität Bayreuth, Universitätsstrasse 30, 95447 Bayreuth,
Germany*

Abstract

Seismic scatterers

ULVZs by seifertite formation?

Geochemistry

Keywords: high pressure, post-stishovite, water, slab, scatterers

1. Introduction

The stishovite structure (tetragonal, $P4_2/mnm$, no. 136) undergoes a weak first-order transition (Andrault et al., 1998; Hemley et al., 2000) with second-order characteristics of Landau/ferroelastic type (Tsuchida and Yagi, 1989; Carpenter et al., 2000) to the CaCl_2 -type structure (orthorhombic, $Pnmm$, no. 58).
This transition occurs at pressures of ~ 50 GPa at room temperature (Kingma et al., 1995; Andrault et al., 1998), increasing with temperature to ~ 70 GPa at 2200 K (Hirose et al., 2005; Nomura et al., 2010). A somewhat higher dT/dP was observed by Ono et al. (2002). The nature of this transition means that the shear modulus $C_{11} - C_{12}$ decreases with increasing pressure and vanishes at the transition pressure, as observed in spectroscopic and high pressure diffraction


*Corresponding author: R. Myhill

Email address: myhill.bob@gmail.com (R. Myhill, D. J. Frost, T. Boffa-Ballaran, H. Bureau, C. Raepsaet)


studies (Kingma et al., 1995; Shieh et al., 2002). The stishovite-post-stishovite transition is therefore of interest to seismologists, as it should produce zones of low shear wave velocity which act as scatterers in the deep mantle.

15 Deviatoric stresses in the diamond anvil cell can cause large variations in the pressure of the stishovite to post-stishovite transition. Nonhydrostaticity greatly decreases the transition pressure from 60 GPa (Andrault et al., 2003; Hemley et al., 2000) to 40 GPa (Kingma et al., 1996; Singh et al., 2012). Molecular and lattice dynamic studies suggest that only 1.5–2.5 GPa differential stress is
20 required to cause this decrease in pressure (Dubrovinsky and Belonoshko, 1996).

Aluminium has a profound effect on the stability of the post-stishovite phase. Al_2O_3 can be incorporated into stishovite by the substitution of Si by Al with the formation of oxygen vacancies (Smyth et al., 1995; Hirose et al., 2005; Bromiley et al., 2006; Lakshtanov et al., 2007a):




25 An alternative mechanism, where charge balance is accomplished by Al occupying the large interstitial sites is probably minor Smyth et al. (1995):



It has been argued that the addition of 4 wt% Al_2O_3 in the absence of any other components causes the transition to shift from 50 to 23 GPa at room temperature by inducing a ‘chemical pressure’ (Bolfan-Casanova et al., 2009). In
30 stishovite, there is one octahedral cation position (Wyckoff notation 2a) located at the origin and the body center of the tetragonal cell. One oxygen at position 4f creates a moderately distorted octahedron with point symmetry mmm. The addition of Al *decreases* distortion of this octahedron (Smyth et al., 1995). The addition of Al should therefore have little effect on the stishovite-post-stishovite
35 transition (Panero, 2006). The results of Bolfan-Casanova et al. (2009) are therefore unexpected; it seems possible that another component (such as H_2O) or deviatoric stress acted to stabilise the CaCl_2 structure.

It has also been suggested that the coupled substitution of Al and H may have a significant effect on the transition pressure. Lakshtanov et al. (2007b) have shown that the addition of 6 wt% Al_2O_3 and 0.24 wt% H_2O reduces the transition pressure to 24 GPa at room temperature. The substitution mechanism in this case is probably



Given that high pressure δ -AlOOH is isostructural with the post-stishovite phase above ca. 19 GPa (Sano-Furukawa et al., 2008; Kuribayashi et al., 2014), after transforming from the $P2_1nm$ structure (no. 31) (Suzuki et al., 2000; Komatsu et al., 2006; Vanpeteghem et al., 2007), it seems sensible to suggest that addition of an AlOOH component lowers the transition pressure. Ab initio calculations support this suggestion (Umemoto et al., 2015). The equilibrium transition for 6.25 mol% AlOOH is at ca. 15 GPa at room temperature, implying an even more marked reduction than observed in the experimental data (Lakshtanov et al., 2007b). This could be due to inaccuracies in the ab-initio data, or metastable preservation of the tetragonal phase in the experiments. Umemoto et al. (2015) also suggest that the larger number of hydrogen sites stabilises the tetragonal structure at high temperatures, changing the slope of the transition. The tetragonal \rightarrow orthorhombic transition in their simulations is associated with splitting of the hydrogen sites on the equatorial oxygens into two groups, one occupied and the other unoccupied. Additionally, the redistribution of hydrogens among equatorial oxygens in the tetragonal \rightarrow orthorhombic transition implies that it is now first order and no longer ferroelastic. Anelasticity by hydrogen hopping seems a plausible alternative to reduce seismic body wave speeds.

Despite this, H_2O contents in tetragonal stishovite measured using the FTIR calibration of Pawley et al. (1993) are less than 20% of that expected from an AlOOH component (Panero et al., 2003; Bromiley et al., 2006; Litasov et al., 2007) and in MORB compositions (Chung and Kagi, 2002).

This study was designed to investigate water solubility in Al-rich stishovite by synthesising crystals at relatively high temperature (Ono, 1999). In doing so, we created and quenched an Al-H rich post-stishovite phase.

1.1. *Hysteresis vs. stabilisation*

70 (Umemoto et al., 2015)

1.2. *Mechanism of shear wave velocity reduction*

Ferroelasticity (Carpenter et al., 2000) Snoek relaxation (anelasticity via H mobility) (Snoek, 1941; Nowick and Berry, 1972; McKnight et al., 2007).

1.3. *Mechanisms of Al-, H incorporation*

75 We found that hydrogen was most stable when bonded to the apical oxygen of the Al-octahedron, with the hydroxyl bond along $\langle 110 \rangle$ and co-planar with Al (Panero and Stixrude, 2004).

The relatively short O-O distance and correspondingly long O-H distance in the stishovite-AlOOH solid solution would lead one to expect a low OH stretching frequency compared to other nominally anhydrous minerals, consistent with
80 the value of 3111 cm^{-1} observed by Pawley et al. (1993).

Symmetric bonding (Panero and Stixrude, 2004)

1.4. *Water concentrations in stishovite*

Pawley et al. (1993) calibration Panero et al. (2003) Bromiley et al. (2006),
85 Litasov et al. (2007), MORB Chung and Kagi (2002),

1.5. *Elastic properties*

Andrault et al. (2003) Lakshtanov et al. (2005) Ono et al. (2002) Sano-Furukawa et al. (2009)

1.6. *Phase relations in the lower mantle*

90 Irifune and Ringwood (1993) Wood (2000) Hirose (2002) Walter et al. (2015) Litasov and Ohtani (2005)

1.7. Datasets and modelling

Holland and Powell (2011) Stixrude and Lithgow-Bertelloni (2011) Cottaar et al. (2014)

95 1.7.1. Elastic properties $AlOOH$

Suzuki (2009) (Sano-Furukawa et al., 2008) Vanpeteghem et al. (2002) Tsuchiya and Tsuchiya (2009) Li et al. (2006)

2. The solid solution $(MgSi, Fe^{2+}Si, Fe_2^{3+}, Al_2)O_4H_2$

2.1. Endmembers

100 2.1.1. SiO_2

2.1.2. $\delta-AlOOH$

2.1.3. $\delta-FeOOH$

Pnma (no.62 α) to $P2_1nm$ (no. 31, ε) (Gleason et al., 2008) to $Pnnm$ (no. 58, high spin) to $Pnnm$ (no. 58, low spin) (Gleason et al., 2013)

105 2.1.4. “Phase H”

The addition of an $MgSiO_4H_2$ component also stabilises the $Pnnm$ structure in stishovite (Komatsu et al., 2011). A previous candidate space group ($Pnn2$; no. 34 Kudoh et al., 2004) is now thought to be unlikely.

Stability (Ohtani et al., 2014) Ab initio $P2/m$ (no. 10) (Tsuchiya, 2013)
110 $Pnnm$ (no. 58) (Bindi et al., 2014) $P2_1nm$ (no. 31) (Nishi et al., 2014)

Kudoh et al. (2004) suggested a hypothetical high pressure form of $Mg(OH)_2$ also in the space group $Pnnm$ from Mg incorporation. Importance of the $AlOOH$ - phase H solid solution Ohira et al. (2014)

3. Other phases

115 Ohtani (2015)

3.1. Water in seifertite

Post-stishovite undergoes a phase transition to the mineral seifertite with the scrutinyite (α -PbO₂) structure (orthorhombic, Pbcn, no. 60 or Pb2n) at about 120-140 GPa and 2000-2500 K (Murakami et al., 2003; Grocholski et al., 2013). The addition of aluminium under dry conditions may stabilise seifertite to 110 GPa at 2000 K (Hirose et al., 2005), although this remains controversial (Grocholski et al., 2013). The effect of hydrogen on the structure is currently unknown. As there are no hydrous phases known with the seifertite structure, the addition of water probably destabilises seifertite relative to post-stishovite. This supposition is supported by the lack of water in the isostructural TiO₂ (II) phase, which contains negligible water (Bromiley et al., 2004).

δ -AlOOH remains stable throughout the lower mantle, and probably transforms to a cubic structure (Pa $\bar{3}$) at core pressures (Tsuchiya and Tsuchiya, 2011).

3.2. δ -Al(OH)₃

P2₁2₁2₁ (no. 19) to Pnma (no. 62) at \sim 67 GPa Matsui et al. (2011). Xue and Kanzaki (2007)

3.3. Ca-perovskite

Tetragonal (probably P4/mmm, no. 123) (Shim et al., 2002) \rightarrow Cubic (Pm3m) for low Al contents. For pure CaSiO₃, transition occurs at ca. 580 K at 50 GPa (Kurashina et al., 2004; Komabayashi et al., 2007)

Orthorhombic (Pbnm) \rightarrow Cubic (Pm3m) for Al-rich CaSiO₃ (Kurashina et al., 2004). At 5.9 wt%, transition occurs at ca. 1840 K at 50 GPa (Kurashina et al., 2004).

In a pyrolitic mantle composition, the CaSiO₃ perovskite contains 1.0–2.3 wt.% Al₂O₃ at upper mantle pressures and somewhat less (0.7–1.6 wt.%) at lower mantle pressures (Kesson et al., 1998; Wood, 2000; Hirose, 2002)

In a MORB composition, 2.0–4.8 wt.% and 1.2–4.5 wt.% Al₂O₃ are included in CaSiO₃-rich perovskite in garnetite and perovskitite lithologies, respectively (Kesson et al., 1994; Irifune and Ringwood, 1993; Hirose and Fei, 2002).

3.4. *Reaction with metal*

(Terasaki et al., 2012)

4. Lower mantle scatterers

Deuss et al. (2013) Kaneshima and Helffrich (1998, 2003) Niu et al. (2003)
150 Krüger et al. (2001) Kaneshima (2009) Bina et al. (2010) Kaneshima et al.
(2010) Bentham and Rost (2014) Asahara et al. (2013) Mookherjee (2011)

5. Experimental and analytical techniques

6. Chemical composition

Figure 1: Single crystal XRD spectra of post-stishovite

7. Conclusions

155 **References**

- Andrault, D., Angel, R.J., Mosenfelder, J.L., Le Bihan, T., 2003. Equation of state of stishovite to lower mantle pressures. *American Mineralogist* 88, 301–307.
- Andrault, D., Angel, R.J., Mosenfelder, J.L., Le Bihan, T., 2003. Equation
160 of state of stishovite to lower mantle pressures. *American Mineralogist* 88, 301–307.
- Andrault, D., Fiquet, G., Guyot, F., Hanfland, M., 1998. Pressure-Induced Landau-Type Transition in Stishovite. *Science* 282, 720.
- Asahara, Y., Hirose, K., Ohishi, Y., Hirao, N., Ozawa, H., Murakami, M.,
165 2013. Acoustic velocity measurements for stishovite across the post-stishovite phase transition under deviatoric stress: Implications for the seismic features of subducting slabs in the mid-mantle. *American Mineralogist* 98, 2053–2062.
- Benthams, H.L.M., Rost, S., 2014. Scattering beneath Western Pacific subduction zones: evidence for oceanic crust in the mid-mantle. *Geophysical Journal International* 197, 1627–1641.
170
- Bina, C.R., Suetsugu, D., Bina, C., Inoue, T., Wiens, D., Jellinek, M., 2010. Scale limits of free-silica seismic scatterers in the lower mantle. *Physics of the Earth and Planetary Interiors* 183, 110–114.
- Bindi, L., Nishi, M., Tsuchiya, J., Irifune, T., 2014. Crystal chemistry of dense
175 hydrous magnesium silicates: The structure of phase H, MgSiH_2O_4 , synthesized at 45 GPa and 1000 °C. *American Mineralogist* 99, 1802–1805.
- Bolfan-Casanova, N., Andrault, D., Amiguet, E., Guignot, N., 2009. Equation of state and post-stishovite transformation of Al-bearing silica up to 100 GPa and 3000 K. *Physics of the Earth and Planetary Interiors* 174, 70–77.
- 180 Bromiley, G., Hilaret, N., McCammon, C., 2004. Solubility of hydrogen and ferric iron in rutile and TiO_2 (II): Implications for phase assemblages during

- ultrahigh-pressure metamorphism and for the stability of silica polymorphs in the lower mantle. *Geophysical Research Letters* 31.
- Bromiley, G.D., Bromiley, F.A., Bromiley, D.W., 2006. On the mechanisms for
185 H and Al incorporation in stishovite. *Physics and Chemistry of Minerals* 33,
613–621.
- Carpenter, M.A., Hemley, R.J., Mao, H.K., 2000. High-pressure elasticity of stishovite and the $P4_2/mnm \rightarrow Pnnm$ phase transition. *Journal of Geophysical Research* 105, 10807.
- 190 Chung, J.I., Kagi, H., 2002. High concentration of water in stishovite in the MORB system. *Geophysical Research Letters* 29, 16–1–16–4. 2020.
- Cottaar, S., Heister, T., Rose, I., Unterborn, C., 2014. BurnMan: A lower mantle mineral physics toolkit. *Geochemistry, Geophysics, Geosystems* 15, 1164–1179.
- 195 Deuss, A., Andrews, J., Day, E., 2013. Seismic Observations of Mantle Discontinuities and Their Mineralogical and Dynamical Interpretation. pp. 295–323.
- Dubrovinsky, L.S., Belonoshko, A.B., 1996. Pressure-induced phase transition and structural changes under deviatoric stress of stishovite to CaCl_2 -like structure. *Geochimica et Cosmochimica Acta* 60, 3657–3663.
- 200 Gleason, A., Jeanloz, R., Kunz, M., 2008. Pressure-temperature stability studies of feooh using x-ray diffraction. *American Mineralogist* 93, 1882–1885.
- Gleason, A., Quiroga, C., Suzuki, A., Pentcheva, R., Mao, W., 2013. Symmetrization driven spin transition in ε -FeOOH at high pressure. *Earth and Planetary Science Letters* 379, 49 – 55.
- 205 Grocholski, B., Shim, S.H., Prakapenka, V.B., 2013. Stability, metastability, and elastic properties of a dense silica polymorph, seifertite. *Journal of Geophysical Research: Solid Earth* 118, 4745–4757.

- Hemley, R.J., Shu, J., Carpenter, M.A., Hu, J., Mao, H.K., Kingma, K.J., 2000. Strain/order parameter coupling in the ferroelastic transition in dense SiO₂. Solid State Communications 114, 527–532.
- 210 Hirose, K., 2002. Phase transitions in pyrolitic mantle around 670-km depth: Implications for upwelling of plumes from the lower mantle. Journal of Geophysical Research (Solid Earth) 107, 2078.
- Hirose, K., Fei, Y., 2002. Subsolvus and melting phase relations of basaltic composition in the uppermost lower mantle. Geochimica et Cosmochimica Acta 66, 2099–2108.
- 215 Hirose, K., Takafuji, N., Sata, N., Ohishi, Y., 2005. Phase transition and density of subducted MORB crust in the lower mantle. Earth and Planetary Science Letters 237, 239–251.
- 220 Holland, T.J.B., Powell, R., 2011. An improved and extended internally consistent thermodynamic dataset for phases of petrological interest, involving a new equation of state for solids. Journal of Metamorphic Geology 29, 333–383.
- Irifune, T., Ringwood, A.E., 1993. Phase transformations in subducted oceanic crust and buoyancy relationships at depths of 600–800 km in the mantle. Earth and Planetary Science Letters 117, 101–110.
- 225 Kaneshima, S., 2009. Seismic scatterers at the shallowest lower mantle beneath subducted slabs. Earth and Planetary Science Letters 286, 304–315.
- Kaneshima, S., Helffrich, G., 1998. Detection of lower mantle scatterers north-east of the Mariana subduction zone using short-period array data. Journal of Geophysical Research 103, 4825–4838.
- 230 Kaneshima, S., Helffrich, G., 2003. Subparallel dipping heterogeneities in the mid-lower mantle. Journal of Geophysical Research (Solid Earth) 108, 2272.
- Kaneshima, S., Helffrich, G., Suetsugu, D., Bina, C., Inoue, T., Wiens, D., Jellinek, M., 2010. Small scale heterogeneity in the mid-lower mantle beneath

- 235 the circum-Pacific area. *Physics of the Earth and Planetary Interiors* 183, 91–103.
- Kesson, S.E., Fitz Gerald, J.D., Shelley, J.M., 1998. Mineralogy and dynamics of a pyrolite lower mantle. *Nature* 393, 252–255.
- Kesson, S.E., Fitz Gerald, J.D., Shelley, J.M.G., 1994. Mineral chemistry and
240 density of subducted basaltic crust at lower-mantle pressures. *Nature* 372, 767–769.
- Kingma, K., Mao, H.K., Hemley, R., 1996. Synchrotron X-ray diffraction of SiO_2 to multimegabar pressures. *High Pressure Research* 14, 363–374.
- Kingma, K.J., Cohen, R.E., Hemley, R.J., Mao, H.K., 1995. Transformation of
245 stishovite to a denser phase at lower-mantle pressures. *Nature* 374, 243–245.
- Komabayashi, T., Hirose, K., Sata, N., Ohishi, Y., Dubrovinsky, L.S., 2007. Phase transition in CaSiO_3 perovskite. *Earth and Planetary Science Letters* 260, 564 – 569.
- Komatsu, K., Kuribayashi, T., Sano, A., Ohtani, E., Kudoh, Y., 2006. Redetermination of the high-pressure modification of AlOOH from single-crystal
250 synchrotron data. *Acta Crystallographica Section E* 62, 216–218.
- Komatsu, K., Sano-Furukawa, A., Kagi, H., 2011. Effects of Mg and Si ions on the symmetry of $\delta\text{-AlOOH}$. *Physics and Chemistry of Minerals* 38, 727–733.
- Krüger, F., Baumann, M., Scherbaum, F., Weber, M., 2001. Mid mantle scatterers near the Mariana Slab detected with a double array method. *Geophysical
255 Research Letters* 28, 667–670.
- Kudoh, Y., Kuribayashi, T., Suzuki, A., Ohtani, E., Kamada, T., 2004. Space group and hydrogen sites of $\delta\text{-AlOOH}$ and implications for a hypothetical high-pressure form of $\text{Mg}(\text{OH})_2$. *Physics and Chemistry of Minerals* 31, 360–
260 364.

- Kurashina, T., Hirose, K., Ono, S., Sata, N., Ohishi, Y., 2004. Phase transition in Al-bearing CaSiO_3 perovskite: implications for seismic discontinuities in the lower mantle. *Physics of the Earth and Planetary Interiors* 145, 67–74.
- Kuribayashi, T., Sano-Furukawa, A., Nagase, T., 2014. Observation of pressure-induced phase transition of $\delta\text{-AlOOH}$ by using single-crystal synchrotron X-ray diffraction method. *Physics and Chemistry of Minerals* 41, 303–312.
- Lakshtanov, D.L., Litasov, K.D., Sinogeikin, S.V., Hellwig, H., Li, J., Ohtani, E., Bass, J.D., 2007a. Effect of Al^{3+} and H^+ on the elastic properties of stishovite. *American Mineralogist* 92, 1026–1030.
- Lakshtanov, D.L., Sinogeikin, S.V., Litasov, K.D., Prakapenka, V.B., Hellwig, H., Wang, J., Sanches-Valle, C., Perrillat, J.P., Chen, B., Somayazulu, M., Li, J., Ohtani, E., Bass, J.D., 2007b. The post-stishovite phase transition in hydrous alumina-bearing SiO_2 in the lower mantle of the Earth. *Proceedings of the National Academy of Sciences* 104, 13588–13590.
- Lakshtanov, D.L., Vanpeteghem, C.B., Jackson, J.M., Bass, J.D., Shen, G., Prakapenka, V.B., Litasov, K., Ohtani, E., 2005. The equation of state of Al,H-bearing SiO_2 stishovite to 58 GPa. *Physics and Chemistry of Minerals* 32, 466–470.
- Li, S., Ahuja, R., Johansson, B., 2006. The elastic and optical properties of the high-pressure hydrous phase $\delta\text{-AlOOH}$. *Solid State Communications* 137, 101–106.
- Litasov, K.D., Kagi, H., Shatskiy, A., Ohtani, E., Lakshtanov, D.L., Bass, J.D., Ito, E., 2007. High hydrogen solubility in Al-rich stishovite and water transport in the lower mantle. *Earth and Planetary Science Letters* 262, 620 – 634.
- Litasov, K.D., Ohtani, E., 2005. Phase relations in hydrous MORB at 18–28 GPa: implications for heterogeneity of the lower mantle. *Physics of the Earth and Planetary Interiors* 150, 239–263.

- 290 Matsui, M., Komatsu, K., Ikeda, E., Sano-Furukawa, A., Gotou, H., Yagi, T.,
2011. The crystal structure of δ -Al(OH)₃: Neutron diffraction measurements
and ab initio calculations. *American Mineralogist* 96, 854–859.
- McKnight, R.E., Carpenter, M.A., Darling, T.W., Buckley, A., Taylor, P.A.,
2007. Acoustic dissipation associated with phase transitions in lawsonite,
CaAl₂Si₂O₇(OH)₂H₂O. *American Mineralogist* 92, 1665–1672.
- 295 Mookherjee, M., 2011. Mid-mantle anisotropy: Elasticity of aluminous phases
in subducted MORB. *Geophysical Research Letters* 38, n/a–n/a. L14302.
- Murakami, M., Hirose, K., Ono, S., Ohishi, Y., 2003. Stability of CaCl₂-type
and α -PbO₂-type SiO₂ at high pressure and temperature determined by in-
situ X-ray measurements. *Geophysical Research Letters* 30. 1207.
- 300 Nishi, M., Irifune, T., Tsuchiya, J., Tange, Y., Nishihara, Y., Fujino, K., Higo,
Y., 2014. Stability of hydrous silicate at high pressures and water transport
to the deep lower mantle. *Nature Geoscience* 7, 224–227.
- Niu, F., Kawakatsu, H., Fukao, Y., 2003. Seismic evidence for a chemical het-
erogeneity in the midmantle: A strong and slightly dipping seismic reflector
305 beneath the Mariana subduction zone. *Journal of Geophysical Research (Solid
Earth)* 108, 2419.
- Nomura, R., Hirose, K., Sata, N., Ohishi, Y., Suetsugu, D., Bina, C., Inoue,
T., Wiens, D., Jellinek, M., 2010. Precise determination of post-stishovite
phase transition boundary and implications for seismic heterogeneities in the
310 mid-lower mantle. *Physics of the Earth and Planetary Interiors* 183, 104–109.
- Nowick, A., Berry, D., 1972. Anelastic relaxation in crystalline solids, acad.
Press, New York .
- Ohira, I., Ohtani, E., Sakai, T., Miyahara, M., Hirao, N., Ohishi, Y., Nishijima,
M., 2014. Stability of a hydrous δ -phase, AlOOH-MgSiO₂(OH)₂, and a mech-
315 anism for water transport into the base of lower mantle. *Earth and Planetary
Science Letters* 401, 12–17.

- Ohtani, E., 2015. Hydrous minerals and the storage of water in the deep mantle. *Chemical Geology* , –.
- Ohtani, E., Amaike, Y., Kamada, S., Sakamaki, T., Hirao, N., 2014. Stability
 320 of hydrous phase H MgSiO_4H_2 under lower mantle conditions. *Geophysical Research Letters* 41, 8283–8287. 2014GL061690.
- Ono, S., 1999. High temperature stability limit of phase egg, $\text{AlSiO}_3(\text{OH})$. *Contributions to Mineralogy and Petrology* 137, 83–89.
- Ono, S., Hirose, K., Murakami, M., Isshiki, M., 2002. Post-stishovite phase
 325 boundary in SiO_2 determined by in situ X-ray observations. *Earth and Planetary Science Letters* 197, 187–192.
- Ono, S., Suto, T., Hirose, K., Kuwayama, Y., Komabayashi, T., Kikegawa, T., 2002. Equation of state of Al-bearing stishovite to 40 GPa at 300 K. *American Mineralogist* 87, 1486–1489.
- 330 Panero, W.R., 2006. Aluminum incorporation in stishovite. *Geophysical Research Letters* 33, n/a–n/a. L20317.
- Panero, W.R., Benedetti, L.R., Jeanloz, R., 2003. Transport of water into the lower mantle: Role of stishovite. *Journal of Geophysical Research (Solid Earth)* 108, 2039.
- 335 Panero, W.R., Stixrude, L.P., 2004. Hydrogen incorporation in stishovite at high pressure and symmetric hydrogen bonding in $\delta\text{-AlOOH}$. *Earth and Planetary Science Letters* 221, 421 – 431.
- Pawley, A.R., McMillan, P.F., Holloway, J.R., 1993. Hydrogen in Stishovite, with Implications for Mantle Water Content. *Science* 261, 1024–1026.
- 340 Sano-Furukawa, A., Kagi, H., Nagai, T., Nakano, S., Fukura, S., Ushijima, D., Iizuka, R., Ohtani, E., Yagi, T., 2009. Change in compressibility of $\delta\text{-AlOOH}$ and $\delta\text{-AlOOD}$ at high pressure: A study of isotope effect and hydrogen-bond symmetrization. *American Mineralogist* 94, 1255–1261.

- Sano-Furukawa, A., Komatsu, K., Vanpeteghem, C.B., Ohtani, E., 2008. Neu-
345 tron diffraction study of δ -AlOOD at high pressure and its implication for
symmetrization of the hydrogen bond. *American Mineralogist* 93, 1558–1567.
- Shieh, S.R., Duffy, T.S., Li, B., 2002. Strength and Elasticity of SiO_2 across the
Stishovite- CaCl_2 -type Structural Phase Boundary. *Physical Review Letters*
89, 255507.
- 350 Shim, S.H., Jeanloz, R., Duffy, T.S., 2002. Tetragonal structure of CaSiO_3
perovskite above 20 GPa. *Geophysical Research Letters* 29, 2166.
- Singh, A.K., Andrault, D., Bouvier, P., 2012. X-ray diffraction from stishovite
under nonhydrostatic compression to 70 GPa: Strength and elasticity across
the tetragonal \rightarrow orthorhombic transition. *Physics of the Earth and Planetary*
355 *Interiors* 208209, 1 – 10.
- Smyth, J.R., Swope, R.J., Pawley, A.R., 1995. H in rutile-type compounds;
II, Crystal chemistry of Al substitution in H-bearing stishovite. *American*
Mineralogist 80, 454–456.
- Snoek, J.L., 1941. Effect of small quantities of carbon and nitrogen on the
360 elastic and plastic properties of iron. *Physica* 8, 711–733.
- Stixrude, L., Lithgow-Bertelloni, C., 2011. Thermodynamics of mantle minerals
- II. Phase equilibria. *Geophysical Journal International* 184, 1180–1213.
- Suzuki, A., 2009. Compressibility of the high-pressure polymorph of AlOOH to
17 GPa. *Mineralogical Magazine* 73, 479–485.
- 365 Suzuki, A., Ohtani, E., Kamada, T., 2000. A new hydrous phase δ - AlOOH
synthesized at 21 GPa and 1000 °C. *Physics and Chemistry of Minerals* 27,
689–693.
- Terasaki, H., Ohtani, E., Sakai, T., Kamada, S., Asanuma, H., Shibazaki, Y.,
Hirao, N., Sata, N., Ohishi, Y., Sakamaki, T., Suzuki, A., ichi Funakoshi, K.,
370 2012. Stability of FeNi hydride after the reaction between Fe-Ni alloy and

- hydrous phase (δ -AlOOH) up to 1.2 Mbar: Possibility of H contribution to the core density deficit. *Physics of the Earth and Planetary Interiors* 194–195, 18–24.
- 375 Tsuchida, Y., Yagi, T., 1989. A new, post-stishovite high-pressure polymorph of silica. *Nature* 340, 217–220.
- Tsuchiya, J., 2013. First principles prediction of a new high-pressure phase of dense hydrous magnesium silicates in the lower mantle. *Geophysical Research Letters* 40, 4570–4573.
- 380 Tsuchiya, J., Tsuchiya, T., 2009. Elastic properties of δ -AlOOH under pressure: First principles investigation. *Physics of the Earth and Planetary Interiors* 174, 122 – 127. *Advances in High Pressure Mineral Physics: from Deep Mantle to the Core*.
- Tsuchiya, J., Tsuchiya, T., 2011. First-principles prediction of a high-pressure hydrous phase of AlOOH. *Phys. Rev. B* 83, 054115.
- 385 Umemoto, K., Kawamura, K., Hirose, K., , Revenaugh, J., Wentzcovitch, R., 2015. Post-stishovite transition in hydrous aluminous SiO_2 . *PNAS* -.
- Vanpeteghem, C., Sano, A., Komatsu, K., Ohtani, E., Suzuki, A., 2007. Neutron diffraction study of aluminous hydroxide δ -AlOOD. *Physics and Chemistry of Minerals* 34, 657–661.
- 390 Vanpeteghem, C.B., Ohtani, E., Kondo, T., 2002. Equation of state of the hydrous phase δ -AlOOH at room temperature up to 22.5 GPa. *Geophysical Research Letters* 29, 1119.
- Walter, M., Thomson, A., Wang, W., Lord, O., Ross, J., McMahon, S., Baron, M., Melekhova, E., Kleppe, A., Kohn, S., 2015. The stability of hydrous silicates in Earth’s lower mantle: Experimental constraints from the systems
395 $\text{MgO-SiO}_2\text{-H}_2\text{O}$ and $\text{MgO-Al}_2\text{O}_3\text{-SiO}_2\text{-H}_2\text{O}$. *Chemical Geology* , -.

- Wood, B.J., 2000. Phase transformations and partitioning relations in peridotite under lower mantle conditions. *Earth and Planetary Science Letters* 174, 341–354.
- 400 Xue, X., Kanzaki, M., 2007. High-Pressure δ -Al(OH)₃ and δ -AlOOH Phases and Isostructural Hydroxides/Oxyhydroxides: New Structural Insights from High-Resolution ¹H and ²⁷Al NMR. *The Journal of Physical Chemistry B* 111, 13156–13166.