
WEEK 8 / PART II / LINEAR PROGRAMMING

Which plan is best?

Keep the allowable choices fixed, state a goal, and compare the consequences. A mathematical optimum answers the question you actually put into the model.



From geometry to a decision

The bike shop's feasible region has corners $(0,0), (6,0), (4,4), (0,8)$. With after-parts contribution $P = 30x + 20y$, scheduling four tune-ups and four quick repairs earns the largest total of \$200. A different goal can favor a different schedule or leave an entire edge tied.

A feasible point represents an allowable plan. A linear objective assigns a score to each plan; examining the feasible region identifies the highest or lowest attainable score, including ties and possible unboundedness.

For the core method you need only substitution and comparison. Equal-score lines and convex mixtures explain why the method works. The whole-number investigation is an extension.

Online reading · Optimizer · Optimization extension

8.6 An objective gives each plan a score

Suppose the shop earns $30 after replacement parts for each tune-up and $20 for each quick repair. The total contribution for a day is

$$P = 30x + 20y.$$

This is the *objective function*: a rule assigning a number to each plan. The letter P names the total contribution in dollars. To *maximize* it means to find the largest value among feasible plans.

Example · Feasible does not mean optimal

Three tune-ups and three quick repairs give $P = 30(3) + 20(3) = 150$ dollars. Four of each give $P = 30(4) + 20(4) = 200$ dollars. Both are possible schedules, but the second earns more.
Five tune-ups and three quick repairs would bring in $210. A larger earnings figure cannot make an impossible schedule workable: it requires more mechanic time than the shop has.

A function such as $30x + 20y$ is *linear*: every additional tune-up adds the same $30 and every quick repair adds the same $20. There is no product xy, squared quantity, or changing rate. A *linear program* combines a linear objective with linear constraints. Here "programming" means planning; writing computer code is not required.

Pause and try 1

At (2,5), what is P=30x+20y? Does knowing the money earned tell you whether the schedule is feasible?

8.7 For this bounded polygon, compare the corners

Our continuous feasible region contains infinitely many points. The *corner-point principle* says that a linear objective on a nonempty, closed, bounded polygonal feasible region attains its maximum and minimum at vertices. "Bounded" means it fits inside some finite box. "Closed" here means the boundary is included.


This guarantees that at least one vertex gives the best value. It does not guarantee a unique best point, or say that every best point must be a vertex. Those distinctions matter when there is a tie.

Feasible vertex	Calculation of P = 30x + 20y	Contribution
(0,0)	30(0)+20(0)	$0
(6,0)	30(6)+20(0)	$180
(4,4)	30(4)+20(4)	$200
(0,8)	30(0)+20(8)	$160

The largest contribution is $200 at (4,4). The shop should schedule four tune-ups and four quick repairs. Report both the schedule and the money earned; they answer different questions.

Why a corner comparison can work

Any point in a filled convex polygon can be expressed as a mixture of its vertices. A linear objective at that point is the same mixture of the vertex scores. An average of scores cannot exceed the largest score or fall below the smallest. This is the geometric reason the table can locate an optimum.



Each dashed line joins plans with an equal objective value. Moving to a parallel line changes that score, not the resource constraints. The star marks the last feasible contact in the direction of increasing score.

In our model, the best plan is also a whole-number plan, so it remains optimal if we restrict production to whole bicycles. A continuous optimum is an upper bound for the integer choices; when a whole-number plan attains it, the integer problem is solved too.


Pause and try 2

Why is checking just the two axis corners insufficient?

8.8 The feasible region stays; the goal can change

Suppose a new service contract favors quick repairs. To see how a change in priorities changes the answer, use the alternative score $R = x + 4y$. At the same four vertices its values are 0, 6, 20, and 32. Now (0,8) is best. The resource limits and region did not change. The objective changed.

Alternatively, suppose the shop wants to return the largest *number of bikes* to customers, regardless of type. Maximize $T = x + y$. The eight-bike limit already tells us $T \leq 8$. Both (4,4) and (0,8) attain 8, and every point on the segment between them also attains 8.



These are multiple optimal solutions. If actual bicycles must be whole, the five plans (0,8), (1,7), (2,6), (3,5), and (4,4) all attain the same largest total.

Example · A tie leaves room for another consideration

If the goal is to help as many customers as possible, (4,4) and (0,8) both serve eight bikes. But the first needs twelve mechanic-hours and the second needs only eight. To save mechanic time, choose (0,8); to earn more after parts, choose (4,4).

A second goal can distinguish plans that tie under the first. We should name that goal rather than pretend the first objective made the whole decision.

Pause and try 3


If two adjacent feasible vertices both give the maximum score for a linear objective, what happens between them? Does the same conclusion follow merely because two vertices have equal scores?

8.9 “At least” often reverses the feasible side

A neighborhood group is organizing a back-to-school supply drive. A store offers two bundles. Bundle A contains 1 notebook and 2 pencils and costs $4. Bundle B contains 1 notebook and 1 pencil and costs $3. The group needs at least six notebooks and eight pencils for the students it is helping. Let x and y be the respective bundle counts.

$$x + y \geq 6, \qquad 2x + y \geq 8, \qquad x, y \geq 0; \qquad \text{minimize } C = 4x + 3y.$$

We can buy extra supplies, so there is no upper limit in this model. The feasible region extends up and to the right. It is *unbounded*, but that does not mean its minimum cost is missing.



The top and right edges of the picture are viewing limits, not extra constraints. The arrow shows the feasible region continuing. Buying more and more bundles raises this positive cost; it does not lower it indefinitely.

Boundary vertex	Notebooks	Pencils	Cost 4x+3y
(0,8)	8	8	$24
(2,4)	6	8	$20
(6,0)	6	12	$24


The group can meet both requirements for $20 with two A-bundles and four B-bundles. For this nonnegative-variable model, the least cost occurs at a feasible vertex. On an unbounded region, we must also check whether some allowed direction keeps improving the objective. Here all costs are positive, so going farther out cannot make the cost decrease forever.

Buying nothing costs zero but would leave students without the supplies. This is why minimizing cost cannot be separated from stating what must be provided. If the only restrictions were upper limits and nonnegativity, doing nothing could indeed be the mathematical minimum-cost plan.

Pause and try 4

Why is (0,6) not feasible for the supply order even though it provides all six notebooks?

8.10 A model can have a best plan, no plan, or no finite best value



Separate the questions: Is any plan feasible? Is the region bounded? Can the objective keep improving without limit? An unbounded region can still have a finite minimum, as the supply-order example showed.

- *Infeasible model:* no point satisfies every constraint. For example, $x + y \leq 4$ and $x + y \geq 6$ cannot both hold. There is no permissible plan to optimize.
- *Unbounded objective:* feasible plans exist, but their scores improve without limit. For example, maximizing $x + y$ with only $x + y \geq 6$ and nonnegativity allows arbitrarily large totals.
- *Finite optimum:* some feasible plan attains the best value. It may be one point, an edge, or a larger set when scores tie.

The optimizer identifies these different outcomes. It computes vertices from the actual constraints; a corner of its graph window is not automatically a real feasible vertex.

Included boundaries matter too

With $0 \leq x < 1$, there is no largest allowable x: any proposed x can be increased a little while staying below 1. The value 1 is a limiting bound, but it is excluded. Our corner method uses non-strict inequalities so the relevant boundaries are available.

Pause and try 5


Does an unbounded feasible region always imply that there is no minimum? Give an example from this reading.

Extension: whole-number plans

Acres, mixture amounts, and hours can often be treated as divisible. Chairs, buses, and repaired bicycles usually cannot. Requiring whole-number decisions adds an *integer constraint*. It changes which points are allowed, even when the drawn continuous region stays the same.

Example · Rounding can fail in two ways

Maximize $S = 5x + 8y$ subject to $x + 2y \leq 8$, $3x + 2y \leq 10$, and nonnegativity. The continuous best point is (1,3.5), with score $5(1) + 8(3.5) = 33$.
Rounding up to (1,4) violates both limits: the totals become 9 and 11. Rounding down to (1,3) is feasible but scores only 29. The whole-number plan (0,4) is feasible and scores 32, so rounding down missed a better plan.



For a small model, list feasible whole-number choices systematically. A fractional continuous optimum bounds the possible whole-number score from above, but it may not itself be usable.

To verify the integer optimum here, try y = 0, 1, 2, 3, 4. The largest allowable whole x-values are 3, 2, 2, 1, 0, giving respective scores 15, 18, 26, 29, 32. This finite check establishes the best whole plan.

Pause and try 6


What does the continuous maximum 33 establish about any feasible whole-number score in this example? Why does it not establish that 33 can be achieved with whole numbers?

8.11 A convex combination is a weighted average of points

Weeks 2–3 used fractions, relative weights, and averages to summarize numbers. We can average points by averaging their coordinates. For points $A = (2, 1)$ and $B = (8, 5)$, use weights 0.75 and 0.25:

$$P = 0.75A + 0.25B = (0.75(2) + 0.25(8),\ 0.75(1) + 0.25(5)) = (3.5, 2).$$

A *convex combination* uses nonnegative weights that add to 1. For two points, all these combinations fill the segment between them. For three noncollinear points, they fill the triangle. The smallest convex set containing the given points is their *convex hull*.



A weight of 0.75 on A places the result nearer A, one quarter of the way from A toward B. The numbers are coordinates in an invented representation; their axes have no assumed physical units.

The AI application isolates one use of this idea: a normalized nonnegative weighted combination of value vectors in attention. The geometric guarantee concerns that one averaging step. Later transformations in a model need not stay in the original hull, and the weights do not certify that an interpretation is true.

Pause and try 7

Why do weights 0.6 and 0.4 define a convex combination, while 1.2 and -0.2 do not?

8.12 A clear recommendation names its goal and assumptions

Linear models assume constant rates: each tune-up takes the same two hours and earns the same \$30 after parts; each quick repair takes one hour and earns \$20. Actual work can involve cancellations, unusually difficult repairs, different mechanics, and changing parts prices. The graph describes the consequences of the assumptions we have made.

Increasing a resource allowance in a "$\leq$" constraint expands the feasible set or leaves it unchanged. Therefore the best maximum score cannot decrease. But it need not improve: another resource may still block every better plan. Tightening a limit can shrink the set or even make it empty.

Example · More mechanic-hours do not create more customers

Keep $x + y \leq 8$, but suppose the shop can now staff sixteen mechanic-hours. It could schedule eight full tune-ups: (8,0) involves eight bikes and sixteen hours and earns \$240 after parts. Adding more hours alone would not increase the maximum while the shop still accepts only eight bikes. At \$30 per tune-up, eight bikes could earn at most \$240.

A practical routine

1. Identify what the people involved can choose, what the units mean, and whether fractions make sense.
2. Translate every resource limit and minimum requirement into a constraint.
3. State the objective and whether to maximize or minimize it.
4. Find all feasible vertices and check candidate intersections.
5. Compare objective values and identify ties or improving directions.
6. Check whole-number restrictions and interpret the plan and value.
7. Ask whom the decision affects, what the model leaves out, and what information could change the recommendation.

Week 9 adds another reason a plan might be impossible: locations and resources must be connected by usable routes. Keep the same distinction there – what is allowed, what is preferred, and what the model leaves out.

Further Worked Models

Furniture: compare every feasible corner

A small furniture maker has 120 standard boards and 70 hours available for finishing. Each chair x needs two boards and one finishing hour; each desk y needs three boards and two finishing hours. After materials, their contributions are \$40 and \$70. The constraints are $2x + 3y \leq 120$ and $x + 2y \leq 70$, with nonnegative counts, and the objective is to maximize $P = 40x + 70y$.

The first boundary has intercepts $(60, 0)$ and $(0, 40)$; the second has $(70, 0)$ and $(0, 35)$. The tighter axis limits give $(60, 0)$ and $(0, 35)$. To find the non-axis intersection, double $x + 2y = 70$ and subtract $2x + 3y = 120$: $y = 20$, then $x = 30$.

Corner	Boards	Finishing hours	Contribution
$(0, 0)$	0	0	\$0
$(60, 0)$	120	60	\$2,400
$(30, 20)$	120	70	\$2,600
$(0, 35)$	105	70	\$2,450

All four corners pass both limits. The maximum is \$2,600 at 30 chairs and 20 desks. The plan and its value are different quantities; both belong in the recommendation.

A resource limit can be redundant

Suppose a different model has limits $x + 2y \leq 20$ and $x + 2y \leq 24$. Every point passing the first also passes the second. The second is redundant. This differs from merely having slack at one plan: a constraint can be slack at a particular optimum yet still exclude other choices elsewhere.

A third real constraint can create a fifth corner

A neighborhood print shop has 60 standard sheets of printing material, room for 40 separate orders, and 70 finishing hours. One poster uses one sheet, one order, and two finishing hours. One banner uses two sheets, one order, and one finishing hour. Its constraints are:

$$x + 2y \leq 60, \quad x + y \leq 40, \quad 2x + y \leq 70, \quad x, y \geq 0.$$

Here x counts poster orders and y banner orders. The shop estimates contributions after materials of \$15 per poster and \$35 per banner. Maximize $P = 15x + 35y$.

Intersect the first two non-axis boundaries to get $(20, 20)$; the third total is 60, so it passes. Intersect the second and third to get $(30, 10)$; the first total is 50, so it passes. The first and third meet at $(80/3, 50/3)$, but this candidate has $x + y = 130/3 > 40$, so it must be discarded.

The x-axis corner is $(35, 0)$: the third constraint is tighter there than the other two. The y-axis corner is $(0, 30)$: the first constraint is tighter there. Together with the origin and the two valid non-axis intersections, these give five corners.

Feasible corner	Calculation	Contribution
$(0, 0)$	$15(0) + 35(0)$	\$0
$(35, 0)$	$15(35) + 35(0)$	\$525
$(30, 10)$	$15(30) + 35(10)$	\$800
$(20, 20)$	$15(20) + 35(20)$	\$1,000
$(0, 30)$	$15(0) + 35(30)$	\$1,050

The maximum is \$1,050 at $(0, 30)$. All three resource constraints matter to this feasible region; none is redundant. A calculation at a visually prominent intersection, such as $(20, 20)$, is not a substitute for comparing all corners.

A useful distinction

A constraint is *binding at a plan* when it holds with equality there. It is *redundant in a model* when removing it leaves the entire feasible set unchanged. These are different claims.

Part II Review

Use nonnegative variables and included boundaries. Show a corner table or explicit calculation and report both a plan and its objective value.

1. For the bike shop's corners $(0,0), (6,0), (4,4), (0,8)$, maximize contribution $P = 30x + 20y$. What is the minimum if the shop is not required to serve anyone that day?
2. On the same bike-shop region, maximize priority score $R = x + 4y$ and then the number of bikes served, $T = x + y$. Describe the optimal edge for T and its whole-number schedules.
3. A library has sixteen materials kits and twenty-four volunteer-hours. Information tables need one kit and three hours and reach five people; reading circles need two kits and two hours and reach eight. Maximize $f = 5x + 8y$ under $x + 2y \le 16$, $3x + 2y \le 24$.
4. A group needs eight folders and ten cards. Bag A has one folder and two cards for \$3; bag B has one folder and one card for \$5. Minimize $C = 3x + 5y$ subject to $x + y \ge 8$, $2x + y \ge 10$. Compare the boundary vertices.
5. For the furniture model in the worked example, compare the best plan with producing only desks. Does the higher per-item contribution determine the answer by itself?
6. For the print shop with limits on material, orders, and finishing hours, why is $(80/3, 50/3)$ not a feasible corner? Which corner maximizes $15x + 35y$?
7. A toy workshop has $3x + y \le 120$, $x + 2y \le 80$, and contribution $P = 8x + 5y$. Find the corners and best plan. Then change the contribution to $5x + 8y$ and compare.
8. A municipality divides an invented \$500-thousand budget between roads x and parks y, measured in thousands of dollars. Require $x \ge 100$, $y \ge 80$, $x + y \le 500$. Maximize an invented score $S = 3x + 4y$. What does the score leave for people to decide?
9. A polygon has corners $(0,0), (8,0), (5,4), (0,6)$. For cost $C = 4x + 6y$, which of the three listed nonzero corners is cheapest? Why is that not the minimum over the full polygon? Give a different positive feasible plan with lower cost.
10. For $x + 2y \le 8$, $3x + 2y \le 10$, maximize $5x + 8y$ continuously and then using whole numbers. Check rounding up and rounding down from the continuous best point.
11. Compute $0.75A + 0.25B$ for $A = (2,1), B = (8,5)$. Add $C = (2,7)$ and normalize amounts $2, 1, 1$. Where do the resulting points lie? Why do weights $1.2, -0.2$ fail the convex-combination test?
12. Explain the difference between an infeasible model and an unbounded objective. Give an example of an unbounded region with a finite minimum. Explain why $0 \le x < 1$ has no largest allowable x.
13. Keep the shop's eight-bike limit and increase mechanic-hours from twelve to sixteen and then twenty. Maximize $30x + 20y$ at each stage. Why do extra hours eventually stop improving contribution?
14. A repair cooperative has 40 restoration hours, can arrange 14 pickups, and earns after-parts contribution $30x + 50y$ from chairs and tables. Recommend a schedule, compare another feasible schedule, and name a missing real-world consideration.

Solutions and Reasoning

Short check 1. P=30(2)+20(5)=160 dollars. The score alone does not prove feasibility; check the constraints. Here the schedule has 7 bikes and uses 9 mechanic-hours, so the plan is feasible.

Short check 2. The best corner can occur where two non-axis boundaries meet. Here (4,4) gives \$200, exceeding the \$180 and \$160 at the axis corners. The method requires every feasible vertex.

Short check 3. Every point on the edge joining two optimal adjacent vertices has the same maximum score. Equal scores alone do not establish optimality: some other vertex might score higher. First compare all feasible corners.

Short check 4. It supplies only 6 pencils, below the minimum 8. Both requirements must hold. A boundary intercept of one requirement may fail another.

Short check 5. No. The supply-order region extends indefinitely, but positive costs give a minimum of \$20 at (2,4). Unbounded choices and an unbounded objective are different properties.

Short check 6. Every whole-number feasible plan is also continuously feasible, so its score is at most 33. But the point giving 33 has a fractional coordinate, so it does not establish an attainable whole-number score.

Short check 7. The first pair is nonnegative and sums to 1. The second also sums to 1 but has a negative weight; it can extrapolate beyond the segment. Both conditions are needed.

Part II Review: solutions

1. Corner contributions are \$0, \$180, \$200, and \$160. The maximum is \$200 at $(4, 4)$: four tune-ups and four quick repairs. With only upper limits, the minimum contribution is zero when no repairs are scheduled; a minimum service requirement would change that decision.

2. R-values are $0, 6, 20, 32$, so $(0, 8)$ is best. T reaches 8 along the edge from $(4, 4)$ to $(0, 8)$. Its whole-number plans are $(0, 8), (1, 7), (2, 6), (3, 5), (4, 4)$.

3. Subtraction gives $x = 4, y = 6$. The corners are $(0, 0), (8, 0), (4, 6), (0, 8)$ and the expected numbers of people reached are 0, 40, 68, 64. Four information tables and six reading circles reach an estimated 68 people, using all sixteen kits and twenty-four volunteer-hours.

4. Corners are $(8, 0), (2, 6), (0, 10)$, with costs $24, 36, 50$. The minimum is 24 at $(8, 0)$. It provides eight folders and sixteen cards, six more cards than required. Under the stated assumptions bag A is less expensive than bag B and includes more cards; a real purchase might also depend on availability or quality. A feasible axis corner can outperform a mixed intersection.

5. $(30, 20)$ gives \$2,600; desks only gives $(0, 35)$ and \$2,450. The best mix depends on both resource limits, not just the larger contribution per item.

6. That intersection has $x + y = 130/3 > 40$ and violates the middle constraint. The maximum is \$1,050 at $(0, 30)$, which passes all three limits.

7. Corners are $(0, 0), (40, 0), (32, 24), (0, 40)$. Original scores are $0, 320, 376, 200$, so $(32, 24)$ is best at 376. Under $5x + 8y$, scores are $0, 200, 352, 320$; the same plan remains best but its score changes. Changing an objective need not always move the optimum.

8. Corners are $(100, 80), (420, 80), (100, 400)$. Scores are $620, 1580, 1900$. The score selects roads \$100,000 and parks \$400,000. Its coefficients and minimum allocations are modeling choices, not

facts establishing what a community ought to value.

9. The three costs are 32,44,36, so $(8, 0)$ is cheapest among those candidates. The full polygon includes $(0, 0)$ at cost 0 and, for example, $(1, 0)$ at cost 4. A real community program needs an explicit minimum service requirement, rather than deleting just one zero-events corner from a comparison.

10. Continuous maximum 33 is at $(1, 3.5)$. Rounding up to $(1, 4)$ gives resource totals 9 and 11, both too large. Rounding down to $(1, 3)$ is feasible but scores 29. The best whole plan is $(0, 4)$ at 32; enumerate y-values 0–4 to verify.

11. The first point is $(3.5, 2)$, on the A–B segment. Normalized amounts give weights $(0.5, 0.25, 0.25)$ and point $(3.5, 3.5)$ inside the triangle. A negative weight violates nonnegativity even if the weights sum to 1.

12. Infeasible means no plan satisfies every condition. Unbounded objective means feasible plans improve the score without finite limit. The supply-order model $x + y \geq 6$, $2x + y \geq 8$ has unbounded choices but minimum $4x + 3y = 20$ at $(2, 4)$. Under $0 \leq x < 1$, any allowed x can be increased while remaining below 1; the limiting bound 1 is excluded.

13. With twelve mechanic-hours, the maximum is \$200 at $(4, 4)$. At sixteen or twenty hours, the maximum is \$240 at $(8, 0)$. The shop can serve at most eight bikes, each worth at most \$30 after parts; beyond sixteen hours extra mechanic time cannot increase this maximum. The time constraint then becomes redundant.

14. Restore eight chairs and six tables: $2(8) + 4(6) = 40$ restoration hours and $8 + 6 = 14$ pickups, yielding \$540 after parts. Ten tables are feasible but yield \$500. The cooperative should check donated-item availability, demand, volunteer schedules, and actual repair times.

Optional primary source for the AI connection

Vaswani et al., *Attention Is All You Need*, Section 3.2. The geometry applies to the normalized weighted value combination at that step; later transformations can move a representation beyond the original hull. The vectors and weights in this unit are constructed examples.