

Chapter 9

Systems of Linear Equations


A single linear equation in two variables describes a line; a pair of them asks where two lines meet. A linear equation in three variables describes a plane; a system of them asks where several planes meet. The solution set of a linear system is the intersection of these flat objects, and that intersection takes one of exactly three forms: a single point, nothing at all, or an entire shared line or plane. The geometry decides which, and substitution finds the answer.

9.1 Two Equations in Two Variables


Each equation $ax + by = c$ graphs as a line in the plane. A solution of the system is a point lying on both lines at once, so the solution set is the intersection of the two lines. Two lines in a plane fall into three positions.

Key idea. Two lines in the plane.


- *Crossing at one point: a unique solution.* The lines have different slopes.
- *Parallel and distinct: no solution.* Same slope, different intercepts; the lines never meet.
- *Coincident (the same line): infinitely many solutions.* Every point of the shared line solves both equations.



unique solution



no solution



infinitely many

Key idea. A linear system has exactly one of three outcomes: a unique solution, no solution, or infinitely many. No linear system has, say, exactly two solutions; flat objects meet in a flat object, and the only flat possibilities are a point, the empty set, or a whole line or plane.

9.2 Solving by Substitution

Substitution solves a system by isolating one variable in one equation and replacing it in the other, reducing two equations in two unknowns to one equation in one. The single equation is solved directly, and the value is fed back to recover the other variable.

Example 9.2.1. (*a unique solution by substitution*) Solve

$$\begin{aligned} 2x + y &= 7, \\ 3x - 2y &= 0. \end{aligned}$$

Isolate y in the first: $y = 7 - 2x$. Substitute into the second: $3x - 2(7 - 2x) = 0$, so $3x - 14 + 4x = 0$, giving $7x = 14$ and $x = 2$. Back-substitute: $y = 7 - 2(2) = 3$. The unique solution is $(2, 3)$, the crossing point of the two lines. Check: $2(2) + 3 = 7$ and $3(2) - 2(3) = 0$.

Example 9.2.2. (*detecting no solution*) Solve

$$\begin{aligned} x - y &= 1, \\ x - y &= 4. \end{aligned}$$

Substituting $x = y + 1$ from the first into the second gives $(y + 1) - y = 4$, that is $1 = 4$, a false statement. A contradiction signals that the lines are parallel and distinct: there is no solution.

Example 9.2.3. (*detecting infinitely many*) Solve

$$\begin{aligned} x + 2y &= 3, \\ 2x + 4y &= 6. \end{aligned}$$

The second equation is twice the first, so it carries no new information. Substituting $x = 3 - 2y$ into the second gives $2(3 - 2y) + 4y = 6$, that is $6 = 6$, always true. Every point on $x + 2y = 3$ solves the system; the solution set is that entire line. Writing $y = t$ as a free parameter, the solutions are $(3 - 2t, t)$ for all t .

Key idea. Substitution reveals the outcome through the equation it produces. A determined value gives a unique solution; a false statement such as $1 = 4$ signals no solution; an identity such as $6 = 6$ signals infinitely many, described by a free


parameter.

9.3 Three Equations in Three Variables

A linear equation $ax + by + cz = d$ in three variables graphs as a plane in space. A system of three such equations asks for the points common to three planes, and the solution set is their intersection. The same three outcomes appear, now staged by planes.

Key idea. Three planes in space.


- *Unique solution:* the three planes meet at a single point, like three walls meeting at a corner.
- *No solution:* the planes have no common point, for instance two parallel planes, or three forming an open triangular prism that never closes.
- *Infinitely many:* the planes share a whole line, or all coincide in a single plane.



unique: a corner



none: parallel



infinite: shared line

9.4 Back-Substitution and Triangular Form

A three-variable system is easiest to solve when it is in *triangular form*: the first equation involves all three variables, the second only two, the third only one. The last equation gives one variable immediately; substituting upward (back- substitution) recovers the others one at a time.

Key idea. A system is in *triangular form* when each successive equation drops a variable, leaving one equation in a single unknown. Solve that equation, then *back-substitute* the value upward to peel off the remaining variables in turn.

Example 9.4.1. (*back-substitution in a triangular system*) Solve

$$\begin{aligned}x + 2y - z &= 3, \\y + 2z &= 4, \\3z &= 9.\end{aligned}$$

The third equation gives $z = 3$. Substitute into the second: $y + 6 = 4$, so $y = -2$. Substitute both into the first: $x + 2(-2) - 3 = 3$, so $x - 7 = 3$ and $x = 10$. The unique solution is $(10, -2, 3)$. Each step uses only values already found, moving upward.

The work of solving a general system is to reach triangular form by eliminating variables. Adding a multiple of one equation to another removes a chosen variable without changing the solution set, because the new equation holds exactly when the old ones do.

Example 9.4.2. (*reducing to triangular form, then back-substituting*) Solve

$$\begin{aligned}x + y + z &= 6, \\2x - y + z &= 3, \\x + 2y - z &= 2.\end{aligned}$$

Subtract twice the first from the second: $(2x - y + z) - 2(x + y + z) = 3 - 12$, giving $-3y - z = -9$. Subtract the first from the third: $y - 2z = -4$. The system is now

$$\begin{aligned}x + y + z &= 6, \\-3y - z &= -9, \\y - 2z &= -4.\end{aligned}$$

From the last two, eliminate y : multiply the third by 3 to get $3y - 6z = -12$ and add to the second, giving $-7z = -21$, so $z = 3$. Back-substitute into $y - 2z = -4$: $y = 2$. Then the first gives $x = 6 - 2 - 3 = 1$. The solution is $(1, 2, 3)$. Check in all three originals: $1 + 2 + 3 = 6$, $2 - 2 + 3 = 3$, $1 + 4 - 3 = 2$.

Common pitfall. Elimination changes the equations but never the solution set, provided each step is reversible: adding a multiple of one equation to another, or scaling an equation by a nonzero constant. Scaling by zero is not allowed, since it discards an equation and can manufacture solutions the original system never had.

9.5 An Application

Example 9.5.1. (*mixing for a target*) A chemist combines three solutions of 10%, 20%, and 50% concentration to make 1 liter of 22% solution, using twice as much of the 10% as the 50%. Let x, y, z be the liters of the 10%, 20%, 50% solutions. The conditions are

$$\begin{aligned}x + y + z &= 1 && \text{(total volume),} \\0.10x + 0.20y + 0.50z &= 0.22 && \text{(total active ingredient),} \\x - 2z &= 0 && \text{(twice as much 10\% as 50\%).}\end{aligned}$$

From the third, $x = 2z$. Substituting into the first two and solving the resulting two-variable system gives $z = 0.2$, $x = 0.4$, and $y = 0.4$ liters. Three plain-language conditions become three linear equations, and their common solution is the recipe.

9.6 Exercises

Exercises 9.6

9.6.1. Describe the geometry and the number of solutions for each: (a) two lines with different slopes; (b) two distinct lines with equal slope; (c) one line written two equivalent ways.

9.6.2. Solve by substitution: $\begin{matrix} x + y = 5 \\ 2x - y = 1 \end{matrix}$ and check both equations.

9.6.3. Solve by substitution and identify the outcome: $\begin{matrix} 3x - y = 2 \\ 6x - 2y = 10 \end{matrix}$

9.6.4. Solve and identify the outcome: $\begin{matrix} x - 2y = 4 \\ -2x + 4y = -8 \end{matrix}$, writing the solution set with a free parameter if appropriate.

9.6.5. Explain why a linear system can never have exactly two solutions, in terms of the geometry of lines and planes.

9.6.6. Describe two distinct geometric arrangements of three planes that each yield no solution.

9.6.7. Solve the triangular system by back-substitution: $\begin{matrix} x - y + 2z = 5 \\ y - z = 1. \\ 4z = 8 \end{matrix}$

9.6.8. Reduce to triangular form and solve: $\begin{matrix} x + y + z = 4 \\ 2x + y - z = 3 \\ x - 2y + 3z = 11 \end{matrix}$, and check your solution.

9.6.9. Solve and classify: $\begin{matrix} x + y + z = 2 \\ 2x + 2y + 2z = 4 \\ x - y + z = 0 \end{matrix}$, describing the solution set geometrically.

9.6.10. A theater sells adult tickets at \$12 and child tickets at \$8. One evening 200 tickets bring in \$2000. Set up and solve a system for the numbers of each kind.

9.6.11. A shop blends coffee at \$8, \$10, and \$14 per pound into 10 pounds selling at \$11 per pound, using equal amounts of the two cheaper beans. Write and solve the system for the three weights.

9.6.12. Explain why adding a multiple of one equation to another leaves the solution set unchanged, and why scaling an equation by zero is forbidden.