Precalculus II

An Original Ten-Week Course Reader

Dr. Lee MATH& 142

Contents

1 Angles, Rotation, and the Unit Circle **2**
- 1.1 Angles measure turning . . . 2
- 1.2 Why radians are natural . . . 2
- 1.3 Arc length and angular speed . . . 2
- 1.4 Rotation becomes coordinates . . . 2
- 1.5 Reference angles and periodicity . . . 3
- 1.6 Exercises . . . 3

2 The Six Trigonometric Functions **4**
- 2.1 From a point to six functions . . . 4
- 2.2 Signs and the quadrants . . . 4
- 2.3 Where are the functions undefined? . . . 4
- 2.4 Right triangles are a special case . . . 4
- 2.5 Special angles without memorizing a disconnected table . . . 5
- 2.6 Applications: orientation and measurement . . . 5
- 2.7 Exercises . . . 5

3 Periodic Functions and Their Graphs **6**

4 Inverse Trigonometry and Identities **7**

5 Trigonometric Equations and Applications **8**

6 Vectors and Geometric Applications **9**

7 Polar Coordinates and Complex Numbers **10**

8 Parametric Equations and Conic Sections **11**

9 Sequences and Series **12**

10 Synthesis: Motion and Patterns **13**

1 Angles, Rotation, and the Unit Circle

This week's central question: How can we turn rotation into coordinates? The answer begins with the natural geometry of a circle.

1.1 Angles measure turning

An angle records a rotation from one ray to another. Counterclockwise rotations are positive, and clockwise rotations are negative. A rotation can exceed one full revolution. For example, 450° and 90° have the same terminal ray.

Definition 1.1. Two angles are *coterminal* if they have the same initial and terminal rays. They differ by an integer multiple of 360°, or 2π radians.

Example 1.1. A positive coterminal angle for -50° is $-50^\circ + 360^\circ = 310^\circ$.

1.2 Why radians are natural

For a circle of radius r with intercepted arc length s, define the radian measure $\theta = s/r$. This is a ratio of lengths. A full circumference is $2\pi r$, so one revolution measures 2π radians:

$$\boxed{\theta = \frac{s}{r}}, \qquad 180^\circ = \pi \text{ radians}.$$

Example 1.2. Convert 150° to radians: $150(\pi/180) = 5\pi/6$. Convert $7\pi/4$ radians to degrees: $(7\pi/4)(180/\pi) = 315^\circ$.

Try it. Convert 225° to radians and $-\pi/3$ radians to degrees, and sketch their terminal rays.

1.3 Arc length and angular speed

Rearrange the definition of radian measure to obtain $s = r\theta$. If rotation through θ takes time t, angular speed is $\omega = \theta/t$ and tangential speed is $v = r\omega$.

Example 1.3. A wheel of radius 0.4 m turns through $3\pi/2$ radians. A point on its rim travels $s = 0.4(3\pi/2) = 0.6\pi$ m.


Try it. A Ferris wheel of radius 12 m makes one revolution every 40 s. Find its angular speed and a passenger's tangential speed.

1.4 Rotation becomes coordinates

Take a circle of radius 1 centered at the origin. The point at angle θ, measured from the positive x-axis, has coordinates

$$\boxed{(x, y) = (\cos\theta, \sin\theta)}.$$

Cosine is the horizontal coordinate and sine is the vertical coordinate. Because the point lies on $x^2 + y^2 = 1$, we get $\cos^2\theta + \sin^2\theta = 1$.



1.5 Reference angles and periodicity

Special angles arise from right triangles:

θ	$\cos\theta$	$\sin\theta$
0	1	0
$\pi/6$	$\sqrt{3}/2$	$1/2$
$\pi/4$	$\sqrt{2}/2$	$\sqrt{2}/2$
$\pi/3$	$1/2$	$\sqrt{3}/2$
$\pi/2$	0	1

The reference angle supplies the absolute values in every quadrant; the quadrant gives the signs.

Example 1.4. At $5\pi/6$, the reference angle is $\pi/6$ and the point is in quadrant II, so $\cos(5\pi/6) = -\sqrt{3}/2$ and $\sin(5\pi/6) = 1/2$.

After one full revolution the point returns to its previous position:

$$\sin(\theta + 2\pi) = \sin\theta, \qquad \cos(\theta + 2\pi) = \cos\theta.$$

This geometric periodicity will become the foundation of sinusoidal graphs.

1.6 Exercises

1. Convert $-135°$ and $300°$ to radians.
2. Convert $11\pi/6$ and $-5\pi/4$ to degrees.
3. Find an angle in $[0, 2\pi)$ coterminal with $19\pi/6$.
4. A circle of radius 8 cm subtends an angle of $5\pi/12$. Find the arc length.
5. A turntable makes $33\frac{1}{3}$ revolutions per minute. Find angular speed in radians per second.
6. Explain why sine and cosine must lie between -1 and 1.
7. Find unit-circle coordinates at $2\pi/3$, $5\pi/4$, and $11\pi/6$.
8. A Ferris wheel has radius 10 m and center $(0, 12)$. Write a rider's coordinates after turning through θ from the rightmost point.

Looking ahead. Next week we extend sine and cosine to the remaining four trigonometric functions.

2 The Six Trigonometric Functions

Central question. How do six functions arise naturally from the same moving point on a circle?

2.1 From a point to six functions

Last week we studied the unit-circle point $P = (x, y) = (\cos\theta, \sin\theta)$. The two coordinates determine four additional ratios whenever the denominators are nonzero:

$$\sin\theta = y, \quad \cos\theta = x, \quad \tan\theta = \frac{y}{x},$$
$$\csc\theta = \frac{1}{y}, \quad \sec\theta = \frac{1}{x}, \quad \cot\theta = \frac{x}{y}.$$

These are not six unrelated measurements. They are six different ways of reading the same geometric information.

Definition 2.1. The *tangent* function is $\tan\theta = \sin\theta/\cos\theta$ where $\cos\theta \neq 0$. The functions csc, sec, cot are the reciprocals of sin, cos, tan respectively, wherever defined.

Example 2.1. At $\theta = \pi/3$ the circle point is $(1/2, \sqrt{3}/2)$. Thus $\tan\theta = \sqrt{3}$, $\sec\theta = 2$, $\csc\theta = 2/\sqrt{3} = 2\sqrt{3}/3$, and $\cot\theta = 1/\sqrt{3} = \sqrt{3}/3$.

2.2 Signs and the quadrants

The coordinate signs determine every trig-function sign. In quadrant I both coordinates are positive; in II, only sine is positive; in III, tangent is positive because both coordinates are negative; and in IV, only cosine is positive. Reciprocal functions have the same signs as their corresponding original functions.

Example 2.2. For $\theta = 5\pi/4$, the reference angle is $\pi/4$ and $P = (-\sqrt{2}/2, -\sqrt{2}/2)$. Therefore $\tan\theta = 1$, $\sec\theta = -\sqrt{2}$, and $\csc\theta = -\sqrt{2}$.

2.3 Where are the functions undefined?

Division by zero has no meaning. In particular, $\tan\theta$ and $\sec\theta$ do not exist where $\cos\theta = 0$, at $\pi/2 + k\pi$. Similarly $\cot\theta$ and $\csc\theta$ do not exist where $\sin\theta = 0$, at $k\pi$, for any integer k. The word *undefined* matters: an undefined value is not the number zero.

Try it. Find all six trig-function values at $\theta = 3\pi/2$, identifying every undefined value.

2.4 Right triangles are a special case

For an acute angle θ in a right triangle, scale the triangle so its hypotenuse has length 1. The opposite and adjacent sides then give the coordinates $\sin\theta$ and $\cos\theta$. Restoring a general hypotenuse h gives

$$\sin\theta = \frac{\text{opposite}}{h}, \quad \cos\theta = \frac{\text{adjacent}}{h}, \quad \tan\theta = \frac{\text{opposite}}{\text{adjacent}}.$$

The unit circle is more general than the right-triangle definitions: it handles negative angles, obtuse angles, and angles exceeding a full revolution.

Example 2.3. A ramp rises 1.5 meters over a horizontal run of 6 meters. Its angle of inclination satisfies $\tan\theta = 1.5/6 = 1/4$. Therefore $\theta = \arctan(1/4) \approx 14.0°$.

2.5 Special angles without memorizing a disconnected table

An equilateral triangle of side 2 splits into two right triangles with sides $1, \sqrt{3}, 2$. This produces $\sin 30^\circ = 1/2$ and $\cos 30^\circ = \sqrt{3}/2$. A square of side 1 has diagonal $\sqrt{2}$, producing $\sin 45^\circ = \cos 45^\circ = \sqrt{2}/2$.



2.6 Applications: orientation and measurement

Suppose an observer stands 40 m from the foot of a vertical tower. The angle of elevation to its top is 35°. The tower's height h satisfies

$$\tan 35^\circ = \frac{h}{40}, \qquad h = 40 \tan 35^\circ \approx 28.0 \text{ m}.$$

Here the tangent function links an angle to a ratio of lengths. Always sketch the physical situation and label which distances are opposite and adjacent before calculating.

2.7 Exercises

1. At $\theta = 2\pi/3$, find all six functions exactly.
2. At $\theta = 7\pi/4$, find $\tan, \cot, \sec, \csc$.
3. Explain geometrically why $\tan(\pi/2)$ is undefined.
4. For which angles in $[0, 2\pi)$ is $\csc\theta$ undefined?
5. A right triangle has opposite side 5 and adjacent side 12. Find all six trig ratios exactly.
6. A ladder leans against a wall, reaching 8 m high. Its foot is 3 m from the wall. Find the angle with the ground.
7. A drone is viewed at an elevation angle of 28° from a point 75 m horizontally from the point below it. Estimate its altitude.
8. Show that $\sin^2\theta + \cos^2\theta = 1$ implies $1 + \tan^2\theta = \sec^2\theta$ wherever $\cos\theta \neq 0$.

3 Periodic Functions and Their Graphs

Full instructional reading in development.

4 Inverse Trigonometry and Identities

Full instructional reading in development.

5 Trigonometric Equations and Applications

Full instructional reading in development.

6 Vectors and Geometric Applications

Full instructional reading in development.

7 Polar Coordinates and Complex Numbers

Full instructional reading in development.

8 Parametric Equations and Conic Sections

Full instructional reading in development.

9 Sequences and Series

Full instructional reading in development.

10 Synthesis: Motion and Patterns

Full instructional reading in development.