1 Angles, Rotation, and the Unit Circle

This week's central question: How can we turn rotation into coordinates? The answer begins with the natural geometry of a circle.

1.1 Angles measure turning

An angle records a rotation from one ray to another. Counterclockwise rotations are positive, and clockwise rotations are negative. A rotation can exceed one full revolution. For example, 450° and 90° have the same terminal ray.

Definition 1.1. Two angles are *coterminal* if they have the same initial and terminal rays. They differ by an integer multiple of 360°, or 2π radians.

Example 1.1. A positive coterminal angle for -50° is $-50^\circ + 360^\circ = 310^\circ$.

1.2 Why radians are natural

For a circle of radius r with intercepted arc length s, define the radian measure $\theta = s/r$. This is a ratio of lengths. A full circumference is $2\pi r$, so one revolution measures 2π radians:

$$\boxed{\theta = \frac{s}{r}}, \qquad 180^\circ = \pi \text{ radians}.$$

Example 1.2. Convert 150° to radians: $150(\pi/180) = 5\pi/6$. Convert $7\pi/4$ radians to degrees: $(7\pi/4)(180/\pi) = 315^\circ$.

Try it. Convert 225° to radians and $-\pi/3$ radians to degrees, and sketch their terminal rays.

1.3 Arc length and angular speed

Rearrange the definition of radian measure to obtain $s = r\theta$. If rotation through θ takes time t, angular speed is $\omega = \theta/t$ and tangential speed is $v = r\omega$.

Example 1.3. A wheel of radius 0.4 m turns through $3\pi/2$ radians. A point on its rim travels $s = 0.4(3\pi/2) = 0.6\pi$ m.


Try it. A Ferris wheel of radius 12 m makes one revolution every 40 s. Find its angular speed and a passenger's tangential speed.

1.4 Rotation becomes coordinates

Take a circle of radius 1 centered at the origin. The point at angle θ, measured from the positive x-axis, has coordinates

$$\boxed{(x, y) = (\cos\theta, \sin\theta)}.$$

Cosine is the horizontal coordinate and sine is the vertical coordinate. Because the point lies on $x^2 + y^2 = 1$, we get $\cos^2\theta + \sin^2\theta = 1$.



1.5 Reference angles and periodicity

Special angles arise from right triangles:

θ	$\cos\theta$	$\sin\theta$
0	1	0
$\pi/6$	$\sqrt{3}/2$	$1/2$
$\pi/4$	$\sqrt{2}/2$	$\sqrt{2}/2$
$\pi/3$	$1/2$	$\sqrt{3}/2$
$\pi/2$	0	1

The reference angle supplies the absolute values in every quadrant; the quadrant gives the signs.

Example 1.4. At $5\pi/6$, the reference angle is $\pi/6$ and the point is in quadrant II, so $\cos(5\pi/6) = -\sqrt{3}/2$ and $\sin(5\pi/6) = 1/2$.

After one full revolution the point returns to its previous position:

$$\sin(\theta + 2\pi) = \sin\theta, \qquad \cos(\theta + 2\pi) = \cos\theta.$$

This geometric periodicity will become the foundation of sinusoidal graphs.

1.6 Exercises

1. Convert $-135°$ and $300°$ to radians.
2. Convert $11\pi/6$ and $-5\pi/4$ to degrees.
3. Find an angle in $[0, 2\pi)$ coterminal with $19\pi/6$.
4. A circle of radius 8 cm subtends an angle of $5\pi/12$. Find the arc length.
5. A turntable makes $33\frac{1}{3}$ revolutions per minute. Find angular speed in radians per second.
6. Explain why sine and cosine must lie between -1 and 1.
7. Find unit-circle coordinates at $2\pi/3$, $5\pi/4$, and $11\pi/6$.
8. A Ferris wheel has radius 10 m and center $(0, 12)$. Write a rider's coordinates after turning through θ from the rightmost point.

Looking ahead. Next week we extend sine and cosine to the remaining four trigonometric functions.