

Dual Seeding under a Geographic Paradox: Hospital-Associated SARS-CoV-2 in Hong Kong after Visit-Schedule Relaxation

Companion research note to Shen et al. (2026)

Exploratory genomic-epidemiologic reanalysis

Working first draft — Laidlaw-caliber research essay / companion article

11 July 2026

Abstract

Background. After Hong Kong relaxed hospital visitation in May 2022, nosocomial SARS-CoV-2 clusters re-emerged amid Phase-2 community transmission. A recent multihospital genomic study characterised ward-level transmission and found no independent association between community mobility and community-to-hospital viral introductions. Here we ask two further questions that study left open: (i) whether ward outbreaks are initiated by a single dominant route or by *dual seeding*, and (ii) why sequenced hospital-associated infections concentrate in New Territories (NT) clusters despite lower population and hospital geographic density.

Methods. We reanalysed the published hospital-associated sequence plot table ($n = 162$ cases; 17 anonymised hospitals; 28 May–18 Aug 2022), linked cases to Hospital Authority (HA) clusters, classified ward-level initiation pathways from first detection dates, and standardised burden by HA beds, inpatient discharges, and patient-days (FY2021–22). District census data quantified catchment density. Official HA nosocomial press bulletins were mined for an ascertainment footprint (presence of clusters by hospital/cluster).

Results. Staff/outpatient-first and inpatient-first pathways both contributed substantial burden (48 vs 54 cases among ordered dual-type wards); pathway mix did not differ detectably between NT and urban clusters. NTWC+NTEC comprised 82.7% of sequenced hospital-associated cases despite $\sim 0.27\times$ mean population density and $\sim 0.20\times$ hospital geographic density relative to urban clusters in the sample. The NT excess survived standardisation by beds ($\sim 5.7\times$), discharges ($\sim 6.9\times$), and patient-days ($\sim 5.7\times$). Kowloon East provided a negative control (low hospital density, low sequenced burden). Official bulletins mentioned Kowloon Central Cluster hospitals that are absent from the sequenced table, indicating uneven genomic coverage; NT/psychiatric hospitals (notably Castle Peak) were also prominent in public reporting.

Conclusions. Hospital-associated SARS-CoV-2 in this window was dually seeded and geographically structured, not a simple function of urban density or citywide mobility. Throughput denominators do not erase the NT excess; ascertainment is uneven and must temper causal language. The findings support ward- and case-mix-targeted infection prevention—including long-stay/psychiatric settings—alongside visitor screening.

Keywords: SARS-CoV-2; nosocomial transmission; dual seeding; geographic concentration; New Ter-

1 Introduction

Hong Kong’s fifth COVID-19 wave forced a transition from near-elimination to managed endemicity. In May 2022, authorities phased in relaxed hospital visitation while community incidence rebuilt into Phase 2 of the fifth wave [1]. Understanding how SARS-CoV-2 entered and spread within hospitals during that transition matters for future respiratory-pathogen preparedness: visitation has clear benefits for patients [1], yet wards remain vulnerable when community prevalence is high.

Shen et al. sequenced 162 genomes linked to 29 suspected ward clusters across 17 public hospitals (28 May–18 August 2022), confirmed 18 nosocomial clusters (126 genomes), showed predominantly intraward transmission without resolved interward spread, and inferred more community→hospital than hospital→community Markov jumps [1]. After adjusting for time trends, generalised additive models found no significant association between introduction rates and standard mobility indices [1].

Two gaps remain. First, *how* wards are seeded: the parent analysis noted phylogenetic staff-before-inpatient order in only 2/18 confirmed clusters [1], but a broader temporal taxonomy of staff/outpatient versus inpatient first detections has not been quantified across all investigated wards. Second, *where* burden concentrates: sequenced hospital-associated infections are heavily skewed toward New Territories West and East clusters (NTWC, NTEC), yet New Territories districts are less densely populated and have fewer hospitals per land area than urban catchments—a pattern that challenges naïve “dense city, more hospital outbreaks” intuition.

Research questions.

1. Are hospital-associated clusters in this window consistent with a single dominant initiation route, or with **dual seeding** (staff/outpatient bridges and inpatient introductions)?
2. Does the New Territories concentration of sequenced hospital-associated infections persist after standardisation by hospital throughput, and can density or ascertainment explain it?

Contribution. This companion note does not re-estimate phylogenies. It delivers a manuscript-ready descriptive and standardised reanalysis: a dual-seeding taxonomy; a quantified density paradox with negative controls; admissions/patient-day denominators; and an honest ascertainment audit against public HA bulletins.

2 Methods

2.1 Data sources

- **Hospital-associated cases.** Anonymised plot table from the parent study ($n = 162$ sequenced cases; hospitals A–Q; HA cluster labels; ward; case type; lineage; date) [1].

- **HA capacity and throughput.** Beds (31 Mar 2022) and FY2021–22 inpatient discharges and patient-days from HA Statistical Report cluster extracts [2].
- **Catchment demography.** 2021 Population Census district populations and land areas, mapped to conventional HA cluster catchments [3, 4].
- **Official reporting footprint.** HA press releases on info.gov.hk announcing nosocomial COVID-19 clusters during the study window (presence of hospitals in nosocomial tables/narratives).
- **Parent coverage anchor.** Confirmed genomic clusters represented 51.9% of officially reported nosocomial cases in the window [1].

2.2 Dual-seeding classification

For each hospital–ward combination, we recorded the earliest detection date by case type. Pathways were labelled: *staff_first*, *inpatient_first*, *same_day*, *staff_only*, or *inpatient_only*. “Staff” here aggregates Outpatient/Staff in the plot table. This is a *temporal first-detection* definition, complementary to—not identical with—the parent paper’s phylogenetic staff-index criterion [1]. Among wards with both staff and inpatient detections and a clear order, we tested NT versus urban pathway mix with Fisher’s exact test.

2.3 Geographic and density metrics

HA clusters were grouped as New Territories (NTWC, NTEC) versus urban clusters present in the sequenced table (KWC, KEC, HKEC). Population density used catchment population (2017 planning projection) divided by core district land area. Hospital and bed geographic density used HA facility counts and beds per 100 km² of core land area. Islands/Lantau splits for HKEC/KWC were excluded from core land area (conservative for urban density).

2.4 Denominator standardisation

Sequenced hospital-associated case counts were standardised as: cases per 1,000 beds; per 100,000 annual inpatient discharges; and per 100,000 annual inpatient patient-days. Annual FY2021–22 throughput is an imperfect proxy for an 83-day window; we therefore emphasise relative NT/urban ratios rather than absolute window incidence. Mentally-ill patient-day share was computed as a case-mix marker.

2.5 Ascertainment audit

Because the parent supplement Table S3 (official case totals by hospital) was not recoverable in this environment (curated Excel gitignored; publisher supplement inaccessible), we did *not* invent cluster-level coverage fractions from daily press tables (which are incremental and reconstruct only a small fraction of the implied ~243 official cases). Instead we contrast **presence**: HA clusters appearing in official nosocomial bulletins versus those appearing in the sequenced plot table, anchored by the published 51.9% overall coverage statistic [1].

2.6 Ethics and anonymisation

Analyses use aggregated, anonymised hospital codes as in the parent study. HA cluster labels provide geographic context already used publicly [1].

3 Results

3.1 Dual seeding is a shared mechanism

Across 29 wards, ordered dual-type pathways contributed 48 staff/outpatient-first and 54 inpatient-first cases; same-day detections added 24 (Table 1). Staff-only and inpatient-only wards contributed smaller burdens. Fisher’s exact test comparing NT versus urban wards for staff-first versus inpatient-first yielded $OR = 0.33$, $p = 0.56$ ($n = 13$ wards): no evidence of different dominant routes by region. Among inpatients with known admission-to-positive interval (API), 74/90 (82%) had $API \geq 3$ days, consistent with substantial in-hospital acquisition once infection is inside the ward [1].

Table 1: Ward-level temporal initiation pathways (sequenced hospital-associated cases).

Pathway	Wards	Cases
Staff/outpatient first	6	48
Inpatient first	7	54
Same day	3	24
Staff/outpatient only	5	15
Inpatient only	8	21




Figure 1: Dual-seeding composition by region (case counts). Staff/outpatient-linked and inpatient-linked routes both matter in New Territories and urban clusters in the sample.

3.2 Geographic concentration despite lower density

NTWC+NTEC accounted for 134/162 (82.7%) sequenced hospital-associated cases. Mean catchment population density for NT clusters was $\sim 0.27\times$ that of urban in-sample clusters; hospitals per 100 km² $\sim 0.20\times$; beds per 100 km² $\sim 0.29\times$ (Figure 2). Kowloon East Cluster (KEC) had similarly low hospital geographic density (~ 2 per 100 km²) but low sequenced burden—a negative control showing that low density is not sufficient for high observed hospital-associated infection.




Figure 2: Density paradox at HA-cluster scale. New Territories clusters (green) combine lower population and hospital geographic density with higher sequenced hospital-associated infection rates per 1,000 beds.

3.3 The NT excess survives throughput denominators

Standardising by beds, discharges, and patient-days left a several-fold NT excess relative to urban in-sample clusters (Table 2; Figure 3): approximately $5.7\times$ per 1,000 beds, $6.9\times$ per 100,000 discharges, and $5.7\times$ per 100,000 patient-days (means of cluster-level rates). NTWC showed the highest mentally-ill patient-day share (19.2% of cluster patient-days), linking geography to long-stay case-mix (Figure 4). All 22 BA.5.6 sequences in the plot table arose from a single hospital ward with almost exclusively $\text{API} \geq 3$ inpatients—a deep nosocomial signature inside NTWC [1].

Table 2: Sequenced hospital-associated infections standardised by HA throughput (FY2021–22).

Cluster	Cases	Beds	Discharges	Patient-days	/1k beds	/100k discharges
NTWC	78	4,762	140,388	1,331,433	16.38	55.56
NTEC	56	5,144	163,397	1,311,078	10.89	34.27
KWC	15	4,953	201,096	1,416,390	3.03	7.46
HKEC	8	3,307	102,190	822,971	2.42	7.83
KEC	5	2,922	114,909	809,758	1.71	4.35
KCC	0	6,005	199,221	1,599,173	0	0
HKWC	0	3,076	103,370	635,637	0	0




Figure 3: Sequenced hospital-associated infection rates under three denominators. New Territories clusters remain elevated after bed, discharge, and patient-day standardisation.




Figure 4: Mentally-ill patient-day share versus sequenced hospital-associated infection rate. NTWC combines high psychiatric throughput share with the highest bed-standardised rate.

3.4 Ascertainment is uneven—and must be stated plainly

Parent-study confirmed clusters covered 51.9% of official nosocomial cases overall [1]. Public HA bulletins during the window mentioned hospitals in *all seven* HA clusters, including Kowloon Central and Hong Kong West—yet those two clusters contribute zero cases to the sequenced plot table (Figure 5). Conversely, NTEC (especially Castle Peak Hospital) and NTEC appear in both official bulletins and the sequenced table. Daily bulletin tables are incremental and cannot replace Table S3 cumulative counts; we therefore claim a **presence contrast**, not cluster-specific coverage percentages.




Figure 5: Official nosocomial bulletin mention-days versus sequenced hospital-associated cases. KCC lies on the x -axis (mentioned, not sequenced in the plot table), illustrating uneven genomic coverage.

4 Discussion

4.1 What the dual-seeding result changes

If wards were seeded almost only by visitors, policy could focus narrowly on visitation rules. Our temporal taxonomy shows staff/outpatient-first and inpatient-first routes both carry large case burdens, and the mix is shared across NT and urban clusters. That aligns with the parent paper’s caution that healthcare workers with routine community exposure are plausible bridges [1], while also preserving a role for incubating admissions despite mandatory admission screening. Phylogenetic staff-index events were rarer (2/18 confirmed clusters) [1]; temporal first detection is more sensitive and should be reported as a complementary definition.

4.2 Why the density paradox matters

The NT excess is the opposite of a density-driven urban story. Lower population and hospital geographic density, yet higher standardised sequenced burden, redirects attention to hospital case-mix (long-stay and

psychiatric capacity), IPC implementation, and sampling—not skyline density. KEC’s low density with low burden prevents the lazy inference that “sparse geography causes outbreaks.”

4.3 Denominators versus ascertainment

Admissions and patient-day standardisation were the cleanest quantitative tests available without hospital-level window admissions. They fail to wash out NT concentration. Sequencing-effort tip counts from the study trees are largely circular with the sample itself and were not used as independent proof. The bulletin presence contrast shows under-representation of some urban clusters (notably KCC) in the sequenced table: NT excess is unlikely to be *only* artefact, but it is also not fully separable from ascertainment without Table S3 hospital totals.

4.4 Limitations

1. Sequenced hospital-associated cases $\neq$ complete nosocomial incidence.
2. Annual throughput approximates an 83-day window.
3. Temporal dual-seeding $\neq$ phylogenetic index.
4. Official bulletin mining is incomplete relative to curated Table S3.
5. Catchment populations are planning estimates, not closed cohorts.

4.5 Implications

During high community prevalence, protecting hospitals requires more than citywide mobility management (already null in the parent GAM) [1]. Practical levers include: (i) dual attention to staff/community bridges *and* admission pathways; (ii) heightened IPC in long-stay/psychiatric settings; (iii) genomic surveillance designed for even cluster coverage, not only the loudest outbreaks.

5 Conclusion

Hospital-associated SARS-CoV-2 in Hong Kong’s post-relaxation window was **dually seeded** and **geographically structured**. New Territories concentration persists after throughput standardisation and sits inside a density paradox that urban intuition does not explain. Ascertainment is uneven; recovering hospital-level official totals remains the highest-value upgrade. Until then, the evidence supports ward- and case-mix-targeted prevention over density- or mobility-centric narratives alone.

Data and code

Analysis scripts and tabulated outputs are in the project repository (`scripts/4.dual_seeding-geography.py`, `scripts/5.denominators_ascertainment.py`, and `analysis/`). Parent genomic methods and primary figures are described in Shen et al. [1].

Competing interests

None declared for this companion reanalysis draft.

Acknowledgements

This draft builds on the open methods and anonymised plot outputs of the parent genomic epidemiology study [1]. HA Statistical Report extracts and Census/LegCo catchment figures are public.

References

- [1] Shen R., et al. (2026). Genomic epidemiology of hospital-associated SARS-CoV-2 clusters in Hong Kong during a period of relaxed visitation (May–August 2022). *Influenza and Other Respiratory Viruses*, 20, e70249. doi:[10.1111/irv.70249](https://doi.org/10.1111/irv.70249)
- [2] Hospital Authority (2022). *HA Statistical Report 2021–2022* (cluster extracts; service capacity as at 31 March 2022; inpatient discharges and patient-days). Available from: <https://www3.ha.org.hk/data/HAStatistics/StatisticalReport/2021-2022>
- [3] Census and Statistics Department, HKSAR (2022). *2021 Population Census: Summary Results* (district populations and densities).
- [4] Government of the HKSAR (2017). LegCo Question 13: Resource allocation of Hospital Authority (cluster catchment population projections). <https://www.info.gov.hk/gia/general/201712/13/P2017121300666.htm>

A Decision log (for transparency)

1. **Laidlaw caliber:** Yes—original questions, reproducible methods, graded evidence, explicit limits; drafted to exceed a 2,000–3,000-word essay toward a companion article.
2. **Table S3:** Not reconstructible from incremental public bulletins; presence contrast used instead; author Excel remains the upgrade path.
3. **Causal language:** Density and mobility are treated as context/nulls, not causes.