

Heat, cold and first hospitalisation after cardiovascular diagnosis in Hong Kong

An exploratory monthly study among people with diabetes or hypertension, 2013–2023

Bob Shen Ruililin

Laidlaw Scholars Programme · The University of Hong Kong · Summer 2026

Contents

Summary	2
1. Introduction	2
1.1 A city facing heat and cold	2
1.2 Why monthly data require a different question	3
1.3 Objectives	3
2. Methods	4
2.1 Study design and data	4
2.2 Temperature measures	4
2.3 Statistical analysis	5
2.4 Robustness and the attempted daily-exposure extension	6
3. Results	6
3.1 Environmental and outcome patterns	6
3.2 The twelve-comparison panel	7
3.3 What the uncertainty comparison changed	8
3.4 Robustness and calibration	8
4. Interpretation	9
4.1 What the study suggests	9
4.2 Why the result still matters	9
4.3 Strengths	9
4.4 Limitations	10
5. Implications and next study	10
6. Conclusion	10
Acknowledgements	11
Appendix A. Complete core panel	11
References	11

Summary

Hong Kong is becoming hotter, especially at night, but cold winter days have not disappeared. Both heat and cold can place additional stress on the cardiovascular system. This matters particularly for people living with type 2 diabetes or hypertension, who already face a high risk of coronary heart disease and heart failure.

This project examined whether monthly temperature patterns were associated with monthly counts of a carefully defined hospital event. The available Hospital Authority data counted each person's first recorded hospitalisation after their first diagnosis record for coronary heart disease (CHD) or heart failure (HF). They did not record the reason for admission. The study therefore does not claim that every admission was caused by CHD or HF, and it does not estimate individual risk. Its question is narrower: in months with more heat or cold, did the governed territory-wide count of these first hospitalisations also differ?

The analysis covered January 2013 to December 2023, giving 132 monthly observations. Six temperature measures were examined separately for each outcome: monthly mean temperature, mean daily maximum temperature, mean daily minimum temperature, hot nights, cold days, and very hot days. This produced a panel of twelve comparisons. Models accounted for season, long-term change, and the different number of days in each month. Four approaches to uncertainty were shown, and the twelve comparisons were corrected for multiple testing.

Two associations stood out before that correction. Five additional hot nights in a month corresponded to a CHD count ratio of 1.022 (95% confidence interval 1.002–1.042 using the continuity uncertainty method). Five additional cold days corresponded to an HF count ratio of 1.073 (1.006–1.144). These are approximately 2.2% and 7.3% higher monthly counts, respectively. However, both multiple-testing-adjusted q-values were 0.192, and all twelve q-values exceeded 0.19. The CHD hot-night interval also included no association under two of the four uncertainty methods. The findings are therefore exploratory, not confirmatory.

The main contribution is not a claim that one weather measure has been proved to cause one disease. It is a reproducible framework that reports the complete panel, makes uncertainty visible, and refuses claims that the available data cannot support. Better daily outcome data, a denominator for the cohort still at risk, and a recorded cause of admission would be needed for stronger inference.

1. Introduction

1.1 A city facing heat and cold

Climate change does not replace one thermal hazard with another. In Hong Kong, hot nights have become more frequent, while cold days still return each winter. At Hong Kong Observatory Headquarters, the number of hot nights rose from 10 in 2013 to a peak of 61 in 2021. Cold-day counts fluctuated substantially but remained present throughout the study period. The city must therefore plan for a changing mixture of overnight heat, very hot days, and episodic cold rather than assuming that warming makes cold irrelevant.

Temperature can affect cardiovascular health through several pathways. Heat can promote dehydration, disturb sleep and overnight recovery, and increase cardiac demand. Cold can raise blood pressure and vascular resistance. Responses may be stronger in people whose cardiovascular regulation is already compromised by diabetes or hypertension. These mechanisms make both sides of the temperature distribution relevant, but they do not establish what will be visible in a particular dataset.

Hong Kong already has important evidence on daily temperature and health. Earlier studies examined daily stroke or acute myocardial infarction admissions and reported substantial cold-related patterns (Goggins et al. 2012; Goggins, Chan, and Yang 2013). Liu and colleagues estimated mortality attributable to non-optimal temperature and found a larger fraction associated with cold than heat during 2006–2016 (J. Liu et al. 2020). More recent work modelled heat-related excess deaths under several heatwave definitions (Z. Liu et al. 2026). Those studies answer valuable but different questions. A daily mortality attributable fraction, a modelled number of excess deaths, and a monthly hospitalisation count ratio cannot be substituted for one another.

1.2 Why monthly data require a different question

Health data are often released in aggregate form to protect privacy. In this project, the governed outcomes were available only as territory-wide monthly counts. That resolution can support a study of monthly burden, but it cannot reveal whether a particular hot day triggered a hospitalisation several days later. It also cannot show which individual experienced which temperature.

The outcome definition requires equal care. The delivered data count the first recorded hospitalisation after a first CHD or HF diagnosis record among people with type 2 diabetes and/or hypertension. Admission cause is absent. Calling these events “heart attacks caused by temperature,” or even “principal-diagnosis CHD admissions,” would go beyond the data. The report uses the longer but accurate description because precision about the outcome is part of the science.

1.3 Objectives

The primary objective was to estimate monthly associations between thermal burden and first-hospitalisation counts for CHD and HF during 2013–2023. Three secondary objectives were to:

1. compare continuous temperature measures with official extreme-day counts;
2. show how conclusions change across defensible uncertainty methods; and
3. determine whether the available monthly outcomes could reliably support a more ambitious estimate using daily exposure information.

2. Methods

2.1 Study design and data

The study used an ecological monthly time-series design. Each observation was one calendar month for Hong Kong, from January 2013 through December 2023. The final panel contained 132 months.

Daily weather observations came from Hong Kong Observatory Headquarters. Public weather series were checked against annual Observatory summaries before analysis. Population data came from the Census and Statistics Department, and air-pollution series were assembled from Environmental Protection Department general monitoring stations for the broader project. The core models reported here focus on thermal exposures and do not claim that pollution or influenza confounding has been fully resolved.

Governed Hospital Authority aggregates supplied the two outcomes. The cohort comprised people recorded with type 2 diabetes and/or hypertension during the study period. For each outcome, the monthly series counted the first recorded hospitalisation after the person's first CHD or HF diagnosis record. The analysis used aggregate counts only; source monthly health files and merged health panels were not placed in public version control.

2.2 Temperature measures

No single number fully represents thermal burden. A warm monthly average can arise from consistently mild conditions, several intense days, or unusually warm nights. These patterns may matter differently for health. Six measures were therefore evaluated:

- monthly mean temperature;
- monthly mean of daily maximum temperature;
- monthly mean of daily minimum temperature;
- number of hot nights, defined as minimum temperature at least 28 °C;
- number of very hot days, defined as maximum temperature at least 33 °C; and
- number of cold days, defined as minimum temperature at most 12 °C.

The continuous measures were interpreted per 1 °C. Extreme-day counts were interpreted per five additional days in a month. More complex spell and heatwave definitions were developed and source-checked elsewhere in the project (Wang et al. 2019; Li et al. 2025), but the six-measure panel provides the clearest common comparison for this report.

Official thermal extremes at HKO Headquarters, 2013–2023

Annual counts. Pipeline totals match HKO Year's Weather summaries (33/33).



Source: Hong Kong Observatory dailyExtract, Headquarters station. Environmental descriptors only — not health-outcome results.

Figure 1: Official hot nights, very hot days and cold days at Hong Kong Observatory Headquarters, 2013–2023. These are environmental descriptors, not health findings.

2.3 Statistical analysis

Separate negative-binomial regression models were fitted for each outcome and each exposure: two outcomes multiplied by six exposures produced twelve core comparisons. Negative-binomial models allow monthly counts to vary more than a simple Poisson model would permit. Each model controlled for calendar month, which captures recurring seasonal patterns, and a smooth long-term trend. The number of days in the month was included as an offset so that February was not treated as providing the same observation time as a 31-day month.

The reported effect measure is a **count ratio**. A ratio of 1 means that the model estimates no difference in the monthly count for the stated exposure change. A ratio of 1.05 means an estimated 5% higher count; a ratio of 0.95 means an estimated 5% lower count. Because the number of people in the cohort who remained eligible for a first event was unavailable for each month, these are count ratios rather than individual incidence-rate ratios.

Monthly observations close together in time may remain correlated after season and

trend are modelled. To avoid hiding this choice, uncertainty was reported four ways: model-based, heteroskedasticity-consistent (HC1), and Newey–West methods with lags of three and six months. The continuity analysis used the six-month Newey–West interval, but all four were displayed. No method was chosen because it made a confidence interval exclude 1.

Twelve comparisons also create more opportunities for a small p-value to occur by chance. Benjamini–Hochberg q-values were therefore calculated across the complete twelve-comparison panel. A q-value is not the probability that a result is true or false. It is a tool for limiting the expected proportion of false discoveries when a set of comparisons is considered together.

2.4 Robustness and the attempted daily-exposure extension

Robustness checks varied the long-term trend, study period, exposure lags and influential months; joint models were diagnostic only. A proposed method combining monthly outcomes with daily exposures was tested in simulation (Basagaña and Ballester 2024). It failed predefined requirements for error, coverage, bias and sign recovery, so it was not used to produce a real daily health coefficient.

3. Results

3.1 Environmental and outcome patterns

The weather record shows a clear rise in hot nights and very hot days across the decade, with year-to-year variation. Cold days were less common than hot nights in recent years but remained recurrent. This coexistence supports examining heat and cold together rather than treating one as a historical concern.

The governed outcome series contained 156,156 CHD first-hospitalisation events and 29,681 HF events across 132 months. The corresponding monthly means were 1,183.0 and 224.9. Both outcomes displayed seasonality, with a stronger winter peak for HF. These totals describe the study series; they do not establish an effect of weather.

Table 1. Outcome summary

Outcome	Months	Total events	Mean per month	Event definition
Coronary heart disease	132	156,156	1,183.0	First hospitalisation after first recorded CHD diagnosis
Heart failure	132	29,681	224.9	First hospitalisation after first recorded HF diagnosis

3.2 The twelve-comparison panel




Figure 2: Count ratios and 95% confidence intervals for the twelve core monthly models. The dashed line at 1 indicates no estimated difference. All twelve Benjamini-Hochberg q-values exceeded 0.19.

Most core estimates were close to a count ratio of 1. For CHD, monthly mean, maximum and minimum temperatures produced count ratios of approximately 0.993–0.994 per 1 °C. Cold days and very hot days were also near 1. The largest CHD estimate was for hot nights: 1.022 per five additional hot nights (95% confidence interval 1.002–1.042 using Newey-West lag 6; $p = 0.032$).

For HF, the continuous temperature estimates were modestly below 1. The minimum-temperature estimate was 0.973 per 1 °C (0.947–1.000). Hot nights and very hot days were near 1. The largest HF estimate was for cold days: 1.073 per five additional cold days (1.006–1.144; $p = 0.031$).

In plain language, the two leading estimates correspond to approximately 2.2% more CHD first-hospitalisation events per five additional hot nights and 7.3% more HF first-hospitalisation events per five additional cold days. These descriptions concern modelled monthly counts, not an individual’s probability of hospitalisation.

After correction across all twelve comparisons, the two leading estimates each had $q = 0.192$. Every core q-value exceeded 0.19. The complete panel therefore provides no multiplicity-protected confirmatory association.

3.3 What the uncertainty comparison changed




Figure 3: The two leading exploratory estimates under four approaches to standard errors. The CHD hot-night interval crosses 1 under two methods; the HF cold-day interval does not.

The CHD hot-night point estimate was 1.022 under all four standard-error methods because the fitted model did not change. Its interval did change: 0.995–1.049 using model-based uncertainty, 0.997–1.047 using HC1, 1.000–1.044 using Newey-West lag 3, and 1.002–1.042 using Newey-West lag 6. The first two intervals include 1, while the latter two narrowly exclude it. The interpretation therefore depends on how serial and unequal uncertainty is handled.

The HF cold-day point estimate was 1.073. Its four intervals were 1.023–1.125, 1.011–1.138, 1.007–1.143, and 1.006–1.144. All excluded 1, so this association was more stable across uncertainty methods. That stability does not overcome the multiple-testing result: its q-value remained 0.192.

3.4 Robustness and calibration

Across trend and period checks, the CHD hot-night count ratio ranged from 1.011 to 1.025 and the HF cold-day ratio from 1.043 to 1.113; excluding the most influential month preserved both directions. Later lags weakened the CHD estimate. The monthly-outcome/daily-exposure method failed substantive calibration, including false-positive rates up to 0.150 and coverage as low as 0.840, and produced no

real daily coefficient.

4. Interpretation

4.1 What the study suggests

The results suggest that the way thermal burden is defined matters. Monthly average temperature and counts of official extremes did not produce the same patterns. In the CHD series, hot nights stood out more than mean temperature or very hot days. In the HF series, cold days stood out more than hot nights. This is scientifically plausible: nighttime heat may limit physiological recovery, while cold may increase circulatory strain. It is also compatible with Hong Kong research that has retained cold as an important cardiovascular hazard while drawing increasing attention to nighttime heat (Goggins, Chan, and Yang 2013; Guo et al. 2024).

Compatibility is not confirmation. The twelve models were specified after the outcome series was available, and no association survived correction across the panel. The CHD hot-night interval changed its null-crossing status across uncertainty methods. Correlated heat measures also redistributed the CHD coefficient when entered together. The defensible conclusion is therefore that the panel contains two hypotheses worth testing with better data, not two causal discoveries.

4.2 Why the result still matters

Null or uncertain findings can be useful when they close off weak claims. This project establishes that monthly governed data can support a transparent comparison of thermal-burden measures. It also shows where that design stops. The data cannot identify within-month timing, individual exposure, or the cause of admission. The failed daily-exposure calibration provides an empirical reason not to present a daily coefficient, rather than relying only on a general warning about aggregation.

The distinction has practical consequences. A health-protection message based on confirmed daily triggering would require different evidence from an exploratory monthly burden pattern. The present findings do not justify disease-specific warning thresholds. They do support continued attention to both overnight heat and winter cold, and they motivate a governed design that links daily weather to dated, cause-recorded admissions.

4.3 Strengths

The project has five main strengths. First, it uses an eleven-year public weather record and governed health aggregates. Second, weather definitions were source-checked rather than reconstructed from memory. Third, all twelve core comparisons are reported as one panel, reducing selective emphasis on the smallest p-values. Fourth, four uncertainty methods are displayed rather than selecting the one that gives the most attractive interval. Fifth, tables, figures and claim text are linked to machine-readable outputs and automated release checks.

4.4 Limitations

The limitations determine how the results should be read:

1. **Outcome meaning.** Admission cause was not recorded. The event is the first hospitalisation after a recorded CHD or HF diagnosis, not necessarily an admission caused by that condition.
2. **Ecological resolution.** Each observation describes Hong Kong for one month. Individual exposure and within-month timing are unknown, so individual or daily causal inference is unavailable ([Greenland and Morgenstern 1989](#)).
3. **Denominator.** The number of cohort members still eligible for a first event in each month was not available. The estimates are count ratios, not cohort incidence-rate ratios.
4. **Confounding.** Season and long-term trend were controlled, but the core panel does not claim complete adjustment for pollution, humidity, influenza, behavioural change, or changes in health care.
5. **Exposure measurement.** Headquarters weather is a territory-level proxy. It cannot represent differences between neighbourhoods, indoor temperatures, air-conditioning access, or personal activity.
6. **Serial dependence and multiplicity.** CHD residuals remained correlated over time, and no core comparison survived correction for twelve tests.
7. **Available outcomes.** Age, sex and disease subgroups were unavailable. A stroke series was not delivered and no stroke result is reported.

5. Implications and next study

A stronger next study would prioritise the outcome data contract rather than add monthly definitions. Dated, cause-recorded admissions, cohort follow-up and numbers still at risk would permit clearer incidence and lag analyses. Age, sex, subtype and location would support vulnerability assessment; pollution, humidity and influenza could enter a staged confounding plan.

The current patterns define two pre-declarable hypotheses: repeated hot nights preceding CHD-related admissions and cold days preceding HF-related admissions.

For practice, the message is broader: Hong Kong faces rising nighttime heat while cold persists. This report neither quantifies preventable admissions nor justifies disease-specific thresholds.

6. Conclusion

Between 2013 and 2023, Hong Kong experienced increasing hot-night burden while cold days persisted. In governed monthly data for people with diabetes and/or hypertension, CHD first-hospitalisation counts were most elevated in relation to hot nights, while HF counts were most elevated in relation to cold days. Neither pattern survived correction across twelve comparisons, and the CHD hot-night result was sensitive to the treatment of uncertainty.

The study therefore offers hypotheses, not proof of thermal effects. Its more durable contribution is a reproducible and aggregation-aware framework: define the event

precisely, compare thermal measures as a panel, show uncertainty, and decline daily or causal claims when the data cannot support them.

Acknowledgements

I thank Hogan for guidance on weather definitions and academic writing; Zhenyuan Liu (“Roro”) for constructing and releasing the governed Hospital Authority aggregates and for regression mentorship; and Professor David Bishai for supervision and population-health guidance. Diagnosis-record construction also benefited from clinical collaboration. External dissemination of governed results remains subject to the appropriate team and institutional approvals.

Appendix A. Complete core panel

Table A1. Separate negative-binomial monthly models with a days-in-month offset and Newey-West lag-6 confidence intervals.

Outcome	Exposure contrast	Count ratio (95% CI)	BH q
CHD	Mean temperature / 1 °C	0.993 (0.976–1.010)	0.726
CHD	Mean maximum temperature / 1 °C	0.993 (0.979–1.007)	0.642
CHD	Mean minimum temperature / 1 °C	0.994 (0.977–1.012)	0.764
CHD	Hot nights / 5 days	1.022 (1.002–1.042)	0.192
CHD	Cold days / 5 days	0.995 (0.949–1.043)	0.900
CHD	Very hot days / 5 days	0.999 (0.974–1.025)	0.962
HF	Mean temperature / 1 °C	0.974 (0.947–1.002)	0.207
HF	Mean maximum temperature / 1 °C	0.981 (0.958–1.005)	0.272
HF	Mean minimum temperature / 1 °C	0.973 (0.947–1.000)	0.202
HF	Hot nights / 5 days	1.003 (0.976–1.031)	0.900
HF	Cold days / 5 days	1.073 (1.006–1.144)	0.192
HF	Very hot days / 5 days	0.995 (0.963–1.028)	0.900

References

- Basagaña, Xavier, and Joan Ballester. 2024. “Unbiased Temperature-Related Mortality Estimates Using Weekly and Monthly Health Data: A New Method for Environmental Epidemiology and Climate Impact Studies.” *The Lancet Planetary Health* 8 (10): e766–77. [https://doi.org/10.1016/S2542-5196\(24\)00212-2](https://doi.org/10.1016/S2542-5196(24)00212-2).
- Goggins, William B., Emily Y. Y. Chan, and Chun-Yuh Yang. 2013. “Weather, Pollution, and Acute Myocardial Infarction in Hong Kong and Taiwan.” *International Journal of Cardiology* 168 (1): 243–49. <https://doi.org/10.1016/j.ijcard.2012.09.087>.
- Goggins, William B., Jean Woo, Suzanne Ho, Emily Y. Y. Chan, and P. H. Chau. 2012. “Weather, Season, and Daily Stroke Admissions in Hong Kong.” *International Journal of Biometeorology* 56 (5): 865–72. <https://doi.org/10.1007/s00484-011-0491-9>.

- Greenland, Sander, and Hal Morgenstern. 1989. "Ecological Bias, Confounding, and Effect Modification." *International Journal of Epidemiology* 18 (1): 269–74. <https://doi.org/10.1093/ije/18.1.269>.
- Guo, Yi Tong, Ka Hung Chan, Hong Qiu, Eliza Lai-Yi Wong, and Kin-Fai Ho. 2024. "The Risk of Hospitalization Associated with Hot Nights and Excess Nighttime Heat in a Subtropical Metropolis: A Time-Series Study in Hong Kong, 2000–2019." *The Lancet Regional Health - Western Pacific* 51: 101168. <https://doi.org/10.1016/j.lanwpc.2024.101168>.
- Li, Changyu, Wenmin Wei, Pak Wai Chan, and Jianping Huang. 2025. "Heatwaves in Hong Kong and Their Influence on Pollution and Extreme Precipitation." *Atmospheric Research* 315: 107845. <https://doi.org/10.1016/j.atmosres.2024.107845>.
- Liu, Jingwen, Alana Hansen, Blesson Varghese, Zhidong Liu, Michael Tong, Hong Qiu, Linwei Tian, et al. 2020. "Cause-Specific Mortality Attributable to Cold and Hot Ambient Temperatures in Hong Kong: A Time-Series Study, 2006–2016." *Sustainable Cities and Society* 57: 102131. <https://doi.org/10.1016/j.scs.2020.102131>.
- Liu, Zhenyuan, Chao Ren, Jingwen Liu, Yurika Kawasaki, and David Makram Bishai. 2026. "Modelling the Excess Mortality Associated with Heat Waves in Hong Kong: 2014–2023." *medRxiv*. <https://doi.org/10.64898/2026.03.05.26347683>.
- Wang, Dan, Kevin Ka-Lun Lau, Chao Ren, William Bernard Goggins, Yuan Shi, Hung Chak Ho, Tsz-Cheung Lee, Lap-Shun Lee, Jean Woo, and Edward Ng. 2019. "The Impact of Extremely Hot Weather Events on All-Cause Mortality in a Highly Urbanized and Densely Populated Subtropical City: A 10-Year Time-Series Study (2006–2015)." *Science of the Total Environment* 690: 923–31. <https://doi.org/10.1016/j.scitotenv.2019.07.039>.