





We thank reviewers [R1](#), [R2](#), [R3](#) for substantial time investment, incisive feedback, and [\[Ba\]](#).

LIMITS. [R1](#) highlights ways our precision must improve. [Thm2](#) states: *“For each d , every non-deg. local min. θ_\star has a nbhd U whose every member θ_0 induces, via [Thm1](#), a sequence $(L_{d,T} : T \in \mathbb{N})$ of truncations, each a degree- d polynomial in η , that converges ptwise as $T \rightarrow \infty$ to some polynomial L_d .”* So if $L_{d,T}(\eta)$ is [Thm1](#)’s truncation, [Thm2](#) controls $L_d(\eta) = \lim_{\tilde{T} \rightarrow \infty} L_{d,\tilde{T}}(\eta)$ but not $L_T(\eta) = \lim_{\tilde{d} \rightarrow \infty} L_{\tilde{d},T}(\eta)$, even for $d, T \gg 1$. **Empirically**, we find that [Thm1/2](#)’s formal power series [\[Wi\]](#) predict SGD-in-practice (w.r.t. which any infinities are idealizations). Regarding our mathematics as but a strong heuristic,¹ we didn’t examine when L_d, L_T agree. Still: **Prop A.** *“Fix $U \subseteq \mathcal{M}$ open, $\theta_\star \in U$ a non-deg. local min. of l . Assume §B.1 and global, prob.-1 bounds $(|l_x(\theta)|, \|\nabla l_x(\theta_\star)\|) < C$. If $\exists Q_-, Q_+ \in \text{SPD}$ bounding the hessian ($Q_- < \nabla^2 l_x(\theta) < Q_+$) on U , then $\forall d$ and $\forall \theta_0$ in some nbhd V_d of θ_\star : $\exists T_0, A, B > 0$ so that $\sup_{T \geq T_0} \text{ReLU}(|L_d(\eta) - L_T(\eta)| - \exp(A - BT))$ exists on some nbhd in SPSD of $\eta = 0$ and is $o(\eta^d)$.”* **SP(S)D** consists of symmetric positive (semi)definites.

SHARP MINIMA. Like us, [R3](#) finds [Cor5](#) counterintuitive.² SGD’s noise consists not of weight **displacements** but of error terms $\nabla l_x - \nabla l$ in the **gradient** estimate; compare [Fig5](#)  to [\[Ke\]](#)’s [Fig1](#). Say θ is 1D with $l(\theta) = a\theta^2/2$ and training loss $\hat{l}(\theta) = l(\theta) + b\theta$. At \hat{l} ’s min. $\theta = -b/a$, $l(\theta) = b^2/(2a)$. So for fixed b , sharp min’a ($a \gg 1$) overfit less ([demo here](#)). C controls b^2 , hence [Cor5](#)’s $C/2H$ factor. Here, opt’z’n to convergence favors sharp min’a (\star); cnv’gnce is slow at flat min’a, so flat min’a also overfit little (\diamond). (Our small- η assumption precludes H from being so sharp that SGD diverges: we treat $\eta H \ll 1$). Prior work ([Pg12Par5](#), e.g. [\[Ke\]](#) and [\[Di\]](#)) supports both pro-flat and pro-sharp intuitions. Recognizing η ’s role in translating gradients to displacements, we account for both (\star, \diamond), unifying existing intuitions (§4.3). It is a merit that our theory makes such counterintuitive phenomena visible.

ODE. [\[Ba\]](#)’s [LemA.3](#) specializes [LemKey](#). In our terms, [\[Ba\]](#)’s [Thm3.1](#) computes η^2 weight displacements using *fuzzless* diagrams (noiseless \equiv cumulants vanish \equiv fuzzy diagrams vanish); see [Tab1](#) for the leading corrections to [\[Ba\]](#) due to noise.. Per §A.6 (fix $E=B=1$), GD displaces θ by $\Delta_{GD}^l(\eta, T) = -T \cdot \frac{1}{2} + \frac{1}{2} \cdot \frac{1}{2} + o(\eta^2)$. Now, [2](#)  $= \nabla^\mu$ , whence arises [\[Ba\]](#)’s $\lambda R = \eta G^2/4 = \frac{1}{2} \cdot \frac{1}{2} / (2 \cdot 2)$. Our [Cor1](#) is analogous to [\[Ba\]](#)’s [Thm3.1](#). See [our analysis](#).

NOTATION. [R3](#) recognizes our expectands as tensor expressions; they are often fully contracted (so scalar) and are always random variables in an \mathbb{R}^k . Per [R2, R3](#), we’ll disemploy ‘Einstein notation’ and cite [\[Cu\]](#) (+ a new §D) for tensor examples. As [R3](#) asked and [R2](#) noted, diagrams organize ([Pg5Par\(-1\)](#)) an otherwise unwieldy ([Pg4Par2](#)) analysis; if advised, we’ll text-ify diagrams: e.g. $[a][ab : c : d][bcd]$ for  (letters name edges). [R3](#), [Pg6Thm2](#) defines ‘non-degenerate’ as ‘ $H \in \text{SPD}$ ’.

ORGANIZATION. [R2, R3](#) stress the paper’s narrative challenge. We’ll arrange the paper (see [revision in progress](#)) into 3 self-contained tracks, each pertinent to a different goal: [TrkA](#) [[Pg1-4](#)], for casual readers, will eschew diagrams, theorems, and §1.1/§2.2’s heavy notations; illustrate Taylor series via §2.1’s proof; identify §3.3’s terms; state [Cor4](#) (w/ §B.1’s assumptions explicit, w/ [PrpA](#)’s precision); explain §4.2’s curl effect. [TrkB](#) [[Pg1-4, 5-12](#)], for seekers of physical intuition, will use [TrkA](#) to motivate (& §A.4 to illustrate) §1.1/§2.2’s def’ns; relegate §2.2/2.1’s [LemKey](#)/discussion to §B; add to §2.3.1 a resum’n cartoon à la [Fig5, 7](#). For space, §C will absorb §4. [TrkC](#) [[Pg5-12, 15-](#)], for our theory’s extenders, will include [PrpA](#) (per [R1](#)) and more explicit statements and proofs throughout.

REFERENCES. [\[Ba\]](#) D.G.Barrett, B.Dherin. *Implicit Gradient Regularization*. ICLR 2021. [\[Cu\]](#) P.McCullagh. *Tensor Methods in Statistics*, §1.1-1.4, §1.8. Dover 2017. [\[Di\]](#) L.Dinh, R.Pascanu, S.Bengio, Y.Bengio. *Sharp Minima Can Generalize for Deep Nets*, §1, §5. ICML 2017. [\[Ke\]](#) N.S.Keskar et alia. *Large-Batch Training for Deep Learning*, §4. ICLR 2017. [\[Wi\]](#) H.Wilf. *Generatingfunctionology*, §2.1-2.3. Academic Press 1994.

1. By cl’cal.mech. (CM) of thermal continua, ice cubes have energy= ∞ [[McQuarrie ’97](#), §1-1]. Still, CM gives insight.

2. i.e.: that **overfitting** ($\triangleq l(\theta_T) - l(\theta_0)$ where θ_0 is a min. of l) has an η^2 term greatest when ηH has moderate eigenvalues.