CAMBRIDGE INTERNATIONAL EXAMINATIONS

Cambridge International General Certificate of Secondary Education

www.PapaCambridge.com MARK SCHEME for the October/November 2014 series

0620 CHEMISTRY

0620/33

Paper 3 (Extended Theory), maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2014 series for most Cambridge IGCSE®, Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.

BB CAMBRIDGE



P	age 2	Mark Scheme	Sylvanoer		
	<u> </u>	Cambridge IGCSE – October/November 2014	Sylvaria per 062 Page 1		
1	(a)	Mark Scheme Cambridge IGCSE – October/November 2014 Bromine Physical: reddish-brown liquid or brown liquid or volatile liquid/low boiling point liquid or poor/non-conductor (of electricity) or soluble in water or soluble in organic/non-polar solvents			
		Chemical: Reacts with water or reacts with iodides (in solution) or dispidence or reacts with alkenes/named alkene/unsaturated hydrocarbons with alkane in UV/named alkane in UV or valency/oxidation state(–)1 or Br or gains or shares 1 electron or combines or reacts with metals/na or combines or reacts with non-metals/named non-metal or oxidising a bleaches litmus paper/indicator paper or corrosive or forms acidic oxide	or reacts r forms med metal gent or		
	(b)	Graphite Physical: (good) conductor (of electricity) or soft or lubricant or high moint/high boiling point or grey black or black solid or slippery or greasy touch) or brittle/breaks when subjected to stress or insoluble in water			
		Chemical: reducing agent or reduces metal oxides/named metal oxide with/burns in air/oxygen or forms an acidic oxide (CO ₂) or valency/oxide of 2 or 4			
	(c)	 (c) Manganese Physical: (good) conductor (of heat/electricity) or high melting point/high boiling point or forms coloured compounds/coloured ions or hard or strong or high density or malleable or ductile or sonorous or shiny Chemical: Variable or different valency/oxidation state/oxidation number or catalytic activity or forms coloured compounds/coloured ions or forms complex ions/complexes or reacts with acids or reducing agent or reacts with non-metals 			
			[Total: 6]		
2	(a)	(i) $(X(s) \leftrightarrow) X(I)$	[1]		
		(ii) melting point/freezing point (of X)	[1]		
	(iii) gas/gaseous or vapour	[1]		
	(iv) not horizontal or line slopes or line is lower	[1]		

www.PapaCambridge.com **Mark Scheme** Page 3 Cambridge IGCSE - October/November 2014

- **(b) (i)** 14.3
 - (ii) $85.7 \div 12$ and $14.3 \div 1$ or 7.14 and 14.3ratio 1:2 CH₂

note: Award all 3 marks for correct answer

allow: alternative working e.g.

$$85.7 \times 84 \div 100$$
 and $14.3 \times 84 \div 100$ or $71.988/72$ and $12/12.012$

6:12 **or** ratio 1:2

CH₂

(iii) C₆H₁₂ [1]

- 3 (a) (i) 3 [1]
 - (ii) 70 [1]
 - **(b)** Add octane (or other liquid hydrocarbon) (to soot) [1]
 - COND(on addition of any solvent) filter (to remove insoluble forms of carbon) [1]
 - (allow to) evaporate **or** heat **or** warm **or** leave in sun(to get crystals of fullerene) [1]
 - (c) (i) graphite [1]
 - (ii) delocalised electrons/free electrons/sea of electrons [1]
 - COND (on electrons) move/mobile/electrons flow [1]
 - [2] (iii) Any **two** from: potassium oxide potassium hydroxide potassium carbonate

potassium hydrogencarbonate (bicarbonate)

(a) carbon dioxide/CO₂ [1]

- **(b)** $2H_2 + O_2 \rightarrow 2H_2O$ [1]
- (c) (i) anode/negative electrode and electrons lost(by hydrogen/H/H₂)/electrons move from this electrode [1]
 - (ii) $H_2 \rightarrow 2H^+ + 2e(^-) / H_2 2e(^-) \rightarrow 2H^+ / H_2 + 2OH^- \rightarrow$ $2H_2O + 2e(^-) / H_2 + 2OH^- - 2e(^-) \rightarrow 2H_2O$ [2] Species (1) Balancing (1)

[Total: 10]

[1]

[1]

[1]

[Total: 9]

Page 4			Mark Scheme Sy.	per
		Cambridg	e IGCSE – October/November 2014 062	ASC.
; ;	Any two from: CELL: SUSTAINABILITY: POLLUTION:		produced conserves a limited resource/petroleum/fossil fuels unlimited supplies of renewable resource(of hydrogen from water) No or less greenhouse effect No or less toxic gases No or less smog No or less C/soot	O ADAC ANDRIGUE
			No or less CO ₂ No or less CO No or less SO ₂ No or less oxides of nitrogen/NO/NO ₂ /N ₂ O ₄ /NO _x No or less (unburnt) hydrocarbons No or less low level ozone H ₂ O is the only product	[2]
				[Total: 7]
5 (a) ((i)	rate decreases concentration of so	odium chlorate ((I))/reactant decreases	[1] [1]
(i	ii)	(initial) gradient gr same final volume	eater/steeper (must start at origin) of oxygen	[1] [1]
(ii	ii)		hemical reaction/(to prevent)reaction catalysed by own or decomposes sodium chlorate((I))	[1]
(i	v)	particles have mor more collisions	re energy/particles move faster/	[1]
		collisions more fre	quent or more often/greater chance of collision/collision re particles have energy to react/more collisions are ctive	[1]
(b)	(i)	$2CT \rightarrow Cl_2 + 2e$	$e(\bar{\ })$ / $2Cl^-$ – $2e(\bar{\ })$ $ ightarrow$ Cl_2	[1]
		$2H^{+} + 2e(^{-}) \rightarrow F$	$H_2 / 2H^+ \rightarrow H_2 - 2e()$	[1]
		hydrogen formed a	at cathode/– and chlorine at anode/+	[1]
		<u>Na[⁺] and OH⁻</u> or so sodium hydroxide	odium <u>ions</u> and hydroxide <u>ions</u> left in solution/form/becom	e [1]
((ii)	Cl_2 + 2NaOH \rightarrow Species (1) Baland	NaClO/NaOCl + NaCl + H₂O cing (1)	[2]
				[Total: 14]

Page 5	Mark Scheme	Syl per
	Cambridge IGCSE – October/November 2014	062

6 (a) Rb loses 1 electron/1 electron in outer shell/1 valency or valence electron

Sr loses 2 electrons/2 electrons in outer shell/2 valency or valence electrons

(b) (i) (mix solutions of) rubidium carbonate/Rb₂CO₃ [1]

strontium chloride/SrC l_2 or strontium nitrate/Sr(NO₃)₂ or strontium sulfate/SrSO₄ or strontium hydroxide/Sr(OH)₂ [1]

COND (on two correct reactants) filter **or** centrifuge **or** decant (the residue) [1]

wash <u>with water</u> **and** dry/press between filter paper/put in (low) oven/put on a (sunny) windowsill/put in sun/heat [1]


(ii) $SrCO_3 \rightarrow SrO + CO_2$ [1]

- (c) (i) rubidium nitr<u>ite</u> or nitr<u>ate(III)</u> [1]
 - (ii) $2Sr(NO_3)_2 \rightarrow 2SrO + 4NO_2 + O_2$ [2] Species (1) Balancing (1)

[Total: 10]

7 (a) (i) butanoic acid/butyric acid [1]

displayed formula below [2]



(ii) any three from:

same or similar chemical properties (same) general (molecular) formula (consecutive members) differ by CH₂ same functional group common methods of preparation physical properties vary in predictable manner/show trends/gradually change **or** example of a physical property variation i.e. melting point/boiling point/volatility

(iii) dissociates/ionises/splits up (into ions) [1]

partially/incompletely/slightly/not fully [1]

(donates) protons/(forms) $H^{+}/H_{3}O^{+}$ (as the only positive ion) [1]

[3]

Page 6		6	Mark Scheme Syl		per
			Cambridge IGCSE – October/November 2014	Sy. 062	OSO I
	(b)	(i)	methyl propanoate		Da Cambridg
			$CH_3CH_2COOCH_3/CH_3CH_2CO_2CH_3/C_2H_5COOCH_3/C_2H_5CO_2CH_3$		196
		(ii)	methyl ethanoate		[1]
	(c)	(i)	$3C_4H_{10} + 5 \frac{1}{2}O_2 \rightarrow 4C_2H_5COOH + 3 H_2O$		[1]
		(ii)	propanol or propan-1-ol or propanal		[1]
					[Total: 14]
8	(a)	(ch	anges from) blue (1) to pink (1)		[2]
	(b)		more (solid) dissolves or no more cobalt(II) carbonate dissolves or rervescence or bubbling or fizzing	no more	[1]
		filte	er(residue)/centrifuge/decant		[1]
		gor	aporate/heat/warm/boil/leave in sun AND until most of the water has ne/some water is left/until it is concentrated/saturation (point)/crystall nt/crystals form on glass rod or microscope slide/crystals start to form		[1]
		wa	ave/allow to cool/allow to crystallise/filter (off crystals)/wash(with distiter)/dry crystals with filter paper/dry crystals in warm place or dry in con windowsill		[1]
	(c)	nur	mber of moles of HC l in 50 cm 3 of acid, concentration 2.2 mol/dm 3 =	0.11	[1]
		ma	ximum number of moles of $CoCl_2.6H_2O$ which could be formed = 0.0	55	[1]
		ma	ss of 1 mole of $CoCl_2.6H_2O = 238g$		
		ma	ximum yield of $CoCl_2.6H_2O = 13.09g$		[1]
		per <u>dp</u>	centage yield = 48.2% or ecf mass of CoC l_2 .6H ₂ O above/13.09 \times 10	00% to <u>1</u>	[1]
					[Total: 10]