

An introduction to the p -adics

Siddharth Bhat

**IIIT Theory group
Seminar Saturday**

October 10th, 2019

Why p -adics?

Analogy between:

- \mathbb{Z} ,

Why p -adics?

Analogy between:

- \mathbb{Z} , where $3, 5, 7, \dots$ are the “primes”

Why p -adics?

Analogy between:

- \mathbb{Z} , where $3, 5, 7, \dots$ are the “primes”
- $\mathbb{C}[X]$,

Why p -adics?

Analogy between:

- \mathbb{Z} , where $3, 5, 7, \dots$ are the “primes”
- $\mathbb{C}[X]$, where $(x - a)$ are the “primes”

What is evaluation?

Remainder when dividing $p(x) = x^3 + x^2 + x + 1$ by $q(x) = x - 1$?

What is evaluation?

Remainder when dividing $p(x) = x^3 + x^2 + x + 1$ by $q(x) = x - 1$?

$$\begin{array}{r}
 X^2 + 2X + 3 \\
 \hline
 X-1) X^2 X + 1 \\
 \underline{-X^3 X^2} X + 1 \\
 2X^2 X + 1 \\
 \underline{-2X^2 + 2X} 1 \\
 3X + 1 \\
 \underline{-3X + 3} \\
 4
 \end{array}$$

What is evaluation?

Remainder when dividing $p(x) = x^3 + x^2 + x + 1$ by $q(x) = x - 1$?

$$\begin{array}{r}
 X^2 + 2X + 3 \\
 \hline
 X-1) X^2 X + 1 \\
 \underline{-X^3 X^2} X + 1 \\
 2X^2 X + 1 \\
 \underline{-2X^2 + 2X} 1 \\
 3X + 1 \\
 \underline{-3X + 3} \\
 4
 \end{array}$$

$$(x^3 + x^2 + x + 1) = (x - 1)(x^2 + 2x + 3) + 4$$

What is evaluation?

Remainder when dividing $p(x) = x^3 + x^2 + x + 1$ by $q(x) = x - 1$?

$$\begin{array}{r}
 X^2 + 2X + 3 \\
 \hline
 X-1) + X^2 + X + 1 \\
 - X^3 + X^2 \\
 \hline
 2X^2 + X \\
 - 2X^2 + 2X \\
 \hline
 3X + 1 \\
 - 3X + 3 \\
 \hline
 4
 \end{array}$$

$$(x^3 + x^2 + x + 1) = (x - 1)(x^2 + 2x + 3) + 4$$

■ $p(1) = 1^3 + 1^2 + 1 + 1 = 4$. Coincidence?

What is evaluation?

Remainder when dividing $p(x) = x^3 + x^2 + x + 1$ by $q(x) = x - 1$?

$$\begin{array}{r}
 X^2 + 2X + 3 \\
 \hline
 X-1) + X^2 + X + 1 \\
 \underline{-X^3 + X^2} \\
 2X^2 + X \\
 \underline{-2X^2 + 2X} \\
 3X + 1 \\
 \underline{-3X + 3} \\
 4
 \end{array}$$

$$(x^3 + x^2 + x + 1) = (x - 1)(x^2 + 2x + 3) + 4$$

- $p(1) = 1^3 + 1^2 + 1 + 1 = 4$. Coincidence?
- Factoring out $q(x) = (x - 1)$

What is evaluation?

Remainder when dividing $p(x) = x^3 + x^2 + x + 1$ by $q(x) = x - 1$?

$$\begin{array}{r}
 X^2 + 2X + 3 \\
 \hline
 X-1) X^2 X + 1 \\
 \underline{-X^3 X^2} X + 1 \\
 2X^2 X 1 \\
 \underline{-2X^2 + 2X} 1 \\
 3X 1 \\
 \underline{-3X + 3} \\
 4
 \end{array}$$

$$(x^3 + x^2 + x + 1) = (x - 1)(x^2 + 2x + 3) + 4$$

- $p(1) = 1^3 + 1^2 + 1 + 1 = 4$. Coincidence?
- Factoring out $q(x) = (x - 1) \simeq$ setting $q(x) = 0$

What is evaluation?

Remainder when dividing $p(x) = x^3 + x^2 + x + 1$ by $q(x) = x - 1$?

$$\begin{array}{r}
 X^2 + 2X + 3 \\
 \hline
 X-1) X^2 X + 1 \\
 \underline{-X^3 X^2} X + 1 \\
 2X^2 X 1 \\
 \underline{-2X^2 + 2X} 1 \\
 3X 1 \\
 \underline{-3X + 3} \\
 4
 \end{array}$$

$$(x^3 + x^2 + x + 1) = (x - 1)(x^2 + 2x + 3) + 4$$

- $p(1) = 1^3 + 1^2 + 1 + 1 = 4$. Coincidence?
- Factoring out $q(x) = (x - 1) \simeq$ setting $q(x) = 0$: remove $q(x)$.

What is evaluation?

Remainder when dividing $p(x) = x^3 + x^2 + x + 1$ by $q(x) = x - 1$?

$$\begin{array}{r}
 X^2 + 2X + 3 \\
 \hline
 X-1) X^2 X + 1 \\
 \underline{-X^3 X^2} X + 1 \\
 2X^2 X + 1 \\
 \underline{-2X^2 + 2X} 1 \\
 3X 1 \\
 \underline{-3X + 3} \\
 4
 \end{array}$$

$$(x^3 + x^2 + x + 1) = (x - 1)(x^2 + 2x + 3) + 4$$

- $p(1) = 1^3 + 1^2 + 1 + 1 = 4$. Coincidence?
- Factoring out $q(x) = (x - 1) \simeq$ setting $q(x) = 0$: remove $q(x)$.
- setting $x - 1 = 0$, or setting $x = 1$

What is evaluation?

Remainder when dividing $p(x) = x^3 + x^2 + x + 1$ by $q(x) = x - 1$?

$$\begin{array}{r}
 X^2 + 2X + 3 \\
 \hline
 X-1) + X^2 + X + 1 \\
 \underline{- X^3 + X^2} \\
 2X^2 + X \\
 \underline{- 2X^2 + 2X} \\
 3X + 1 \\
 \underline{- 3X + 3} \\
 4
 \end{array}$$

$$(x^3 + x^2 + x + 1) = (x - 1)(x^2 + 2x + 3) + 4$$

- $p(1) = 1^3 + 1^2 + 1 + 1 = 4$. Coincidence?
- Factoring out $q(x) = (x - 1) \simeq$ setting $q(x) = 0$: remove $q(x)$.
- setting $x - 1 = 0$, or setting $x = 1$
- Substituting $x = 1$: $p(1) = 1^3 + 1^2 + 1 + 1 = 4$

What is evaluation?

Remainder when dividing $p(x) = x^3 + x^2 + x + 1$ by $q(x) = x - 1$?

$$\begin{array}{r}
 X^2 + 2X + 3 \\
 X-1 \overline{) X^3 + X^2 + X + 1} \\
 \underline{-X^3 + X^2} \\
 2X^2 + X \\
 \underline{-2X^2 + 2X} \\
 3X + 1 \\
 \underline{-3X + 3} \\
 4
 \end{array}$$

$$(x^3 + x^2 + x + 1) = (x - 1)(x^2 + 2x + 3) + 4$$

- $p(1) = 1^3 + 1^2 + 1 + 1 = 4$. Coincidence?
- Factoring out $q(x) = (x - 1) \simeq$ setting $q(x) = 0$: remove $q(x)$.
- setting $x - 1 = 0$, or setting $x = 1$
- Substituting $x = 1$: $p(1) = 1^3 + 1^2 + 1 + 1 = 4$

remainder of $p(x)$ on factoring $(x - a) \simeq$ evaluation of $p(x_0)$ at $x_0 = a$

What is evaluation in \mathbb{Z} ?

remainder of $p(x)$ on factoring $(x - a) \simeq$ evaluation of $p(x_0)$ at $x_0 = a$

What is evaluation in \mathbb{Z} ?

remainder of $p(x)$ on factoring $(x - a) \simeq$ evaluation of $p(x_0)$ at $x_0 = a$

evaluation of $p(x_0)$ at $x_0 = a \simeq$ remainder of $p(x)$ on factoring $(x - a)$

What is evaluation in \mathbb{Z} ?

remainder of $p(x)$ on factoring $(x - a) \simeq$ evaluation of $p(x_0)$ at $x_0 = a$

evaluation of $p(x_0)$ at $x_0 = a \simeq$ remainder of $p(x)$ on factoring $(x - a)$

■ 10(2)

What is evaluation in \mathbb{Z} ?

remainder of $p(x)$ on factoring $(x - a) \simeq$ evaluation of $p(x_0)$ at $x_0 = a$

evaluation of $p(x_0)$ at $x_0 = a \simeq$ remainder of $p(x)$ on factoring $(x - a)$

- $10(2) =$ remainder of 10 when factored by 2;

What is evaluation in \mathbb{Z} ?

remainder of $p(x)$ on factoring $(x - a) \simeq$ evaluation of $p(x_0)$ at $x_0 = a$

evaluation of $p(x_0)$ at $x_0 = a \simeq$ remainder of $p(x)$ on factoring $(x - a)$

- $10(2) =$ remainder of 10 when factored by 2; $10 = 2 \cdot 5 + 0$

What is evaluation in \mathbb{Z} ?

remainder of $p(x)$ on factoring $(x - a) \simeq$ evaluation of $p(x_0)$ at $x_0 = a$

evaluation of $p(x_0)$ at $x_0 = a \simeq$ remainder of $p(x)$ on factoring $(x - a)$

- $10(2) =$ remainder of 10 when factored by 2; $10 = 2 \cdot 5 + 0$; $10(2) = 0$

What is evaluation in \mathbb{Z} ?

remainder of $p(x)$ on factoring $(x - a) \simeq$ evaluation of $p(x_0)$ at $x_0 = a$

evaluation of $p(x_0)$ at $x_0 = a \simeq$ remainder of $p(x)$ on factoring $(x - a)$

- $10(2)$ = remainder of 10 when factored by 2; $10 = 2 \cdot 5 + 0$; $10(2) = 0$
- $10(3)$

What is evaluation in \mathbb{Z} ?

remainder of $p(x)$ on factoring $(x - a) \simeq$ evaluation of $p(x_0)$ at $x_0 = a$

evaluation of $p(x_0)$ at $x_0 = a \simeq$ remainder of $p(x)$ on factoring $(x - a)$

- $10(2) =$ remainder of 10 when factored by 2; $10 = 2 \cdot 5 + 0$; $10(2) = 0$
- $10(3) =$ remainder of 10 when factored by 3;

What is evaluation in \mathbb{Z} ?

remainder of $p(x)$ on factoring $(x - a) \simeq$ evaluation of $p(x_0)$ at $x_0 = a$

evaluation of $p(x_0)$ at $x_0 = a \simeq$ remainder of $p(x)$ on factoring $(x - a)$

- $10(2) =$ remainder of 10 when factored by 2; $10 = 2 \cdot 5 + 0$; $10(2) = 0$
- $10(3) =$ remainder of 10 when factored by 3; $10 = 3 \cdot 3 + 1$

What is evaluation in \mathbb{Z} ?

remainder of $p(x)$ on factoring $(x - a) \simeq$ evaluation of $p(x_0)$ at $x_0 = a$

evaluation of $p(x_0)$ at $x_0 = a \simeq$ remainder of $p(x)$ on factoring $(x - a)$

- $10(2) =$ remainder of 10 when factored by 2; $10 = 2 \cdot 5 + 0$; $10(2) = 0$
- $10(3) =$ remainder of 10 when factored by 3; $10 = 3 \cdot 3 + 1$; $10(3) = 1$

What is evaluation in \mathbb{Z} ?

remainder of $p(x)$ on factoring $(x - a) \simeq$ evaluation of $p(x_0)$ at $x_0 = a$

evaluation of $p(x_0)$ at $x_0 = a \simeq$ remainder of $p(x)$ on factoring $(x - a)$

- $10(2) =$ remainder of 10 when factored by 2; $10 = 2 \cdot 5 + 0$; $10(2) = 0$
- $10(3) =$ remainder of 10 when factored by 3; $10 = 3 \cdot 3 + 1$; $10(3) = 1$
- $10(5)$

What is evaluation in \mathbb{Z} ?

remainder of $p(x)$ on factoring $(x - a) \simeq$ evaluation of $p(x_0)$ at $x_0 = a$

evaluation of $p(x_0)$ at $x_0 = a \simeq$ remainder of $p(x)$ on factoring $(x - a)$

- $10(2) =$ remainder of 10 when factored by 2; $10 = 2 \cdot 5 + 0$; $10(2) = 0$
- $10(3) =$ remainder of 10 when factored by 3; $10 = 3 \cdot 3 + 1$; $10(3) = 1$
- $10(5) =$ remainder of 10 when factored by 5;

What is evaluation in \mathbb{Z} ?

remainder of $p(x)$ on factoring $(x - a) \simeq$ evaluation of $p(x_0)$ at $x_0 = a$

evaluation of $p(x_0)$ at $x_0 = a \simeq$ remainder of $p(x)$ on factoring $(x - a)$

- $10(2) =$ remainder of 10 when factored by 2; $10 = 2 \cdot 5 + 0$; $10(2) = 0$
- $10(3) =$ remainder of 10 when factored by 3; $10 = 3 \cdot 3 + 1$; $10(3) = 1$
- $10(5) =$ remainder of 10 when factored by 5; $10 = 5 \cdot 2 + 0$

What is evaluation in \mathbb{Z} ?

remainder of $p(x)$ on factoring $(x - a) \simeq$ evaluation of $p(x_0)$ at $x_0 = a$

evaluation of $p(x_0)$ at $x_0 = a \simeq$ remainder of $p(x)$ on factoring $(x - a)$

- $10(2) =$ remainder of 10 when factored by 2; $10 = 2 \cdot 5 + 0$; $10(2) = 0$
- $10(3) =$ remainder of 10 when factored by 3; $10 = 3 \cdot 3 + 1$; $10(3) = 1$
- $10(5) =$ remainder of 10 when factored by 5; $10 = 5 \cdot 2 + 0$; $10(5) = 0$

What is evaluation in \mathbb{Z} ?

remainder of $p(x)$ on factoring $(x - a) \simeq$ evaluation of $p(x_0)$ at $x_0 = a$

evaluation of $p(x_0)$ at $x_0 = a \simeq$ remainder of $p(x)$ on factoring $(x - a)$

- $10(2) =$ remainder of 10 when factored by 2; $10 = 2 \cdot 5 + 0$; $10(2) = 0$
- $10(3) =$ remainder of 10 when factored by 3; $10 = 3 \cdot 3 + 1$; $10(3) = 1$
- $10(5) =$ remainder of 10 when factored by 5; $10 = 5 \cdot 2 + 0$; $10(5) = 0$
- $10(7)$

What is evaluation in \mathbb{Z} ?

remainder of $p(x)$ on factoring $(x - a) \simeq$ evaluation of $p(x_0)$ at $x_0 = a$

evaluation of $p(x_0)$ at $x_0 = a \simeq$ remainder of $p(x)$ on factoring $(x - a)$

- $10(2) =$ remainder of 10 when factored by 2; $10 = 2 \cdot 5 + 0$; $10(2) = 0$
- $10(3) =$ remainder of 10 when factored by 3; $10 = 3 \cdot 3 + 1$; $10(3) = 1$
- $10(5) =$ remainder of 10 when factored by 5; $10 = 5 \cdot 2 + 0$; $10(5) = 0$
- $10(7) =$ remainder of 10 when factored by 7;

What is evaluation in \mathbb{Z} ?

remainder of $p(x)$ on factoring $(x - a) \simeq$ evaluation of $p(x_0)$ at $x_0 = a$

evaluation of $p(x_0)$ at $x_0 = a \simeq$ remainder of $p(x)$ on factoring $(x - a)$

- $10(2) =$ remainder of 10 when factored by 2; $10 = 2 \cdot 5 + 0$; $10(2) = 0$
- $10(3) =$ remainder of 10 when factored by 3; $10 = 3 \cdot 3 + 1$; $10(3) = 1$
- $10(5) =$ remainder of 10 when factored by 5; $10 = 5 \cdot 2 + 0$; $10(5) = 0$
- $10(7) =$ remainder of 10 when factored by 7; $10 = 7 \cdot 1 + 3$

What is evaluation in \mathbb{Z} ?

remainder of $p(x)$ on factoring $(x - a) \simeq$ evaluation of $p(x_0)$ at $x_0 = a$

evaluation of $p(x_0)$ at $x_0 = a \simeq$ remainder of $p(x)$ on factoring $(x - a)$

- $10(2) =$ remainder of 10 when factored by 2; $10 = 2 \cdot 5 + 0$; $10(2) = 0$
- $10(3) =$ remainder of 10 when factored by 3; $10 = 3 \cdot 3 + 1$; $10(3) = 1$
- $10(5) =$ remainder of 10 when factored by 5; $10 = 5 \cdot 2 + 0$; $10(5) = 0$
- $10(7) =$ remainder of 10 when factored by 7; $10 = 7 \cdot 1 + 3$; $10(7) = 3$

What is evaluation in \mathbb{Z} ?

remainder of $p(x)$ on factoring $(x - a) \simeq$ evaluation of $p(x_0)$ at $x_0 = a$

evaluation of $p(x_0)$ at $x_0 = a \simeq$ remainder of $p(x)$ on factoring $(x - a)$


- $10(2) =$ remainder of 10 when factored by 2; $10 = 2 \cdot 5 + 0$; $10(2) = 0$
- $10(3) =$ remainder of 10 when factored by 3; $10 = 3 \cdot 3 + 1$; $10(3) = 1$
- $10(5) =$ remainder of 10 when factored by 5; $10 = 5 \cdot 2 + 0$; $10(5) = 0$
- $10(7) =$ remainder of 10 when factored by 7; $10 = 7 \cdot 1 + 3$; $10(7) = 3$

What is evaluation in \mathbb{Z} ?

remainder of $p(x)$ on factoring $(x - a) \simeq$ evaluation of $p(x_0)$ at $x_0 = a$

evaluation of $p(x_0)$ at $x_0 = a \simeq$ remainder of $p(x)$ on factoring $(x - a)$

- $10(2) =$ remainder of 10 when factored by 2; $10 = 2 \cdot 5 + 0$; $10(2) = 0$
- $10(3) =$ remainder of 10 when factored by 3; $10 = 3 \cdot 3 + 1$; $10(3) = 1$
- $10(5) =$ remainder of 10 when factored by 5; $10 = 5 \cdot 2 + 0$; $10(5) = 0$
- $10(7) =$ remainder of 10 when factored by 7; $10 = 7 \cdot 1 + 3$; $10(7) = 3$



Why $n(p)$: only primes?

- $10(5)$ = remainder of 10 when factored by 5; $10 = 5 \cdot 2 + 0$; $10(5) = 0$
- $10(7)$ = remainder of 10 when factored by 7; $10 = 7 \cdot 1 + 3$; $10(7) = 3$
- $p(x) = (x^2 - 15x + 50)$.

Why $n(p)$: only primes?

- $10(5)$ = remainder of 10 when factored by 5; $10 = 5 \cdot 2 + 0$; $10(5) = 0$
- $10(7)$ = remainder of 10 when factored by 7; $10 = 7 \cdot 1 + 3$; $10(7) = 3$
- $p(x) = (x^2 - 15x + 50)$.
- $p(5) =$

Why $n(p)$: only primes?

- $10(5) =$ remainder of 10 when factored by 5; $10 = 5 \cdot 2 + 0$; $10(5) = 0$
- $10(7) =$ remainder of 10 when factored by 7; $10 = 7 \cdot 1 + 3$; $10(7) = 3$
- $p(x) = (x^2 - 15x + 50)$.
- $p(5) =$ remainder of $p(x)$ when factored by $(x - 5)$;

Why $n(p)$: only primes?

- $10(5) =$ remainder of 10 when factored by 5; $10 = 5 \cdot 2 + 0$; $10(5) = 0$
- $10(7) =$ remainder of 10 when factored by 7; $10 = 7 \cdot 1 + 3$; $10(7) = 3$
- $p(x) = (x^2 - 15x + 50)$.
- $p(5) =$ remainder of $p(x)$ when factored by $(x - 5)$;
- $p(x) = (x - 5)(x - 10) + 0$;

Why $n(p)$: only primes?

- $10(5) =$ remainder of 10 when factored by 5; $10 = 5 \cdot 2 + 0$; $10(5) = 0$
- $10(7) =$ remainder of 10 when factored by 7; $10 = 7 \cdot 1 + 3$; $10(7) = 3$
- $p(x) = (x^2 - 15x + 50)$.
- $p(5) =$ remainder of $p(x)$ when factored by $(x - 5)$;
- $p(x) = (x - 5)(x - 10) + 0$; $p(5) = 0$

Why $n(p)$: only primes?

- $10(5) =$ remainder of 10 when factored by 5; $10 = 5 \cdot 2 + 0$; $10(5) = 0$
- $10(7) =$ remainder of 10 when factored by 7; $10 = 7 \cdot 1 + 3$; $10(7) = 3$
- $p(x) = (x^2 - 15x + 50)$.
- $p(5) =$ remainder of $p(x)$ when factored by $(x - 5)$;
- $p(x) = (x - 5)(x - 10) + 0$; $p(5) = 0$
- $p(1) =$

Why $n(p)$: only primes?

- $10(5) =$ remainder of 10 when factored by 5; $10 = 5 \cdot 2 + 0$; $10(5) = 0$
- $10(7) =$ remainder of 10 when factored by 7; $10 = 7 \cdot 1 + 3$; $10(7) = 3$
- $p(x) = (x^2 - 15x + 50)$.
- $p(5) =$ remainder of $p(x)$ when factored by $(x - 5)$;
- $p(x) = (x - 5)(x - 10) + 0$; $p(5) = 0$
- $p(1) =$ remainder of $p(x)$ when factored by $(x - 1)$;

Why $n(p)$: only primes?

- $10(5) =$ remainder of 10 when factored by 5; $10 = 5 \cdot 2 + 0$; $10(5) = 0$
- $10(7) =$ remainder of 10 when factored by 7; $10 = 7 \cdot 1 + 3$; $10(7) = 3$
- $p(x) = (x^2 - 15x + 50)$.
- $p(5) =$ remainder of $p(x)$ when factored by $(x - 5)$;
- $p(x) = (x - 5)(x - 10) + 0$; $p(5) = 0$
- $p(1) =$ remainder of $p(x)$ when factored by $(x - 1)$;
- $p(x) = (x - 1)(x - 14) + 36$;

Why $n(p)$: only primes?

- $10(5) =$ remainder of 10 when factored by 5; $10 = 5 \cdot 2 + 0$; $10(5) = 0$
- $10(7) =$ remainder of 10 when factored by 7; $10 = 7 \cdot 1 + 3$; $10(7) = 3$
- $p(x) = (x^2 - 15x + 50)$.
- $p(5) =$ remainder of $p(x)$ when factored by $(x - 5)$;
- $p(x) = (x - 5)(x - 10) + 0$; $p(5) = 0$
- $p(1) =$ remainder of $p(x)$ when factored by $(x - 1)$;
- $p(x) = (x - 1)(x - 14) + 36$; $p(1) = 36$

Why $n(p)$: only primes?

- $10(5) =$ remainder of 10 when factored by 5; $10 = 5 \cdot 2 + 0$; $10(5) = 0$
- $10(7) =$ remainder of 10 when factored by 7; $10 = 7 \cdot 1 + 3$; $10(7) = 3$
- $p(x) = (x^2 - 15x + 50)$.
- $p(5) =$ remainder of $p(x)$ when factored by $(x - 5)$;
- $p(x) = (x - 5)(x - 10) + 0$; $p(5) = 0$
- $p(1) =$ remainder of $p(x)$ when factored by $(x - 1)$;
- $p(x) = (x - 1)(x - 14) + 36$; $p(1) = 36$

Why $n(p)$: only primes?

- $10(5) =$ remainder of 10 when factored by 5; $10 = 5 \cdot 2 + 0$; $10(5) = 0$
- $10(7) =$ remainder of 10 when factored by 7; $10 = 7 \cdot 1 + 3$; $10(7) = 3$
- $p(x) = (x^2 - 15x + 50)$.
- $p(5) =$ remainder of $p(x)$ when factored by $(x - 5)$;
- $p(x) = (x - 5)(x - 10) + 0$; $p(5) = 0$
- $p(1) =$ remainder of $p(x)$ when factored by $(x - 1)$;
- $p(x) = (x - 1)(x - 14) + 36$; $p(1) = 36$

Theorem (Fundamental theorem of algebra)

Every nonconstant polynomial $p(x) \in \mathbb{C}[X]$ can be written uniquely (upto reordering) as a product of monic irreducibles of the form $(x - z_i)$ for $z_i \in \mathbb{C}[X]$.

$$p(x) = \pm 1 \prod_i (x - z_i)$$

Why $n(p)$: only primes?

- $10(5) =$ remainder of 10 when factored by 5; $10 = 5 \cdot 2 + 0$; $10(5) = 0$
- $10(7) =$ remainder of 10 when factored by 7; $10 = 7 \cdot 1 + 3$; $10(7) = 3$
- $p(x) = (x^2 - 15x + 50)$.
- $p(5) =$ remainder of $p(x)$ when factored by $(x - 5)$;
- $p(x) = (x - 5)(x - 10) + 0$; $p(5) = 0$
- $p(1) =$ remainder of $p(x)$ when factored by $(x - 1)$;
- $p(x) = (x - 1)(x - 14) + 36$; $p(1) = 36$

Theorem (Fundamental theorem of algebra)

Every nonconstant polynomial $p(x) \in \mathbb{C}[X]$ can be written uniquely (upto reordering) as a product of monic irreducibles of the form $(x - z_i)$ for $z_i \in \mathbb{C}[X]$.

$$p(x) = \pm 1 \prod_i (x - z_i)$$

Theorem (Fundamental theorem of arithmetic)

Every non-zero integer can be written uniquely (upto reordering) as a product of primes

$$n = \pm 1 \prod_i p_i$$

Cheap trick?

- What are the complex numbers?

Cheap trick?

- What are the complex numbers?
- \mathbb{R} with i : $i^2 = -1$.

Cheap trick?

- What are the complex numbers?
- \mathbb{R} with i : $i^2 = -1$. That is, $i^2 + 1 = 0$.

Cheap trick?

- What are the complex numbers?
- \mathbb{R} with i : $i^2 = -1$. That is, $i^2 + 1 = 0$.
- Equivalently: $\mathbb{R}[X]$

Cheap trick?

- What are the complex numbers?
- \mathbb{R} with i : $i^2 = -1$. That is, $i^2 + 1 = 0$.
- Equivalently: $\mathbb{R}[X]$ divided by $q(x) = x^2 + 1$.

Cheap trick?

- What are the complex numbers?
- \mathbb{R} with i : $i^2 = -1$. That is, $i^2 + 1 = 0$.
- Equivalently: $\mathbb{R}[X]$ divided by $q(x) = x^2 + 1$.
- Left with only linear polynomials.

Cheap trick?

- What are the complex numbers?
- \mathbb{R} with i : $i^2 = -1$. That is, $i^2 + 1 = 0$.
- Equivalently: $\mathbb{R}[X]$ divided by $q(x) = x^2 + 1$.
- Left with only linear polynomials.
- All higher power polynomials $h(x)$ are $h(x) = p(x) \cdot q(x) + r(x)$

Cheap trick?

- What are the complex numbers?
- \mathbb{R} with i : $i^2 = -1$. That is, $i^2 + 1 = 0$.
- Equivalently: $\mathbb{R}[X]$ divided by $q(x) = x^2 + 1$.
- Left with only linear polynomials.
- All higher power polynomials $h(x)$ are $h(x) = p(x) \cdot q(x) + r(x)$ $\text{degree}(r) \leq 1$.

Cheap trick?

- What are the complex numbers?
- \mathbb{R} with i : $i^2 = -1$. That is, $i^2 + 1 = 0$.
- Equivalently: $\mathbb{R}[X]$ divided by $q(x) = x^2 + 1$.
- Left with only linear polynomials.
- All higher power polynomials $h(x)$ are $h(x) = p(x) \cdot q(x) + r(x)$ $\text{degree}(r) \leq 1$.
- Example: $7x^2 + 5 = 7(x^2 + 1) - 2$

Cheap trick?

- What are the complex numbers?
- \mathbb{R} with i : $i^2 = -1$. That is, $i^2 + 1 = 0$.
- Equivalently: $\mathbb{R}[X]$ divided by $q(x) = x^2 + 1$.
- Left with only linear polynomials.
- All higher power polynomials $h(x)$ are $h(x) = p(x) \cdot q(x) + r(x)$ $\text{degree}(r) \leq 1$.
- Example: $7x^2 + 5 = 7(x^2 + 1) - 2$
- Sum of linear polynomials: $(a + xb) + (c + xd) = (a + c) + x(b + d)$

Cheap trick?

- What are the complex numbers?
- \mathbb{R} with i : $i^2 = -1$. That is, $i^2 + 1 = 0$.
- Equivalently: $\mathbb{R}[X]$ divided by $q(x) = x^2 + 1$.
- Left with only linear polynomials.
- All higher power polynomials $h(x)$ are $h(x) = p(x) \cdot q(x) + r(x)$ $\text{degree}(r) \leq 1$.
- Example: $7x^2 + 5 = 7(x^2 + 1) - 2$
- Sum of linear polynomials: $(a + xb) + (c + xd) = (a + c) + x(b + d)$
- Product of linear polynomials: $(a + xb) \cdot (c + xd) = ac + x(ad + bc) + bdx^2$

Cheap trick?

- What are the complex numbers?
- \mathbb{R} with $i: i^2 = -1$. That is, $i^2 + 1 = 0$.
- Equivalently: $\mathbb{R}[X]$ divided by $q(x) = x^2 + 1$.
- Left with only linear polynomials.
- All higher power polynomials $h(x)$ are $h(x) = p(x) \cdot q(x) + r(x)$ $\text{degree}(r) \leq 1$.
- Example: $7x^2 + 5 = 7(x^2 + 1) - 2$
- Sum of linear polynomials: $(a + xb) + (c + xd) = (a + c) + x(b + d)$
- Product of linear polynomials: $(a + xb) \cdot (c + xd) = ac + x(ad + bc) + bdx^2$
- dividing product by $q(x) = x^2 + 1$:

[illegible]

Taylor series

- Taylor series of $q(x) \in \mathbb{C}[X]$ at $x = x_0$: $q(x) = \sum_i a_i (x - x_0)^i$

Taylor series

- Taylor series of $q(x) \in \mathbb{C}[X]$ at $x = x_0$: $q(x) = \sum_i a_i (x - x_0)^i$
- Taylor series of $n \in \mathbb{Z}$ at p prime: $n = \sum_i b_i p^i$.

Taylor series

- Taylor series of $q(x) \in \mathbb{C}[X]$ at $x = x_0$: $q(x) = \sum_i a_i (x - x_0)^i$
- Taylor series of $n \in \mathbb{Z}$ at p prime: $n = \sum_i b_i p^i$.

Definition

The p -adic expansion of a natural number n is the unique decomposition $n = \sum_i b_i p^i$ for $0 \leq b_i < p$.

- Taylor series of $q(x) = x^3 - 7x^2 + 15x - 9$

Taylor series

- Taylor series of $q(x) \in \mathbb{C}[X]$ at $x = x_0$: $q(x) = \sum_i a_i (x - x_0)^i$
- Taylor series of $n \in \mathbb{Z}$ at p prime: $n = \sum_i b_i p^i$.

Definition

The p -adic expansion of a natural number n is the unique decomposition $n = \sum_i b_i p^i$ for $0 \leq b_i < p$.

- Taylor series of $q(x) = x^3 - 7x^2 + 15x - 9$ at $x = 3$:

Taylor series

- Taylor series of $q(x) \in \mathbb{C}[X]$ at $x = x_0$: $q(x) = \sum_i a_i (x - x_0)^i$
- Taylor series of $n \in \mathbb{Z}$ at p prime: $n = \sum_i b_i p^i$.

Definition

The p -adic expansion of a natural number n is the unique decomposition $n = \sum_i b_i p^i$ for $0 \leq b_i < p$.

- Taylor series of $q(x) = x^3 - 7x^2 + 15x - 9$ at $x = 3$: $q(x) = 2(x - 3)^2 + (x - 3)^3$

Taylor series

- Taylor series of $q(x) \in \mathbb{C}[X]$ at $x = x_0$: $q(x) = \sum_i a_i (x - x_0)^i$
- Taylor series of $n \in \mathbb{Z}$ at p prime: $n = \sum_i b_i p^i$.

Definition

The p -adic expansion of a natural number n is the unique decomposition $n = \sum_i b_i p^i$ for $0 \leq b_i < p$.

- Taylor series of $q(x) = x^3 - 7x^2 + 15x - 9$ at $x = 3$: $q(x) = 2(x - 3)^2 + (x - 3)^3$
- $q(x) = (x - 3)^2(2 + (x - 3))$

Taylor series

- Taylor series of $q(x) \in \mathbb{C}[X]$ at $x = x_0$: $q(x) = \sum_i a_i (x - x_0)^i$
- Taylor series of $n \in \mathbb{Z}$ at p prime: $n = \sum_i b_i p^i$.

Definition

The p -adic expansion of a natural number n is the unique decomposition $n = \sum_i b_i p^i$ for $0 \leq b_i < p$.

- Taylor series of $q(x) = x^3 - 7x^2 + 15x - 9$ at $x = 3$: $q(x) = 2(x - 3)^2 + (x - 3)^3$
- $q(x) = (x - 3)^2(2 + (x - 3)) = (x - 3)^2(x - 1)$

Taylor series

- Taylor series of $q(x) \in \mathbb{C}[X]$ at $x = x_0$: $q(x) = \sum_i a_i (x - x_0)^i$
- Taylor series of $n \in \mathbb{Z}$ at p prime: $n = \sum_i b_i p^i$.

Definition

The p -adic expansion of a natural number n is the unique decomposition $n = \sum_i b_i p^i$ for $0 \leq b_i < p$.

- Taylor series of $q(x) = x^3 - 7x^2 + 15x - 9$ at $x = 3$: $q(x) = 2(x - 3)^2 + (x - 3)^3$
- $q(x) = (x - 3)^2(2 + (x - 3)) = (x - 3)^2(x - 1)$
- $x^3 - 7x^2 + 15x - 9$ has a root at 3 of order 2

Taylor series

- Taylor series of $q(x) \in \mathbb{C}[X]$ at $x = x_0$: $q(x) = \sum_i a_i (x - x_0)^i$
- Taylor series of $n \in \mathbb{Z}$ at p prime: $n = \sum_i b_i p^i$.

Definition

The p -adic expansion of a natural number n is the unique decomposition $n = \sum_i b_i p^i$ for $0 \leq b_i < p$.

- Taylor series of $q(x) = x^3 - 7x^2 + 15x - 9$ at $x = 3$: $q(x) = 2(x - 3)^2 + (x - 3)^3$
- $q(x) = (x - 3)^2(2 + (x - 3)) = (x - 3)^2(x - 1)$
- $x^3 - 7x^2 + 15x - 9$ has a root at 3 of order 2
- Taylor series/ p -adic expansion of 72 at $p = 3$:

Taylor series

- Taylor series of $q(x) \in \mathbb{C}[X]$ at $x = x_0$: $q(x) = \sum_i a_i (x - x_0)^i$
- Taylor series of $n \in \mathbb{Z}$ at p prime: $n = \sum_i b_i p^i$.

Definition

The p -adic expansion of a natural number n is the unique decomposition $n = \sum_i b_i p^i$ for $0 \leq b_i < p$.

- Taylor series of $q(x) = x^3 - 7x^2 + 15x - 9$ at $x = 3$: $q(x) = 2(x - 3)^2 + (x - 3)^3$
- $q(x) = (x - 3)^2(2 + (x - 3)) = (x - 3)^2(x - 1)$
- $x^3 - 7x^2 + 15x - 9$ has a root at 3 of order 2
- Taylor series/ p -adic expansion of 72 at $p = 3$: $72 = 0 \cdot 3 + 2 \cdot 3^2 + 2 \cdot 3^3$

Taylor series

- Taylor series of $q(x) \in \mathbb{C}[X]$ at $x = x_0$: $q(x) = \sum_i a_i (x - x_0)^i$
- Taylor series of $n \in \mathbb{Z}$ at p prime: $n = \sum_i b_i p^i$.

Definition

The p -adic expansion of a natural number n is the unique decomposition $n = \sum_i b_i p^i$ for $0 \leq b_i < p$.

- Taylor series of $q(x) = x^3 - 7x^2 + 15x - 9$ at $x = 3$: $q(x) = 2(x - 3)^2 + (x - 3)^3$
- $q(x) = (x - 3)^2(2 + (x - 3)) = (x - 3)^2(x - 1)$
- $x^3 - 7x^2 + 15x - 9$ has a root at 3 of order 2
- Taylor series/ p -adic expansion of 72 at $p = 3$: $72 = 0 \cdot 3 + 2 \cdot 3^2 + 2 \cdot 3^3$
- $72 = 3^2 * (2 + 2 \cdot 3) = 3^2 * 2^3$

Taylor series

- Taylor series of $q(x) \in \mathbb{C}[X]$ at $x = x_0$: $q(x) = \sum_i a_i (x - x_0)^i$
- Taylor series of $n \in \mathbb{Z}$ at p prime: $n = \sum_i b_i p^i$.

Definition

The p -adic expansion of a natural number n is the unique decomposition $n = \sum_i b_i p^i$ for $0 \leq b_i < p$.

- Taylor series of $q(x) = x^3 - 7x^2 + 15x - 9$ at $x = 3$: $q(x) = 2(x - 3)^2 + (x - 3)^3$
- $q(x) = (x - 3)^2(2 + (x - 3)) = (x - 3)^2(x - 1)$
- $x^3 - 7x^2 + 15x - 9$ has a root at 3 of order 2
- Taylor series/ p -adic expansion of 72 at $p = 3$: $72 = 0 \cdot 3 + 2 \cdot 3^2 + 2 \cdot 3^3$
- $72 = 3^2 * (2 + 2 \cdot 3) = 3^2 * 2^3$
- 72 has a root at $p = 3$ of order 2

Taylor series

- Taylor series of $q(x) \in \mathbb{C}[X]$ at $x = x_0$: $q(x) = \sum_i a_i (x - x_0)^i$
- Taylor series of $n \in \mathbb{Z}$ at p prime: $n = \sum_i b_i p^i$.

Definition

The p -adic expansion of a natural number n is the unique decomposition $n = \sum_i b_i p^i$ for $0 \leq b_i < p$.

- Taylor series of $q(x) = x^3 - 7x^2 + 15x - 9$ at $x = 3$: $q(x) = 2(x - 3)^2 + (x - 3)^3$
- $q(x) = (x - 3)^2(2 + (x - 3)) = (x - 3)^2(x - 1)$
- $x^3 - 7x^2 + 15x - 9$ has a root at 3 of order 2
- Taylor series/ p -adic expansion of 72 at $p = 3$: $72 = 0 \cdot 3 + 2 \cdot 3^2 + 2 \cdot 3^3$
- $72 = 3^2 * (2 + 2 \cdot 3) = 3^2 * 2^3$
- 72 has a root at $p = 3$ of order 2

Extending to the negatives

- Consider -1 .

Extending to the negatives

- Consider -1 .
- Goal: write $-1 = a_0 3^0 + a_1 3^1 + a_2 3^2 + a_3 3^3 + \dots$.
- $-1 \equiv -1 + 3 - 3$.

Extending to the negatives

- Consider -1 .
- Goal: write $-1 = a_0 3^0 + a_1 3^1 + a_2 3^2 + a_3 3^3 + \dots$.
- $-1 \equiv -1 + 3 - 3$.
- $-1 \equiv 2 - 3$

Extending to the negatives

- Consider -1 .
- Goal: write $-1 = a_0 3^0 + a_1 3^1 + a_2 3^2 + a_3 3^3 + \dots$.
- $-1 \equiv -1 + 3 - 3$.
- $-1 \equiv 2 - 3$
- $-1 \equiv 2 - 3 + 9 - 9$

Extending to the negatives

- Consider -1 .
- Goal: write $-1 = a_0 3^0 + a_1 3^1 + a_2 3^2 + a_3 3^3 + \dots$.
- $-1 \equiv -1 + 3 - 3$.
- $-1 \equiv 2 - 3$
- $-1 \equiv 2 - 3 + 9 - 9$
- $-1 \equiv 2 + (9 - 3) - 9$

Extending to the negatives

- Consider -1 .
- Goal: write $-1 = a_0 3^0 + a_1 3^1 + a_2 3^2 + a_3 3^3 + \dots$.
- $-1 \equiv -1 + 3 - 3$.
- $-1 \equiv 2 - 3$
- $-1 \equiv 2 - 3 + 9 - 9$
- $-1 \equiv 2 + (9 - 3) - 9$
- $-1 \equiv 2 + 6 - 9$

Extending to the negatives

- Consider -1 .
- Goal: write $-1 = a_0 3^0 + a_1 3^1 + a_2 3^2 + a_3 3^3 + \dots$.
- $-1 \equiv -1 + 3 - 3$.
- $-1 \equiv 2 - 3$
- $-1 \equiv 2 - 3 + 9 - 9$
- $-1 \equiv 2 + (9 - 3) - 9$
- $-1 \equiv 2 + 6 - 9$
- $-1 \equiv 2 + 6 - 9 + 27 - 27$

Extending to the negatives

- Consider -1 .
- Goal: write $-1 = a_0 3^0 + a_1 3^1 + a_2 3^2 + a_3 3^3 + \dots$.
- $-1 \equiv -1 + 3 - 3$.
- $-1 \equiv 2 - 3$
- $-1 \equiv 2 - 3 + 9 - 9$
- $-1 \equiv 2 + (9 - 3) - 9$
- $-1 \equiv 2 + 6 - 9$
- $-1 \equiv 2 + 6 - 9 + 27 - 27$
- $-1 \equiv 2 + 6 + (27 - 9) - 125$

Extending to the negatives

- Consider -1 .
- Goal: write $-1 = a_0 3^0 + a_1 3^1 + a_2 3^2 + a_3 3^3 + \dots$.
- $-1 \equiv -1 + 3 - 3$.
- $-1 \equiv 2 - 3$
- $-1 \equiv 2 - 3 + 9 - 9$
- $-1 \equiv 2 + (9 - 3) - 9$
- $-1 \equiv 2 + 6 - 9$
- $-1 \equiv 2 + 6 - 9 + 27 - 27$
- $-1 \equiv 2 + 6 + (27 - 9) - 125$
- $-1 \equiv 2 + 6 + 100 - 125$

Extending to the negatives

- Consider -1 .
- Goal: write $-1 = a_0 3^0 + a_1 3^1 + a_2 3^2 + a_3 3^3 + \dots$.
- $-1 \equiv -1 + 3 - 3$.
- $-1 \equiv 2 - 3$
- $-1 \equiv 2 - 3 + 9 - 9$
- $-1 \equiv 2 + (9 - 3) - 9$
- $-1 \equiv 2 + 6 - 9$
- $-1 \equiv 2 + 6 - 9 + 27 - 27$
- $-1 \equiv 2 + 6 + (27 - 9) - 125$
- $-1 \equiv 2 + 6 + 100 - 125$
- $-1 \equiv 2 \cdot 3^0 + 2 \cdot 3^1 + 2 \cdot 3^2 + \dots$.

Extending to the negatives

- Consider -1 .
- Goal: write $-1 = a_0 3^0 + a_1 3^1 + a_2 3^2 + a_3 3^3 + \dots$.
- $-1 \equiv -1 + 3 - 3$.
- $-1 \equiv 2 - 3$
- $-1 \equiv 2 - 3 + 9 - 9$
- $-1 \equiv 2 + (9 - 3) - 9$
- $-1 \equiv 2 + 6 - 9$
- $-1 \equiv 2 + 6 - 9 + 27 - 27$
- $-1 \equiv 2 + 6 + (27 - 9) - 125$
- $-1 \equiv 2 + 6 + 100 - 125$
- $-1 \equiv 2 \cdot 3^0 + 2 \cdot 3^1 + 2 \cdot 3^2 + \dots$.

Checking our math: $-1 + 1$

$$\blacksquare -1 \equiv 2 \cdot 3^0 + 2 \cdot 3^1 + 2 \cdot 3^2 + 2 \cdot 3^3 + \dots$$

Checking our math: $-1 + 1$

- $-1 \equiv 2 \cdot 3^0 + 2 \cdot 3^1 + 2 \cdot 3^2 + 2 \cdot 3^3 + \dots$
- $-1 + 1 = \mathbf{1} + \mathbf{2 \cdot 3^0} + 2 \cdot 3^1 + 2 \cdot 3^2 + 2 \cdot 3^3 + \dots$

Checking our math: $-1 + 1$

- $-1 \equiv 2 \cdot 3^0 + 2 \cdot 3^1 + 2 \cdot 3^2 + 2 \cdot 3^3 + \dots$
- $-1 + 1 = \mathbf{1} + \mathbf{2 \cdot 3^0} + + 2 \cdot 3^1 + 2 \cdot 3^2 + 2 \cdot 3^3 + \dots$
- $-1 + 1 = \mathbf{1 \cdot 3^1} + \mathbf{2 \cdot 3^1} + 2 \cdot 3^2 + 2 \cdot 3^3 + \dots$

Checking our math: $-1 + 1$

- $-1 \equiv 2 \cdot 3^0 + 2 \cdot 3^1 + 2 \cdot 3^2 + 2 \cdot 3^3 + \dots$.
- $-1 + 1 = \mathbf{1} + \mathbf{2 \cdot 3^0} + 2 \cdot 3^1 + 2 \cdot 3^2 + 2 \cdot 3^3 + \dots$.
- $-1 + 1 = \mathbf{1 \cdot 3^1} + \mathbf{2 \cdot 3^1} + 2 \cdot 3^2 + 2 \cdot 3^3 + \dots$.
- $-1 + 1 = \mathbf{1 \cdot 3^2} + \mathbf{2 \cdot 3^2} + 2 \cdot 3^3 + \dots$.

Checking our math: $-1 + 1$

- $-1 \equiv 2 \cdot 3^0 + 2 \cdot 3^1 + 2 \cdot 3^2 + 2 \cdot 3^3 + \dots$.
- $-1 + 1 = \mathbf{1} + \mathbf{2 \cdot 3^0} + 2 \cdot 3^1 + 2 \cdot 3^2 + 2 \cdot 3^3 + \dots$.
- $-1 + 1 = \mathbf{1 \cdot 3^1} + \mathbf{2 \cdot 3^1} + 2 \cdot 3^2 + 2 \cdot 3^3 + \dots$.
- $-1 + 1 = \mathbf{1 \cdot 3^2} + \mathbf{2 \cdot 3^2} + 2 \cdot 3^3 + \dots$.
- $-1 + 1 = \mathbf{1 \cdot 3^3} + \mathbf{2 \cdot 3^3} + \dots$.

Checking our math: $-1 + 1$

- $-1 \equiv 2 \cdot 3^0 + 2 \cdot 3^1 + 2 \cdot 3^2 + 2 \cdot 3^3 + \dots$.
- $-1 + 1 = \mathbf{1} + \mathbf{2 \cdot 3^0} + 2 \cdot 3^1 + 2 \cdot 3^2 + 2 \cdot 3^3 + \dots$.
- $-1 + 1 = \mathbf{1 \cdot 3^1} + \mathbf{2 \cdot 3^1} + 2 \cdot 3^2 + 2 \cdot 3^3 + \dots$.
- $-1 + 1 = \mathbf{1 \cdot 3^2} + \mathbf{2 \cdot 3^2} + 2 \cdot 3^3 + \dots$.
- $-1 + 1 = \mathbf{1 \cdot 3^3} + \mathbf{2 \cdot 3^3} + \dots$.
- $-1 + 1 = \dots$

Checking our math: $-1 + 1$

- $-1 \equiv 2 \cdot 3^0 + 2 \cdot 3^1 + 2 \cdot 3^2 + 2 \cdot 3^3 + \dots$.
- $-1 + 1 = \mathbf{1} + \mathbf{2 \cdot 3^0} + + 2 \cdot 3^1 + 2 \cdot 3^2 + 2 \cdot 3^3 + \dots$.
- $-1 + 1 = \mathbf{1 \cdot 3^1} + \mathbf{2 \cdot 3^1} + 2 \cdot 3^2 + 2 \cdot 3^3 + \dots$.
- $-1 + 1 = \mathbf{1 \cdot 3^2} + \mathbf{2 \cdot 3^2} + 2 \cdot 3^3 + \dots$.
- $-1 + 1 = \mathbf{1 \cdot 3^3} + \mathbf{2 \cdot 3^3} + \dots$.
- $-1 + 1 = \dots$
- $-1 + 1 = 0$.

Positional notation

$$\begin{array}{r} \dots 22222 \\ \dots 00001 + \\ \hline \dots ????? \\ \hline \end{array}$$

Positional notation

$$\begin{array}{r}
 \dots 22222 \\
 \dots 00001 \quad + \\
 \hline
 \dots ?????? \\
 \hline
 \end{array}$$

$$\begin{array}{r}
 1 \\
 \dots 22222 \\
 \dots 00001 \quad + \\
 \hline
 0 \\
 \hline
 \end{array}$$

Positional notation

$$\begin{array}{r}
 \dots 22222 \\
 \dots 00001 \quad + \\
 \hline
 \dots ????? \\
 \hline
 \end{array}$$

$$\begin{array}{r}
 1 \\
 \dots 22222 \\
 \dots 00001 \quad + \\
 \hline
 0 \\
 \hline
 \end{array}$$

$$\begin{array}{r}
 1 \\
 \dots 22222 \\
 \dots 00001 \quad + \\
 \hline
 00 \\
 \hline
 \end{array}$$

Positional notation

$$\begin{array}{r}
 \dots 22222 \\
 \dots 00001 + \\
 \hline
 \dots ?????? \\
 \hline
 \end{array}$$

$$\begin{array}{r}
 1 \\
 \dots 22222 \\
 \dots 00001 + \\
 \hline
 0 \\
 \hline
 \end{array}$$

$$\begin{array}{r}
 1 \\
 \dots 22222 \\
 \dots 00001 + \\
 \hline
 00 \\
 \hline
 \end{array}$$

$$\begin{array}{r}
 \dots 22222 \\
 \dots 00001 + \\
 \hline
 \dots 00000 \\
 \hline
 \end{array}$$

Positional notation

$$\begin{array}{r}
 \dots 22222 \\
 \dots 00001 + \\
 \hline
 \dots ????? \\
 \hline
 \end{array}$$

$$\begin{array}{r}
 1 \\
 \dots 22222 \\
 \dots 00001 + \\
 \hline
 0 \\
 \hline
 \end{array}$$

$$\begin{array}{r}
 1 \\
 \dots 22222 \\
 \dots 00001 + \\
 \hline
 00 \\
 \hline
 \end{array}$$

$$\begin{array}{r}
 \dots 22222 \\
 \dots 00001 + \\
 \hline
 \dots 00000 \\
 \hline
 \end{array}$$

- What is -1 is 2 - adically?

Positional notation

$$\begin{array}{r}
 \dots 22222 \\
 \dots 00001 + \\
 \hline
 \dots ????? \\
 \hline
 \end{array}$$

$$\begin{array}{r}
 1 \\
 \dots 22222 \\
 \dots 00001 + \\
 \hline
 0 \\
 \hline
 \end{array}$$

$$\begin{array}{r}
 1 \\
 \dots 22222 \\
 \dots 00001 + \\
 \hline
 00 \\
 \hline
 \end{array}$$

$$\begin{array}{r}
 \dots 22222 \\
 \dots 00001 + \\
 \hline
 \dots 00000 \\
 \hline
 \end{array}$$

- What is -1 is 2 - adically?
- $-1 = \dots 11111$.

Positional notation

$$\begin{array}{r}
 \dots 22222 \\
 \dots 00001 + \\
 \hline
 \dots ????? \\
 \hline
 \end{array}$$

$$\begin{array}{r}
 1 \\
 \dots 22222 \\
 \dots 00001 + \\
 \hline
 0 \\
 \hline
 \end{array}$$

$$\begin{array}{r}
 1 \\
 \dots 22222 \\
 \dots 00001 + \\
 \hline
 00 \\
 \hline
 \end{array}$$

$$\begin{array}{r}
 \dots 22222 \\
 \dots 00001 + \\
 \hline
 \dots 00000 \\
 \hline
 \end{array}$$

- What is -1 is 2 - adically?
- $-1 = \dots 11111$.
- Same as 2's complement!

Rationals

- Evaluate $1/4$ in the 3-adic system.

Rationals

- Evaluate $1/4$ in the 3-adic system.
- $1/4$

Rationals

- Evaluate $1/4$ in the 3-adic system.
- $1/4 = 1/(1 + 3)$

Rationals

- Evaluate $1/4$ in the 3-adic system.
- $1/4 = 1/(1+3) = 1 - 3 + 3^2 - 3^3 + 3^4 + \dots$

Rationals

- Evaluate $1/4$ in the 3-adic system.
- $1/4 = 1/(1+3) = 1 - 3 + 3^2 - 3^3 + 3^4 + \dots$
- What is -3 ? that's not allowed!

Rationals

- Evaluate $1/4$ in the 3-adic system.
- $1/4 = 1/(1+3) = 1 - 3 + 3^2 - 3^3 + 3^4 + \dots$
- What is -3 ? that's not allowed!
- $3^2 = 3 \cdot 3$

Rationals

- Evaluate $1/4$ in the 3-adic system.
- $1/4 = 1/(1+3) = 1 - 3 + 3^2 - 3^3 + 3^4 + \dots$
- What is -3 ? that's not allowed!
- $3^2 = 3 \cdot 3$
- $1/4 = 1 - \mathbf{3} + \mathbf{3 \cdot 3} - 3^3 + 3^4 + \dots$

Rationals

- Evaluate $1/4$ in the 3-adic system.
- $1/4 = 1/(1+3) = 1 - 3 + 3^2 - 3^3 + 3^4 + \dots$
- What is -3 ? that's not allowed!
- $3^2 = 3 \cdot 3$
- $1/4 = 1 - \mathbf{3} + \mathbf{3} \cdot \mathbf{3} - 3^3 + 3^4 + \dots$
- $1/4 = 1 + \mathbf{2} \cdot \mathbf{3} - 3^3 + 3^4 + \dots$

Rationals

- Evaluate $1/4$ in the 3-adic system.
- $1/4 = 1/(1+3) = 1 - 3 + 3^2 - 3^3 + 3^4 + \dots$
- What is -3 ? that's not allowed!
- $3^2 = 3 \cdot 3$
- $1/4 = 1 - \mathbf{3} + \mathbf{3} \cdot \mathbf{3} - 3^3 + 3^4 + \dots$
- $1/4 = 1 + \mathbf{2} \cdot \mathbf{3} - 3^3 + 3^4 + \dots$
- $1/4 = 1 + 2 \cdot 3 - \mathbf{3}^3 + \mathbf{3}^4 + \dots$

Rationals

- Evaluate $1/4$ in the 3-adic system.
- $1/4 = 1/(1+3) = 1 - 3 + 3^2 - 3^3 + 3^4 + \dots$
- What is -3 ? that's not allowed!
- $3^2 = 3 \cdot 3$
- $1/4 = 1 - \mathbf{3} + \mathbf{3} \cdot \mathbf{3} - 3^3 + 3^4 + \dots$
- $1/4 = 1 + \mathbf{2} \cdot \mathbf{3} - 3^3 + 3^4 + \dots$
- $1/4 = 1 + 2 \cdot 3 - \mathbf{3}^3 + \mathbf{3}^4 + \dots$
- $1/4 = 1 + 2 \cdot 3 - \mathbf{3}^3 + \mathbf{3} \cdot \mathbf{3}^3 + \dots$

Rationals

- Evaluate $1/4$ in the 3-adic system.
- $1/4 = 1/(1+3) = 1 - 3 + 3^2 - 3^3 + 3^4 + \dots$
- What is -3 ? that's not allowed!
- $3^2 = 3 \cdot 3$
- $1/4 = 1 - \mathbf{3} + \mathbf{3} \cdot \mathbf{3} - 3^3 + 3^4 + \dots$
- $1/4 = 1 + \mathbf{2} \cdot \mathbf{3} - 3^3 + 3^4 + \dots$
- $1/4 = 1 + 2 \cdot 3 - \mathbf{3}^3 + \mathbf{3}^4 + \dots$
- $1/4 = 1 + 2 \cdot 3 - \mathbf{3}^3 + \mathbf{3} \cdot \mathbf{3}^3 + \dots$
- $1/4 = 1 + 2 \cdot 3 + 2 \cdot 3^3 + \dots$

Rationals

- Evaluate $1/4$ in the 3-adic system.
- $1/4 = 1/(1+3) = 1 - 3 + 3^2 - 3^3 + 3^4 + \dots$
- What is -3 ? that's not allowed!
- $3^2 = 3 \cdot 3$
- $1/4 = 1 - \mathbf{3} + \mathbf{3} \cdot \mathbf{3} - 3^3 + 3^4 + \dots$
- $1/4 = 1 + \mathbf{2} \cdot \mathbf{3} - 3^3 + 3^4 + \dots$
- $1/4 = 1 + 2 \cdot 3 - \mathbf{3}^3 + \mathbf{3}^4 + \dots$
- $1/4 = 1 + 2 \cdot 3 - \mathbf{3}^3 + \mathbf{3} \cdot \mathbf{3}^3 + \dots$
- $1/4 = 1 + 2 \cdot 3 + 2 \cdot 3^3 + \dots$
- $1/4 = 1 + 2 \cdot 3 + 2 \cdot 3^3 + 2 \cdot 3^5 + 2 \cdot 3^7 + \dots$

Rationals

- Evaluate $1/4$ in the 3-adic system.
- $1/4 = 1/(1+3) = 1 - 3 + 3^2 - 3^3 + 3^4 + \dots$
- What is -3 ? that's not allowed!
- $3^2 = 3 \cdot 3$
- $1/4 = 1 - \mathbf{3} + \mathbf{3} \cdot \mathbf{3} - 3^3 + 3^4 + \dots$
- $1/4 = 1 + \mathbf{2} \cdot \mathbf{3} - 3^3 + 3^4 + \dots$
- $1/4 = 1 + 2 \cdot 3 - \mathbf{3}^3 + \mathbf{3}^4 + \dots$
- $1/4 = 1 + 2 \cdot 3 - \mathbf{3}^3 + \mathbf{3} \cdot \mathbf{3}^3 + \dots$
- $1/4 = 1 + 2 \cdot 3 + 2 \cdot 3^3 + \dots$
- $1/4 = 1 + 2 \cdot 3 + 2 \cdot 3^3 + 2 \cdot 3^5 + 2 \cdot 3^7 + \dots$
- Similar cleverness produces $1/p$ for any rational.

Rationals

- Evaluate $1/4$ in the 3-adic system.
- $1/4 = 1/(1+3) = 1 - 3 + 3^2 - 3^3 + 3^4 + \dots$
- What is -3 ? that's not allowed!
- $3^2 = 3 \cdot 3$
- $1/4 = 1 - \mathbf{3} + \mathbf{3} \cdot \mathbf{3} - 3^3 + 3^4 + \dots$
- $1/4 = 1 + \mathbf{2} \cdot \mathbf{3} - 3^3 + 3^4 + \dots$
- $1/4 = 1 + 2 \cdot 3 - \mathbf{3}^3 + \mathbf{3}^4 + \dots$
- $1/4 = 1 + 2 \cdot 3 - \mathbf{3}^3 + \mathbf{3} \cdot \mathbf{3}^3 + \dots$
- $1/4 = 1 + 2 \cdot 3 + 2 \cdot 3^3 + \dots$
- $1/4 = 1 + 2 \cdot 3 + 2 \cdot 3^3 + 2 \cdot 3^5 + 2 \cdot 3^7 + \dots$
- Similar cleverness produces $1/p$ for any rational.

A step back

- Let $x = a_0 + a_1p + a_2p^2 + \dots$

A step back

- Let $x = a_0 + a_1p + a_2p^2 + \dots$
- $x \equiv a_0 \pmod{p}$

A step back

- Let $x = a_0 + a_1p + a_2p^2 + \dots$
- $x \equiv a_0 \pmod{p}$
- $x \equiv a_0 + a_1p \pmod{p^2}$

A step back

- Let $x = a_0 + a_1p + a_2p^2 + \dots$
- $x \equiv a_0 \pmod{p}$
- $x \equiv a_0 + a_1p \pmod{p^2}$
- $x - a_0 \equiv a_1p \pmod{p^2}$

A step back

- Let $x = a_0 + a_1p + a_2p^2 + \dots$
- $x \equiv a_0 \pmod{p}$
- $x \equiv a_0 + a_1p \pmod{p^2}$
- $x - a_0 \equiv a_1p \pmod{p^2}$
- $x \equiv a_0 + a_1p + a_2p^2 \pmod{p^3}$

A step back

- Let $x = a_0 + a_1p + a_2p^2 + \dots$
- $x \equiv a_0 \pmod{p}$
- $x \equiv a_0 + a_1p \pmod{p^2}$
- $x - a_0 \equiv a_1p \pmod{p^2}$
- $x \equiv a_0 + a_1p + a_2p^2 \pmod{p^3}$
- $x - a_0 - a_1p \equiv a_2p^2 \pmod{p^3}$

A step back

- Let $-1 = \sum_i a_i 3^i$

- Let $x = a_0 + a_1 p + a_2 p^2 + \dots$

- $x \equiv a_0 \pmod{p}$

- $x \equiv a_0 + a_1 p \pmod{p^2}$

- $x - a_0 \equiv a_1 p \pmod{p^2}$

- $x \equiv a_0 + a_1 p + a_2 p^2 \pmod{p^3}$

- $x - a_0 - a_1 p \equiv a_2 p^2 \pmod{p^3}$

A step back

- Let $-1 = \sum_i a_i 3^i$
 - Let $-1 = a_0 \pmod{3}$
-
- Let $x = a_0 + a_1 p + a_2 p^2 + \dots$
 - $x \equiv a_0 \pmod{p}$
 - $x \equiv a_0 + a_1 p \pmod{p^2}$
 - $x - a_0 \equiv a_1 p \pmod{p^2}$
 - $x \equiv a_0 + a_1 p + a_2 p^2 \pmod{p^3}$
 - $x - a_0 - a_1 p \equiv a_2 p^2 \pmod{p^3}$

A step back

- Let $-1 = \sum_i a_i 3^i$
 - Let $-1 = a_0 \pmod{3}$; $a_0 = 2 \pmod{3}$
-
- Let $x = a_0 + a_1 p + a_2 p^2 + \dots$
 - $x \equiv a_0 \pmod{p}$
 - $x \equiv a_0 + a_1 p \pmod{p^2}$
 - $x - a_0 \equiv a_1 p \pmod{p^2}$
 - $x \equiv a_0 + a_1 p + a_2 p^2 \pmod{p^3}$
 - $x - a_0 - a_1 p \equiv a_2 p^2 \pmod{p^3}$

A step back

- Let $x = a_0 + a_1p + a_2p^2 + \dots$
 - $x \equiv a_0 \pmod{p}$
 - $x \equiv a_0 + a_1p \pmod{p^2}$
 - $x - a_0 \equiv a_1p \pmod{p^2}$
 - $x \equiv a_0 + a_1p + a_2p^2 \pmod{p^3}$
 - $x - a_0 - a_1p \equiv a_2p^2 \pmod{p^3}$
- Let $-1 = \sum_i a_i 3^i$
 - Let $-1 = a_0 \pmod{3}$; $a_0 = 2 \pmod{3}$
 - Let $-1 = 2 + a_1 \cdot 3 \pmod{9}$;

A step back

- Let $x = a_0 + a_1p + a_2p^2 + \dots$
 - $x \equiv a_0 \pmod{p}$
 - $x \equiv a_0 + a_1p \pmod{p^2}$
 - $x - a_0 \equiv a_1p \pmod{p^2}$
 - $x \equiv a_0 + a_1p + a_2p^2 \pmod{p^3}$
 - $x - a_0 - a_1p \equiv a_2p^2 \pmod{p^3}$
- Let $-1 = \sum_i a_i 3^i$
 - Let $-1 = a_0 \pmod{3}$; $a_0 = 2 \pmod{3}$
 - Let $-1 = 2 + a_1 \cdot 3 \pmod{9}$;
 $-3 = a_1 \cdot 3 \pmod{9}$;

A step back

- Let $x = a_0 + a_1p + a_2p^2 + \dots$
 - $x \equiv a_0 \pmod{p}$
 - $x \equiv a_0 + a_1p \pmod{p^2}$
 - $x - a_0 \equiv a_1p \pmod{p^2}$
 - $x \equiv a_0 + a_1p + a_2p^2 \pmod{p^3}$
 - $x - a_0 - a_1p \equiv a_2p^2 \pmod{p^3}$
- Let $-1 = \sum_i a_i 3^i$
 - Let $-1 = a_0 \pmod{3}$; $a_0 = 2 \pmod{3}$
 - Let $-1 = 2 + a_1 \cdot 3 \pmod{9}$;
 $-3 = a_1 \cdot 3 \pmod{9}$; $6 = a_1 \cdot 3 \pmod{9}$;

A step back

- Let $x = a_0 + a_1p + a_2p^2 + \dots$
- $x \equiv a_0 \pmod{p}$
- $x \equiv a_0 + a_1p \pmod{p^2}$
- $x - a_0 \equiv a_1p \pmod{p^2}$
- $x \equiv a_0 + a_1p + a_2p^2 \pmod{p^3}$
- $x - a_0 - a_1p \equiv a_2p^2 \pmod{p^3}$

- Let $-1 = \sum_i a_i 3^i$
- Let $-1 = a_0 \pmod{3}$; $a_0 = 2 \pmod{3}$
- Let $-1 = 2 + a_1 \cdot 3 \pmod{9}$;
 $-3 = a_1 \cdot 3 \pmod{9}$; $6 = a_1 \cdot 3 \pmod{9}$;
 $a_1 = 2$
- Let $-1 = 2 + 2 \cdot 3 + 2 \cdot 3^2 + \dots$

-
- Let $1/4 = \sum_i a_i 3^i$

A step back

- Let $x = a_0 + a_1p + a_2p^2 + \dots$
- $x \equiv a_0 \pmod{p}$
- $x \equiv a_0 + a_1p \pmod{p^2}$
- $x - a_0 \equiv a_1p \pmod{p^2}$
- $x \equiv a_0 + a_1p + a_2p^2 \pmod{p^3}$
- $x - a_0 - a_1p \equiv a_2p^2 \pmod{p^3}$

- Let $-1 = \sum_i a_i 3^i$
- Let $-1 = a_0 \pmod{3}$; $a_0 = 2 \pmod{3}$
- Let $-1 = 2 + a_1 \cdot 3 \pmod{9}$;
 $-3 = a_1 \cdot 3 \pmod{9}$; $6 = a_1 \cdot 3 \pmod{9}$;
 $a_1 = 2$
- Let $-1 = 2 + 2 \cdot 3 + 2 \cdot 3^2 + \dots$

-
- Let $1/4 = \sum_i a_i 3^i$
 - What defines $1/4$?

A step back

- Let $x = a_0 + a_1p + a_2p^2 + \dots$
- $x \equiv a_0 \pmod{p}$
- $x \equiv a_0 + a_1p \pmod{p^2}$
- $x - a_0 \equiv a_1p \pmod{p^2}$
- $x \equiv a_0 + a_1p + a_2p^2 \pmod{p^3}$
- $x - a_0 - a_1p \equiv a_2p^2 \pmod{p^3}$

- Let $-1 = \sum_i a_i 3^i$
- Let $-1 = a_0 \pmod{3}$; $a_0 = 2 \pmod{3}$
- Let $-1 = 2 + a_1 \cdot 3 \pmod{9}$;
 $-3 = a_1 \cdot 3 \pmod{9}$; $6 = a_1 \cdot 3 \pmod{9}$;
 $a_1 = 2$
- Let $-1 = 2 + 2 \cdot 3 + 2 \cdot 3^2 + \dots$

-
- Let $1/4 = \sum_i a_i 3^i$
 - What defines $1/4$? The equation $1/4 \cdot 4 = 1$.

A step back

- Let $x = a_0 + a_1p + a_2p^2 + \dots$
- $x \equiv a_0 \pmod{p}$
- $x \equiv a_0 + a_1p \pmod{p^2}$
- $x - a_0 \equiv a_1p \pmod{p^2}$
- $x \equiv a_0 + a_1p + a_2p^2 \pmod{p^3}$
- $x - a_0 - a_1p \equiv a_2p^2 \pmod{p^3}$

- Let $-1 = \sum_i a_i 3^i$
- Let $-1 = a_0 \pmod{3}$; $a_0 = 2 \pmod{3}$
- Let $-1 = 2 + a_1 \cdot 3 \pmod{9}$;
 $-3 = a_1 \cdot 3 \pmod{9}$; $6 = a_1 \cdot 3 \pmod{9}$;
 $a_1 = 2$
- Let $-1 = 2 + 2 \cdot 3 + 2 \cdot 3^2 + \dots$

-
- Let $1/4 = \sum_i a_i 3^i$
 - What defines $1/4$? The equation $1/4 \cdot 4 = 1$.
 - $(a_0 + 3a_1 + 9a_2 + \dots)(1 + 3 + 0 \cdot 9 + \dots) = 1$

A step back

- Let $x = a_0 + a_1p + a_2p^2 + \dots$
- $x \equiv a_0 \pmod{p}$
- $x \equiv a_0 + a_1p \pmod{p^2}$
- $x - a_0 \equiv a_1p \pmod{p^2}$
- $x \equiv a_0 + a_1p + a_2p^2 \pmod{p^3}$
- $x - a_0 - a_1p \equiv a_2p^2 \pmod{p^3}$

- Let $-1 = \sum_i a_i 3^i$
- Let $-1 = a_0 \pmod{3}$; $a_0 = 2 \pmod{3}$
- Let $-1 = 2 + a_1 \cdot 3 \pmod{9}$;
 $-3 = a_1 \cdot 3 \pmod{9}$; $6 = a_1 \cdot 3 \pmod{9}$;
 $a_1 = 2$
- Let $-1 = 2 + 2 \cdot 3 + 2 \cdot 3^2 + \dots$

-
- Let $1/4 = \sum_i a_i 3^i$
 - What defines $1/4$? The equation $1/4 \cdot 4 = 1$.
 - $(a_0 + 3a_1 + 9a_2 + \dots)(1 + 3 + 0 \cdot 9 + \dots) = 1$
 - $a_0 \cdot 1 \equiv 1 \pmod{3}$

A step back

- Let $x = a_0 + a_1p + a_2p^2 + \dots$
- $x \equiv a_0 \pmod{p}$
- $x \equiv a_0 + a_1p \pmod{p^2}$
- $x - a_0 \equiv a_1p \pmod{p^2}$
- $x \equiv a_0 + a_1p + a_2p^2 \pmod{p^3}$
- $x - a_0 - a_1p \equiv a_2p^2 \pmod{p^3}$

- Let $-1 = \sum_i a_i 3^i$
- Let $-1 = a_0 \pmod{3}$; $a_0 = 2 \pmod{3}$
- Let $-1 = 2 + a_1 \cdot 3 \pmod{9}$;
 $-3 = a_1 \cdot 3 \pmod{9}$; $6 = a_1 \cdot 3 \pmod{9}$;
 $a_1 = 2$
- Let $-1 = 2 + 2 \cdot 3 + 2 \cdot 3^2 + \dots$

-
- Let $1/4 = \sum_i a_i 3^i$
 - What defines $1/4$? The equation $1/4 \cdot 4 = 1$.
 - $(a_0 + 3a_1 + 9a_2 + \dots)(1 + 3 + 0 \cdot 9 + \dots) = 1$
 - $a_0 \cdot 1 \equiv 1 \pmod{3}$ $a_0 = 1$.

A step back

- Let $x = a_0 + a_1p + a_2p^2 + \dots$
- $x \equiv a_0 \pmod{p}$
- $x \equiv a_0 + a_1p \pmod{p^2}$
- $x - a_0 \equiv a_1p \pmod{p^2}$
- $x \equiv a_0 + a_1p + a_2p^2 \pmod{p^3}$
- $x - a_0 - a_1p \equiv a_2p^2 \pmod{p^3}$

- Let $-1 = \sum_i a_i 3^i$
- Let $-1 = a_0 \pmod{3}$; $a_0 = 2 \pmod{3}$
- Let $-1 = 2 + a_1 \cdot 3 \pmod{9}$;
 $-3 = a_1 \cdot 3 \pmod{9}$; $6 = a_1 \cdot 3 \pmod{9}$;
 $a_1 = 2$
- Let $-1 = 2 + 2 \cdot 3 + 2 \cdot 3^2 + \dots$

-
- Let $1/4 = \sum_i a_i 3^i$
 - What defines $1/4$? The equation $1/4 \cdot 4 = 1$.
 - $(a_0 + 3a_1 + 9a_2 + \dots)(1 + 3 + 0 \cdot 9 + \dots) = 1$
 - $a_0 \cdot 1 \equiv 1 \pmod{3}$ $a_0 = 1$.
 - $(1 + 3a_1)(1 + 3) \equiv 1 \pmod{9}$

A step back

- Let $x = a_0 + a_1p + a_2p^2 + \dots$
- $x \equiv a_0 \pmod{p}$
- $x \equiv a_0 + a_1p \pmod{p^2}$
- $x - a_0 \equiv a_1p \pmod{p^2}$
- $x \equiv a_0 + a_1p + a_2p^2 \pmod{p^3}$
- $x - a_0 - a_1p \equiv a_2p^2 \pmod{p^3}$

- Let $-1 = \sum_i a_i 3^i$
- Let $-1 = a_0 \pmod{3}$; $a_0 = 2 \pmod{3}$
- Let $-1 = 2 + a_1 \cdot 3 \pmod{9}$;
 $-3 = a_1 \cdot 3 \pmod{9}$; $6 = a_1 \cdot 3 \pmod{9}$;
 $a_1 = 2$
- Let $-1 = 2 + 2 \cdot 3 + 2 \cdot 3^2 + \dots$

-
- Let $1/4 = \sum_i a_i 3^i$
 - What defines $1/4$? The equation $1/4 \cdot 4 = 1$.
 - $(a_0 + 3a_1 + 9a_2 + \dots)(1 + 3 + 0 \cdot 9 + \dots) = 1$
 - $a_0 \cdot 1 \equiv 1 \pmod{3}$ $a_0 = 1$.
 - $(1 + 3a_1)(1 + 3) \equiv 1 \pmod{9}$ $4 + 12a_1 \equiv 1 \pmod{9}$

A step back

- Let $x = a_0 + a_1p + a_2p^2 + \dots$
- $x \equiv a_0 \pmod{p}$
- $x \equiv a_0 + a_1p \pmod{p^2}$
- $x - a_0 \equiv a_1p \pmod{p^2}$
- $x \equiv a_0 + a_1p + a_2p^2 \pmod{p^3}$
- $x - a_0 - a_1p \equiv a_2p^2 \pmod{p^3}$

- Let $-1 = \sum_i a_i 3^i$
- Let $-1 = a_0 \pmod{3}$; $a_0 = 2 \pmod{3}$
- Let $-1 = 2 + a_1 \cdot 3 \pmod{9}$;
 $-3 = a_1 \cdot 3 \pmod{9}$; $6 = a_1 \cdot 3 \pmod{9}$;
 $a_1 = 2$
- Let $-1 = 2 + 2 \cdot 3 + 2 \cdot 3^2 + \dots$

-
- Let $1/4 = \sum_i a_i 3^i$
 - What defines $1/4$? The equation $1/4 \cdot 4 = 1$.
 - $(a_0 + 3a_1 + 9a_2 + \dots)(1 + 3 + 0 \cdot 9 + \dots) = 1$
 - $a_0 \cdot 1 \equiv 1 \pmod{3}$ $a_0 = 1$.
 - $(1 + 3a_1)(1 + 3) \equiv 1 \pmod{9}$ $4 + 12a_1 \equiv 1 \pmod{9}$
 - $3a_1 \equiv -3 \equiv 6 \pmod{9}$

A step back

- Let $x = a_0 + a_1p + a_2p^2 + \dots$
- $x \equiv a_0 \pmod{p}$
- $x \equiv a_0 + a_1p \pmod{p^2}$
- $x - a_0 \equiv a_1p \pmod{p^2}$
- $x \equiv a_0 + a_1p + a_2p^2 \pmod{p^3}$
- $x - a_0 - a_1p \equiv a_2p^2 \pmod{p^3}$

- Let $-1 = \sum_i a_i 3^i$
- Let $-1 = a_0 \pmod{3}$; $a_0 = 2 \pmod{3}$
- Let $-1 = 2 + a_1 \cdot 3 \pmod{9}$;
 $-3 = a_1 \cdot 3 \pmod{9}$; $6 = a_1 \cdot 3 \pmod{9}$;
 $a_1 = 2$
- Let $-1 = 2 + 2 \cdot 3 + 2 \cdot 3^2 + \dots$

-
- Let $1/4 = \sum_i a_i 3^i$
 - What defines $1/4$? The equation $1/4 \cdot 4 = 1$.
 - $(a_0 + 3a_1 + 9a_2 + \dots)(1 + 3 + 0 \cdot 9 + \dots) = 1$
 - $a_0 \cdot 1 \equiv 1 \pmod{3}$ $a_0 = 1$.
 - $(1 + 3a_1)(1 + 3) \equiv 1 \pmod{9}$ $4 + 12a_1 \equiv 1 \pmod{9}$
 - $3a_1 \equiv -3 \equiv 6 \pmod{9}$
 - $a_1 \equiv 2 \pmod{9}$

A step back

- Let $x = a_0 + a_1p + a_2p^2 + \dots$
- $x \equiv a_0 \pmod{p}$
- $x \equiv a_0 + a_1p \pmod{p^2}$
- $x - a_0 \equiv a_1p \pmod{p^2}$
- $x \equiv a_0 + a_1p + a_2p^2 \pmod{p^3}$
- $x - a_0 - a_1p \equiv a_2p^2 \pmod{p^3}$

- Let $-1 = \sum_i a_i 3^i$
- Let $-1 = a_0 \pmod{3}$; $a_0 = 2 \pmod{3}$
- Let $-1 = 2 + a_1 \cdot 3 \pmod{9}$;
 $-3 = a_1 \cdot 3 \pmod{9}$; $6 = a_1 \cdot 3 \pmod{9}$;
 $a_1 = 2$
- Let $-1 = 2 + 2 \cdot 3 + 2 \cdot 3^2 + \dots$

-
- Let $1/4 = \sum_i a_i 3^i$
 - What defines $1/4$? The equation $1/4 \cdot 4 = 1$.
 - $(a_0 + 3a_1 + 9a_2 + \dots)(1 + 3 + 0 \cdot 9 + \dots) = 1$
 - $a_0 \cdot 1 \equiv 1 \pmod{3}$ $a_0 = 1$.
 - $(1 + 3a_1)(1 + 3) \equiv 1 \pmod{9}$ $4 + 12a_1 \equiv 1 \pmod{9}$
 - $3a_1 \equiv -3 \equiv 6 \pmod{9}$
 - $a_1 \equiv 2 \pmod{9}$
 - $1/4 = 1 + 2 \cdot 3 + 2 \cdot 3^3 + \dots$

Irrationals

- Let's solve $X^2 = 2$ in the 7-adics.

Irrationals

- Let's solve $X^2 = 2$ in the 7-adics.
- Such a solution does not "really exist" in the rationals or the integers.

Irrationals

- Let's solve $X^2 = 2$ in the 7-adics.
- Such a solution does not "really exist" in the rationals or the integers.
- Start with $X \equiv 3 \pmod{7}$, $X \equiv 4 \equiv -3 \pmod{7}$.

Convergence

- Intuition: higher powers of p should become “smaller” for convergence!

Convergence

- Intuition: higher powers of p should become “smaller” for convergence!
- $|a|_p = p^{-1 \cdot \text{highest power of } p \text{ which divides } a}$

Convergence

- Intuition: higher powers of p should become “smaller” for convergence!
- $|a|_p = p^{-1 \cdot \text{highest power of } p \text{ which divides } a}$
- $|10|_3 = 3^{-0} = 1$

Convergence

- Intuition: higher powers of p should become “smaller” for convergence!
- $|a|_p = p^{-1 \cdot \text{highest power of } p \text{ which divides } a}$
- $|10|_3 = 3^{-0} = 1$
- $|3|_3 = 3^{-1} = 1/3$

Convergence

- Intuition: higher powers of p should become “smaller” for convergence!
- $|a|_p = p^{-1 \cdot \text{highest power of } p \text{ which divides } a}$
- $|10|_3 = 3^{-0} = 1$
- $|3|_3 = 3^{-1} = 1/3$
- $|9|_3 = 3^{-2} = 1/9$

Convergence

- Intuition: higher powers of p should become “smaller” for convergence!
- $|a|_p = p^{-1 \cdot \text{highest power of } p \text{ which divides } a}$
- $|10|_3 = 3^{-0} = 1$
- $|3|_3 = 3^{-1} = 1/3$
- $|9|_3 = 3^{-2} = 1/9$
- $|90|_3 = 3^{-2} = 1/9$

Convergence

- Intuition: higher powers of p should become “smaller” for convergence!
- $|a|_p = p^{-1 \cdot \text{highest power of } p \text{ which divides } a}$
- $|10|_3 = 3^{-0} = 1$
- $|3|_3 = 3^{-1} = 1/3$
- $|9|_3 = 3^{-2} = 1/9$
- $|90|_3 = 3^{-2} = 1/9$
- $|27|_3 = 3^{-3} = 1/27$

Convergence

- Intuition: higher powers of p should become “smaller” for convergence!
- $|a|_p = p^{-1 \cdot \text{highest power of } p \text{ which divides } a}$
- $|10|_3 = 3^{-0} = 1$
- $|3|_3 = 3^{-1} = 1/3$
- $|9|_3 = 3^{-2} = 1/9$
- $|90|_3 = 3^{-2} = 1/9$
- $|27|_3 = 3^{-3} = 1/27$
- $|a + b|_p \leq |a|_p + |b|_p?$

Convergence

- Intuition: higher powers of p should become “smaller” for convergence!
- $|a|_p = p^{-1 \cdot \text{highest power of } p \text{ which divides } a}$
- $|10|_3 = 3^{-0} = 1$
- $|3|_3 = 3^{-1} = 1/3$
- $|9|_3 = 3^{-2} = 1/9$
- $|90|_3 = 3^{-2} = 1/9$
- $|27|_3 = 3^{-3} = 1/27$
- $|a + b|_p \leq |a|_p + |b|_p?$

Scam v/s non-scam

- Solve $x = 1 + 3x$

Scam v/s non-scam

- Solve $x = 1 + 3x$
- **Non scam:** $-2x = 1$

Scam v/s non-scam

- Solve $x = 1 + 3x$
- **Non scam:** $-2x = 1$; $x = -1/2$
- $x_0 = 1$

Scam v/s non-scam

- Solve $x = 1 + 3x$
- **Non scam:** $-2x = 1$; $x = -1/2$
- $x_0 = 1$
- $x_1 = 1 + 3$
- $x_2 = 1 + 3x_1 = 1 + 3(1 + 3) = 1 + 3 + 3^2$

Scam v/s non-scam

- Solve $x = 1 + 3x$
- **Non scam:** $-2x = 1$; $x = -1/2$
- $x_0 = 1$
- $x_1 = 1 + 3$
- $x_2 = 1 + 3x_1 = 1 + 3(1 + 3) = 1 + 3 + 3^2$
- $x_i = 1 + 3 + 3^2 + \cdots + 3^i$

Scam v/s non-scam

- Solve $x = 1 + 3x$
- **Non scam:** $-2x = 1$; $x = -1/2$
- $x_0 = 1$
- $x_1 = 1 + 3$
- $x_2 = 1 + 3x_1 = 1 + 3(1 + 3) = 1 + 3 + 3^2$
- $x_i = 1 + 3 + 3^2 + \cdots + 3^i$
- $x_\infty = 1/(1 - 3) = -1/2$
- Converges? We need $|3| < 1$


Scam v/s non-scam

- Solve $x = 1 + 3x$
- **Non scam:** $-2x = 1$; $x = -1/2$
- $x_0 = 1$
- $x_1 = 1 + 3$
- $x_2 = 1 + 3x_1 = 1 + 3(1 + 3) = 1 + 3 + 3^2$
- $x_i = 1 + 3 + 3^2 + \dots + 3^i$
- $x_\infty = 1/(1 - 3) = -1/2$
- Converges? We need $|3| < 1$
- But it is! $|3|_3 < 1$


Scam v/s non-scam

- Solve $x = 1 + 3x$
- **Non scam:** $-2x = 1$; $x = -1/2$
- $x_0 = 1$
- $x_1 = 1 + 3$
- $x_2 = 1 + 3x_1 = 1 + 3(1 + 3) = 1 + 3 + 3^2$
- $x_i = 1 + 3 + 3^2 + \dots + 3^i$
- $x_\infty = 1/(1 - 3) = -1/2$
- Converges? We need $|3| < 1$
- But it is! $|3|_3 < 1$


Why only primes? Geometry of functions




$f(x_0) \neq 0 \Rightarrow f$ invertible (locally)
 $f' \cdot (x - x_0) \neq 0 \Rightarrow f$ invertible (locally)



Why only primes? Geometry of functions




$f(x_0) \neq 0 \Rightarrow f$ invertible (locally)
 $f' \cdot (x - x_0) \neq 0 \Rightarrow f$ invertible (locally)




- $f(x)$: continuous, *non-zero* at $x = x_0$.

Why only primes? Geometry of functions




$$f(x_0) \neq 0 \Rightarrow f \text{ invertible (locally)}$$

$$f' \cdot (x - x_0) \neq 0 \Rightarrow f \text{ invertible (locally)}$$




- $f(x)$: continuous, *non-zero* at $x = x_0$.
- $f(x)$: *locally invertible* at $x = x_0$.

Why only primes? Geometry of functions




$f(x_0) \neq 0 \Rightarrow f$ invertible (locally)
 $f'(x_0) \neq 0 \Rightarrow f$ invertible (locally)




- $f(x)$: continuous, *non-zero* at $x = x_0$.
- $f(x)$: *locally invertible* at $x = x_0$.

Why only primes? Geometry of numbers




- consider 4.

Why only primes? Geometry of numbers




- consider 4.
- nonzero at $a_0 = 6$: $4 \simeq 4 \pmod{6}$

Why only primes? Geometry of numbers




- consider 4.
- nonzero at $a_0 = 6$: $4 \simeq 4 \pmod{6}$
- *not invertible* modulo 6:
 $[0, 1, 2, 3, 4, 5] \times 4 \equiv [0, 4, 8, 12, 16, 20] \equiv [0, 4, 2, 0, 4, 2] \pmod{6}$

Why only primes? Geometry of numbers



- consider 4.
- nonzero at $a_0 = 6$: $4 \simeq 4 \pmod{6}$
- *not invertible* modulo 6:
 $[0, 1, 2, 3, 4, 5] \times 4 \equiv [0, 4, 8, 12, 16, 20] \equiv [0, 4, 2, 0, 4, 2] \pmod{6}$
- If we want 4 to be a *continuous* function

Why only primes? Geometry of numbers



- consider 4.
- nonzero at $a_0 = 6$: $4 \simeq 4 \pmod{6}$
- *not invertible* modulo 6:
 $[0, 1, 2, 3, 4, 5] \times 4 \equiv [0, 4, 8, 12, 16, 20] \equiv [0, 4, 2, 0, 4, 2] \pmod{6}$
- If we want 4 to be a *continuous* function
- then 6 should not be a point!