

# Limits and Colimits as universal cones

Siddharth Bhat

`##harmless` **Category Theory in Context**

Sun 20, June 2021

## Building objects from other ones


$$\mathbb{R} \uparrow \times \xrightarrow{\mathbb{R}} = \mathbb{I} \text{ (isomorphism)} : \mathbb{R} \times \mathbb{R} = \mathbb{R}^2$$

$$\mathbb{R}^2 \xrightarrow{+} x^2 + y^2 - 1 = 0 = \left\{ \mathbb{R} \times \mathbb{R} ; \begin{matrix} x \\ y \end{matrix} \right\} : \left\{ \mathbb{R} \times \mathbb{R} ; x^2 + y^2 - 1 = 0 \right\} = S^1$$


$D^2 / S^1 = S^2$  (QUOTIENT)

  $S^1 \times S^1 = T^2$

## Cone over a diagram (Picture)




## Cone over a diagram (formally)




- Given: (1) a diagram category  $J$ , (2) a target category  $C$ , (3) a functor  $F: J \rightarrow C$ , (4) a choice of apex  $c_* \in C$ .

## Cone over a diagram (formally)



- Given: (1) a diagram category  $J$ , (2) a target category  $C$ , (3) a functor  $F : J \rightarrow C$ , (4) a choice of apex  $c_* \in C$ .
- The cone is: A natural transformation  $\eta$  between the constant functor  $\Delta_{c_*} : J \rightarrow C$  (defined by the eqn  $\Delta_{c_*}(\_) \equiv c_*$ ) and the given  $F : J \rightarrow C$ .

## Cone over a diagram (formally)




- Given: (1) a diagram category  $J$ , (2) a target category  $C$ , (3) a functor  $F: J \rightarrow C$ , (4) a choice of apex  $c_* \in C$ .
- The cone is: A natural transformation  $\eta$  between the constant functor  $\Delta_{c_*}: J \rightarrow C$  (defined by the eqn  $\Delta_{c_*}(\_) \equiv c_*$ ) and the given  $F: J \rightarrow C$ .
- At each component, we have:

$$\eta: \Delta_{c_*} \Rightarrow F$$

$$\eta_j: \Delta_{c_*}(j) \rightarrow F(j)$$

$$\eta_j: c_* \rightarrow F(j)$$



- Given: (1) a diagram category  $J$ , (2) a target category  $C$ , (3) a functor  $F : J \rightarrow C$ , (4) a choice of apex  $c_* \in C$ .
- The cone is: A natural transformation  $\eta$  between the constant functor  $\Delta_{c_*} : J \rightarrow C$  (defined by the eqn  $\Delta_{c_*}(\_) \equiv c_*$ ) and the given  $F : J \rightarrow C$ .
- At each component, we have:
 
$$\eta : \Delta_{c_*} \Rightarrow F$$

$$\eta_j : \Delta_{c_*}(j) \rightarrow F(j)$$


$$\eta_j : c_* \rightarrow F(j)$$

## Example Cone 1




## Example Cone 2


## Definition of a Limit 1




- For any diagram  $F: J \rightarrow C$ , there is a functor:  $\text{Cone}(-, F): C \rightarrow \text{Set}$  which sends a given object  $c_* \in C$  to the set of cones over  $F$  with summit  $c_*$ .




## Definition of a Limit 1




- For any diagram  $F : J \rightarrow C$ , there is a functor:  
 $\text{Cone}(-, F) : C \rightarrow \text{Set}$  which sends a given object  $c_* \in C$  to the set of cones over  $F$  with summit  $c_*$ .
- A limit is a representation of  $\text{Cone}(-, F)$ .




## Definition of a Limit 1



- For any diagram  $F: J \rightarrow C$ , there is a functor:  $\text{Cone}(-, F): C \rightarrow \text{Set}$  which sends a given object  $c_* \in C$  to the set of cones over  $F$  with summit  $c_*$ .
- A limit is a representation of  $\text{Cone}(-, F)$ .
- So we have an object  $\lim F$  whose Hom functor represents  $\text{Cone}(-, F)$ . This is given by  $\eta: \text{Hom}_C(-, \lim F) \simeq \text{Cone}(-, F)$ .




## Definition of a Limit 1




- For any diagram  $F: J \rightarrow C$ , there is a functor:  $\text{Cone}(-, F): C \rightarrow \text{Set}$  which sends a given object  $c_* \in C$  to the set of cones over  $F$  with summit  $c_*$ .
- A limit is a representation of  $\text{Cone}(-, F)$ .
- So we have an object  $\text{lim } F$  whose Hom functor represents  $\text{Cone}(-, F)$ . This is given by  $\eta: \text{Hom}_C(-, \text{lim } F) \simeq \text{Cone}(-, F)$ .
- By Yoneda, such a natural transformation is determined entirely by an element of  $\text{Cone}(-, F)(\text{lim } F)$ , or  $\text{Cone}(\text{lim } F, F)$ .

## Definition of a Limit 1



- For any diagram  $F : J \rightarrow C$ , there is a functor:  $\text{Cone}(-, F) : C \rightarrow \text{Set}$  which sends a given object  $c_* \in C$  to the set of cones over  $F$  with summit  $c_*$ .
- A limit is a representation of  $\text{Cone}(-, F)$ .
- So we have an object  $\lim F$  whose Hom functor represents  $\text{Cone}(-, F)$ . This is given by  $\eta : \text{Hom}_C(-, \lim F) \simeq \text{Cone}(-, F)$ .
- By Yoneda, such a natural transformation is determined entirely by an element of  $\text{Cone}(-, F)(\lim F)$ , or  $\text{Cone}(\lim F, F)$ .
- Call this (universal) element  $\lambda \in \text{Cone}(\lim F, F)$ . This is a natural transformation  $\lambda : \Delta_{\lim F} \Rightarrow F$ .


## Definition of a Limit 1



- For any diagram  $F : J \rightarrow C$ , there is a functor:  $\text{Cone}(-, F) : C \rightarrow \text{Set}$  which sends a given object  $c_* \in C$  to the set of cones over  $F$  with summit  $c_*$ .
- A limit is a representation of  $\text{Cone}(-, F)$ .
- So we have an object  $\lim F$  whose Hom functor represents  $\text{Cone}(-, F)$ . This is given by  $\eta : \text{Hom}_C(-, \lim F) \simeq \text{Cone}(-, F)$ .
- By Yoneda, such a natural transformation is determined entirely by an element of  $\text{Cone}(-, F)(\lim F)$ , or  $\text{Cone}(\lim F, F)$ .
- Call this (universal) element  $\lambda \in \text{Cone}(\lim F, F)$ . This is a natural transformation  $\lambda : \Delta_{\lim F} \Rightarrow F$ .<sup>a</sup>

<sup>a</sup>Riehl writes  $\lambda : \lim F \Rightarrow F$  which does not type-check for me.

## Definition of a Limit 1: Natural Iso



- An object  $\lim F$  such that  $\eta : \text{Hom}_C(-, \lim F) \simeq \text{Cone}(-, F)$ .
- By Yoneda,  $\eta$  is determined entirely by an element of  $\text{Cone}(\lim F, F)$ .
- This Universal element is  $\lambda \in \text{Cone}(\lim F, F)$ . I.e., a natural transformation  $\lambda : \Delta_{\lim F} \Rightarrow F$ .