

Porosphaera globularis (Phillips, 1829) (Porifera, Calcarea) from the Maastrichtian of the Farokhi Formation of Central Iran

Markus Wilmsen^{a,*}, Franz Theodor Fürsich^b, Mahmoud Reza Majidifard^c

^aSenckenberg Naturhistorische Sammlungen Dresden, Museum für Mineralogie und Geologie, Sektion Paläozoologie, Königsbrücker Landstr. 159, D-01109 Dresden, Germany

^bGeoZentrum Nordbayern, Fachgruppe PaläoUmwelt, Friedrich-Alexander-Universität Erlangen-Nürnberg, Loewenichstr. 28, D-91054 Erlangen, Germany

^cGeological Survey of Iran, Box 131851-1494, Tehran, Iran

ARTICLE INFO

Article history:

Received 6 December 2010

Accepted in revised form 6 September 2011

Available online 21 September 2011

Keywords:

Upper Cretaceous

Iran

Porifera

Minchinellidae

First record

ABSTRACT

A small faunule of the calcareous sponge *Porosphaera globularis* (Phillips, 1829) (Calcarea, Minchinellidae) is described and illustrated from the Maastrichtian Farokhi Formation of Central Iran. The shape of the sponges varies from near-spherical to pyriform with diameters ranging between 5 and 13 mm. The surfaces show small, equidistant apertures corresponding to radiating oscular tubes within a choanosomal skeleton consisting of large tetractines fused to a rigid framework. Radially arranged superficial grooves may also be present in some specimens. External attachment scars have not been observed, but perforations in two specimens may represent sipuncular worm borings. A quiet and fairly deep-water environment can be inferred as the habitat of the sponges, corresponding to data in the literature. So far, *P. globularis* was known only from the Upper Cretaceous (Cenomanian–Maastrichtian) of the Boreal Realm. The record from Central Iran is the southernmost occurrence of *P. globularis* (palaeo-latitude ca. 20°N) known to date and is the first description of the species from Iran and the Middle East.

© 2011 Elsevier Ltd. All rights reserved.

1. Introduction

Cretaceous strata in Central Iran are superbly exposed and relatively fossiliferous. However, their detailed stratigraphy (apart from gross lithostratigraphical mapping) and palaeontology are poorly known, with the exception of ammonite faunas, which have been recorded and described from numerous localities in Central Iran (e.g., Seyed-Emami, 1977, 1982; Kennedy et al., 1979; Seyed-Emami and Immel, 1995, 1996; Wilmsen et al., 2005). In contrast, data on benthic macro-invertebrates are scarce and only a few outdated monographs (e.g., on echinoids; Cotteau and Gauthier, 1895) exist. Here, we record the calcareous sponge *Porosphaera globularis* from the Khur area of Central Iran, which is the first documentation of the species from Iran and the Middle East.

2. Geological setting

The study area belongs to the Central-East Iranian Micro-continent (CEIM), which forms the core of the so-called Iran Plate.

The CEIM consists, from east to west, of three structural units, the Lut, Tabas and Yazd blocks (see Berberian and King, 1981 and Davoudzadeh, 1997 for overview). While Jurassic strata are well exposed (and have been well studied) on the Tabas Block (see Fürsich et al., 2003; Wilmsen et al., 2003, 2009a, 2010), Cretaceous strata are widespread and very thick in the western part of the CEIM (Yazd Block). The Cretaceous successions of the Yazd Block document, in excellent outcrops, the development of a mid- to Late Cretaceous (Late Barremian–Maastrichtian) carbonate platform–basin system. In terms of geodynamic evolution, the area is of great importance for the Middle East as the CEIM formed a separate microplate during Cretaceous times which was separated from the nearby Eurasian margin by small oceanic basins (Fig. 1; e.g., Dercourt et al., 1986; Philip and Floquet, 2000). These oceanic basins started to open in Early Cretaceous ('Neocomian') times and may have been related to an inferred post-Triassic rotation of the CEIM around a vertical axis of about 135° with respect to Eurasia (e.g., Davoudzadeh et al., 1981; Soffel et al., 1996). The various small oceans (Nain-Baft Ocean in the west, Sabzevar Ocean in the north, and Sistan Ocean in the east; Fig. 1) were subsequently closed during the later part of the Early to Late Cretaceous and in the Palaeocene, in connection with the advance of the Arabian Plate and the closure of the Neotethys (e.g., Tirrul et al., 1983; Dercourt et al., 1986; Stampfli and Borel,

* Corresponding author. Tel.: +49 351 7958414273; fax: +49 351 8926404.

E-mail address: markus.wilmsen@senckenberg.de (M. Wilmsen).




Fig. 1. Upper Cretaceous palaeogeographic map of the Middle East (A, modified after Philip and Floquet, 2000) and structural map of Iran (B) with indication of the Khur area, north of Yazd, in Central Iran (modified from Wilmsen et al., 2009b).

2002; Rosetti et al., 2010). Of special interest here are the Haftoman and Farokhi formations, which comprise the middle and upper part of the Upper Cretaceous succession in the Khur area (Fig. 2).

The section studied at Kuh-e-Honu, south of Khur (Figs. 1 and 3; N 33°43.355', E 55°00.869') starts with shallow-water carbonates of the Haftoman Formation, which rest unconformably on the Albian–Cenomanian Debarsu Formation. The age of the lower part of the up to 1000 m-thick Haftoman Formation is Early Coniacian, based on the occurrence of the inoceramid bivalve *Tethyoceras* ex gr. *wandereri* (Andert). In the upper part of the Haftoman Formation, a deepening trend is indicated by intercalations of marly limestones and marls with more open marine faunas (echinoids, inoceramid bivalves and ammonites). The heteromorph ammonite *Bostrychoceras polyplacum* (Roemer) indicates an early Late Campanian age. Up-section, 100–200 m of unfossiliferous limestone follow until, at a sharp, karstic contact with reworked pebbles of the Haftoman Formation, the marls of the Farokhi Formation commence (Fig. 3). The Farokhi Formation is 21.5 m thick and consists of soft, grey marl. In a slightly more calcareous and bioclastic bed, ca. 8 m above the base of the formation, numerous specimens of the calcareous sponge *P. globularis* (Phillips) occur and a small number has been collected. The Farokhi Formation is unconformably overlain by the Palaeocene Chupanan Formation (Fig. 2; see Aistov et al., 1984 for details). The age of the Farokhi Formation at Kuh-e-Honu is thus late Late Campanian–Maastrichtian with an exclusively Maastrichtian age being more likely on the basis of stratigraphic and sedimentological data (early Late Campanian index fossils occur a considerable distance below the lower boundary of the Farokhi Formation which furthermore represents a subaerial unconformity).

3. Taxonomy

We here follow the classification of the Porifera as proposed by Hooper and van Soest (2002) and Finks et al. (2004). According to Vacelet et al. (2002), fossil Minchinellidae (to which *Porosphaera* belongs) are in urgent need of revision. Authors of higher rank taxonomic categories are not listed in the references.

Class: Calcarea Bowerbank, 1864
Order: Lithonida Doederlein, 1892
Family: Minchinellidae Dendy and Row, 1913
Genus *Porosphaera* Steinmann, 1878

Type species: *Millepora globularis* Phillips, 1829, p. 186, pl. 1, fig. 12, by original designation.

Porosphaera globularis (Phillips, 1829)
Figs. 4 and 5

- 1829 *Millepora globularis* Phillips, p. 186, pl. 1, fig. 12.
- 1878 *Porosphaera globularis* (Phillips); Steinmann, p. 102, pl. 13, figs 8–12.
- 1904 *Porosphaera globularis* (Phillips); Hinde, p. 18, pl. 1, figs 1–10, pl. 2, figs 1–3, 6–10. [with synonymy]
- 1961 *Porosphaera globularis* Phillips; Nestler, p. 39, pl. 10, pl. 11, figs 1–8. [with synonymy]
- 1970 *Porosphaera globularis* Phillips; Müller, text-fig. 1, pls 1, 2.
- 1985a *Porosphaera globularis* (Phillips); Termier and Termier, p. 153, fig. 4e.
- 1985b *Porosphaera globularis* (Phillips) Steinmann; Termier and Termier, p. 54, pl. 5, figs 1–3, pl. 6, figs 1–3.




Fig. 2. Lithostratigraphy of the Upper Cretaceous in the Khur area, Yazd Block, Central Iran, after our own observations.

- 1989 *Porosphaera globularis* (Phillips); Małeck, p. 206, text-figs 1, 2, pls 1–4,
 1992 *Porosphaera globularis* (Phillips); Reitner, p. 112, pl. 6, figs 1–3.
 2002 *Porosphaera globularis* Phillips; Nestler, p. 38, fig. 36.
 2002 *Porosphaera globularis* (Phillips); Wood, p. 35, pl. 4, fig. 3.
 2002 *Porosphaera globularis* Phillips; Reich and Frenzel, p. 126, pl. 13, fig. 3.
 2008 *Porosphaera globularis* with *Trypanites mobilis* isp. n.; Neumann et al., fig. 5.
 2009 *Porosphaera globularis*; Rigaud et al., figs 2, 3, 7, 8.

Material. Fourteen specimens from the Farokhi Formation of Kuh-e-Honu (N 33°43.355', E 55°00.869' at 1.300 m altitude), ca. 20 km south of Khur, Central Iran (sample no. 091014-18, stored at the Museum für Mineralogie und Geologie der Senckenberg Naturhistorischen Sammlungen Dresden).

Measurements.

091014-18	max. Ø [mm]	min. Ø [mm]
–1	12.9	11.0
–2	11.9	8.0
–3	10.9	9.8
–4	9.7	7.8
–5	9.6	8.9
–6	9.2	7.9
–7	9.1	8.0
–8	8.9	8.1
–9	8.8	7.8
–10	8.6	8.0
–11	7.7	7.5
–12	6.9	6.3
–13	5.7	5.3
–14	5.6	5.5
Mean:	9.0	7.9

Description. The specimens are roughly spherical, but ovoid, flattened or pea-shaped forms also occur (Fig. 4). One specimen shows a flattened base and a tapering opposite end, resulting in a somewhat pyriform shape (Fig. 4H, I). Diameters range from 5.3 to 12.9 mm with a mean maximum diameter of 9.0 and a mean minimum diameter of 7.9 mm. The smaller forms are usually more spherical than the larger ones. The surface of all specimens shows small, equidistant apertures (ca. 0.1–0.2 mm in diameter). Radiating superficial grooves are sometimes also present (Fig. 4J, K) and two specimens show perforations (Fig. 4L, M). In thin section (Fig. 5), the radiating main oscular canals are clearly visible. The choanosomal skeleton consists of large tetractines fused to a rigid framework via cemented bridges. Skeletal interspaces are filled by a (late) diagenetic sparry calcite.

Discussion. The specimens from the Farokhi Formation do not differ significantly from *P. globularis* as described and illustrated in the literature. The mean diameter of 9.0 mm is comparable to that of a large population from the Lower Maastrichtian chalk of Rügen, Germany (2734 specimens with a mean diameter of 10 mm; Nestler, 1961). However, larger forms up to 50 mm in diameter as in Rügen have not been observed. External perforations are common in *P. globularis* and are interpreted as borings (Nestler, 1961; Müller, 1970; Neumann et al., 2008). The occasionally visible radiating superficial grooves almost certainly represent the traces of the oscular tubes in a zone between the dermal layer and the basal skeleton.

The internal structure of the skeleton shows the typical radiating oscular canals within a framework of mutually cemented tetractines (Fig. 5), also illustrated by Termier and Termier (1985b, pl. 5) and Reitner (1992, pl. 6, figs 1, 2). However, large diactines as relics of the non-cemented ectosomal spicular skeleton have not been observed (compare Reitner, 1992).

According to Nestler (1961, 1965), *P. globularis* from the Early Maastrichtian of Rügen thrived at sub-photic depths (ca. 100–300 m), and also Reitner (1992) regarded the sponge as a deep-water species adapted to low-energy conditions. The sponge may have been anchored to the substrate by a non-calcified dermal tissue, which could explain the lack of external attachment scars (Reitner, 1992). The occurrence of *P. globularis* in the Farokhi Formation is not in contradiction with this inferred mode of life as the facies also indicates a relatively low-energy environment below storm wave base.




Fig. 3. Section of the Farokhi Formation at Kuh-e-Honu, Khur area, Central Iran (N 33°43.355', E 55°00.869').

The large perforations in two specimens (Fig. 4L, M) represent sipunculan worm borings named *Trypanites mobilis* by Neumann et al. (2008). The bored sponges are interpreted as mobile shelters for the worms. Neumann et al. (2008, table 1) found that ca. 11% of specimens in *P. globularis* populations were bored, which roughly (i.e., in the absence of a statistically significant population) matches the two bored specimens out of 14 (~14%) from the Farokhi Formation.

The form of *P. globularis*, albeit generally spherical, may be very variable: pea-, pear-, marble-, oval-, loaf- and cushion-shaped specimens occur (e.g., Nestler, 1961, p. 39; Wood, 2002). The different forms and the presence of superficial grooves have also been used to separate species within the genus *Porosphaera* [e.g., *Porosphaera patelliformis* Hinde, 1904, *Porosphaera nuciformis* (von Hagenow, 1839), *Porosphaera plana* Stolley, 1892]. According to this approach, the specimens illustrated in Fig. 4H–K are close to *P. nuciformis*, which is pyriform and in some cases shows superficial grooves radiating from a centre, the latter carrying a tubercle (not seen in the Iranian material and not present either in many *P. nuciformis* illustrated in the literature). However, there seem to be a gradual morphological transitions from globular to pyriform shapes in the sparse material from Iran as weak grooves are also present in some other specimens (e.g., Fig. 4C, D). Furthermore, the internal skeletal structure and surface perforations are identical to those of *P. globularis* (Nestler, 1961, p. 43), while authors such as Steinmann (1878) and Malecki (1989) doubted the validity of von Hagenow's species and regarded it as junior synonym. It is noteworthy that the stratigraphic ranges and distribution of the numerous species of *Porosphaera* recognized seem to be identical to those of the type species and that these taxa tend to be very rare (Nestler, 1961; Termier and Termier, 1985b; Malecki, 1989; Wood, 2002; Jagt and Snellings, 2009). Many of these 'species' may merely represent different growth or preservational forms. However, a taxonomic revision of the genus is beyond the scope of the present paper.

Occurrence. *P. globularis* occurs in the Farokhi Formation of Kuh-e-Honu, ca. 20 km south of Khur, Central Iran (N 33°43.355', E 55°00.869'). Based on the presence of the Late Campanian heteromorph ammonoid *B. polyplacum* a considerable distance below in the underlying Haftoman Formation and the superposition by the Palaeocene Chupanan Formation, the Farokhi Formation at Kuh-e-Honu can be dated as late Late Campanian to Maastrichtian (an exclusively Maastrichtian age being more likely). *P. globularis* is mainly known from the Upper Cretaceous (Cenomanian–Maastrichtian) of Boreal Europe (Nestler, 1961; Termier and Termier, 1985a, b; Reitner, 1992; Wood, 2002). The type locality of *P. globularis* appears to be Danes Dyke in Yorkshire (Hinde, 1904, p. 3), from which a *Marsupites* Zone horizon (latest Santonian) can be inferred (pers. commun. C.J. Wood, 08/2011). Interestingly, the lowermost Maastrichtian strata in the Chalk of Norfolk were termed the *Porosphaera* Beds by Brydone (1908) because of the abundance in them of *Porosphaera*. However, Mortimore et al. (2001, 369) found this idea of a particular abundance difficult to understand but it is well possible that the find layer of *Porosphaera* from the Farokhi Formation is likewise of Early Maastrichtian age.

The record of *P. globularis* from Central Iran represents the southernmost known palaeo-latitude occurrence to date (ca. 15–20°N; Fig. 1) and it is the first mention of the species from the Middle East. However, it has been noted from the Asian part of the former Soviet Union (e.g., Pervushov, 1998). The present record from subtropical settings (rudists are common in the underlying Haftoman Formation) may be explained by the deep-water habitat of the sponge which lived at comparatively cool levels in the water column.




Fig. 4. *Porosphaera globularis* (Phillips, 1829) from the Farokhi Formation of Central Iran. All specimens $\times 3.0$. A, 091014-18/1. B, 091014-18/2. C, 091014-18/7. D, 091014-18/8. E, 091014-18/13. F, 091014-18/14. G, 091014-18/4. H, I, 091014-18/4. J, 091014-18/9. K, 091014-18/10. L, 091014-18/5. M, 091014-18/12.




Fig. 5. Thin sections of *Porosphaera globularis* (Phillips, 1829) from the Farokhi Formation of Central Iran (plane-polarized light). A–C, 091014-18/6. D–F, 091014-18/11. A, near-median cross-section showing the radiating oscular canals ($\times 6.0$). B, close-up of the central part of the choanosomal skeleton (width of photomicrograph 7.5 mm). C, close-up of the marginal part of the choanosomal skeleton (width of photomicrograph 3.75 mm). D, tangential cross-section ($\times 5.0$). E, F close-up of the choanosomal skeleton (widths of photomicrographs 6 and 3.75 mm respectively). G, H, photomicrographs showing fused tetractines of the choanosomal skeleton (widths of photomicrographs 300 and 200 μm respectively).

Acknowledgements

We thank J.W.M Jagt (Maastricht) and C.J. Wood (Minehead) for their helpful reviews and D.J. Batten (Manchester) for his editorial work. Financial support by the Darius Programme is gratefully acknowledged. We thank M. Kaden (MMG, Dresden) for preparing the thin sections, R. Winkler (MMG, Dresden) for photographic work, and K.-A. Tröger (Freiberg) for identification of the inoceramid bivalve. I. Walaszczyk (Warszawa), J.W.M Jagt (Maastricht) and E.A. Jagt-Yazykova (Opole) helped to access obscure literature, while B. Senowbari-Daryan (Erlangen) is thanked for discussion.

References

- Aistov, L., Melnikov, B., Krivyakin, B., Morozov, L., Kiristaev, V., 1984. Geology of the Khur Area (Central Iran). Explanatory Text of the Khur Quadrangle Map 1:250,000, V/O Technoexport, Report 20, pp. 1–130.
- Berberian, M., King, G.C.P., 1981. Towards a palaeogeography and tectonic evolution of Iran. Canadian Journal of Earth Sciences 18, 210–265.
- Brydone, R.M., 1908. On the subdivisions of the chalk of Trimmingham (Norfolk). Quarterly Journal of the Geological Society of London 54, 401–412.
- Cotteau, G., Gauthier, V., 1895. Echinides fossils. In: Morgan, J. de (Ed.), Mission scientifique en Perse. Paléontologie, vol. 3(1), pp. 1–142, 16 pls.
- Davoudzadeh, M., 1997. Iran. In: Moores, E.M., Fairbridge, R.M. (Eds.), Encyclopedia of European and Asian Regional Geology. Encyclopedia of Earth Sciences Series. Chapman & Hall, London, pp. 384–405.

- Davoudzadeh, M., Soffel, H., Schmidt, K., 1981. On the rotation of the central-east Iran microplate. *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte* 1981, 180–192.
- Dercourt, J., Zonenshain, L.P., Ricou, L.-E., Kazmin, V.G., Le Pichon, X., Knipper, A.L., Grandjacquet, C., Sbertshikov, I.M., Geyssant, J., Lepvrier, C., Pechersky, D.H., Boulain, J., Sibuet, J.-C., Savostin, L.A., Sorokhtin, O., Westphal, M., Bazhenov, M.L., Lauer, J.P., Biju-Duval, B., 1986. Geological evolution of the Tethys belt from the Atlantic to the Pamir since the Lias. *Tectonophysics* 123, 241–315.
- Finks, R.M., Reid, R.E.H., Rigby, J.K., 2004. Porifera (Demospongia, Hexactinellida, Heteractinia, Calcarea). In: Moore, R.C., Kaesler, R.L. (Eds.), *Treatise on Invertebrate Palaeontology, Part E Porifera, Revised*, vol. 3. Geological Society of America, Boulder, pp. 1–872.
- Fürsich, F.T., Wilmsen, M., Seyed-Emami, K., Schairer, G., Majidifard, M.R., 2003. Platform/basin transect of a large-scale Middle-Late Jurassic carbonate platform system (Shotori Mountains, Tabas area, east-central Iran). *Facies* 48, 171–198.
- von Hagenow, F., 1839–40. Monographie der Kreide-Versteinerungen Neu-vorpommerns und Rügens. *Neues Jahrbuch für Mineralogie, Geologie und Paläontologie* 1839–40 (Separatdruck 1842), 253–296.
- Hinde, G.J., 1904. On the structure and affinities of the genus *Porosphaera*, Steinmann. *Journal of the Royal Microscopical Society London* 1904, 1–25, 2 pls.
- Hooper, J.N.A., van Soest, R.W.M., 2002. *Systema Porifera: A Guide for the Classifications of Sponges*, vols. 1+2. Plenum Press, New York, pp. i–xlvi, 1–1708.
- Jagt, J.W.M., Snellings, J., 2009. Opmerkelijke Luiks-Limbursse Krijtfofossielen. *Natuurhistorisch Maandblad* 98 (10), 203–205.
- Kennedy, W.J., Chahida, M.R., Djafarian, M.A., 1979. Cenomanian cephalopods from the Glauconitic Limestone southeast of Esfahan, Iran. *Acta Palaeontologica Polonica* 14, 3–50.
- Malecki, J., 1989. The genus *Porosphaera* (Spongiae) from the Lower Campanian sediments in the vicinity of Krakow. *Bulletin of the Polish Academy of Sciences, Earth Sciences* 37, 205–217.
- Müller, A.H., 1970. Über *Porosphaera* (Porifera, Calcarea) und ihr Endolithion. *Monatsberichte der deutschen Akademie der Wissenschaften Berlin* 12, 708–720.
- Mortimore, R.N., Wood, C.J., Gallois, R.W., 2001. British Upper Cretaceous stratigraphy. *Geological Conservation Review Series* 23, 1–558.
- Nestler, H., 1961. Spongien aus der weißen Schreibkreide (Unt. Maastricht) der Insel Rügen (Ostsee). *Paläontologische Abhandlungen* 1, 1–70, pls 1–12.
- Nestler, H., 1965. Die Rekonstruktion des Lebensraumes der Rügener Schreibkreide-Fauna (Unter-Maastricht) mit Hilfe der Paläoökologie und Paläobiologie. *Beihefte zur Zeitschrift Geologie* 49, 1–147.
- Nestler, H., 2002. Die Fossilien der Rügener Schreibkreide. In: *Westarp Wissenschaften, Hohenwarsleben, Die Neue Brehm-Bücherei*, Band 486, pp. 1–129.
- Neumann, C., Wisshak, M., Bromley, R.G., 2008. Boring a mobile domicile: an alternative to the conchicolous life habit. In: Wisshak, M., Tapanila, L. (Eds.), *Current Developments in Bioerosion*. Springer, Berlin, Heidelberg, New York, pp. 307–328.
- Pervushov, E.M., 1998. Pozdneemelovye ventrikulitidnye gubki Povolzh'ya. *Tom 2. Geologii SGU, Saratov; GosUNTs Kolledzh*, pp. 1–168, 42 pls.
- Phillips, J., 1829. *Illustrations of the Geology of Yorkshire. Part 1: The Yorkshire Coast*, Printed for the author by T. Wilson, York, pp. 1–192.
- Philip, J., Floquet, M., 2000. Late Cenomanian. In: Dercourt, J., Gaetani, M., Vrielynck, B., Barrier, E., Biju-Duval, B., Brunet, M.F., Cadet, J.P., Crasquin, S., Sandulescu, M. (Eds.), *Atlas Peri-Tethys Palaeogeographical Maps. CCGM/CGMW*, Paris, pp. 129–136.
- Reich, M., Frenzel, P., 2002. Die Fauna und Flora der Rügener Schreibkreide (Maastrichtium, Ostsee). *Archiv für Geschichtskunde* 3, 73–284.
- Reitner, J., 1992. "Coralline Spongien". *Der Versuch einer phylogenetisch-taxonomischen Analyse. Berliner Geowissenschaftliche Abhandlungen E1*, 1–352, pls 1–62.
- Rigaud, S., d'Errico, F., Vanhaeren, M., Vanhaeren, M., Neumann, C., 2009. Critical reassessment of putative Acheulean *Porosphaera globularis* beads. *Journal of Archaeological Science* 36 (1), 25–34.
- Rosetti, F., Nasrabad, M., Vignaroli, G., Theye, T., Gerdes, A., Razavi, M.H., Vaziri, M., 2010. Early Cretaceous migmatitic granulites from the Sabzevar range (NE Iran): implications for the closure of the Mesozoic peri-Tethyan oceans in central Iran. *Terra Nova* 22, 26–34.
- Seyed-Emami, K., 1977. Scaphiten aus dem Oberen Alb und Cenoman der Umgebung von Esfahan (Zentral-Iran). *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und historische Geologie* 17, 125–136.
- Seyed-Emami, K., 1982. Turrilitidae (Ammonoidea) aus dem Glaukonitkalk bei Esfahan (Zentraliran). *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 163, 417–434.
- Seyed-Emami, K., Immel, H., 1995. Ammoniten aus dem Alb (Kreide) von Shir-Kuh (N Yazd, Zentraliran). *Paläontologische Zeitschrift* 69, 377–399.
- Seyed-Emami, K., Immel, H., 1996. Ammoniten aus dem Alb (höhere Unterkreide) des Zentralirans. *Palaeontographica* A241, 1–26.
- Soffel, H., Davoudzadeh, M., Rolf, C., Schmidt, S., 1996. New palaeomagnetic data from Central Iran and a Triassic palaeoreconstruction. *Geologische Rundschau* 85, 293–302.
- Stampfli, G.M., Borel, G.D., 2002. A plate tectonic model for the Paleozoic and Mesozoic constrained by dynamic plate boundaries and restored synthetic oceanic isochrones. *Earth and Planetary Science Letters* 196, 17–33.
- Steinmann, G., 1878. Über fossile Hydrozoen aus der Familie der Coryniden. *Palaeontographica* A25, 101–124.
- Stolley, E., 1892. Die Kreide Schleswig-Holsteins. *Mitteilungen aus dem Mineralogischen Institut der Universität Kiel* 1, 191–309.
- Termier, H., Termier, G., 1985a. Sponges of Santonian and Campanian age along motorway-A10 (France). *Cretaceous Research* 6, 143–155.
- Termier, H., Termier, G., 1985b. Les Spongiaires du Crétacé Normand. 1ère Partie, Généralités sur les Spongiaires et Ischyrosponges. *Bulletin trimestriel de la Société Géologique de Normandie* 132 (3), 8–89.
- Tirul, R., Bell, I.R., Griffis, R.J., Camp, V.E., 1983. The Sistan suture zone of eastern Iran. *Geological Society of America Bulletin* 94, 134–150.
- Vacelet, J., Borjovic, R., Boury-Esnault, N., Manuel, M., 2002. Order Lithonida Vacelet, 1981, Recent. In: Hooper, J.N.A., van Soest, R.W.M. (Eds.), *Systema Porifera: A Guide for the Classifications of Sponges*, vol. 2. Plenum Press, New York, pp. 1185–1192.
- Wilmsen, M., Fürsich, F.T., Seyed-Emami, K., 2003. Revised lithostratigraphy of the Middle and Upper Jurassic Magu Group of the northern Tabas Block, east-central Iran. *Newsletters on Stratigraphy* 39, 143–156.
- Wilmsen, M., Wiese, F., Seyed-Emami, K., Fürsich, F.T., 2005. First record and significance of Cretaceous (Turonian) ammonites from the Shotori Mountains, east-central Iran. *Cretaceous Research* 26, 181–195.
- Wilmsen, M., Fürsich, F.T., Seyed-Emami, K., Majidifard, M.R., 2009a. An overview of the lithostratigraphy and facies development of the Jurassic System on the Tabas Block, east-central Iran. In: Brunet, M.-F., Wilmsen, M., Granath, J. (Eds.), *South Caspian to Central Iran basins. Geological Society, London, Special Publications* 312, pp. 323–344.
- Wilmsen, M., Fürsich, F.T., Seyed-Emami, K., Majidifard, M.R., Taheri, J., 2009b. The Cimmarian orogeny in northern Iran: tectono-stratigraphic evidence from the foreland. *Terra Nova* 21, 211–218.
- Wilmsen, M., Fürsich, F.T., Seyed-Emami, K., Majidifard, M.R., Zamani-Pedram, M., 2010. Facies analysis of a large-scale Jurassic shelf-lagoon: The Kamar-e-Mehdi Formation of east-central Iran. *Facies* 56, 59–87.
- Wood, R., 2002. Sponges. In: Smith, A.B., Batten, D.J. (Eds.), *Fossils of the Chalk Palaeontological Association Field Guides to Fossils*, second ed., vol. 2, pp. 27–41.