

Statistical Methods for Real-Time Monitoring of Physical Disability in Multiple Sclerosis

Vadim Zipunnikov (JHSPH)


Kathleen Zackowski (KKI)

Actigraphy studies


- ▶ Physical Activity and Aging
- ▶ Heart Failure Hospitalizations: real-time monitoring
- ▶ Aortic Valve Replacement: pre/post-operation
- ▶ Mental Health: Mood Disorders and EMA
- ▶ Physical Activity & Location (GPS)
- ▶ Parkinson Disease: cross-species studies
- ▶ Pediatric Surgery: early dispatch
- ▶ Sleep-wake vs Brain Atrophy & $A\beta$ -amyloid



- Quantified-self (self-chronometry)



Stability/Disruptions in Circadian Rhythms



Wearables

Research



Consumer




What do research grade sensors offer?

- Dense measurements of Physical Activity
 - G's (mg)
 - Activity Counts (per minute)
 - Steps (per minute)
 - Gait (temporal gait asymmetry, stride variability)
 - Energy Expenditure (calories, ...)
 - Sleep (duration, the number of wakes, ...)
- Light, Temperature, others
- Heart Rate (ECG, BPM)
- Voice (Mood, Progression of Disease)
- App-based surveys (2-4 times a day)
- GPS

Accelerometers


- Use sensors to detect accelerations in one-to-three orthogonal planes

- Anteroposterior
- Mediolateral
- Vertical




- Small size: wireless, non-invasive
- Long battery life, can record activity for up to 45 days (and down to 14 days)
- Provides an objective, comprehensive assessment of free-living physical activity

Value of the Raw data



- 400m walk: currently only time (gait speed)



Temporal Gait Parameters





- quality of steps (AM vs PM)
- temporal gait parameters:
 - step asymmetry
 - stride variability
 - instantaneous walking speed

Urbanek, et al, arxiv.org

Minute-level activity count



Minute-level activity profiles



Epidemiological Studies and Clinical/Interventional Trials


- Epidemiological studies:
 - Cross-sectional/one visit: 7 -14 days per visit
 - Age, Sex, BMI,...
 - Nutrition, Heart Diseases, Mood Disorders, ...
- Clinical trials:
 - two visits, multiple visits, continuous monitoring
 - mobile monitoring:
 - comparative effectiveness, pre-/post- intervention
 - progression, recovery
 - early detection (Congestive Heart Failure, Multiple Sclerosis)
 - part of the treatment
 - compliance to treatment
 - FDA: to define endpoints at Clinical Trial of 2020

Multiple Sclerosis (MS)

- Multiple Sclerosis
 - affects physical and cognitive function
 - slow accumulation of physical disability
- Expanded Disability Status Scale (EDSS)
 - non-linear scale
 - poor reliability and poor responsiveness
- **Our proposal:**
 - normalized physical disability score (real-life, accelerometry-based)


Preliminary Data

- Collected data on 10 subjects (1-7 days)
- Patient 1 (red) is a 61 year old female with an EDSS of 6.0
- Patient 2 (blue) is a 54 year old male with an EDSS of 4.0




Preliminary Data

- Accelerometry vs EDSS




- Accelerometry vs t25fw



Preliminary Data

- Slow t25fw (red) vs Fast t25fw (blue)



Our approach

Step 1: Feature Extraction













Step 2: Feature Selection

Step 3: Creating normalized physical-disability score


Step 1: Feature Extraction

- Traditional time-invariant summaries (# of active minutes, total activity, max10, etc.)
- Time-dependent summaries (multiresolution: minute, hour)
- Temporal modeling of activity bouts (duration, frequency, magnitude), switching, fragmentation, stability/variability
- Temporal architecture of PA (functional PCs, landmarks of daily profiles)

Step 2: Feature Selection



Step 3: Creating normalized physical-disability score



Zipunnikov, et al, unpublished data

Challenges

- Main Challenge: Steps 2 and 3
 - identifying features responsive to disability
 - no well-defined “events” (surgery/treatment, hospitalization)

Next Steps

- Collection data on 20 subjects (different levels of disability, 2 weeks each)
- Working out the main challenge