

First Name: _____ **Last Name:** _____

Quiz 4

-
- 5 minute individual quiz;
 - Answer the questions in the space provided. If you run out of space, continue onto the back of the page. Additional space is provided at the end;
 - **Show and explain all work;**
 - **Underline** the answer of each steps;
 - The use of books, personal notes, **calculator**, cellphone, laptop, and communication with others is forbidden;
 - By taking this quiz, you agree to follow the university's code of academic integrity.
-

Exercise 1 40%

Consider the following set of vectors

$$\begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix}, \quad \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \quad \begin{pmatrix} -1 \\ 2 \\ 0 \end{pmatrix}$$

and determine whether they are linearly independent.

Exercise 2 60%

Find *one* eigenvalues and one associated eigenvector of the matrix

$$\begin{pmatrix} -3 & 3/4 \\ -5 & 1 \end{pmatrix}.$$

Quiz 4: solutions

Exercise 1 40%

To check independence, we compute

$$\det \begin{pmatrix} 2 & 1 & -1 \\ 1 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix} = \begin{pmatrix} 2 & 1 & -1 \\ 0 & \frac{1}{2} & \frac{5}{2} \\ 0 & 0 & 0 \end{pmatrix} = 0.$$

We could also have used the row formula

$$\det(A) = \sum_{i=1}^3 (-1)^{i+3} a_{3,i} \det(A(3,i)) = 0$$

because $a_{3,i} = 0$, $i = 1, 2, 3$.

Notice that we can therefore find nonzero $\alpha_i = 1, 2, 3$ such that

$$\alpha_1 \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix} + \alpha_2 \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} + \alpha_3 \begin{pmatrix} -1 \\ 2 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}.$$

Indeed, this is equivalent to

$$\begin{cases} 2\alpha_1 - \alpha_3 = 0 \\ \alpha_1 + \alpha_2 + 2\alpha_3 = 0 \\ 0 = 0 \end{cases}.$$

For instance $\alpha_1 = 1$, $\alpha_2 = -5$, $\alpha_3 = 2$ is a non trivial solution. The vectors are linearly dependent and

$$\begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix} = 5 \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} - 2 \begin{pmatrix} -1 \\ 2 \\ 0 \end{pmatrix}.$$

Exercise 2 60%

The eigenvalues are given by

$$\det(A - \lambda I) = 0.$$

This is

$$\det \begin{pmatrix} -3 - \lambda & 3/4 \\ -5 & 1 - \lambda \end{pmatrix} = 0$$

or computing the determinant

$$\lambda^2 + 2\lambda + 3/4 = 0$$

The solutions of the above equation are given by

$$\lambda_1 = -\frac{1}{2} \quad \lambda_2 = -\frac{3}{2}.$$

The eigenvectors $\xi^1 = (\xi_1, \xi_2)^t$ associated with λ_1 satisfy

$$-5\xi_1 + \frac{3}{2}\xi_2 = 0$$

or

$$\xi^1 = \alpha \begin{pmatrix} 1 \\ -\frac{3}{10} \end{pmatrix}$$

for any constant α . For instance

$$\xi^1 = \begin{pmatrix} 10 \\ -3 \end{pmatrix}$$

The eigenvectors $\xi^2 = (\xi_1, \xi_2)^t$ associated with λ_2 satisfy

$$-3\xi_1 + \frac{3}{2}\xi_2 = 0$$

or

$$\xi^2 = \alpha \begin{pmatrix} 1 \\ \frac{1}{2} \end{pmatrix}$$

for any constant α . For instance,

$$\xi^2 = \begin{pmatrix} 2 \\ 1 \end{pmatrix}.$$