





Meggerdorf

Peatland rewetting – Eider-Treene-Sorge test area

Farmed peat: 1,558 ha
grassland 1,556 ha · cropland 2 ha
Current emissions: ≈ 49,394 t CO2-eq/yr
Saving if all rewetted: ≈ 40,826 t CO2-eq/yr
Farmed peat in chosen catchments: 755 ha (48 %)

Legend

Sub-catchments

- Catchment chosen for rewetting
- Other sub-catchment


Farmed peat


- Grassland on peat
- Cropland on peat

Municipalities (atlas)

- Municipality
- Neighbouring municipality
- Test area

Data: organic soils © Thünen-Institut (Wittnebel et al. 2023, CC BY 4.0); Feldblöcke © Land Schleswig-Holstein 2026 (dl-de/zero-2.0); DGM1 © GeoBasis-DE/LVermGeo SH (CC BY 4.0); VG250 © GeoBasis-DE/BKG (dl-de/by-2.0); background © OpenStreetMap contributors.
Emission factors: Tiemeyer et al. 2020. All emission figures are ESTIMATES (area × national emission factors).





Bergenhusen

Peatland rewetting – Eider-Treene-Sorge test area

Farmed peat: 938 ha
grassland 928 ha · cropland 9 ha
Current emissions: ≈ 29,809 t CO2-eq/yr
Saving if all rewetted: ≈ 24,651 t CO2-eq/yr
Farmed peat in chosen catchments: 344 ha (37 %)

Legend

Sub-catchments

- Catchment chosen for rewetting
- Other sub-catchment


Farmed peat


- Grassland on peat
- Cropland on peat

Municipalities (atlas)

- Municipality
- Neighbouring municipality
- Test area

Data: organic soils © Thünen-Institut (Wittnebel et al. 2023, CC BY 4.0); Feldblöcke © Land Schleswig-Holstein 2026 (dl-de/zero-2.0); DGM1 © GeoBasis-DE/LVermGeo SH (CC BY 4.0); VG250 © GeoBasis-DE/BKG (dl-de/by-2.0); background © OpenStreetMap contributors.
Emission factors: Tiemeyer et al. 2020. All emission figures are ESTIMATES (area × national emission factors).





Wohlde

Peatland rewetting – Eider-Treene-Sorge test area

Farmed peat: 809 ha
grassland 769 ha · cropland 40 ha
Current emissions: ≈ 25,985 t CO2-eq/yr
Saving if all rewetted: ≈ 21,537 t CO2-eq/yr
Farmed peat in chosen catchments: 139 ha (17 %)

Legend

Sub-catchments

- Catchment chosen for rewetting
- Other sub-catchment


Farmed peat


- Grassland on peat
- Cropland on peat

Municipalities (atlas)

- Municipality
- Neighbouring municipality
- Test area

Data: organic soils © Thünen-Institut (Wittnebel et al. 2023, CC BY 4.0); Feldblöcke © Land Schleswig-Holstein 2026 (dl-de/zero-2.0); DGM1 © GeoBasis-DE/LVermGeo SH (CC BY 4.0); VG250 © GeoBasis-DE/BKG (dl-de/by-2.0); background © OpenStreetMap contributors.
Emission factors: Tiemeyer et al. 2020. All emission figures are ESTIMATES (area × national emission factors).





Stapel

Peatland rewetting – Eider-Treene-Sorge test area

Farmed peat: 693 ha
grassland 690 ha · cropland 3 ha
Current emissions: ≈ 21,995 t CO2-eq/yr
Saving if all rewetted: ≈ 18,183 t CO2-eq/yr
Farmed peat in chosen catchments: 75 ha (11 %)

Legend

Sub-catchments

- Catchment chosen for rewetting
- Other sub-catchment


Farmed peat


- Grassland on peat
- Cropland on peat

Municipalities (atlas)

- Municipality
- Neighbouring municipality
- Test area

Data: organic soils © Thünen-Institut (Wittnebel et al. 2023, CC BY 4.0); Feldblöcke © Land Schleswig-Holstein 2026 (dl-de/zero-2.0); DGM1 © GeoBasis-DE/LVermGeo SH (CC BY 4.0); VG250 © GeoBasis-DE/BKG (dl-de/by-2.0); background © OpenStreetMap contributors.
Emission factors: Tiemeyer et al. 2020. All emission figures are ESTIMATES (area × national emission factors).





Börm

Peatland rewetting – Eider-Treene-Sorge test area

Farmed peat: 638 ha
grassland 609 ha · cropland 29 ha
Current emissions: ≈ 20,490 t CO2-eq/yr
Saving if all rewetted: ≈ 16,979 t CO2-eq/yr
Farmed peat in chosen catchments: 177 ha (28 %)

Legend

Sub-catchments

- Catchment chosen for rewetting
- Other sub-catchment


Farmed peat


- Grassland on peat
- Cropland on peat

Municipalities (atlas)

- Municipality
- Neighbouring municipality
- Test area

Data: organic soils © Thünen-Institut (Wittnebel et al. 2023, CC BY 4.0); Feldblöcke © Land Schleswig-Holstein 2026 (dl-de/zero-2.0); DGM1 © GeoBasis-DE/LVermGeo SH (CC BY 4.0); VG250 © GeoBasis-DE/BKG (dl-de/by-2.0); background © OpenStreetMap contributors.
Emission factors: Tiemeyer et al. 2020. All emission figures are ESTIMATES (area × national emission factors).





Hohn

Peatland rewetting – Eider-Treene-Sorge test area

Farmed peat: 637 ha
grassland 620 ha · cropland 17 ha
Current emissions: ≈ 20,332 t CO2-eq/yr
Saving if all rewetted: ≈ 16,830 t CO2-eq/yr
Farmed peat in chosen catchments: 167 ha (26 %)

Legend

Sub-catchments

- Catchment chosen for rewetting
- Other sub-catchment


Farmed peat


- Grassland on peat
- Cropland on peat

Municipalities (atlas)

- Municipality
- Neighbouring municipality
- Test area

Data: organic soils © Thünen-Institut (Wittnebel et al. 2023, CC BY 4.0); Feldblöcke © Land Schleswig-Holstein 2026 (dl-de/zero-2.0); DGM1 © GeoBasis-DE/LVermGeo SH (CC BY 4.0); VG250 © GeoBasis-DE/BKG (dl-de/by-2.0); background © OpenStreetMap contributors.
Emission factors: Tiemeyer et al. 2020. All emission figures are ESTIMATES (area × national emission factors).





Erfde

Peatland rewetting – Eider-Treene-Sorge test area

Farmed peat: 620 ha
grassland 617 ha · cropland 3 ha
Current emissions: ≈ 19,669 t CO2-eq/yr
Saving if all rewetted: ≈ 16,260 t CO2-eq/yr
Farmed peat in chosen catchments: 81 ha (13 %)

Legend

Sub-catchments

- Catchment chosen for rewetting
- Other sub-catchment


Farmed peat


- Grassland on peat
- Cropland on peat

Municipalities (atlas)

- Municipality
- Neighbouring municipality
- Test area

Data: organic soils © Thünen-Institut (Wittnebel et al. 2023, CC BY 4.0); Feldblöcke © Land Schleswig-Holstein 2026 (dl-de/zero-2.0); DGM1 © GeoBasis-DE/LVermGeo SH (CC BY 4.0); VG250 © GeoBasis-DE/BKG (dl-de/by-2.0); background © OpenStreetMap contributors.
Emission factors: Tiemeyer et al. 2020. All emission figures are ESTIMATES (area × national emission factors).





Tetenhusen

Peatland rewetting – Eider-Treene-Sorge test area

Farmed peat: 603 ha
grassland 596 ha · cropland 6 ha
Current emissions: ≈ 19,156 t CO2-eq/yr
Saving if all rewetted: ≈ 15,842 t CO2-eq/yr
Farmed peat in chosen catchments: 144 ha (24 %)

Legend

Sub-catchments

- Catchment chosen for rewetting
- Other sub-catchment


Farmed peat


- Grassland on peat
- Cropland on peat

Municipalities (atlas)

- Municipality
- Neighbouring municipality
- Test area

Data: organic soils © Thünen-Institut (Wittnebel et al. 2023, CC BY 4.0); Feldblöcke © Land Schleswig-Holstein 2026 (dl-de/zero-2.0); DGM1 © GeoBasis-DE/LVermGeo SH (CC BY 4.0); VG250 © GeoBasis-DE/BKG (dl-de/by-2.0); background © OpenStreetMap contributors.
Emission factors: Tiemeyer et al. 2020. All emission figures are ESTIMATES (area × national emission factors).





Ostenfeld (Husum)

Peatland rewetting – Eider-Treene-Sorge test area

Farmed peat: 563 ha
grassland 561 ha · cropland 2 ha
Current emissions: ≈ 17,868 t CO2-eq/yr
Saving if all rewetted: ≈ 14,771 t CO2-eq/yr
Farmed peat in chosen catchments: 258 ha (46 %)

Legend

Sub-catchments

- Catchment chosen for rewetting
- Other sub-catchment


Farmed peat


- Grassland on peat
- Cropland on peat

Municipalities (atlas)

- Municipality
- Neighbouring municipality
- Test area

Data: organic soils © Thünen-Institut (Wittnebel et al. 2023, CC BY 4.0); Feldblöcke © Land Schleswig-Holstein 2026 (dl-de/zero-2.0); DGM1 © GeoBasis-DE/LVermGeo SH (CC BY 4.0); VG250 © GeoBasis-DE/BKG (dl-de/by-2.0); background © OpenStreetMap contributors.
Emission factors: Tiemeyer et al. 2020. All emission figures are ESTIMATES (area × national emission factors).





Königshügel

Peatland rewetting – Eider-Treene-Sorge test area

Farmed peat: 490 ha
grassland 490 ha · cropland 0 ha
Current emissions: ≈ 15,546 t CO2-eq/yr
Saving if all rewetted: ≈ 12,849 t CO2-eq/yr
Farmed peat in chosen catchments: 225 ha (46 %)

Legend

Sub-catchments

- Catchment chosen for rewetting
- Other sub-catchment


Farmed peat


- Grassland on peat
- Cropland on peat

Municipalities (atlas)

- Municipality
- Neighbouring municipality
- Test area

Data: organic soils © Thünen-Institut (Wittnebel et al. 2023, CC BY 4.0); Feldblöcke © Land Schleswig-Holstein 2026 (dl-de/zero-2.0); DGM1 © GeoBasis-DE/LVermGeo SH (CC BY 4.0); VG250 © GeoBasis-DE/BKG (dl-de/by-2.0); background © OpenStreetMap contributors.
Emission factors: Tiemeyer et al. 2020. All emission figures are ESTIMATES (area × national emission factors).





Dörpstedt

Peatland rewetting – Eider-Treene-Sorge test area

Farmed peat: 441 ha
grassland 392 ha · cropland 48 ha
Current emissions: ≈ 14,395 t CO2-eq/yr
Saving if all rewetted: ≈ 11,970 t CO2-eq/yr
No catchment chosen at the 75,000 t target

Legend

Sub-catchments

- Catchment chosen for rewetting
- Other sub-catchment


Farmed peat


- Grassland on peat
- Cropland on peat

Municipalities (atlas)

- Municipality
- Neighbouring municipality
- Test area

Data: organic soils © Thünen-Institut (Wittnebel et al. 2023, CC BY 4.0); Feldblöcke © Land Schleswig-Holstein 2026 (dl-de/zero-2.0); DGM1 © GeoBasis-DE/LVermGeo SH (CC BY 4.0); VG250 © GeoBasis-DE/BKG (dl-de/by-2.0); background © OpenStreetMap contributors.
Emission factors: Tiemeyer et al. 2020. All emission figures are ESTIMATES (area × national emission factors).





Friedrichsholm

Peatland rewetting – Eider-Treene-Sorge test area

Farmed peat: 428 ha
grassland 420 ha · cropland 7 ha
Current emissions: ≈ 13,624 t CO2-eq/yr
Saving if all rewetted: ≈ 11,271 t CO2-eq/yr
Farmed peat in chosen catchments: 19 ha (4 %)

Legend

Sub-catchments

- Catchment chosen for rewetting
- Other sub-catchment


Farmed peat


- Grassland on peat
- Cropland on peat

Municipalities (atlas)

- Municipality
- Neighbouring municipality
- Test area

Data: organic soils © Thünen-Institut (Wittnebel et al. 2023, CC BY 4.0); Feldblöcke © Land Schleswig-Holstein 2026 (dl-de/zero-2.0); DGM1 © GeoBasis-DE/LVermGeo SH (CC BY 4.0); VG250 © GeoBasis-DE/BKG (dl-de/by-2.0); background © OpenStreetMap contributors.
Emission factors: Tiemeyer et al. 2020. All emission figures are ESTIMATES (area × national emission factors).





Alt Bennebek

Peatland rewetting – Eider-Treene-Sorge test area

Farmed peat: 396 ha
grassland 390 ha · cropland 6 ha
Current emissions: ≈ 12,590 t CO2-eq/yr
Saving if all rewetted: ≈ 10,415 t CO2-eq/yr
Farmed peat in chosen catchments: 64 ha (16 %)

Legend

Sub-catchments

- Catchment chosen for rewetting
- Other sub-catchment


Farmed peat


- Grassland on peat
- Cropland on peat

Municipalities (atlas)

- Municipality
- Neighbouring municipality
- Test area

Data: organic soils © Thünen-Institut (Wittnebel et al. 2023, CC BY 4.0); Feldblöcke © Land Schleswig-Holstein 2026 (dl-de/zero-2.0); DGM1 © GeoBasis-DE/LVermGeo SH (CC BY 4.0); VG250 © GeoBasis-DE/BKG (dl-de/by-2.0); background © OpenStreetMap contributors.
Emission factors: Tiemeyer et al. 2020. All emission figures are ESTIMATES (area × national emission factors).





Klein Bennebek

Peatland rewetting – Eider-Treene-Sorge test area

Farmed peat: 343 ha
grassland 334 ha · cropland 9 ha
Current emissions: ≈ 10,962 t CO2-eq/yr
Saving if all rewetted: ≈ 9,074 t CO2-eq/yr
Farmed peat in chosen catchments: 96 ha (28 %)

Legend

Sub-catchments

- Catchment chosen for rewetting
- Other sub-catchment


Farmed peat


- Grassland on peat
- Cropland on peat

Municipalities (atlas)

- Municipality
- Neighbouring municipality
- Test area

Data: organic soils © Thünen-Institut (Wittnebel et al. 2023, CC BY 4.0); Feldblöcke © Land Schleswig-Holstein 2026 (dl-de/zero-2.0); DGM1 © GeoBasis-DE/LVermGeo SH (CC BY 4.0); VG250 © GeoBasis-DE/BKG (dl-de/by-2.0); background © OpenStreetMap contributors.
Emission factors: Tiemeyer et al. 2020. All emission figures are ESTIMATES (area × national emission factors).





Christiansholm

Peatland rewetting – Eider-Treene-Sorge test area

Farmed peat: 310 ha
grassland 304 ha · cropland 7 ha
Current emissions: ≈ 9,895 t CO2-eq/yr
Saving if all rewetted: ≈ 8,188 t CO2-eq/yr
Farmed peat in chosen catchments: 53 ha (17 %)

Legend

Sub-catchments

- Catchment chosen for rewetting
- Other sub-catchment


Farmed peat


- Grassland on peat
- Cropland on peat

Municipalities (atlas)

- Municipality
- Neighbouring municipality
- Test area

Data: organic soils © Thünen-Institut (Wittnebel et al. 2023, CC BY 4.0); Feldblöcke © Land Schleswig-Holstein 2026 (dl-de/zero-2.0); DGM1 © GeoBasis-DE/LVermGeo SH (CC BY 4.0); VG250 © GeoBasis-DE/BKG (dl-de/by-2.0); background © OpenStreetMap contributors.
Emission factors: Tiemeyer et al. 2020. All emission figures are ESTIMATES (area × national emission factors).





Friedrichsgraben

Peatland rewetting – Eider-Treene-Sorge test area

Farmed peat: 231 ha
grassland 230 ha · cropland 1 ha
Current emissions: ≈ 7,330 t CO2-eq/yr
Saving if all rewetted: ≈ 6,060 t CO2-eq/yr
Farmed peat in chosen catchments: 14 ha (6 %)

Legend

Sub-catchments

- Catchment chosen for rewetting
- Other sub-catchment


Farmed peat


- Grassland on peat
- Cropland on peat

Municipalities (atlas)

- Municipality
- Neighbouring municipality
- Test area

Data: organic soils © Thünen-Institut (Wittnebel et al. 2023, CC BY 4.0); Feldblöcke © Land Schleswig-Holstein 2026 (dl-de/zero-2.0); DGM1 © GeoBasis-DE/LVermGeo SH (CC BY 4.0); VG250 © GeoBasis-DE/BKG (dl-de/by-2.0); background © OpenStreetMap contributors.
Emission factors: Tiemeyer et al. 2020. All emission figures are ESTIMATES (area × national emission factors).





Winnert

Peatland rewetting – Eider-Treene-Sorge test area

Farmed peat: 221 ha
grassland 220 ha · cropland 1 ha
Current emissions: ≈ 7,012 t CO2-eq/yr
Saving if all rewetted: ≈ 5,796 t CO2-eq/yr
Farmed peat in chosen catchments: 112 ha (51 %)

Legend

Sub-catchments

- Catchment chosen for rewetting
- Other sub-catchment


Farmed peat


- Grassland on peat
- Cropland on peat

Municipalities (atlas)

- Municipality
- Neighbouring municipality
- Test area

Data: organic soils © Thünen-Institut (Wittnebel et al. 2023, CC BY 4.0); Feldblöcke © Land Schleswig-Holstein 2026 (dl-de/zero-2.0); DGM1 © GeoBasis-DE/LVermGeo SH (CC BY 4.0); VG250 © GeoBasis-DE/BKG (dl-de/by-2.0); background © OpenStreetMap contributors.
Emission factors: Tiemeyer et al. 2020. All emission figures are ESTIMATES (area × national emission factors).





Lohe-Föhrden

Peatland rewetting – Eider-Treene-Sorge test area

Farmed peat: 217 ha
grassland 204 ha · cropland 13 ha
Current emissions: ≈ 6,992 t CO2-eq/yr
Saving if all rewetted: ≈ 5,799 t CO2-eq/yr
Farmed peat in chosen catchments: 50 ha (23 %)

Legend

Sub-catchments

- Catchment chosen for rewetting
- Other sub-catchment


Farmed peat


- Grassland on peat
- Cropland on peat

Municipalities (atlas)

- Municipality
- Neighbouring municipality
- Test area

Data: organic soils © Thünen-Institut (Wittnebel et al. 2023, CC BY 4.0); Feldblöcke © Land Schleswig-Holstein 2026 (dl-de/zero-2.0); DGM1 © GeoBasis-DE/LVermGeo SH (CC BY 4.0); VG250 © GeoBasis-DE/BKG (dl-de/by-2.0); background © OpenStreetMap contributors.
Emission factors: Tiemeyer et al. 2020. All emission figures are ESTIMATES (area × national emission factors).





Schwabstedt

Peatland rewetting – Eider-Treene-Sorge test area

Farmed peat: 206 ha
grassland 186 ha · cropland 21 ha
Current emissions: ≈ 6,712 t CO2-eq/yr
Saving if all rewetted: ≈ 5,578 t CO2-eq/yr
Farmed peat in chosen catchments: 87 ha (42 %)

Legend

Sub-catchments

- Catchment chosen for rewetting
- Other sub-catchment


Farmed peat


- Grassland on peat
- Cropland on peat

Municipalities (atlas)

- Municipality
- Neighbouring municipality
- Test area

Data: organic soils © Thünen-Institut (Wittnebel et al. 2023, CC BY 4.0); Feldblöcke © Land Schleswig-Holstein 2026 (dl-de/zero-2.0); DGM1 © GeoBasis-DE/LVermGeo SH (CC BY 4.0); VG250 © GeoBasis-DE/BKG (dl-de/by-2.0); background © OpenStreetMap contributors.
Emission factors: Tiemeyer et al. 2020. All emission figures are ESTIMATES (area × national emission factors).





Kropp

Peatland rewetting – Eider-Treene-Sorge test area

Farmed peat: 203 ha
grassland 198 ha · cropland 5 ha
Current emissions: ≈ 6,471 t CO2-eq/yr
Saving if all rewetted: ≈ 5,355 t CO2-eq/yr
No catchment chosen at the 75,000 t target

Legend

Sub-catchments

- Catchment chosen for rewetting
- Other sub-catchment


Farmed peat


- Grassland on peat
- Cropland on peat

Municipalities (atlas)

- Municipality
- Neighbouring municipality
- Test area

Data: organic soils © Thünen-Institut (Wittnebel et al. 2023, CC BY 4.0); Feldblöcke © Land Schleswig-Holstein 2026 (dl-de/zero-2.0); DGM1 © GeoBasis-DE/LVermGeo SH (CC BY 4.0); VG250 © GeoBasis-DE/BKG (dl-de/by-2.0); background © OpenStreetMap contributors.
Emission factors: Tiemeyer et al. 2020. All emission figures are ESTIMATES (area × national emission factors).





Hollingstedt

Peatland rewetting – Eider-Treene-Sorge test area

Farmed peat: 163 ha
grassland 129 ha · cropland 34 ha
Current emissions: ≈ 5,447 t CO2-eq/yr
Saving if all rewetted: ≈ 4,553 t CO2-eq/yr
No catchment chosen at the 75,000 t target

Legend

Sub-catchments

- Catchment chosen for rewetting
- Other sub-catchment


Farmed peat


- Grassland on peat
- Cropland on peat

Municipalities (atlas)

- Municipality
- Neighbouring municipality
- Test area

Data: organic soils © Thünen-Institut (Wittnebel et al. 2023, CC BY 4.0); Feldblöcke © Land Schleswig-Holstein 2026 (dl-de/zero-2.0); DGM1 © GeoBasis-DE/LVermGeo SH (CC BY 4.0); VG250 © GeoBasis-DE/BKG (dl-de/by-2.0); background © OpenStreetMap contributors.
Emission factors: Tiemeyer et al. 2020. All emission figures are ESTIMATES (area × national emission factors).





Tielen

Peatland rewetting – Eider-Treene-Sorge test area

Farmed peat: 94 ha
grassland 94 ha · cropland 0 ha
Current emissions: ≈ 2,967 t CO2-eq/yr
Saving if all rewetted: ≈ 2,452 t CO2-eq/yr
No catchment chosen at the 75,000 t target

Legend

Sub-catchments

- Catchment chosen for rewetting
- Other sub-catchment


Farmed peat


- Grassland on peat
- Cropland on peat

Municipalities (atlas)

- Municipality
- Neighbouring municipality
- Test area

Data: organic soils © Thünen-Institut (Wittnebel et al. 2023, CC BY 4.0); Feldblöcke © Land Schleswig-Holstein 2026 (dl-de/zero-2.0); DGM1 © GeoBasis-DE/LVermGeo SH (CC BY 4.0); VG250 © GeoBasis-DE/BKG (dl-de/by-2.0); background © OpenStreetMap contributors.
Emission factors: Tiemeyer et al. 2020. All emission figures are ESTIMATES (area × national emission factors).





Fresendelf

Peatland rewetting – Eider-Treene-Sorge test area

Farmed peat: 85 ha
grassland 76 ha · cropland 9 ha
Current emissions: ≈ 2,776 t CO2-eq/yr
Saving if all rewetted: ≈ 2,308 t CO2-eq/yr
No catchment chosen at the 75,000 t target

Legend

Sub-catchments

- Catchment chosen for rewetting
- Other sub-catchment


Farmed peat


- Grassland on peat
- Cropland on peat

Municipalities (atlas)

- Municipality
- Neighbouring municipality
- Test area

Data: organic soils © Thünen-Institut (Wittnebel et al. 2023, CC BY 4.0); Feldblöcke © Land Schleswig-Holstein 2026 (dl-de/zero-2.0); DGM1 © GeoBasis-DE/LVermGeo SH (CC BY 4.0); VG250 © GeoBasis-DE/BKG (dl-de/by-2.0); background © OpenStreetMap contributors.
Emission factors: Tiemeyer et al. 2020. All emission figures are ESTIMATES (area × national emission factors).





Sophienhamm

Peatland rewetting – Eider-Treene-Sorge test area

Farmed peat: 81 ha
grassland 81 ha · cropland 0 ha
Current emissions: ≈ 2,570 t CO2-eq/yr
Saving if all rewetted: ≈ 2,124 t CO2-eq/yr
No catchment chosen at the 75,000 t target

Legend

Sub-catchments

- Catchment chosen for rewetting
- Other sub-catchment

Farmed peat

- Grassland on peat
- Cropland on peat

Municipalities (atlas)

- Municipality
- Neighbouring municipality
- Test area

Data: organic soils © Thünen-Institut (Wittnebel et al. 2023, CC BY 4.0); Feldblöcke © Land Schleswig-Holstein 2026 (dl-de/zero-2.0); DGM1 © GeoBasis-DE/LVermGeo SH (CC BY 4.0); VG250 © GeoBasis-DE/BKG (dl-de/by-2.0); background © OpenStreetMap contributors.
Emission factors: Tiemeyer et al. 2020. All emission figures are ESTIMATES (area × national emission factors).