

Thermodynamics & Statistical Physics

Chapter 10. Fluctuation

Yuan-Chuan Zou
zouyc@hust.edu.cn

School of Physics, Huazhong University of Science and Technology

December 30, 2013

Table of contents

1 §10. Fluctuation

- 10.1 Quasi-thermodynamics of fluctuation
- 10.5 Brownian motion
- 10.6 Diffusion and temporal correlation of Brownian particle's momentum
- 10.7 Examples of Brownian motion

§10.1 Quasi-thermodynamics of fluctuation

§10.1 Quasi-thermodynamics of fluctuation

- Statistical average: $\overline{B} = \sum_s B_s \rho_s$.

§10.1 Quasi-thermodynamics of fluctuation

- Statistical average: $\overline{B} = \sum_s B_s \rho_s$.
- Deviation to the average: $B - \overline{B}$;

§10.1 Quasi-thermodynamics of fluctuation

- Statistical average: $\overline{B} = \sum_s B_s \rho_s$.
- Deviation to the average: $B - \overline{B}$;
Average of the deviation: $\overline{B - \overline{B}}$

§10.1 Quasi-thermodynamics of fluctuation

- Statistical average: $\overline{B} = \sum_s B_s \rho_s$.
- Deviation to the average: $B - \overline{B}$;
Average of the deviation: $\overline{B - \overline{B}} = \sum_s (B - \overline{B}) \rho_s$

§10.1 Quasi-thermodynamics of fluctuation

- Statistical average: $\overline{B} = \sum_s B_s \rho_s$.

- Deviation to the average: $B - \overline{B}$;

Average of the deviation: $\overline{B - \overline{B}} = \sum_s (B - \overline{B}) \rho_s$

$$= \sum_s B_s \rho_s - \overline{B} \sum_s \rho_s$$

§10.1 Quasi-thermodynamics of fluctuation

- Statistical average: $\overline{B} = \sum_s B_s \rho_s$.

- Deviation to the average: $B - \overline{B}$;

Average of the deviation: $\overline{B - \overline{B}} = \sum_s (B - \overline{B}) \rho_s$

$$= \sum_s B_s \rho_s - \overline{B} \sum_s \rho_s = \overline{B} - \overline{B} = 0.$$

§10.1 Quasi-thermodynamics of fluctuation

- Statistical average: $\overline{B} = \sum_s B_s \rho_s$.
- Deviation to the average: $B - \overline{B}$;
Average of the deviation: $\overline{B - \overline{B}} = \sum_s (B - \overline{B}) \rho_s$
 $= \sum_s B_s \rho_s - \overline{B} \sum_s \rho_s = \overline{B} - \overline{B} = 0$.
- Use $\overline{(B - \overline{B})^2}$ to indicate the fluctuation.

§10.1 Quasi-thermodynamics of fluctuation

- Statistical average: $\overline{B} = \sum_s B_s \rho_s$.
- Deviation to the average: $B - \overline{B}$;
Average of the deviation: $\overline{B - \overline{B}} = \sum_s (B - \overline{B}) \rho_s$
$$= \sum_s B_s \rho_s - \overline{B} \sum_s \rho_s = \overline{B} - \overline{B} = 0.$$
- Use $\overline{(B - \overline{B})^2}$ to indicate the fluctuation.
$$\overline{(B - \overline{B})^2} = \sum_s \rho_s (B_s - \overline{B})^2$$

§10.1 Quasi-thermodynamics of fluctuation

- Statistical average: $\overline{B} = \sum_s B_s \rho_s$.
- Deviation to the average: $B - \overline{B}$;
Average of the deviation: $\overline{B - \overline{B}} = \sum_s (B - \overline{B}) \rho_s$
$$= \sum_s B_s \rho_s - \overline{B} \sum_s \rho_s = \overline{B} - \overline{B} = 0.$$
- Use $\overline{(B - \overline{B})^2}$ to indicate the fluctuation.
$$\overline{(B - \overline{B})^2} = \sum_s \rho_s (B_s - \overline{B})^2$$
$$= \sum_s \rho_s B_s^2 - 2\overline{B} \sum_s \rho_s B_s + \overline{B}^2$$

§10.1 Quasi-thermodynamics of fluctuation

- Statistical average: $\overline{B} = \sum_s B_s \rho_s$.
- Deviation to the average: $B - \overline{B}$;
Average of the deviation: $\overline{B - \overline{B}} = \sum_s (B - \overline{B}) \rho_s$
$$= \sum_s B_s \rho_s - \overline{B} \sum_s \rho_s = \overline{B} - \overline{B} = 0.$$
- Use $\overline{(B - \overline{B})^2}$ to indicate the fluctuation.
$$\overline{(B - \overline{B})^2} = \sum_s \rho_s (B_s - \overline{B})^2$$
$$= \sum_s \rho_s B_s^2 - 2\overline{B} \sum_s \rho_s B_s + \overline{B}^2 = \overline{B^2} - \overline{B}^2.$$

§10.1 Quasi-thermodynamics of fluctuation

- Statistical average: $\overline{B} = \sum_s B_s \rho_s$.
- Deviation to the average: $B - \overline{B}$;
Average of the deviation: $\overline{B - \overline{B}} = \sum_s (B - \overline{B}) \rho_s$

$$= \sum_s B_s \rho_s - \overline{B} \sum_s \rho_s = \overline{B} - \overline{B} = 0.$$
- Use $\overline{(B - \overline{B})^2}$ to indicate the fluctuation.

$$\overline{(B - \overline{B})^2} = \sum_s \rho_s (B_s - \overline{B})^2$$

$$= \sum_s \rho_s B_s^2 - 2\overline{B} \sum_s \rho_s B_s + \overline{B}^2 = \overline{B^2} - \overline{B}^2.$$
- Fluctuation of thermal parameters: $N, V, E; S, T$.

§10.1 Quasi-thermodynamics of fluctuation

- Statistical average: $\overline{B} = \sum_s B_s \rho_s$.
- Deviation to the average: $B - \overline{B}$;
Average of the deviation: $\overline{B - \overline{B}} = \sum_s (B - \overline{B}) \rho_s$

$$= \sum_s B_s \rho_s - \overline{B} \sum_s \rho_s = \overline{B} - \overline{B} = 0.$$
- Use $\overline{(B - \overline{B})^2}$ to indicate the fluctuation.

$$\overline{(B - \overline{B})^2} = \sum_s \rho_s (B_s - \overline{B})^2$$

$$= \sum_s \rho_s B_s^2 - 2\overline{B} \sum_s \rho_s B_s + \overline{B}^2 = \overline{B^2} - \overline{B}^2.$$
- Fluctuation of thermal parameters: $N, V, E; S, T$.
For S (or T), means $\overline{[S(N, V, E) - S(\overline{N}, \overline{V}, \overline{E})]^2}$.

Fluctuation of canonical ensemble

Fluctuation of canonical ensemble

- Canonical ensemble:

Fluctuation of canonical ensemble

- Canonical ensemble:

Consider the combined system, it has:

$$\Delta E + \Delta E_r = 0, \Delta V + \Delta V_r = 0.$$

Fluctuation of canonical ensemble

- Canonical ensemble:

Consider the combined system, it has:

$$\Delta E + \Delta E_r = 0, \Delta V + \Delta V_r = 0.$$

- (\bar{E}, \bar{V}) is the most probable value, the corresponding number of states of the combined system: $\bar{\Omega}^{(0)}$;

Fluctuation of canonical ensemble

- Canonical ensemble:

Consider the combined system, it has:

$$\Delta E + \Delta E_r = 0, \Delta V + \Delta V_r = 0.$$

- (\bar{E}, \bar{V}) is the most probable value, the corresponding number of states of the combined system: $\bar{\Omega}^{(0)}$; the corresponding entropy: $\bar{S}^{(0)} = k \ln \bar{\Omega}^{(0)}$.

Fluctuation of canonical ensemble

- Canonical ensemble:

Consider the combined system, it has:

$$\Delta E + \Delta E_r = 0, \Delta V + \Delta V_r = 0.$$

- (\bar{E}, \bar{V}) is the most probable value, the corresponding number of states of the combined system: $\bar{\Omega}^{(0)}$; the corresponding entropy: $\bar{S}^{(0)} = k \ln \bar{\Omega}^{(0)}$.
- Any state: (E, V) , the corresponding number of states of the combined system: $\Omega^{(0)}$,

Fluctuation of canonical ensemble

- Canonical ensemble:

Consider the combined system, it has:

$$\Delta E + \Delta E_r = 0, \Delta V + \Delta V_r = 0.$$

- (\bar{E}, \bar{V}) is the most probable value, the corresponding number of states of the combined system: $\bar{\Omega}^{(0)}$; the corresponding entropy: $\bar{S}^{(0)} = k \ln \bar{\Omega}^{(0)}$.
- Any state: (E, V) , the corresponding number of states of the combined system: $\Omega^{(0)}$, and entropy: $S^{(0)}(E, V) = k \ln \Omega^{(0)}$.

Fluctuation of canonical ensemble

- Canonical ensemble:

Consider the combined system, it has:

$$\Delta E + \Delta E_r = 0, \Delta V + \Delta V_r = 0.$$

- (\bar{E}, \bar{V}) is the most probable value, the corresponding number of states of the combined system: $\bar{\Omega}^{(0)}$; the corresponding entropy: $\bar{S}^{(0)} = k \ln \bar{\Omega}^{(0)}$.
- Any state: (E, V) , the corresponding number of states of the combined system: $\Omega^{(0)}$, and entropy: $S^{(0)}(E, V) = k \ln \Omega^{(0)}$.
- The relative probability for (E, V) comparing to (\bar{E}, \bar{V}) : $w = \frac{W}{W_{\max}} = \frac{\Omega^{(0)}}{\bar{\Omega}^{(0)}} = e^{\frac{\Delta S^{(0)}}{k}}$

Fluctuation of canonical ensemble

- Canonical ensemble:

Consider the combined system, it has:

$$\Delta E + \Delta E_r = 0, \Delta V + \Delta V_r = 0.$$

- (\bar{E}, \bar{V}) is the most probable value, the corresponding number of states of the combined system: $\bar{\Omega}^{(0)}$; the corresponding entropy: $\bar{S}^{(0)} = k \ln \bar{\Omega}^{(0)}$.
- Any state: (E, V) , the corresponding number of states of the combined system: $\Omega^{(0)}$, and entropy: $S^{(0)}(E, V) = k \ln \Omega^{(0)}$.
- The relative probability for (E, V) comparing to (\bar{E}, \bar{V}) : $w = \frac{W}{W_{\max}} = \frac{\Omega^{(0)}}{\bar{\Omega}^{(0)}} = e^{\frac{\Delta S^{(0)}}{k}} \Rightarrow W \propto e^{\frac{\Delta S^{(0)}}{k}}$.

Fluctuation of canonical ensemble

- Canonical ensemble:

Consider the combined system, it has:

$$\Delta E + \Delta E_r = 0, \Delta V + \Delta V_r = 0.$$

- (\bar{E}, \bar{V}) is the most probable value, the corresponding number of states of the combined system: $\bar{\Omega}^{(0)}$; the corresponding entropy: $\bar{S}^{(0)} = k \ln \bar{\Omega}^{(0)}$.
- Any state: (E, V) , the corresponding number of states of the combined system: $\Omega^{(0)}$, and entropy: $S^{(0)}(E, V) = k \ln \Omega^{(0)}$.
- The relative probability for (E, V) comparing to (\bar{E}, \bar{V}) : $w = \frac{W}{W_{\max}} = \frac{\Omega^{(0)}}{\bar{\Omega}^{(0)}} = e^{\frac{\Delta S^{(0)}}{k}} \Rightarrow W \propto e^{\frac{\Delta S^{(0)}}{k}}$.

Fluctuation of canonical ensemble

$$W \propto e^{\frac{\Delta S^{(0)}}{k}}$$

- 1. Extensive property of the entropy:

$$\Delta S^{(0)} = \Delta S + \Delta S_r,$$

$$\text{where } \Delta S = S - \overline{S}, \Delta S_r = S_r - \overline{S_r}.$$

Fluctuation of canonical ensemble

$$W \propto e^{\frac{\Delta S^{(0)}}{k}}$$

- 1. Extensive property of the entropy:

$$\Delta S^{(0)} = \Delta S + \Delta S_r,$$

$$\text{where } \Delta S = S - \overline{S}, \Delta S_r = S_r - \overline{S}_r.$$

- 2. The basic equation: $\Delta S_r = \frac{\Delta E_r + p \Delta V_r}{T}$

Fluctuation of canonical ensemble

$$W \propto e^{\frac{\Delta S^{(0)}}{k}}$$

- 1. Extensive property of the entropy:

$$\Delta S^{(0)} = \Delta S + \Delta S_r,$$

$$\text{where } \Delta S = S - \overline{S}, \Delta S_r = S_r - \overline{S}_r.$$

- 2. The basic equation: $\Delta S_r = \frac{\Delta E_r + p \Delta V_r}{T} = -\frac{\Delta E + p \Delta V}{T}.$

Fluctuation of canonical ensemble

$$W \propto e^{\frac{\Delta S^{(0)}}{k}}$$

- 1. Extensive property of the entropy:

$$\Delta S^{(0)} = \Delta S + \Delta S_r,$$

$$\text{where } \Delta S = S - \overline{S}, \Delta S_r = S_r - \overline{S_r}.$$

- 2. The basic equation: $\Delta S_r = \frac{\Delta E_r + p \Delta V_r}{T} = -\frac{\Delta E + p \Delta V}{T}.$
- $\Rightarrow W \propto e^{\frac{\Delta S^{(0)}}{k}}$

Fluctuation of canonical ensemble

$$W \propto e^{\frac{\Delta S^{(0)}}{k}}$$

- 1. Extensive property of the entropy:

$$\Delta S^{(0)} = \Delta S + \Delta S_r,$$

$$\text{where } \Delta S = S - \overline{S}, \Delta S_r = S_r - \overline{S}_r.$$

- 2. The basic equation: $\Delta S_r = \frac{\Delta E_r + p \Delta V_r}{T} = -\frac{\Delta E + p \Delta V}{T}.$
- $\Rightarrow W \propto e^{\frac{\Delta S^{(0)}}{k}} \propto e^{\frac{\Delta S + \Delta S_r}{k}}$

Fluctuation of canonical ensemble

$$W \propto e^{\frac{\Delta S^{(0)}}{k}}$$

- 1. Extensive property of the entropy:

$$\Delta S^{(0)} = \Delta S + \Delta S_r,$$

$$\text{where } \Delta S = S - \bar{S}, \Delta S_r = S_r - \bar{S}_r.$$

- 2. The basic equation: $\Delta S_r = \frac{\Delta E_r + p \Delta V_r}{T} = -\frac{\Delta E + p \Delta V}{T}.$
- $\Rightarrow W \propto e^{\frac{\Delta S^{(0)}}{k}} \propto e^{\frac{\Delta S + \Delta S_r}{k}} \propto e^{\frac{\Delta S - \frac{\Delta E + p \Delta V}{T}}{k}}$

Fluctuation of canonical ensemble

$$W \propto e^{\frac{\Delta S^{(0)}}{k}}$$

- 1. Extensive property of the entropy:

$$\Delta S^{(0)} = \Delta S + \Delta S_r,$$

$$\text{where } \Delta S = S - \bar{S}, \Delta S_r = S_r - \bar{S}_r.$$

- 2. The basic equation: $\Delta S_r = \frac{\Delta E_r + p \Delta V_r}{T} = -\frac{\Delta E + p \Delta V}{T}.$
- $\Rightarrow W \propto e^{\frac{\Delta S^{(0)}}{k}} \propto e^{\frac{\Delta S + \Delta S_r}{k}} \propto e^{\frac{\Delta S - \frac{\Delta E + p \Delta V}{T}}{k}}$
 $\propto e^{-\frac{\Delta E - T \Delta S + p \Delta V}{kT}}.$

Fluctuation of canonical ensemble

$$W \propto e^{\frac{\Delta S^{(0)}}{k}}$$

- 1. Extensive property of the entropy:

$$\Delta S^{(0)} = \Delta S + \Delta S_r,$$

$$\text{where } \Delta S = S - \bar{S}, \Delta S_r = S_r - \bar{S}_r.$$

- 2. The basic equation: $\Delta S_r = \frac{\Delta E_r + p \Delta V_r}{T} = -\frac{\Delta E + p \Delta V}{T}.$
- $\Rightarrow W \propto e^{\frac{\Delta S^{(0)}}{k}} \propto e^{\frac{\Delta S + \Delta S_r}{k}} \propto e^{\frac{\Delta S - \frac{\Delta E + p \Delta V}{T}}{k}}$
 $\propto e^{-\frac{\Delta E - T \Delta S + p \Delta V}{kT}}.$
- Take E as $E(S, V)$, expand at (\bar{S}, \bar{V}) (labeled as 0):

Fluctuation of canonical ensemble

$$W \propto e^{\frac{\Delta S^{(0)}}{k}}$$

- 1. Extensive property of the entropy:

$$\Delta S^{(0)} = \Delta S + \Delta S_r,$$

$$\text{where } \Delta S = S - \bar{S}, \Delta S_r = S_r - \bar{S}_r.$$

- 2. The basic equation: $\Delta S_r = \frac{\Delta E_r + p \Delta V_r}{T} = -\frac{\Delta E + p \Delta V}{T}.$

$$\Rightarrow W \propto e^{\frac{\Delta S^{(0)}}{k}} \propto e^{\frac{\Delta S + \Delta S_r}{k}} \propto e^{\frac{\Delta S - \frac{\Delta E + p \Delta V}{T}}{k}} \\ \propto e^{-\frac{\Delta E - T \Delta S + p \Delta V}{kT}}.$$

- Take E as $E(S, V)$, expand at (\bar{S}, \bar{V}) (labeled as 0):

$$E = \bar{E} + \left(\frac{\partial E}{\partial S}\right)_0 \Delta S + \left(\frac{\partial E}{\partial V}\right)_0 \Delta V + \frac{1}{2} \left[\left(\frac{\partial^2 E}{\partial S^2}\right)_0 (\Delta S)^2 + 2 \left(\frac{\partial^2 E}{\partial S \partial V}\right)_0 \Delta S \Delta V + \left(\frac{\partial^2 E}{\partial V^2}\right)_0 (\Delta V)^2 \right].$$

Fluctuation of canonical ensemble

$$W \propto e^{-\frac{\Delta E - T\Delta S + p\Delta V}{kT}}$$

- $dE = TdS - pdV$, $\Rightarrow (\frac{\partial E}{\partial S})_0 = T$, $(\frac{\partial E}{\partial V})_0 = -p$, then

Fluctuation of canonical ensemble

$$W \propto e^{-\frac{\Delta E - T\Delta S + p\Delta V}{kT}}$$

- $dE = TdS - pdV$, $\Rightarrow (\frac{\partial E}{\partial S})_0 = T$, $(\frac{\partial E}{\partial V})_0 = -p$, then

$$E = \bar{E} + T\Delta S - p\Delta V + \frac{1}{2} \left[\frac{\partial}{\partial S} \left(\frac{\partial E}{\partial S} \right) (\Delta S)^2 + \frac{\partial}{\partial S} \left(\frac{\partial E}{\partial V} \right) \Delta S \Delta V + \frac{\partial}{\partial V} \left(\frac{\partial E}{\partial S} \right) \Delta S \Delta V + \frac{\partial}{\partial V} \left(\frac{\partial E}{\partial V} \right) (\Delta V)^2 \right]$$

Fluctuation of canonical ensemble

$$W \propto e^{-\frac{\Delta E - T\Delta S + p\Delta V}{kT}}$$

- $dE = TdS - pdV$, $\Rightarrow (\frac{\partial E}{\partial S})_0 = T$, $(\frac{\partial E}{\partial V})_0 = -p$, then

$$E = \bar{E} + T\Delta S - p\Delta V + \frac{1}{2} \left[\frac{\partial}{\partial S} \left(\frac{\partial E}{\partial S} \right) (\Delta S)^2 + \frac{\partial}{\partial S} \left(\frac{\partial E}{\partial V} \right) \Delta S \Delta V + \frac{\partial}{\partial V} \left(\frac{\partial E}{\partial S} \right) \Delta S \Delta V + \frac{\partial}{\partial V} \left(\frac{\partial E}{\partial V} \right) (\Delta V)^2 \right]$$

$$= \bar{E} + T\Delta S - p\Delta V + \frac{1}{2} \left[\frac{\partial T}{\partial S} (\Delta S)^2 + \frac{\partial(-p)}{\partial S} \Delta S \Delta V + \frac{\partial T}{\partial V} \Delta S \Delta V + \frac{\partial(-p)}{\partial V} (\Delta V)^2 \right]$$

Fluctuation of canonical ensemble

$$W \propto e^{-\frac{\Delta E - T\Delta S + p\Delta V}{kT}}$$

- $dE = TdS - pdV, \Rightarrow (\frac{\partial E}{\partial S})_0 = T, (\frac{\partial E}{\partial V})_0 = -p$, then

$$\begin{aligned}
 E &= \bar{E} + T\Delta S - p\Delta V + \frac{1}{2} \left[\frac{\partial}{\partial S} \left(\frac{\partial E}{\partial S} \right) (\Delta S)^2 + \right. \\
 &\quad \left. \frac{\partial}{\partial S} \left(\frac{\partial E}{\partial V} \right) \Delta S \Delta V + \frac{\partial}{\partial V} \left(\frac{\partial E}{\partial S} \right) \Delta S \Delta V + \frac{\partial}{\partial V} \left(\frac{\partial E}{\partial V} \right) (\Delta V)^2 \right] \\
 &= \bar{E} + T\Delta S - p\Delta V + \frac{1}{2} \left[\frac{\partial T}{\partial S} (\Delta S)^2 + \frac{\partial(-p)}{\partial S} \Delta S \Delta V + \right. \\
 &\quad \left. \frac{\partial T}{\partial V} \Delta S \Delta V + \frac{\partial(-p)}{\partial V} (\Delta V)^2 \right] \\
 &= \bar{E} + T\Delta S - p\Delta V + \frac{1}{2} \left[\left(\frac{\partial T}{\partial S} \Delta S + \frac{\partial T}{\partial V} \Delta V \right) \Delta S - \right. \\
 &\quad \left. \left(\frac{\partial p}{\partial S} \Delta S + \frac{\partial p}{\partial V} \Delta V \right) \Delta V \right]
 \end{aligned}$$

Fluctuation of canonical ensemble

$$W \propto e^{-\frac{\Delta E - T\Delta S + p\Delta V}{kT}}$$

- $dE = TdS - pdV$, $\Rightarrow (\frac{\partial E}{\partial S})_0 = T$, $(\frac{\partial E}{\partial V})_0 = -p$, then

$$\begin{aligned}
 E &= \bar{E} + T\Delta S - p\Delta V + \frac{1}{2} \left[\frac{\partial}{\partial S} \left(\frac{\partial E}{\partial S} \right) (\Delta S)^2 + \right. \\
 &\quad \left. \frac{\partial}{\partial S} \left(\frac{\partial E}{\partial V} \right) \Delta S \Delta V + \frac{\partial}{\partial V} \left(\frac{\partial E}{\partial S} \right) \Delta S \Delta V + \frac{\partial}{\partial V} \left(\frac{\partial E}{\partial V} \right) (\Delta V)^2 \right] \\
 &= \bar{E} + T\Delta S - p\Delta V + \frac{1}{2} \left[\frac{\partial T}{\partial S} (\Delta S)^2 + \frac{\partial(-p)}{\partial S} \Delta S \Delta V + \right. \\
 &\quad \left. \frac{\partial T}{\partial V} \Delta S \Delta V + \frac{\partial(-p)}{\partial V} (\Delta V)^2 \right] \\
 &= \bar{E} + T\Delta S - p\Delta V + \frac{1}{2} \left[\left(\frac{\partial T}{\partial S} \Delta S + \frac{\partial T}{\partial V} \Delta V \right) \Delta S - \right. \\
 &\quad \left. \left(\frac{\partial p}{\partial S} \Delta S + \frac{\partial p}{\partial V} \Delta V \right) \Delta V \right] \\
 &= \bar{E} + T\Delta S - p\Delta V + \frac{1}{2} (\Delta T \Delta S - \Delta p \Delta V).
 \end{aligned}$$

Fluctuation of canonical ensemble

$$W \propto e^{-\frac{\Delta E - T\Delta S + p\Delta V}{kT}}$$

- $dE = TdS - pdV$, $\Rightarrow (\frac{\partial E}{\partial S})_0 = T$, $(\frac{\partial E}{\partial V})_0 = -p$, then

$$\begin{aligned}
 E &= \bar{E} + T\Delta S - p\Delta V + \frac{1}{2} \left[\frac{\partial}{\partial S} \left(\frac{\partial E}{\partial S} \right) (\Delta S)^2 + \frac{\partial}{\partial S} \left(\frac{\partial E}{\partial V} \right) \Delta S \Delta V + \frac{\partial}{\partial V} \left(\frac{\partial E}{\partial S} \right) \Delta S \Delta V + \frac{\partial}{\partial V} \left(\frac{\partial E}{\partial V} \right) (\Delta V)^2 \right] \\
 &= \bar{E} + T\Delta S - p\Delta V + \frac{1}{2} \left[\frac{\partial T}{\partial S} (\Delta S)^2 + \frac{\partial(-p)}{\partial S} \Delta S \Delta V + \frac{\partial T}{\partial V} \Delta S \Delta V + \frac{\partial(-p)}{\partial V} (\Delta V)^2 \right] \\
 &= \bar{E} + T\Delta S - p\Delta V + \frac{1}{2} \left[\left(\frac{\partial T}{\partial S} \Delta S + \frac{\partial T}{\partial V} \Delta V \right) \Delta S - \left(\frac{\partial p}{\partial S} \Delta S + \frac{\partial p}{\partial V} \Delta V \right) \Delta V \right] \\
 &= \bar{E} + T\Delta S - p\Delta V + \frac{1}{2} (\Delta T \Delta S - \Delta p \Delta V).
 \end{aligned}$$
- $\Rightarrow \Delta E - T\Delta S + p\Delta V = \frac{1}{2} (\Delta T \Delta S - \Delta p \Delta V)$

Fluctuation of canonical ensemble

$$W \propto e^{-\frac{\Delta E - T\Delta S + p\Delta V}{kT}}$$

- $dE = TdS - pdV$, $\Rightarrow (\frac{\partial E}{\partial S})_0 = T$, $(\frac{\partial E}{\partial V})_0 = -p$, then

$$\begin{aligned}
 E &= \bar{E} + T\Delta S - p\Delta V + \frac{1}{2} \left[\frac{\partial}{\partial S} \left(\frac{\partial E}{\partial S} \right) (\Delta S)^2 + \frac{\partial}{\partial S} \left(\frac{\partial E}{\partial V} \right) \Delta S \Delta V + \frac{\partial}{\partial V} \left(\frac{\partial E}{\partial S} \right) \Delta S \Delta V + \frac{\partial}{\partial V} \left(\frac{\partial E}{\partial V} \right) (\Delta V)^2 \right] \\
 &= \bar{E} + T\Delta S - p\Delta V + \frac{1}{2} \left[\frac{\partial T}{\partial S} (\Delta S)^2 + \frac{\partial(-p)}{\partial S} \Delta S \Delta V + \frac{\partial T}{\partial V} \Delta S \Delta V + \frac{\partial(-p)}{\partial V} (\Delta V)^2 \right] \\
 &= \bar{E} + T\Delta S - p\Delta V + \frac{1}{2} \left[\left(\frac{\partial T}{\partial S} \Delta S + \frac{\partial T}{\partial V} \Delta V \right) \Delta S - \left(\frac{\partial p}{\partial S} \Delta S + \frac{\partial p}{\partial V} \Delta V \right) \Delta V \right] \\
 &= \bar{E} + T\Delta S - p\Delta V + \frac{1}{2} (\Delta T \Delta S - \Delta p \Delta V).
 \end{aligned}$$
- $\Rightarrow \Delta E - T\Delta S + p\Delta V = \frac{1}{2} (\Delta T \Delta S - \Delta p \Delta V)$
- $\Rightarrow W \propto e^{-\frac{\Delta T \Delta S - \Delta p \Delta V}{2kT}}.$

Fluctuation of canonical ensemble

$$W \propto e^{-\frac{\Delta T \Delta S - \Delta p \Delta V}{2kT}}$$

- If take (T, V) as variable,

Fluctuation of canonical ensemble

$$W \propto e^{-\frac{\Delta T \Delta S - \Delta p \Delta V}{2kT}}$$

- If take (T, V) as variable,
$$\Delta S = \left(\frac{\partial S}{\partial T}\right)_V \Delta T + \left(\frac{\partial S}{\partial V}\right)_T \Delta V$$

Fluctuation of canonical ensemble

$$W \propto e^{-\frac{\Delta T \Delta S - \Delta p \Delta V}{2kT}}$$

- If take (T, V) as variable,

$$\Delta S = \left(\frac{\partial S}{\partial T}\right)_V \Delta T + \left(\frac{\partial S}{\partial V}\right)_T \Delta V = \frac{C_V}{T} \Delta T + \left(\frac{\partial p}{\partial T}\right)_V \Delta V;$$

Fluctuation of canonical ensemble

$$W \propto e^{-\frac{\Delta T \Delta S - \Delta p \Delta V}{2kT}}$$

- If take (T, V) as variable,

$$\Delta S = \left(\frac{\partial S}{\partial T}\right)_V \Delta T + \left(\frac{\partial S}{\partial V}\right)_T \Delta V = \frac{C_V}{T} \Delta T + \left(\frac{\partial p}{\partial T}\right)_V \Delta V;$$

$$\Delta p = \left(\frac{\partial p}{\partial T}\right)_V \Delta T + \left(\frac{\partial p}{\partial V}\right)_T \Delta V.$$

Fluctuation of canonical ensemble

$$W \propto e^{-\frac{\Delta T \Delta S - \Delta p \Delta V}{2kT}}$$

- If take (T, V) as variable,

$$\Delta S = \left(\frac{\partial S}{\partial T}\right)_V \Delta T + \left(\frac{\partial S}{\partial V}\right)_T \Delta V = \frac{C_V}{T} \Delta T + \left(\frac{\partial p}{\partial T}\right)_V \Delta V;$$

$$\Delta p = \left(\frac{\partial p}{\partial T}\right)_V \Delta T + \left(\frac{\partial p}{\partial V}\right)_T \Delta V.$$

$$\therefore \Delta T \Delta S - \Delta p \Delta V = \frac{C_V}{T} (\Delta T)^2 + \left(\frac{\partial p}{\partial T}\right)_V \Delta T \Delta V - \left(\frac{\partial p}{\partial T}\right)_V \Delta T \Delta V - \left(\frac{\partial p}{\partial V}\right)_T (\Delta V)^2$$

Fluctuation of canonical ensemble

$$W \propto e^{-\frac{\Delta T \Delta S - \Delta p \Delta V}{2kT}}$$

- If take (T, V) as variable,

$$\Delta S = \left(\frac{\partial S}{\partial T}\right)_V \Delta T + \left(\frac{\partial S}{\partial V}\right)_T \Delta V = \frac{C_V}{T} \Delta T + \left(\frac{\partial p}{\partial T}\right)_V \Delta V;$$

$$\Delta p = \left(\frac{\partial p}{\partial T}\right)_V \Delta T + \left(\frac{\partial p}{\partial V}\right)_T \Delta V.$$

$$\begin{aligned} \therefore \Delta T \Delta S - \Delta p \Delta V &= \frac{C_V}{T} (\Delta T)^2 + \left(\frac{\partial p}{\partial T}\right)_V \Delta T \Delta V - \\ &\quad \left(\frac{\partial p}{\partial T}\right)_V \Delta T \Delta V - \left(\frac{\partial p}{\partial V}\right)_T (\Delta V)^2 \\ &= \frac{C_V}{T} (\Delta T)^2 - \left(\frac{\partial p}{\partial V}\right)_T (\Delta V)^2 \end{aligned}$$

Fluctuation of canonical ensemble

$$W \propto e^{-\frac{\Delta T \Delta S - \Delta p \Delta V}{2kT}}$$

- If take (T, V) as variable,

$$\Delta S = \left(\frac{\partial S}{\partial T}\right)_V \Delta T + \left(\frac{\partial S}{\partial V}\right)_T \Delta V = \frac{C_V}{T} \Delta T + \left(\frac{\partial p}{\partial T}\right)_V \Delta V;$$

$$\Delta p = \left(\frac{\partial p}{\partial T}\right)_V \Delta T + \left(\frac{\partial p}{\partial V}\right)_T \Delta V.$$

$$\begin{aligned} \therefore \Delta T \Delta S - \Delta p \Delta V &= \frac{C_V}{T} (\Delta T)^2 + \left(\frac{\partial p}{\partial T}\right)_V \Delta T \Delta V - \\ &\quad \left(\frac{\partial p}{\partial T}\right)_V \Delta T \Delta V - \left(\frac{\partial p}{\partial V}\right)_T (\Delta V)^2 \\ &= \frac{C_V}{T} (\Delta T)^2 - \left(\frac{\partial p}{\partial V}\right)_T (\Delta V)^2 \end{aligned}$$

- $W \propto e^{-\frac{C_V}{2kT^2} (\Delta T)^2 + \frac{1}{2kT} \left(\frac{\partial p}{\partial V}\right)_T (\Delta V)^2},$

Fluctuation of canonical ensemble

$$W \propto e^{-\frac{\Delta T \Delta S - \Delta p \Delta V}{2kT}}$$

- If take (T, V) as variable,

$$\Delta S = \left(\frac{\partial S}{\partial T}\right)_V \Delta T + \left(\frac{\partial S}{\partial V}\right)_T \Delta V = \frac{C_V}{T} \Delta T + \left(\frac{\partial p}{\partial T}\right)_V \Delta V;$$

$$\Delta p = \left(\frac{\partial p}{\partial T}\right)_V \Delta T + \left(\frac{\partial p}{\partial V}\right)_T \Delta V.$$

$$\begin{aligned} \therefore \Delta T \Delta S - \Delta p \Delta V &= \frac{C_V}{T} (\Delta T)^2 + \left(\frac{\partial p}{\partial T}\right)_V \Delta T \Delta V - \\ &\quad \left(\frac{\partial p}{\partial T}\right)_V \Delta T \Delta V - \left(\frac{\partial p}{\partial V}\right)_T (\Delta V)^2 \\ &= \frac{C_V}{T} (\Delta T)^2 - \left(\frac{\partial p}{\partial V}\right)_T (\Delta V)^2 \end{aligned}$$

- $W \propto e^{-\frac{C_V}{2kT^2} (\Delta T)^2 + \frac{1}{2kT} \left(\frac{\partial p}{\partial V}\right)_T (\Delta V)^2}$, the probability for the system with deviation ΔT and ΔV .

Fluctuation of canonical ensemble

$$W \propto e^{-\frac{\Delta T \Delta S - \Delta p \Delta V}{2kT}}$$

- If take (T, V) as variable,

$$\Delta S = \left(\frac{\partial S}{\partial T}\right)_V \Delta T + \left(\frac{\partial S}{\partial V}\right)_T \Delta V = \frac{C_V}{T} \Delta T + \left(\frac{\partial p}{\partial T}\right)_V \Delta V;$$

$$\Delta p = \left(\frac{\partial p}{\partial T}\right)_V \Delta T + \left(\frac{\partial p}{\partial V}\right)_T \Delta V.$$

$$\begin{aligned} \therefore \Delta T \Delta S - \Delta p \Delta V &= \frac{C_V}{T} (\Delta T)^2 + \left(\frac{\partial p}{\partial T}\right)_V \Delta T \Delta V - \\ &\quad \left(\frac{\partial p}{\partial T}\right)_V \Delta T \Delta V - \left(\frac{\partial p}{\partial V}\right)_T (\Delta V)^2 \\ &= \frac{C_V}{T} (\Delta T)^2 - \left(\frac{\partial p}{\partial V}\right)_T (\Delta V)^2 \end{aligned}$$

- $W \propto e^{-\frac{C_V}{2kT^2} (\Delta T)^2 + \frac{1}{2kT} \left(\frac{\partial p}{\partial V}\right)_T (\Delta V)^2}$, the probability for the system with deviation ΔT and ΔV .
- Two independent Gaussian distributions indicates ΔT and ΔV are independent,

Fluctuation of canonical ensemble

$$W \propto e^{-\frac{\Delta T \Delta S - \Delta p \Delta V}{2kT}}$$

- If take (T, V) as variable,

$$\Delta S = \left(\frac{\partial S}{\partial T}\right)_V \Delta T + \left(\frac{\partial S}{\partial V}\right)_T \Delta V = \frac{C_V}{T} \Delta T + \left(\frac{\partial p}{\partial T}\right)_V \Delta V;$$

$$\Delta p = \left(\frac{\partial p}{\partial T}\right)_V \Delta T + \left(\frac{\partial p}{\partial V}\right)_T \Delta V.$$

$$\begin{aligned} \therefore \Delta T \Delta S - \Delta p \Delta V &= \frac{C_V}{T} (\Delta T)^2 + \left(\frac{\partial p}{\partial T}\right)_V \Delta T \Delta V - \\ &\quad \left(\frac{\partial p}{\partial T}\right)_V \Delta T \Delta V - \left(\frac{\partial p}{\partial V}\right)_T (\Delta V)^2 \\ &= \frac{C_V}{T} (\Delta T)^2 - \left(\frac{\partial p}{\partial V}\right)_T (\Delta V)^2 \end{aligned}$$

- $W \propto e^{-\frac{C_V}{2kT^2} (\Delta T)^2 + \frac{1}{2kT} \left(\frac{\partial p}{\partial V}\right)_T (\Delta V)^2}$, the probability for the system with deviation ΔT and ΔV .
- Two independent Gaussian distributions indicates ΔT and ΔV are independent, and $\overline{(\Delta T)^2} = \frac{kT^2}{C_V}$,

Fluctuation of canonical ensemble

$$W \propto e^{-\frac{\Delta T \Delta S - \Delta p \Delta V}{2kT}}$$

- If take (T, V) as variable,

$$\Delta S = \left(\frac{\partial S}{\partial T}\right)_V \Delta T + \left(\frac{\partial S}{\partial V}\right)_T \Delta V = \frac{C_V}{T} \Delta T + \left(\frac{\partial p}{\partial T}\right)_V \Delta V;$$

$$\Delta p = \left(\frac{\partial p}{\partial T}\right)_V \Delta T + \left(\frac{\partial p}{\partial V}\right)_T \Delta V.$$

$$\begin{aligned} \therefore \Delta T \Delta S - \Delta p \Delta V &= \frac{C_V}{T} (\Delta T)^2 + \left(\frac{\partial p}{\partial T}\right)_V \Delta T \Delta V - \\ &\quad \left(\frac{\partial p}{\partial T}\right)_V \Delta T \Delta V - \left(\frac{\partial p}{\partial V}\right)_T (\Delta V)^2 \\ &= \frac{C_V}{T} (\Delta T)^2 - \left(\frac{\partial p}{\partial V}\right)_T (\Delta V)^2 \end{aligned}$$

- $W \propto e^{-\frac{C_V}{2kT^2} (\Delta T)^2 + \frac{1}{2kT} \left(\frac{\partial p}{\partial V}\right)_T (\Delta V)^2}$, the probability for the system with deviation ΔT and ΔV .
- Two independent Gaussian distributions indicates ΔT and ΔV are independent,
and $\overline{(\Delta T)^2} = \frac{kT^2}{C_V}$, $\overline{(\Delta V)^2} = -kT \left(\frac{\partial V}{\partial p}\right)_T = kTV \kappa_T$.

Fluctuation of grand canonical ensemble

$$W \propto e^{-\frac{\Delta T \Delta S - \Delta p \Delta V}{2kT}}, \quad W \propto e^{-\frac{C_V}{2kT^2}(\Delta T)^2 + \frac{1}{2kT}\left(\frac{\partial p}{\partial V}\right)_T(\Delta V)^2}$$

- For magnetic medium, the work part (1.4.8):

$$\delta W = \mu_0 V H \delta M = \mu_0 H \delta m.$$

Fluctuation of grand canonical ensemble

$$W \propto e^{-\frac{\Delta T \Delta S - \Delta p \Delta V}{2kT}}, \quad W \propto e^{-\frac{C_V}{2kT^2}(\Delta T)^2 + \frac{1}{2kT}\left(\frac{\partial p}{\partial V}\right)_T(\Delta V)^2}$$

- For magnetic medium, the work part (1.4.8):

$$\delta W = \mu_0 V H \delta M = \mu_0 H \delta m.$$

$-p \rightarrow \mu_0 H$, $V \rightarrow m$, then

Fluctuation of grand canonical ensemble

$$W \propto e^{-\frac{\Delta T \Delta S - \Delta p \Delta V}{2kT}}, \quad W \propto e^{-\frac{C_V}{2kT^2}(\Delta T)^2 + \frac{1}{2kT}\left(\frac{\partial p}{\partial V}\right)_T(\Delta V)^2}$$

- For magnetic medium, the work part (1.4.8):

$$\delta W = \mu_0 V H \delta M = \mu_0 H \delta m.$$

$-p \rightarrow \mu_0 H$, $V \rightarrow m$, then

$$W \propto e^{-\frac{C_m}{2kT^2}(\Delta T)^2 - \frac{\mu_0}{2kT}\left(\frac{\partial H}{\partial m}\right)_T(\Delta m)^2}$$

Fluctuation of grand canonical ensemble

$$W \propto e^{-\frac{\Delta T \Delta S - \Delta p \Delta V}{2kT}}, \quad W \propto e^{-\frac{C_V}{2kT^2}(\Delta T)^2 + \frac{1}{2kT}(\frac{\partial p}{\partial V})_T(\Delta V)^2}$$

- For magnetic medium, the work part (1.4.8):

$$\delta W = \mu_0 V H \delta M = \mu_0 H \delta m.$$

$-p \rightarrow \mu_0 H$, $V \rightarrow m$, then

$$W \propto e^{-\frac{C_m}{2kT^2}(\Delta T)^2 - \frac{\mu_0}{2kT}(\frac{\partial H}{\partial m})_T(\Delta m)^2}$$

- For grand canonical ensemble:

Fluctuation of grand canonical ensemble

$$W \propto e^{-\frac{\Delta T \Delta S - \Delta p \Delta V}{2kT}}, \quad W \propto e^{-\frac{C_V}{2kT^2}(\Delta T)^2 + \frac{1}{2kT}\left(\frac{\partial p}{\partial V}\right)_T(\Delta V)^2}$$

- For magnetic medium, the work part (1.4.8):

$$\delta W = \mu_0 V H \delta M = \mu_0 H \delta m.$$

$-p \rightarrow \mu_0 H$, $V \rightarrow m$, then

$$W \propto e^{-\frac{C_m}{2kT^2}(\Delta T)^2 - \frac{\mu_0}{2kT}\left(\frac{\partial H}{\partial m}\right)_T(\Delta m)^2}$$

- For grand canonical ensemble:

$$W \propto e^{-\frac{\Delta T \Delta S - \Delta p \Delta V + \Delta \mu \Delta N}{2kT}}.$$

Fluctuation of grand canonical ensemble

$$W \propto e^{-\frac{\Delta T \Delta S - \Delta p \Delta V}{2kT}}, \quad W \propto e^{-\frac{C_V}{2kT^2}(\Delta T)^2 + \frac{1}{2kT}\left(\frac{\partial p}{\partial V}\right)_T(\Delta V)^2}$$

- For magnetic medium, the work part (1.4.8):

$$\delta W = \mu_0 V H \delta M = \mu_0 H \delta m.$$

$-p \rightarrow \mu_0 H$, $V \rightarrow m$, then

$$W \propto e^{-\frac{C_m}{2kT^2}(\Delta T)^2 - \frac{\mu_0}{2kT}\left(\frac{\partial H}{\partial m}\right)_T(\Delta m)^2}$$

- For grand canonical ensemble:

$$W \propto e^{-\frac{\Delta T \Delta S - \Delta p \Delta V + \Delta \mu \Delta N}{2kT}}.$$

- If take (N, V, T) as variables, but keep V fixed,

Fluctuation of grand canonical ensemble

$$W \propto e^{-\frac{\Delta T \Delta S - \Delta p \Delta V}{2kT}}, \quad W \propto e^{-\frac{C_V}{2kT^2}(\Delta T)^2 + \frac{1}{2kT}\left(\frac{\partial p}{\partial V}\right)_T(\Delta V)^2}$$

- For magnetic medium, the work part (1.4.8):

$$\delta W = \mu_0 V H \delta M = \mu_0 H \delta m.$$

$-p \rightarrow \mu_0 H$, $V \rightarrow m$, then

$$W \propto e^{-\frac{C_m}{2kT^2}(\Delta T)^2 - \frac{\mu_0}{2kT}\left(\frac{\partial H}{\partial m}\right)_T(\Delta m)^2}$$

- For grand canonical ensemble:

$$W \propto e^{-\frac{\Delta T \Delta S - \Delta p \Delta V + \Delta \mu \Delta N}{2kT}}.$$

- If take (N, V, T) as variables, but keep V fixed,

$$\Delta S = \left(\frac{\partial S}{\partial T}\right)_{V,N} \Delta T + \left(\frac{\partial S}{\partial N}\right)_{V,T} \Delta N$$

Fluctuation of grand canonical ensemble

$$W \propto e^{-\frac{\Delta T \Delta S - \Delta p \Delta V}{2kT}}, \quad W \propto e^{-\frac{C_V}{2kT^2}(\Delta T)^2 + \frac{1}{2kT}\left(\frac{\partial p}{\partial V}\right)_T(\Delta V)^2}$$

- For magnetic medium, the work part (1.4.8):

$$\delta W = \mu_0 V H \delta M = \mu_0 H \delta m.$$

$-p \rightarrow \mu_0 H$, $V \rightarrow m$, then

$$W \propto e^{-\frac{C_m}{2kT^2}(\Delta T)^2 - \frac{\mu_0}{2kT}\left(\frac{\partial H}{\partial m}\right)_T(\Delta m)^2}$$

- For grand canonical ensemble:

$$W \propto e^{-\frac{\Delta T \Delta S - \Delta p \Delta V + \Delta \mu \Delta N}{2kT}}.$$

- If take (N, V, T) as variables, but keep V fixed,

$$\begin{aligned} \Delta S &= \left(\frac{\partial S}{\partial T}\right)_{V,N} \Delta T + \left(\frac{\partial S}{\partial N}\right)_{V,T} \Delta N \\ &= \frac{C_V}{T} \Delta T - \left(\frac{\partial \mu}{\partial T}\right)_{V,T} \Delta N; \end{aligned}$$

Fluctuation of grand canonical ensemble

$$W \propto e^{-\frac{\Delta T \Delta S - \Delta p \Delta V}{2kT}}, \quad W \propto e^{-\frac{C_V}{2kT^2}(\Delta T)^2 + \frac{1}{2kT}\left(\frac{\partial p}{\partial V}\right)_T(\Delta V)^2}$$

- For magnetic medium, the work part (1.4.8):

$$\delta W = \mu_0 V H \delta M = \mu_0 H \delta m.$$

$-p \rightarrow \mu_0 H$, $V \rightarrow m$, then

$$W \propto e^{-\frac{C_m}{2kT^2}(\Delta T)^2 - \frac{\mu_0}{2kT}\left(\frac{\partial H}{\partial m}\right)_T(\Delta m)^2}$$

- For grand canonical ensemble:

$$W \propto e^{-\frac{\Delta T \Delta S - \Delta p \Delta V + \Delta \mu \Delta N}{2kT}}.$$

- If take (N, V, T) as variables, but keep V fixed,

$$\Delta S = \left(\frac{\partial S}{\partial T}\right)_{V,N} \Delta T + \left(\frac{\partial S}{\partial N}\right)_{V,T} \Delta N$$

$$= \frac{C_V}{T} \Delta T - \left(\frac{\partial \mu}{\partial T}\right)_{V,T} \Delta N;$$

$$\Delta \mu = \left(\frac{\partial \mu}{\partial T}\right)_{V,N} \Delta T + \left(\frac{\partial \mu}{\partial N}\right)_{V,T} \Delta N.$$

Fluctuation of grand canonical ensemble

$$W \propto e^{-\frac{\Delta T \Delta S - \Delta p \Delta V}{2kT}}, \quad W \propto e^{-\frac{C_V}{2kT^2}(\Delta T)^2 + \frac{1}{2kT}\left(\frac{\partial p}{\partial V}\right)_T(\Delta V)^2}$$

- For magnetic medium, the work part (1.4.8):

$$\delta W = \mu_0 V H \delta M = \mu_0 H \delta m.$$

$-p \rightarrow \mu_0 H$, $V \rightarrow m$, then

$$W \propto e^{-\frac{C_m}{2kT^2}(\Delta T)^2 - \frac{\mu_0}{2kT}\left(\frac{\partial H}{\partial m}\right)_T(\Delta m)^2}$$

- For grand canonical ensemble:

$$W \propto e^{-\frac{\Delta T \Delta S - \Delta p \Delta V + \Delta \mu \Delta N}{2kT}}.$$

- If take (N, V, T) as variables, but keep V fixed,

$$\Delta S = \left(\frac{\partial S}{\partial T}\right)_{V,N} \Delta T + \left(\frac{\partial S}{\partial N}\right)_{V,T} \Delta N$$

$$= \frac{C_V}{T} \Delta T - \left(\frac{\partial \mu}{\partial T}\right)_{V,T} \Delta N;$$

$$\Delta \mu = \left(\frac{\partial \mu}{\partial T}\right)_{V,N} \Delta T + \left(\frac{\partial \mu}{\partial N}\right)_{V,T} \Delta N.$$

Fluctuation of grand canonical ensemble

- $W \propto e^{-\frac{C_V}{2kT^2}(\Delta T)^2 - \frac{1}{2kT}\left(\frac{\partial \mu}{\partial N}\right)_{V,T}(\Delta N)^2}$.

Fluctuation of grand canonical ensemble

- $W \propto e^{-\frac{C_V}{2kT^2}(\Delta T)^2 - \frac{1}{2kT}(\frac{\partial \mu}{\partial N})_{V,T}(\Delta N)^2}$.
- $\overline{(\Delta T)^2} = \frac{kT^2}{C_V}$, $\overline{\Delta N \Delta T} = 0$, $\overline{(\Delta N)^2} = kT(\frac{\partial N}{\partial \mu})_{V,T}$.
- For the energy: $\Delta E = (\frac{\partial E}{\partial T})_{V,N} \Delta T + (\frac{\partial E}{\partial N})_{V,T} \Delta N$.

Fluctuation of grand canonical ensemble

- $W \propto e^{-\frac{C_V}{2kT^2}(\Delta T)^2 - \frac{1}{2kT}(\frac{\partial \mu}{\partial N})_{V,T}(\Delta N)^2}$.
- $\overline{(\Delta T)^2} = \frac{kT^2}{C_V}$, $\overline{\Delta N \Delta T} = 0$, $\overline{(\Delta N)^2} = kT(\frac{\partial N}{\partial \mu})_{V,T}$.
- For the energy: $\Delta E = (\frac{\partial E}{\partial T})_{V,N} \Delta T + (\frac{\partial E}{\partial N})_{V,T} \Delta N$.
- $\overline{(\Delta E)^2} = (\frac{\partial E}{\partial T})_{V,N}^2 \overline{(\Delta T)^2} + 2(\frac{\partial E}{\partial T})_{V,N}(\frac{\partial E}{\partial N})_{V,T} \overline{\Delta N \Delta T} + (\frac{\partial E}{\partial N})_{V,T}^2 \overline{(\Delta N)^2}$

Fluctuation of grand canonical ensemble

- $W \propto e^{-\frac{C_V}{2kT^2}(\Delta T)^2 - \frac{1}{2kT}(\frac{\partial \mu}{\partial N})_{V,T}(\Delta N)^2}$.
- $\overline{(\Delta T)^2} = \frac{kT^2}{C_V}$, $\overline{\Delta N \Delta T} = 0$, $\overline{(\Delta N)^2} = kT(\frac{\partial N}{\partial \mu})_{V,T}$.
- For the energy: $\Delta E = (\frac{\partial E}{\partial T})_{V,N} \Delta T + (\frac{\partial E}{\partial N})_{V,T} \Delta N$.
- $$\begin{aligned} \overline{(\Delta E)^2} &= (\frac{\partial E}{\partial T})_{V,N}^2 \overline{(\Delta T)^2} + 2(\frac{\partial E}{\partial T})_{V,N}(\frac{\partial E}{\partial N})_{V,T} \overline{\Delta N \Delta T} + \\ &\quad (\frac{\partial E}{\partial N})_{V,T}^2 \overline{(\Delta N)^2} \\ &= kT^2 C_V + (\frac{\partial E}{\partial N})_{V,T}^2 \overline{(\Delta N)^2}. \end{aligned}$$

Table of contents

1 §10. Fluctuation

- 10.1 Quasi-thermodynamics of fluctuation
- 10.5 Brownian motion
- 10.6 Diffusion and temporal correlation of Brownian particle's momentum
- 10.7 Examples of Brownian motion

§10.5 Brownian motion

§10.5 Brownian motion

- Microscopic essential: the Brownian particle is collided randomly by the molecules.

§10.5 Brownian motion

- Microscopic essential: the Brownian particle is collided randomly by the molecules.
- For the particle, it obeys Newton's 2nd law (only x):
$$m \frac{d^2 x}{dt^2} = f(t) + \mathcal{F}(t),$$
 where $f(t)$ is a random force, and $\mathcal{F}(t)$ is the stable external force.

§10.5 Brownian motion

- Microscopic essential: the Brownian particle is collided randomly by the molecules.
- For the particle, it obeys Newton's 2nd law (only x):
$$m \frac{d^2 x}{dt^2} = f(t) + \mathcal{F}(t),$$
 where $f(t)$ is a random force, and $\mathcal{F}(t)$ is the stable external force.
- $f(t)$ consists by two parts: damping the movement, called viscous force $-\alpha v$;

§10.5 Brownian motion

- Microscopic essential: the Brownian particle is collided randomly by the molecules.
- For the particle, it obeys Newton's 2nd law (only x):
$$m \frac{d^2 x}{dt^2} = f(t) + \mathcal{F}(t),$$
 where $f(t)$ is a random force, and $\mathcal{F}(t)$ is the stable external force.
- $f(t)$ consists by two parts: damping the movement, called viscous force $-\alpha v$; and randomly fluctuated force $F(t)$, with $\overline{F}(t) = 0$.

§10.5 Brownian motion

- Microscopic essential: the Brownian particle is collided randomly by the molecules.
- For the particle, it obeys Newton's 2nd law (only x): $m \frac{d^2x}{dt^2} = f(t) + \mathcal{F}(t)$, where $f(t)$ is a random force, and $\mathcal{F}(t)$ is the stable external force.
- $f(t)$ consists by two parts: damping the movement, called viscous force $-\alpha v$; and randomly fluctuated force $F(t)$, with $\overline{F}(t) = 0$.
- The equation of motion: $m \frac{d^2x}{dt^2} = -\alpha \frac{dx}{dt} + F(t) + \mathcal{F}(t)$, called Langevin's equation.

§10.5 Brownian motion

- Microscopic essential: the Brownian particle is collided randomly by the molecules.
- For the particle, it obeys Newton's 2nd law (only x): $m \frac{d^2x}{dt^2} = f(t) + \mathcal{F}(t)$, where $f(t)$ is a random force, and $\mathcal{F}(t)$ is the stable external force.
- $f(t)$ consists by two parts: damping the movement, called viscous force $-\alpha v$; and randomly fluctuated force $F(t)$, with $\overline{F}(t) = 0$.
- The equation of motion: $m \frac{d^2x}{dt^2} = -\alpha \frac{dx}{dt} + F(t) + \mathcal{F}(t)$, called Langevin's equation.
- When $\mathcal{F}(t) = 0$, $m \frac{d^2x}{dt^2} = -\alpha \frac{dx}{dt} + F(t)$.

Brownian motion

$$m \frac{d^2 x}{dt^2} = -\alpha \frac{dx}{dt} + F(t)$$

- As $x\ddot{x} = \frac{d(x\dot{x})}{dt} - \dot{x}^2$

Brownian motion

$$m \frac{d^2 x}{dt^2} = -\alpha \frac{dx}{dt} + F(t)$$

- As $x\ddot{x} = \frac{d(x\dot{x})}{dt} - \dot{x}^2 = \frac{1}{2} \frac{d^2(x^2)}{dt^2} - \dot{x}^2$.

Brownian motion

$$m \frac{d^2 x}{dt^2} = -\alpha \frac{dx}{dt} + F(t)$$

- As $x\ddot{x} = \frac{d(x\dot{x})}{dt} - \dot{x}^2 = \frac{1}{2} \frac{d^2(x^2)}{dt^2} - \dot{x}^2$.
- Introduce $\frac{1}{2} \frac{d^2(mx^2)}{dt^2} - m\dot{x}^2$

Brownian motion

$$m \frac{d^2 x}{dt^2} = -\alpha \frac{dx}{dt} + F(t)$$

- As $x\ddot{x} = \frac{d(x\dot{x})}{dt} - \dot{x}^2 = \frac{1}{2} \frac{d^2(x^2)}{dt^2} - \dot{x}^2$.
- Introduce $\frac{1}{2} \frac{d^2(mx^2)}{dt^2} - m\dot{x}^2 = mx\ddot{x}$

Brownian motion

$$m \frac{d^2 x}{dt^2} = -\alpha \frac{dx}{dt} + F(t)$$

- As $x\ddot{x} = \frac{d(x\dot{x})}{dt} - \dot{x}^2 = \frac{1}{2} \frac{d^2(x^2)}{dt^2} - \dot{x}^2$.
- Introduce $\frac{1}{2} \frac{d^2(mx^2)}{dt^2} - m\dot{x}^2 = mx\ddot{x} = x[-\alpha\dot{x} + F(t)]$

Brownian motion

$$m \frac{d^2 x}{dt^2} = -\alpha \frac{dx}{dt} + F(t)$$

- As $x\ddot{x} = \frac{d(x\dot{x})}{dt} - \dot{x}^2 = \frac{1}{2} \frac{d^2(x^2)}{dt^2} - \dot{x}^2$.
- Introduce $\frac{1}{2} \frac{d^2(mx^2)}{dt^2} - m\dot{x}^2 = mx\ddot{x} = x[-\alpha\dot{x} + F(t)]$
 $= -\frac{1}{2}\alpha \frac{dx^2}{dt} + xF(t)$.

Brownian motion

$$m \frac{d^2 x}{dt^2} = -\alpha \frac{dx}{dt} + F(t)$$

- As $x\ddot{x} = \frac{d(x\dot{x})}{dt} - \dot{x}^2 = \frac{1}{2} \frac{d^2(x^2)}{dt^2} - \dot{x}^2$.
- Introduce $\frac{1}{2} \frac{d^2(mx^2)}{dt^2} - m\dot{x}^2 = mx\ddot{x} = x[-\alpha\dot{x} + F(t)]$
 $= -\frac{1}{2}\alpha \frac{dx^2}{dt} + xF(t)$.
- Consider average for each item: $\overline{\frac{dx^2}{dt}} = \frac{d\overline{x^2}}{dt}$;
 $\overline{xF(t)} = \overline{x}\overline{F(t)} = 0$. And $\frac{1}{2}\overline{m\dot{x}^2} = \frac{1}{2}kT$.

Brownian motion

$$m \frac{d^2 x}{dt^2} = -\alpha \frac{dx}{dt} + F(t)$$

- As $x\ddot{x} = \frac{d(x\dot{x})}{dt} - \dot{x}^2 = \frac{1}{2} \frac{d^2(x^2)}{dt^2} - \dot{x}^2$.
- Introduce $\frac{1}{2} \frac{d^2(mx^2)}{dt^2} - m\dot{x}^2 = mx\ddot{x} = x[-\alpha\dot{x} + F(t)]$
 $= -\frac{1}{2}\alpha \frac{dx^2}{dt} + xF(t)$.
- Consider average for each item: $\overline{\frac{dx^2}{dt}} = \frac{d\overline{x^2}}{dt}$;
 $\overline{xF(t)} = \overline{x}\overline{F(t)} = 0$. And $\frac{1}{2}\overline{m\dot{x}^2} = \frac{1}{2}kT$.

Brownian motion

$$m \frac{d^2 x}{dt^2} = -\alpha \frac{dx}{dt} + F(t)$$

- As $x\ddot{x} = \frac{d(x\dot{x})}{dt} - \dot{x}^2 = \frac{1}{2} \frac{d^2(x^2)}{dt^2} - \dot{x}^2$.
- Introduce $\frac{1}{2} \frac{d^2(mx^2)}{dt^2} - m\dot{x}^2 = mx\ddot{x} = x[-\alpha\dot{x} + F(t)]$
 $= -\frac{1}{2}\alpha \frac{dx^2}{dt} + xF(t)$.
- Consider average for each item: $\overline{\frac{dx^2}{dt}} = \frac{d\overline{x^2}}{dt}$;
 $\overline{x F(t)} = \overline{x} \overline{F(t)} = 0$. And $\frac{1}{2} \overline{m\dot{x}^2} = \frac{1}{2} kT$.
 $\Rightarrow \frac{1}{2} m \frac{d^2}{dt^2} \overline{x^2} - m \overline{\dot{x}^2} = -\frac{\alpha}{2} \frac{d}{dt} \overline{x^2} + \overline{x F(t)}$,

Brownian motion

$$m \frac{d^2 x}{dt^2} = -\alpha \frac{dx}{dt} + F(t)$$

- As $x\ddot{x} = \frac{d(x\dot{x})}{dt} - \dot{x}^2 = \frac{1}{2} \frac{d^2(x^2)}{dt^2} - \dot{x}^2$.
- Introduce $\frac{1}{2} \frac{d^2(mx^2)}{dt^2} - m\dot{x}^2 = mx\ddot{x} = x[-\alpha\dot{x} + F(t)]$
 $= -\frac{1}{2}\alpha \frac{dx^2}{dt} + xF(t)$.
- Consider average for each item: $\overline{\frac{dx^2}{dt}} = \frac{d\overline{x^2}}{dt}$;
 $\overline{xF(t)} = \overline{x}\overline{F(t)} = 0$. And $\frac{1}{2}\overline{m\dot{x}^2} = \frac{1}{2}kT$.
 $\Rightarrow \frac{1}{2}m \frac{d^2}{dt^2} \overline{x^2} - m\overline{\dot{x}^2} = -\frac{\alpha}{2} \frac{d}{dt} \overline{x^2} + \overline{xF(t)}$,
 $\Rightarrow \frac{1}{2}m \frac{d^2}{dt^2} \overline{x^2} - kT = -\frac{\alpha}{2} \frac{d}{dt} \overline{x^2}$,

Brownian motion

$$m \frac{d^2 x}{dt^2} = -\alpha \frac{dx}{dt} + F(t)$$

- As $x\ddot{x} = \frac{d(x\dot{x})}{dt} - \dot{x}^2 = \frac{1}{2} \frac{d^2(x^2)}{dt^2} - \dot{x}^2$.
- Introduce $\frac{1}{2} \frac{d^2(mx^2)}{dt^2} - m\dot{x}^2 = mx\ddot{x} = x[-\alpha\dot{x} + F(t)]$
 $= -\frac{1}{2}\alpha \frac{dx^2}{dt} + xF(t)$.
- Consider average for each item: $\overline{\frac{dx^2}{dt}} = \frac{d\overline{x^2}}{dt}$;
 $\overline{x F(t)} = \overline{x} \overline{F(t)} = 0$. And $\frac{1}{2} \overline{m\dot{x}^2} = \frac{1}{2} kT$.
 $\Rightarrow \frac{1}{2} m \frac{d^2}{dt^2} \overline{x^2} - m \overline{\dot{x}^2} = -\frac{\alpha}{2} \frac{d}{dt} \overline{x^2} + \overline{x F(t)}$,
 $\Rightarrow \frac{1}{2} m \frac{d^2}{dt^2} \overline{x^2} - kT = -\frac{\alpha}{2} \frac{d}{dt} \overline{x^2}$,
 $\Rightarrow \frac{d^2}{dt^2} \overline{x^2} + \frac{\alpha}{m} \frac{d}{dt} \overline{x^2} - \frac{2}{m} kT = 0$.

Brownian motion

$$m \frac{d^2 x}{dt^2} = -\alpha \frac{dx}{dt} + F(t)$$

- As $x\ddot{x} = \frac{d(x\dot{x})}{dt} - \dot{x}^2 = \frac{1}{2} \frac{d^2(x^2)}{dt^2} - \dot{x}^2$.
- Introduce $\frac{1}{2} \frac{d^2(mx^2)}{dt^2} - m\dot{x}^2 = mx\ddot{x} = x[-\alpha\dot{x} + F(t)]$
 $= -\frac{1}{2}\alpha \frac{dx^2}{dt} + xF(t)$.
- Consider average for each item: $\overline{\frac{dx^2}{dt}} = \frac{d\overline{x^2}}{dt}$;
 $\overline{x F(t)} = \overline{x} \overline{F(t)} = 0$. And $\frac{1}{2} \overline{m\dot{x}^2} = \frac{1}{2} kT$.
 $\Rightarrow \frac{1}{2} m \frac{d^2}{dt^2} \overline{x^2} - m \overline{\dot{x}^2} = -\frac{\alpha}{2} \frac{d}{dt} \overline{x^2} + \overline{x F(t)}$,
 $\Rightarrow \frac{1}{2} m \frac{d^2}{dt^2} \overline{x^2} - kT = -\frac{\alpha}{2} \frac{d}{dt} \overline{x^2}$,
 $\Rightarrow \frac{d^2}{dt^2} \overline{x^2} + \frac{\alpha}{m} \frac{d}{dt} \overline{x^2} - \frac{2}{m} kT = 0$.
- The solution: $\overline{x^2} = \frac{2kT}{\alpha} t + C_1 e^{-\frac{\alpha}{m} t} + C_2$.

Brownian motion

$$m \frac{d^2 x}{dt^2} = -\alpha \frac{dx}{dt} + F(t)$$

- As $x\ddot{x} = \frac{d(x\dot{x})}{dt} - \dot{x}^2 = \frac{1}{2} \frac{d^2(x^2)}{dt^2} - \dot{x}^2$.
- Introduce $\frac{1}{2} \frac{d^2(mx^2)}{dt^2} - m\dot{x}^2 = mx\ddot{x} = x[-\alpha\dot{x} + F(t)]$
 $= -\frac{1}{2}\alpha \frac{dx^2}{dt} + xF(t)$.
- Consider average for each item: $\overline{\frac{dx^2}{dt}} = \frac{d\overline{x^2}}{dt}$;
 $\overline{xF(t)} = \overline{x}\overline{F(t)} = 0$. And $\frac{1}{2}\overline{m\dot{x}^2} = \frac{1}{2}kT$.
 $\Rightarrow \frac{1}{2}m \frac{d^2}{dt^2} \overline{x^2} - m\overline{\dot{x}^2} = -\frac{\alpha}{2} \frac{d}{dt} \overline{x^2} + \overline{xF(t)}$,
 $\Rightarrow \frac{1}{2}m \frac{d^2}{dt^2} \overline{x^2} - kT = -\frac{\alpha}{2} \frac{d}{dt} \overline{x^2}$,
 $\Rightarrow \frac{d^2}{dt^2} \overline{x^2} + \frac{\alpha}{m} \frac{d}{dt} \overline{x^2} - \frac{2}{m} kT = 0$.
- The solution: $\overline{x^2} = \frac{2kT}{\alpha} t + C_1 e^{-\frac{\alpha}{m}t} + C_2$.
- As $t = 0$, $x = 0$. $\Rightarrow C_2 = -C_1$.

Brownian motion

$$\overline{x^2} = \frac{2kT}{\alpha}t + C_1 e^{-\frac{\alpha}{m}t} - C_1$$

Brownian motion

$$\overline{x^2} = \frac{2kT}{\alpha}t + C_1 e^{-\frac{\alpha}{m}t} - C_1$$

- As $t = 0$, $\dot{x} = 0$. $\Rightarrow 0 = \dot{\overline{x^2}} = \frac{2kT}{\alpha} - C_1 \frac{\alpha}{m} e^{-\frac{\alpha}{m}0}$

Brownian motion

$$\overline{x^2} = \frac{2kT}{\alpha}t + C_1 e^{-\frac{\alpha}{m}t} - C_1$$

- As $t = 0$, $\dot{x} = 0$. $\Rightarrow 0 = \dot{\overline{x^2}} = \frac{2kT}{\alpha} - C_1 \frac{\alpha}{m} e^{-\frac{\alpha}{m}0}$
 $\Rightarrow C_1 = \frac{2mkT}{\alpha^2}$.

Brownian motion

$$\overline{x^2} = \frac{2kT}{\alpha}t + C_1 e^{-\frac{\alpha}{m}t} - C_1$$

- As $t = 0$, $\dot{x} = 0$. $\Rightarrow 0 = \dot{\overline{x^2}} = \frac{2kT}{\alpha} - C_1 \frac{\alpha}{m} e^{-\frac{\alpha}{m}0}$
 $\Rightarrow C_1 = \frac{2mkT}{\alpha^2}$.
 $\therefore \overline{x^2} = \frac{2kT}{\alpha}t + \frac{2mkT}{\alpha^2}e^{-\frac{\alpha}{m}t} - \frac{2mkT}{\alpha^2}$.

Brownian motion

$$\overline{x^2} = \frac{2kT}{\alpha}t + C_1 e^{-\frac{\alpha}{m}t} - C_1$$

- As $t = 0$, $\dot{x} = 0$. $\Rightarrow 0 = \dot{\overline{x^2}} = \frac{2kT}{\alpha} - C_1 \frac{\alpha}{m} e^{-\frac{\alpha}{m}0}$
 $\Rightarrow C_1 = \frac{2mkT}{\alpha^2}$.
 $\therefore \overline{x^2} = \frac{2kT}{\alpha}t + \frac{2mkT}{\alpha^2}e^{-\frac{\alpha}{m}t} - \frac{2mkT}{\alpha^2}$.
- Exists typical time $\tau = \frac{m}{\alpha}$. For $\tau \ll 1$, $C_1 \rightarrow 0$.

Brownian motion

$$\overline{x^2} = \frac{2kT}{\alpha}t + C_1 e^{-\frac{\alpha}{m}t} - C_1$$

- As $t = 0$, $\dot{x} = 0$. $\Rightarrow 0 = \dot{\overline{x^2}} = \frac{2kT}{\alpha} - C_1 \frac{\alpha}{m} e^{-\frac{\alpha}{m}0}$
 $\Rightarrow C_1 = \frac{2mkT}{\alpha^2}$.
 $\therefore \overline{x^2} = \frac{2kT}{\alpha}t + \frac{2mkT}{\alpha^2}e^{-\frac{\alpha}{m}t} - \frac{2mkT}{\alpha^2}$.
- Exists typical time $\tau = \frac{m}{\alpha}$. For $\tau \ll 1$, $C_1 \rightarrow 0$.
- 1. $t \ll \tau$,
 $\overline{x^2} \simeq \frac{2kT}{\alpha}t + \frac{2mkT}{\alpha^2}(1 - \frac{\alpha}{m}t + \frac{1}{2}\frac{\alpha^2}{m^2}t^2) - \frac{2mkT}{\alpha^2} \simeq \frac{kT}{m}t^2$,

Brownian motion

$$\overline{x^2} = \frac{2kT}{\alpha}t + C_1 e^{-\frac{\alpha}{m}t} - C_1$$

- As $t = 0$, $\dot{x} = 0$. $\Rightarrow 0 = \dot{\overline{x^2}} = \frac{2kT}{\alpha} - C_1 \frac{\alpha}{m} e^{-\frac{\alpha}{m}0}$
 $\Rightarrow C_1 = \frac{2mkT}{\alpha^2}$.
 $\therefore \overline{x^2} = \frac{2kT}{\alpha}t + \frac{2mkT}{\alpha^2}e^{-\frac{\alpha}{m}t} - \frac{2mkT}{\alpha^2}$.
- Exists typical time $\tau = \frac{m}{\alpha}$. For $\tau \ll 1$, $C_1 \rightarrow 0$.
- 1. $t \ll \tau$,
 $\overline{x^2} \simeq \frac{2kT}{\alpha}t + \frac{2mkT}{\alpha^2}(1 - \frac{\alpha}{m}t + \frac{1}{2}\frac{\alpha^2}{m^2}t^2) - \frac{2mkT}{\alpha^2} \simeq \frac{kT}{m}t^2$,
 like a uniform motion.

Brownian motion

$$\overline{x^2} = \frac{2kT}{\alpha}t + C_1 e^{-\frac{\alpha}{m}t} - C_1$$

- As $t = 0$, $\dot{x} = 0$. $\Rightarrow 0 = \dot{\overline{x^2}} = \frac{2kT}{\alpha} - C_1 \frac{\alpha}{m} e^{-\frac{\alpha}{m}0}$
 $\Rightarrow C_1 = \frac{2mkT}{\alpha^2}$.
 $\therefore \overline{x^2} = \frac{2kT}{\alpha}t + \frac{2mkT}{\alpha^2}e^{-\frac{\alpha}{m}t} - \frac{2mkT}{\alpha^2}$.
- Exists typical time $\tau = \frac{m}{\alpha}$. For $\tau \ll 1$, $C_1 \rightarrow 0$.
- 1. $t \ll \tau$,
 $\overline{x^2} \simeq \frac{2kT}{\alpha}t + \frac{2mkT}{\alpha^2}(1 - \frac{\alpha}{m}t + \frac{1}{2}\frac{\alpha^2}{m^2}t^2) - \frac{2mkT}{\alpha^2} \simeq \frac{kT}{m}t^2$,
 like a uniform motion.
- 2. $t \gg \tau$, $\overline{x^2} \simeq \frac{2kT}{\alpha}t \propto t$,

Brownian motion

$$\overline{x^2} = \frac{2kT}{\alpha}t + C_1 e^{-\frac{\alpha}{m}t} - C_1$$

- As $t = 0$, $\dot{x} = 0$. $\Rightarrow 0 = \dot{\overline{x^2}} = \frac{2kT}{\alpha} - C_1 \frac{\alpha}{m} e^{-\frac{\alpha}{m}0}$
 $\Rightarrow C_1 = \frac{2mkT}{\alpha^2}$.
 $\therefore \overline{x^2} = \frac{2kT}{\alpha}t + \frac{2mkT}{\alpha^2}e^{-\frac{\alpha}{m}t} - \frac{2mkT}{\alpha^2}$.
- Exists typical time $\tau = \frac{m}{\alpha}$. For $\tau \ll 1$, $C_1 \rightarrow 0$.
- 1. $t \ll \tau$,
 $\overline{x^2} \simeq \frac{2kT}{\alpha}t + \frac{2mkT}{\alpha^2}(1 - \frac{\alpha}{m}t + \frac{1}{2}\frac{\alpha^2}{m^2}t^2) - \frac{2mkT}{\alpha^2} \simeq \frac{kT}{m}t^2$,
 like a uniform motion.
- 2. $t \gg \tau$, $\overline{x^2} \simeq \frac{2kT}{\alpha}t \propto t$, then $|x| \propto t^{\frac{1}{2}}$.

Brownian motion, in the view of diffusion

$$\overline{x^2} \simeq \frac{2kT}{\alpha} t$$

Brownian motion, in the view of diffusion

$$\overline{x^2} \simeq \frac{2kT}{\alpha} t$$

- In the view of diffusion: $\vec{J} = -D \nabla n$, where $J(x, t)$ is the number of Brownian particle crossing unit area in unit time.

Brownian motion, in the view of diffusion

$$\overline{x^2} \simeq \frac{2kT}{\alpha} t$$

- In the view of diffusion: $\vec{J} = -D \nabla n$, where $J(x, t)$ is the number of Brownian particle crossing unit area in unit time.
- Equation of continuity: $\frac{\partial n}{\partial t} + \nabla \cdot \vec{J} = 0$,

Brownian motion, in the view of diffusion

$$\overline{x^2} \simeq \frac{2kT}{\alpha} t$$

- In the view of diffusion: $\vec{J} = -D \nabla n$, where $J(x, t)$ is the number of Brownian particle crossing unit area in unit time.
- Equation of continuity: $\frac{\partial n}{\partial t} + \nabla \cdot \vec{J} = 0$, then $\frac{\partial n}{\partial t} - D \nabla^2 n = 0$.

Brownian motion, in the view of diffusion

$$\overline{x^2} \simeq \frac{2kT}{\alpha} t$$

- In the view of diffusion: $\vec{J} = -D \nabla n$, where $J(x, t)$ is the number of Brownian particle crossing unit area in unit time.
- Equation of continuity: $\frac{\partial n}{\partial t} + \nabla \cdot \vec{J} = 0$, then $\frac{\partial n}{\partial t} - D \nabla^2 n = 0$.
- For 1-D, $\frac{\partial n}{\partial t} = D \frac{\partial^2 n}{\partial x^2}$.

Brownian motion, in the view of diffusion

$$\overline{x^2} \simeq \frac{2kT}{\alpha} t$$

- In the view of diffusion: $\vec{J} = -D \nabla n$, where $J(x, t)$ is the number of Brownian particle crossing unit area in unit time.
- Equation of continuity: $\frac{\partial n}{\partial t} + \nabla \cdot \vec{J} = 0$, then $\frac{\partial n}{\partial t} - D \nabla^2 n = 0$.
- For 1-D, $\frac{\partial n}{\partial t} = D \frac{\partial^2 n}{\partial x^2}$.
- Suppose $t = 0$, all particles at $x = 0$, i.e., $n(x, 0) = N\delta(x)$,

Brownian motion, in the view of diffusion

$$\overline{x^2} \simeq \frac{2kT}{\alpha} t$$

- In the view of diffusion: $\vec{J} = -D \nabla n$, where $J(x, t)$ is the number of Brownian particle crossing unit area in unit time.
- Equation of continuity: $\frac{\partial n}{\partial t} + \nabla \cdot \vec{J} = 0$, then $\frac{\partial n}{\partial t} - D \nabla^2 n = 0$.
- For 1-D, $\frac{\partial n}{\partial t} = D \frac{\partial^2 n}{\partial x^2}$.
- Suppose $t = 0$, all particles at $x = 0$, i.e., $n(x, 0) = N\delta(x)$, the solution: $n(x, t) = \frac{N}{2\sqrt{\pi Dt}} e^{-\frac{x^2}{4Dt}}$.

Brownian motion, in the view of diffusion

$$\overline{x^2} \simeq \frac{2kT}{\alpha} t$$

- In the view of diffusion: $\vec{J} = -D \nabla n$, where $J(x, t)$ is the number of Brownian particle crossing unit area in unit time.
- Equation of continuity: $\frac{\partial n}{\partial t} + \nabla \cdot \vec{J} = 0$, then $\frac{\partial n}{\partial t} - D \nabla^2 n = 0$.
- For 1-D, $\frac{\partial n}{\partial t} = D \frac{\partial^2 n}{\partial x^2}$.
- Suppose $t = 0$, all particles at $x = 0$, i.e., $n(x, 0) = N\delta(x)$, the solution: $n(x, t) = \frac{N}{2\sqrt{\pi Dt}} e^{-\frac{x^2}{4Dt}}$.
- $\overline{x^2(t)} = \frac{1}{N} \int_{-\infty}^{\infty} x^2 n(x, t) dx = 2Dt$,

Brownian motion, in the view of diffusion

$$\overline{x^2} \simeq \frac{2kT}{\alpha} t$$

- In the view of diffusion: $\vec{J} = -D \nabla n$, where $J(x, t)$ is the number of Brownian particle crossing unit area in unit time.
- Equation of continuity: $\frac{\partial n}{\partial t} + \nabla \cdot \vec{J} = 0$, then $\frac{\partial n}{\partial t} - D \nabla^2 n = 0$.
- For 1-D, $\frac{\partial n}{\partial t} = D \frac{\partial^2 n}{\partial x^2}$.
- Suppose $t = 0$, all particles at $x = 0$, i.e., $n(x, 0) = N\delta(x)$, the solution: $n(x, t) = \frac{N}{2\sqrt{\pi Dt}} e^{-\frac{x^2}{4Dt}}$.
- $\overline{x^2(t)} = \frac{1}{N} \int_{-\infty}^{\infty} x^2 n(x, t) dx = 2Dt$, should be the same as (10.5.8),

Brownian motion, in the view of diffusion

$$\overline{x^2} \simeq \frac{2kT}{\alpha} t$$

- In the view of diffusion: $\vec{J} = -D \nabla n$, where $J(x, t)$ is the number of Brownian particle crossing unit area in unit time.
- Equation of continuity: $\frac{\partial n}{\partial t} + \nabla \cdot \vec{J} = 0$, then $\frac{\partial n}{\partial t} - D \nabla^2 n = 0$.
- For 1-D, $\frac{\partial n}{\partial t} = D \frac{\partial^2 n}{\partial x^2}$.
- Suppose $t = 0$, all particles at $x = 0$, i.e., $n(x, 0) = N\delta(x)$, the solution: $n(x, t) = \frac{N}{2\sqrt{\pi Dt}} e^{-\frac{x^2}{4Dt}}$.
- $\overline{x^2(t)} = \frac{1}{N} \int_{-\infty}^{\infty} x^2 n(x, t) dx = 2Dt$, should be the same as (10.5.8), then, $D = \frac{kT}{\alpha}$.

Table of contents

1 §10. Fluctuation

- 10.1 Quasi-thermodynamics of fluctuation
- 10.5 Brownian motion
- 10.6 Diffusion and temporal correlation of Brownian particle's momentum
- 10.7 Examples of Brownian motion

§10.6 Diffusion and temporal correlation of Brownian particle's momentum

§10.6 Diffusion and temporal correlation of Brownian particle's momentum

- Assuming an initial momentum $p(0)$, consider the momentum evolution of Brownian particle.

§10.6 Diffusion and temporal correlation of Brownian particle's momentum

- Assuming an initial momentum $p(0)$, consider the momentum evolution of Brownian particle.
- Only average momentum is available $\bar{p}(t)$.

§10.6 Diffusion and temporal correlation of Brownian particle's momentum

- Assuming an initial momentum $p(0)$, consider the momentum evolution of Brownian particle.
- Only average momentum is available $\bar{p}(t)$.
- Langevin's equation (10.5.4): $m \frac{d^2x}{dt^2} = -\alpha \frac{dx}{dt} + F(t)$.

§10.6 Diffusion and temporal correlation of Brownian particle's momentum

- Assuming an initial momentum $p(0)$, consider the momentum evolution of Brownian particle.
- Only average momentum is available $\bar{p}(t)$.
- Langevin's equation (10.5.4): $m \frac{d^2x}{dt^2} = -\alpha \frac{dx}{dt} + F(t)$.
- Convert to momentum: $\frac{dp}{dt} = -\gamma p + F(t)$, with $\gamma = \frac{\alpha}{m}$.

§10.6 Diffusion and temporal correlation of Brownian particle's momentum

- Assuming an initial momentum $p(0)$, consider the momentum evolution of Brownian particle.
- Only average momentum is available $\bar{p}(t)$.
- Langevin's equation (10.5.4): $m \frac{d^2 x}{dt^2} = -\alpha \frac{dx}{dt} + F(t)$.
- Convert to momentum: $\frac{dp}{dt} = -\gamma p + F(t)$, with $\gamma = \frac{\alpha}{m}$.
- Before to get \bar{p} , turn to the temporal correlation function: $\overline{F(t)F(t+\tau)} \equiv \frac{1}{N} \sum_{i=1}^N F_i(t)F_i(t+\tau)$.

§10.6 Diffusion and temporal correlation of Brownian particle's momentum

- Assuming an initial momentum $p(0)$, consider the momentum evolution of Brownian particle.
- Only average momentum is available $\bar{p}(t)$.
- Langevin's equation (10.5.4): $m \frac{d^2x}{dt^2} = -\alpha \frac{dx}{dt} + F(t)$.
- Convert to momentum: $\frac{dp}{dt} = -\gamma p + F(t)$, with $\gamma = \frac{\alpha}{m}$.
- Before to get \bar{p} , turn to the temporal correlation function: $\overline{F(t)F(t+\tau)} \equiv \frac{1}{N} \sum_{i=1}^N F_i(t)F_i(t+\tau)$.
- If the duration (τ_c) of the fluctuating force is short enough, $\overline{F(t)F(t+\tau)} = 2D_p\delta(\tau)$;

§10.6 Diffusion and temporal correlation of Brownian particle's momentum

- Assuming an initial momentum $p(0)$, consider the momentum evolution of Brownian particle.
- Only average momentum is available $\bar{p}(t)$.
- Langevin's equation (10.5.4): $m \frac{d^2x}{dt^2} = -\alpha \frac{dx}{dt} + F(t)$.
- Convert to momentum: $\frac{dp}{dt} = -\gamma p + F(t)$, with $\gamma = \frac{\alpha}{m}$.
- Before to get \bar{p} , turn to the temporal correlation function: $\overline{F(t)F(t+\tau)} \equiv \frac{1}{N} \sum_{i=1}^N F_i(t)F_i(t+\tau)$.
- If the duration (τ_c) of the fluctuating force is short enough, $\overline{F(t)F(t+\tau)} = 2D_p\delta(\tau)$; and $2D_p = \overline{F^2(t)}$.

Diffusion and temporal correlation of Brownian particle's momentum

$$\frac{dp}{dt} = -\gamma p + F(t)$$

Diffusion and temporal correlation of Brownian particle's momentum

$$\frac{dp}{dt} = -\gamma p + F(t)$$

- Back to the Langevin's equation:

$$e^{\gamma t} \frac{dp}{dt} = -e^{\gamma t} \gamma p + e^{\gamma t} F(t),$$

Diffusion and temporal correlation of Brownian particle's momentum

$$\frac{dp}{dt} = -\gamma p + F(t)$$

- Back to the Langevin's equation:

$$e^{\gamma t} \frac{dp}{dt} = -e^{\gamma t} \gamma p + e^{\gamma t} F(t),$$

$$\Rightarrow \frac{d(pe^{\gamma t})}{dt} = e^{\gamma t} F(t),$$

Diffusion and temporal correlation of Brownian particle's momentum

$$\frac{dp}{dt} = -\gamma p + F(t)$$

- Back to the Langevin's equation:

$$e^{\gamma t} \frac{dp}{dt} = -e^{\gamma t} \gamma p + e^{\gamma t} F(t),$$

$$\Rightarrow \frac{d(pe^{\gamma t})}{dt} = e^{\gamma t} F(t),$$

$$\Rightarrow p(t) = p(0)e^{-\gamma t} + e^{-\gamma t} \int_0^t F(\xi) e^{\gamma \xi} d\xi.$$

Diffusion and temporal correlation of Brownian particle's momentum

$$\frac{dp}{dt} = -\gamma p + F(t)$$

- Back to the Langevin's equation:

$$e^{\gamma t} \frac{dp}{dt} = -e^{\gamma t} \gamma p + e^{\gamma t} F(t),$$

$$\Rightarrow \frac{d(pe^{\gamma t})}{dt} = e^{\gamma t} F(t),$$

$$\Rightarrow p(t) = p(0)e^{-\gamma t} + e^{-\gamma t} \int_0^t F(\xi) e^{\gamma \xi} d\xi.$$

- Take the average, noticing $\overline{F(\xi)} = 0$,

$$\Rightarrow \bar{p}(t) = p(0)e^{-\gamma t},$$

Diffusion and temporal correlation of Brownian particle's momentum

$$\frac{dp}{dt} = -\gamma p + F(t)$$

- Back to the Langevin's equation:

$$e^{\gamma t} \frac{dp}{dt} = -e^{\gamma t} \gamma p + e^{\gamma t} F(t),$$

$$\Rightarrow \frac{d(pe^{\gamma t})}{dt} = e^{\gamma t} F(t),$$

$$\Rightarrow p(t) = p(0)e^{-\gamma t} + e^{-\gamma t} \int_0^t F(\xi) e^{\gamma \xi} d\xi.$$

- Take the average, noticing $\overline{F(\xi)} = 0$,

$$\Rightarrow \bar{p}(t) = p(0)e^{-\gamma t},$$

where $\frac{1}{\gamma}$ is the time scale for a notable change of momentum (larger than τ_c).

Diffusion of Brownian particle's momentum

$$p(t) = p(0)e^{-\gamma t} + e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$$

- $\overline{(\Delta p)^2} = \overline{[p(t) - \bar{p}(t)]^2}$

Diffusion of Brownian particle's momentum

$$p(t) = p(0)e^{-\gamma t} + e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$$

- $$\begin{aligned} \overline{(\Delta p)^2} &= \overline{[p(t) - \bar{p}(t)]^2} \\ &= \overline{[e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi]^2} \end{aligned}$$

Diffusion of Brownian particle's momentum

$$p(t) = p(0)e^{-\gamma t} + e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$$

- $$\begin{aligned} \overline{(\Delta p)^2} &= \overline{[p(t) - \bar{p}(t)]^2} \\ &= \overline{[e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi]^2} = \overline{[\int_0^t F(\xi)e^{-\gamma(t-\xi)} d\xi]^2} \end{aligned}$$

Diffusion of Brownian particle's momentum

$$p(t) = p(0)e^{-\gamma t} + e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$$

- $$\begin{aligned} \bullet \quad \overline{(\Delta p)^2} &= \overline{[p(t) - \bar{p}(t)]^2} \\ &= \overline{[e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi]^2} = \overline{[\int_0^t F(\xi)e^{-\gamma(t-\xi)} d\xi]^2} \\ &= \int_0^t d\xi \int_0^t d\xi' F(\xi)F(\xi')e^{-\gamma(t-\xi)}e^{-\gamma(t-\xi')} \end{aligned}$$

Diffusion of Brownian particle's momentum

$$p(t) = p(0)e^{-\gamma t} + e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$$

- $$\begin{aligned}
 \bullet \quad \overline{(\Delta p)^2} &= \overline{[p(t) - \bar{p}(t)]^2} \\
 &= \overline{[e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi]^2} = \overline{[\int_0^t F(\xi)e^{-\gamma(t-\xi)} d\xi]^2} \\
 &= \int_0^t d\xi \int_0^t d\xi' F(\xi)F(\xi')e^{-\gamma(t-\xi)}e^{-\gamma(t-\xi')} \\
 &= \int_0^t d\xi \int_0^t d\xi' \overline{F(\xi)F(\xi')}e^{-\gamma(t-\xi)}e^{-\gamma(t-\xi')}
 \end{aligned}$$

Diffusion of Brownian particle's momentum

$$p(t) = p(0)e^{-\gamma t} + e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$$

$$\begin{aligned}
 \bullet \quad \overline{(\Delta p)^2} &= \overline{[p(t) - \bar{p}(t)]^2} \\
 &= \overline{[e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi]^2} = \overline{[\int_0^t F(\xi)e^{-\gamma(t-\xi)} d\xi]^2} \\
 &= \int_0^t d\xi \int_0^t d\xi' F(\xi)F(\xi')e^{-\gamma(t-\xi)}e^{-\gamma(t-\xi')} \\
 &= \int_0^t d\xi \int_0^t d\xi' \overline{F(\xi)F(\xi')}e^{-\gamma(t-\xi)}e^{-\gamma(t-\xi')} \\
 &= \int_0^t d\xi \int_0^t d\xi' 2D_p \delta(\xi - \xi')e^{-\gamma(t-\xi)}e^{-\gamma(t-\xi')}
 \end{aligned}$$

Diffusion of Brownian particle's momentum

$$p(t) = p(0)e^{-\gamma t} + e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$$

$$\begin{aligned}
 \bullet \quad \overline{(\Delta p)^2} &= \overline{[p(t) - \bar{p}(t)]^2} \\
 &= \overline{[e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi]^2} = \overline{[\int_0^t F(\xi)e^{-\gamma(t-\xi)} d\xi]^2} \\
 &= \int_0^t d\xi \int_0^t d\xi' F(\xi)F(\xi')e^{-\gamma(t-\xi)}e^{-\gamma(t-\xi')} \\
 &= \int_0^t d\xi \int_0^t d\xi' \overline{F(\xi)F(\xi')}e^{-\gamma(t-\xi)}e^{-\gamma(t-\xi')} \\
 &= \int_0^t d\xi \int_0^t d\xi' 2D_p \delta(\xi - \xi')e^{-\gamma(t-\xi)}e^{-\gamma(t-\xi')} \\
 &= \int_0^t d\xi 2D_p e^{-2\gamma(t-\xi)}
 \end{aligned}$$

Diffusion of Brownian particle's momentum

$$p(t) = p(0)e^{-\gamma t} + e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$$

$$\begin{aligned}
 \bullet \quad \overline{(\Delta p)^2} &= \overline{[p(t) - \bar{p}(t)]^2} \\
 &= \overline{[e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi]^2} = \overline{[\int_0^t F(\xi)e^{-\gamma(t-\xi)} d\xi]^2} \\
 &= \int_0^t d\xi \int_0^t d\xi' F(\xi)F(\xi')e^{-\gamma(t-\xi)}e^{-\gamma(t-\xi')} \\
 &= \int_0^t d\xi \int_0^t d\xi' \overline{F(\xi)F(\xi')}e^{-\gamma(t-\xi)}e^{-\gamma(t-\xi')} \\
 &= \int_0^t d\xi \int_0^t d\xi' 2D_p \delta(\xi - \xi')e^{-\gamma(t-\xi)}e^{-\gamma(t-\xi')} \\
 &= \int_0^t d\xi 2D_p e^{-2\gamma(t-\xi)} = \frac{D_p}{\gamma}(1 - e^{-2\gamma t}).
 \end{aligned}$$

Diffusion of Brownian particle's momentum

$$p(t) = p(0)e^{-\gamma t} + e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$$

- $$\begin{aligned} \bullet \quad \overline{(\Delta p)^2} &= \overline{[p(t) - \bar{p}(t)]^2} \\ &= \overline{[e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi]^2} = \overline{[\int_0^t F(\xi)e^{-\gamma(t-\xi)} d\xi]^2} \\ &= \int_0^t d\xi \int_0^t d\xi' F(\xi)F(\xi')e^{-\gamma(t-\xi)}e^{-\gamma(t-\xi')} \\ &= \int_0^t d\xi \int_0^t d\xi' \overline{F(\xi)F(\xi')}e^{-\gamma(t-\xi)}e^{-\gamma(t-\xi')} \\ &= \int_0^t d\xi \int_0^t d\xi' 2D_p \delta(\xi - \xi')e^{-\gamma(t-\xi)}e^{-\gamma(t-\xi')} \\ &= \int_0^t d\xi 2D_p e^{-2\gamma(t-\xi)} = \frac{D_p}{\gamma}(1 - e^{-2\gamma t}). \end{aligned}$$
- $$\bullet \quad 1. \quad \tau_c \ll t \ll \frac{1}{\gamma}, \quad \overline{(\Delta p)^2} = 2D_p t;$$

Diffusion of Brownian particle's momentum

$$p(t) = p(0)e^{-\gamma t} + e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$$

- $$\begin{aligned} \bullet \quad \overline{(\Delta p)^2} &= \overline{[p(t) - \bar{p}(t)]^2} \\ &= \overline{[e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi]^2} = \overline{[\int_0^t F(\xi)e^{-\gamma(t-\xi)} d\xi]^2} \\ &= \int_0^t d\xi \int_0^t d\xi' F(\xi)F(\xi')e^{-\gamma(t-\xi)}e^{-\gamma(t-\xi')} \\ &= \int_0^t d\xi \int_0^t d\xi' \overline{F(\xi)F(\xi')}e^{-\gamma(t-\xi)}e^{-\gamma(t-\xi')} \\ &= \int_0^t d\xi \int_0^t d\xi' 2D_p \delta(\xi - \xi')e^{-\gamma(t-\xi)}e^{-\gamma(t-\xi')} \\ &= \int_0^t d\xi 2D_p e^{-2\gamma(t-\xi)} = \frac{D_p}{\gamma}(1 - e^{-2\gamma t}). \end{aligned}$$
- $$\bullet \quad 1. \quad \tau_c \ll t \ll \frac{1}{\gamma}, \quad \overline{(\Delta p)^2} = 2D_p t;$$
- $$\bullet \quad 2. \quad t \gg \frac{1}{\gamma}, \quad \bar{p}(t) = 0, \text{ so } \overline{p^2} = \overline{(\Delta p)^2} = \frac{D_p}{\gamma}.$$

Diffusion of Brownian particle's momentum

$$p(t) = p(0)e^{-\gamma t} + e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$$

- $$\begin{aligned} \bullet \quad \overline{(\Delta p)^2} &= \overline{[p(t) - \bar{p}(t)]^2} \\ &= \overline{[e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi]^2} = \overline{[\int_0^t F(\xi)e^{-\gamma(t-\xi)} d\xi]^2} \\ &= \int_0^t d\xi \int_0^t d\xi' F(\xi)F(\xi')e^{-\gamma(t-\xi)}e^{-\gamma(t-\xi')} \\ &= \int_0^t d\xi \int_0^t d\xi' \overline{F(\xi)F(\xi')}e^{-\gamma(t-\xi)}e^{-\gamma(t-\xi')} \\ &= \int_0^t d\xi \int_0^t d\xi' 2D_p \delta(\xi - \xi')e^{-\gamma(t-\xi)}e^{-\gamma(t-\xi')} \\ &= \int_0^t d\xi 2D_p e^{-2\gamma(t-\xi)} = \frac{D_p}{\gamma}(1 - e^{-2\gamma t}). \end{aligned}$$
- 1. $\tau_c \ll t \ll \frac{1}{\gamma}$, $\overline{(\Delta p)^2} = 2D_p t$;
- 2. $t \gg \frac{1}{\gamma}$, $\bar{p}(t) = 0$, so $\overline{p^2} = \overline{(\Delta p)^2} = \frac{D_p}{\gamma}$.

In equilibrium, $\frac{\overline{p^2}}{2m} = \frac{1}{2}kT$, $\therefore D_p = m\gamma kT = \alpha kT$.

Temporal correlation of the momentum

$$p(t) = p(0)e^{-\gamma t} + e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$$

Temporal correlation of the momentum

$$p(t) = p(0)e^{-\gamma t} + e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$$

- For $t > \frac{1}{\gamma}$, $e^{-\gamma t} \rightarrow 0$, then $p(t) = e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$.

Temporal correlation of the momentum

$$p(t) = p(0)e^{-\gamma t} + e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$$

- For $t > \frac{1}{\gamma}$, $e^{-\gamma t} \rightarrow 0$, then $p(t) = e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$.
- The temporal correlation:

$$\overline{p(t)p(t')}$$

Temporal correlation of the momentum

$$p(t) = p(0)e^{-\gamma t} + e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$$

- For $t > \frac{1}{\gamma}$, $e^{-\gamma t} \rightarrow 0$, then $p(t) = e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$.
- The temporal correlation:

$$\overline{p(t)p(t')} = \overline{e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi \cdot e^{-\gamma t'} \int_0^{t'} F(\xi')e^{\gamma \xi'} d\xi'}$$

Temporal correlation of the momentum

$$p(t) = p(0)e^{-\gamma t} + e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$$

- For $t > \frac{1}{\gamma}$, $e^{-\gamma t} \rightarrow 0$, then $p(t) = e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$.
- The temporal correlation:

$$\begin{aligned} \overline{p(t)p(t')} &= \overline{e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi \cdot e^{-\gamma t'} \int_0^{t'} F(\xi')e^{\gamma \xi'} d\xi'} \\ &= e^{-\gamma(t+t')} \int_0^t d\xi \int_0^{t'} d\xi' \overline{F(\xi)F(\xi')} e^{\gamma(\xi+\xi')} \end{aligned}$$

Temporal correlation of the momentum

$$p(t) = p(0)e^{-\gamma t} + e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$$

- For $t > \frac{1}{\gamma}$, $e^{-\gamma t} \rightarrow 0$, then $p(t) = e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$.
- The temporal correlation:

$$\begin{aligned} \overline{p(t)p(t')} &= \overline{e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi \cdot e^{-\gamma t'} \int_0^{t'} F(\xi')e^{\gamma \xi'} d\xi'} \\ &= e^{-\gamma(t+t')} \int_0^t d\xi \int_0^{t'} d\xi' \overline{F(\xi)F(\xi')} e^{\gamma(\xi+\xi')} \\ &= e^{-\gamma(t+t')} \int_0^t d\xi \int_0^{t'} d\xi' 2D_p \delta(\xi - \xi') e^{\gamma(\xi+\xi')} \end{aligned}$$

Temporal correlation of the momentum

$$p(t) = p(0)e^{-\gamma t} + e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$$

- For $t > \frac{1}{\gamma}$, $e^{-\gamma t} \rightarrow 0$, then $p(t) = e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$.
- The temporal correlation:

$$\begin{aligned} \overline{p(t)p(t')} &= \overline{e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi \cdot e^{-\gamma t'} \int_0^{t'} F(\xi')e^{\gamma \xi'} d\xi'} \\ &= e^{-\gamma(t+t')} \int_0^t d\xi \int_0^{t'} d\xi' \overline{F(\xi)F(\xi')} e^{\gamma(\xi+\xi')} \\ &= e^{-\gamma(t+t')} \int_0^t d\xi \int_0^{t'} d\xi' 2D_p \delta(\xi - \xi') e^{\gamma(\xi+\xi')} \end{aligned}$$

- 1. If $t < t'$, $\overline{p(t)p(t')} = e^{-\gamma(t+t')} \int_0^t d\xi 2D_p e^{2\gamma \xi}$

Temporal correlation of the momentum

$$p(t) = p(0)e^{-\gamma t} + e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$$

- For $t > \frac{1}{\gamma}$, $e^{-\gamma t} \rightarrow 0$, then $p(t) = e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$.
- The temporal correlation:

$$\begin{aligned} \overline{p(t)p(t')} &= \overline{e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi \cdot e^{-\gamma t'} \int_0^{t'} F(\xi')e^{\gamma \xi'} d\xi'} \\ &= e^{-\gamma(t+t')} \int_0^t d\xi \int_0^{t'} d\xi' \overline{F(\xi)F(\xi')} e^{\gamma(\xi+\xi')} \\ &= e^{-\gamma(t+t')} \int_0^t d\xi \int_0^{t'} d\xi' 2D_p \delta(\xi - \xi') e^{\gamma(\xi+\xi')} \end{aligned}$$

- 1. If $t < t'$, $\overline{p(t)p(t')} = e^{-\gamma(t+t')} \int_0^t d\xi 2D_p e^{2\gamma \xi}$
 $= 2D_p e^{-\gamma(t+t')} \frac{1}{2\gamma} (e^{2\gamma t} - 1)$

Temporal correlation of the momentum

$$p(t) = p(0)e^{-\gamma t} + e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$$

- For $t > \frac{1}{\gamma}$, $e^{-\gamma t} \rightarrow 0$, then $p(t) = e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$.
- The temporal correlation:

$$\begin{aligned} \overline{p(t)p(t')} &= \overline{e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi \cdot e^{-\gamma t'} \int_0^{t'} F(\xi')e^{\gamma \xi'} d\xi'} \\ &= e^{-\gamma(t+t')} \int_0^t d\xi \int_0^{t'} d\xi' \overline{F(\xi)F(\xi')} e^{\gamma(\xi+\xi')} \\ &= e^{-\gamma(t+t')} \int_0^t d\xi \int_0^{t'} d\xi' 2D_p \delta(\xi - \xi') e^{\gamma(\xi+\xi')} \end{aligned}$$

- 1. If $t < t'$, $\overline{p(t)p(t')} = e^{-\gamma(t+t')} \int_0^t d\xi 2D_p e^{2\gamma \xi}$
 $= 2D_p e^{-\gamma(t+t')} \frac{1}{2\gamma} (e^{2\gamma t} - 1) = \frac{D_p}{\gamma} [e^{-\gamma(t'-t)} - e^{-\gamma(t+t')}]$.

Temporal correlation of the momentum

$$p(t) = p(0)e^{-\gamma t} + e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$$

- For $t > \frac{1}{\gamma}$, $e^{-\gamma t} \rightarrow 0$, then $p(t) = e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$.
- The temporal correlation:

$$\begin{aligned} \overline{p(t)p(t')} &= \overline{e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi \cdot e^{-\gamma t'} \int_0^{t'} F(\xi')e^{\gamma \xi'} d\xi'} \\ &= e^{-\gamma(t+t')} \int_0^t d\xi \int_0^{t'} d\xi' \overline{F(\xi)F(\xi')} e^{\gamma(\xi+\xi')} \\ &= e^{-\gamma(t+t')} \int_0^t d\xi \int_0^{t'} d\xi' 2D_p \delta(\xi - \xi') e^{\gamma(\xi+\xi')} \end{aligned}$$

- 1. If $t < t'$, $\overline{p(t)p(t')} = e^{-\gamma(t+t')} \int_0^t d\xi 2D_p e^{2\gamma \xi}$
 $= 2D_p e^{-\gamma(t+t')} \frac{1}{2\gamma} (e^{2\gamma t} - 1) = \frac{D_p}{\gamma} [e^{-\gamma(t'-t)} - e^{-\gamma(t+t')}]$.
- 2. Similarly $t > t'$, $\overline{p(t)p(t')} = \frac{D_p}{\gamma} [e^{-\gamma(t-t')} - e^{-\gamma(t+t')}]$.

Temporal correlation of the momentum

$$p(t) = p(0)e^{-\gamma t} + e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$$

- For $t > \frac{1}{\gamma}$, $e^{-\gamma t} \rightarrow 0$, then $p(t) = e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi$.
- The temporal correlation:

$$\begin{aligned} \overline{p(t)p(t')} &= \overline{e^{-\gamma t} \int_0^t F(\xi)e^{\gamma \xi} d\xi \cdot e^{-\gamma t'} \int_0^{t'} F(\xi')e^{\gamma \xi'} d\xi'} \\ &= e^{-\gamma(t+t')} \int_0^t d\xi \int_0^{t'} d\xi' \overline{F(\xi)F(\xi')} e^{\gamma(\xi+\xi')} \\ &= e^{-\gamma(t+t')} \int_0^t d\xi \int_0^{t'} d\xi' 2D_p \delta(\xi - \xi') e^{\gamma(\xi+\xi')} \end{aligned}$$

- 1. If $t < t'$, $\overline{p(t)p(t')} = e^{-\gamma(t+t')} \int_0^t d\xi 2D_p e^{2\gamma \xi}$
 $= 2D_p e^{-\gamma(t+t')} \frac{1}{2\gamma} (e^{2\gamma t} - 1) = \frac{D_p}{\gamma} [e^{-\gamma(t'-t)} - e^{-\gamma(t+t')}]$.
- 2. Similarly $t > t'$, $\overline{p(t)p(t')} = \frac{D_p}{\gamma} [e^{-\gamma(t-t')} - e^{-\gamma(t+t')}]$.
- The combination: $\overline{p(t)p(t')} = \frac{D_p}{\gamma} [e^{-\gamma|t-t'|} - e^{-\gamma(t+t')}]$.

Temporal correlation of the momentum

$$\overline{p(t)p(t')} = \frac{D_p}{\gamma} [e^{-\gamma|t-t'|} - e^{-\gamma(t+t')}]$$

Temporal correlation of the momentum

$$\overline{p(t)p(t')} = \frac{D_p}{\gamma} [e^{-\gamma|t-t'|} - e^{-\gamma(t+t')}]$$

- For $t, t' > \frac{1}{\gamma}$ (normal), $e^{-\gamma(t+t')} \rightarrow 0$;

Temporal correlation of the momentum

$$\overline{p(t)p(t')} = \frac{D_p}{\gamma} [e^{-\gamma|t-t'|} - e^{-\gamma(t+t')}]$$

- For $t, t' > \frac{1}{\gamma}$ (normal), $e^{-\gamma(t+t')} \rightarrow 0$; then

$$\overline{p(t)p(t')} = \frac{D_p}{\gamma} e^{-\gamma|t-t'|} = mkTe^{-\gamma|t-t'|}.$$

Temporal correlation of the momentum

$$\overline{p(t)p(t')} = \frac{D_p}{\gamma} [e^{-\gamma|t-t'|} - e^{-\gamma(t+t')}]$$

- For $t, t' > \frac{1}{\gamma}$ (normal), $e^{-\gamma(t+t')} \rightarrow 0$; then
$$\overline{p(t)p(t')} = \frac{D_p}{\gamma} e^{-\gamma|t-t'|} = mkTe^{-\gamma|t-t'|}.$$
- Back to $\overline{x^2(t)}$ again.

Temporal correlation of the momentum

$$\overline{p(t)p(t')} = \frac{D_p}{\gamma} [e^{-\gamma|t-t'|} - e^{-\gamma(t+t')}]$$

- For $t, t' > \frac{1}{\gamma}$ (normal), $e^{-\gamma(t+t')} \rightarrow 0$; then

$$\overline{p(t)p(t')} = \frac{D_p}{\gamma} e^{-\gamma|t-t'|} = mkTe^{-\gamma|t-t'|}.$$

- Back to $\overline{x^2(t)}$ again.

$$x(t) = \int_0^t \dot{x}(\xi) d\xi$$

Temporal correlation of the momentum

$$\overline{p(t)p(t')} = \frac{D_p}{\gamma} [e^{-\gamma|t-t'|} - e^{-\gamma(t+t')}]$$

- For $t, t' > \frac{1}{\gamma}$ (normal), $e^{-\gamma(t+t')} \rightarrow 0$; then

$$\overline{p(t)p(t')} = \frac{D_p}{\gamma} e^{-\gamma|t-t'|} = mkTe^{-\gamma|t-t'|}.$$

- Back to $\overline{x^2(t)}$ again.

$$x(t) = \int_0^t \dot{x}(\xi) d\xi = \frac{1}{m} \int_0^t p(\xi) d\xi.$$

Temporal correlation of the momentum

$$\overline{p(t)p(t')} = \frac{D_p}{\gamma} [e^{-\gamma|t-t'|} - e^{-\gamma(t+t')}]$$

- For $t, t' > \frac{1}{\gamma}$ (normal), $e^{-\gamma(t+t')} \rightarrow 0$; then

$$\overline{p(t)p(t')} = \frac{D_p}{\gamma} e^{-\gamma|t-t'|} = mkTe^{-\gamma|t-t'|}.$$

- Back to $\overline{x^2(t)}$ again.

$$x(t) = \int_0^t \dot{x}(\xi) d\xi = \frac{1}{m} \int_0^t p(\xi) d\xi.$$

$$\overline{x^2(t)} = \overline{\frac{1}{m} \int_0^t p(\xi) d\xi \cdot \frac{1}{m} \int_0^t p(\xi') d\xi'}$$

Temporal correlation of the momentum

$$\overline{p(t)p(t')} = \frac{D_p}{\gamma} [e^{-\gamma|t-t'|} - e^{-\gamma(t+t')}]$$

- For $t, t' > \frac{1}{\gamma}$ (normal), $e^{-\gamma(t+t')} \rightarrow 0$; then

$$\overline{p(t)p(t')} = \frac{D_p}{\gamma} e^{-\gamma|t-t'|} = mkTe^{-\gamma|t-t'|}.$$

- Back to $\overline{x^2(t)}$ again.

$$x(t) = \int_0^t \dot{x}(\xi) d\xi = \frac{1}{m} \int_0^t p(\xi) d\xi.$$

$$\begin{aligned} \overline{x^2(t)} &= \overline{\frac{1}{m} \int_0^t p(\xi) d\xi \cdot \frac{1}{m} \int_0^t p(\xi') d\xi'} \\ &= \frac{1}{m^2} \int_0^t d\xi \int_0^t d\xi' \overline{p(\xi)p(\xi')} \end{aligned}$$

Temporal correlation of the momentum

$$\overline{p(t)p(t')} = \frac{D_p}{\gamma} [e^{-\gamma|t-t'|} - e^{-\gamma(t+t')}]$$

- For $t, t' > \frac{1}{\gamma}$ (normal), $e^{-\gamma(t+t')} \rightarrow 0$; then

$$\overline{p(t)p(t')} = \frac{D_p}{\gamma} e^{-\gamma|t-t'|} = mkTe^{-\gamma|t-t'|}.$$

- Back to $\overline{x^2(t)}$ again.

$$x(t) = \int_0^t \dot{x}(\xi) d\xi = \frac{1}{m} \int_0^t p(\xi) d\xi.$$

$$\begin{aligned} \overline{x^2(t)} &= \frac{1}{m} \int_0^t p(\xi) d\xi \cdot \frac{1}{m} \int_0^t p(\xi') d\xi' \\ &= \frac{1}{m^2} \int_0^t d\xi \int_0^t d\xi' \overline{p(\xi)p(\xi')} \\ &= \frac{kT}{m} \int_0^t d\xi \int_0^t d\xi' e^{-\gamma|\xi-\xi'|} \end{aligned}$$

Temporal correlation of the momentum

$$\overline{p(t)p(t')} = \frac{D_p}{\gamma} [e^{-\gamma|t-t'|} - e^{-\gamma(t+t')}]$$

- For $t, t' > \frac{1}{\gamma}$ (normal), $e^{-\gamma(t+t')} \rightarrow 0$; then

$$\overline{p(t)p(t')} = \frac{D_p}{\gamma} e^{-\gamma|t-t'|} = mkT e^{-\gamma|t-t'|}.$$

- Back to $\overline{x^2(t)}$ again.

$$x(t) = \int_0^t \dot{x}(\xi) d\xi = \frac{1}{m} \int_0^t p(\xi) d\xi.$$

$$\begin{aligned} \overline{x^2(t)} &= \overline{\frac{1}{m} \int_0^t p(\xi) d\xi \cdot \frac{1}{m} \int_0^t p(\xi') d\xi'} \\ &= \frac{1}{m^2} \int_0^t d\xi \int_0^t d\xi' \overline{p(\xi)p(\xi')} \\ &= \frac{kT}{m} \int_0^t d\xi \int_0^t d\xi' e^{-\gamma|\xi-\xi'|} \\ &= \frac{kT}{m} \int_0^t d\xi \left[\int_0^\xi d\xi' e^{-\gamma(\xi-\xi')} + \int_\xi^t d\xi' e^{-\gamma(\xi'-\xi)} \right] \end{aligned}$$

Temporal correlation of the momentum

$$\overline{p(t)p(t')} = \frac{D_p}{\gamma} [e^{-\gamma|t-t'|} - e^{-\gamma(t+t')}]$$

- For $t, t' > \frac{1}{\gamma}$ (normal), $e^{-\gamma(t+t')} \rightarrow 0$; then

$$\overline{p(t)p(t')} = \frac{D_p}{\gamma} e^{-\gamma|t-t'|} = mkTe^{-\gamma|t-t'|}.$$

- Back to $\overline{x^2(t)}$ again.

$$x(t) = \int_0^t \dot{x}(\xi) d\xi = \frac{1}{m} \int_0^t p(\xi) d\xi.$$

$$\begin{aligned} \overline{x^2(t)} &= \overline{\frac{1}{m} \int_0^t p(\xi) d\xi \cdot \frac{1}{m} \int_0^t p(\xi') d\xi'} \\ &= \frac{1}{m^2} \int_0^t d\xi \int_0^t d\xi' \overline{p(\xi)p(\xi')} \\ &= \frac{kT}{m} \int_0^t d\xi \int_0^t d\xi' e^{-\gamma|\xi-\xi'|} \\ &= \frac{kT}{m} \int_0^t d\xi \left[\int_0^\xi d\xi' e^{-\gamma(\xi-\xi')} + \int_\xi^t d\xi' e^{-\gamma(\xi'-\xi)} \right] \\ &= \frac{kT}{m} \int_0^t d\xi \left[\int_0^\xi d\xi' e^{\gamma(\xi'-\xi)} + \int_\xi^t d\xi' e^{\gamma(\xi-\xi')} \right] \end{aligned}$$

Temporal correlation of the momentum

$$\overline{x^2(t)} = \frac{kT}{m} \int_0^t d\xi \left[\int_0^\xi d\xi' e^{\gamma(\xi' - \xi)} + \int_\xi^t d\xi' e^{\gamma(\xi - \xi')} \right]$$

Temporal correlation of the momentum

$$\begin{aligned}\overline{x^2(t)} &= \frac{kT}{m} \int_0^t d\xi \left[\int_0^\xi d\xi' e^{\gamma(\xi'-\xi)} + \int_\xi^t d\xi' e^{\gamma(\xi-\xi')} \right] \\ &= \frac{kT}{m} \int_0^t d\xi \left[\frac{1}{\gamma} \int_0^\xi d\xi' e^{\gamma(\xi'-\xi)} - \frac{1}{\gamma} \int_\xi^t d\xi' e^{\gamma(\xi-\xi')} \right]\end{aligned}$$

Temporal correlation of the momentum

$$\begin{aligned}\overline{x^2(t)} &= \frac{kT}{m} \int_0^t d\xi \left[\int_0^\xi d\xi' e^{\gamma(\xi'-\xi)} + \int_\xi^t d\xi' e^{\gamma(\xi-\xi')} \right] \\ &= \frac{kT}{m} \int_0^t d\xi \left[\frac{1}{\gamma} \int_0^\xi d\xi' e^{\gamma(\xi'-\xi)} - \frac{1}{\gamma} \int_\xi^t d\xi' e^{\gamma(\xi-\xi')} \right] \\ &= \frac{kT}{m} \int_0^t d\xi \left[\frac{1}{\gamma} (1 - e^{-\gamma\xi}) - \frac{1}{\gamma} (e^{\gamma(\xi-t)} - 1) \right]\end{aligned}$$

Temporal correlation of the momentum

$$\begin{aligned}\overline{x^2(t)} &= \frac{kT}{m} \int_0^t d\xi \left[\int_0^\xi d\xi' e^{\gamma(\xi'-\xi)} + \int_\xi^t d\xi' e^{\gamma(\xi-\xi')} \right] \\ &= \frac{kT}{m} \int_0^t d\xi \left[\frac{1}{\gamma} \int_0^\xi d\xi' e^{\gamma(\xi'-\xi)} - \frac{1}{\gamma} \int_\xi^t d\xi' e^{\gamma(\xi-\xi')} \right] \\ &= \frac{kT}{m} \int_0^t d\xi \left[\frac{1}{\gamma} (1 - e^{-\gamma\xi}) - \frac{1}{\gamma} (e^{\gamma(\xi-t)} - 1) \right] \\ &= \frac{kT}{\gamma m} \int_0^t d\xi \left[2 - e^{-\gamma\xi} - e^{\gamma(\xi-t)} \right]\end{aligned}$$

Temporal correlation of the momentum

$$\begin{aligned}\overline{x^2(t)} &= \frac{kT}{m} \int_0^t d\xi \left[\int_0^\xi d\xi' e^{\gamma(\xi'-\xi)} + \int_\xi^t d\xi' e^{\gamma(\xi-\xi')} \right] \\ &= \frac{kT}{m} \int_0^t d\xi \left[\frac{1}{\gamma} \int_0^\xi d\xi' e^{\gamma(\xi'-\xi)} - \frac{1}{\gamma} \int_\xi^t d\xi' e^{\gamma(\xi-\xi')} \right] \\ &= \frac{kT}{m} \int_0^t d\xi \left[\frac{1}{\gamma} (1 - e^{-\gamma\xi}) - \frac{1}{\gamma} (e^{\gamma(\xi-t)} - 1) \right] \\ &= \frac{kT}{\gamma m} \int_0^t d\xi [2 - e^{-\gamma\xi} - e^{\gamma(\xi-t)}] \\ &= \frac{kT}{\gamma m} \left[2t + \frac{1}{\gamma} \int_0^t d\xi e^{-\gamma\xi} - \frac{1}{\gamma} \int_0^t d\xi e^{\gamma(\xi-t)} \right]\end{aligned}$$

Temporal correlation of the momentum

$$\begin{aligned}\overline{x^2(t)} &= \frac{kT}{m} \int_0^t d\xi \left[\int_0^\xi d\xi' e^{\gamma(\xi'-\xi)} + \int_\xi^t d\xi' e^{\gamma(\xi-\xi')} \right] \\ &= \frac{kT}{m} \int_0^t d\xi \left[\frac{1}{\gamma} \int_0^\xi d\xi' e^{\gamma(\xi'-\xi)} - \frac{1}{\gamma} \int_\xi^t d\xi' e^{\gamma(\xi-\xi')} \right] \\ &= \frac{kT}{m} \int_0^t d\xi \left[\frac{1}{\gamma} (1 - e^{-\gamma\xi}) - \frac{1}{\gamma} (e^{\gamma(\xi-t)} - 1) \right] \\ &= \frac{kT}{\gamma m} \int_0^t d\xi [2 - e^{-\gamma\xi} - e^{\gamma(\xi-t)}] \\ &= \frac{kT}{\gamma m} \left[2t + \frac{1}{\gamma} \int_0^t d\xi e^{-\gamma\xi} - \frac{1}{\gamma} \int_0^t d\xi e^{\gamma(\xi-t)} \right] \\ &= \frac{kT}{\gamma m} \left[2t + \frac{1}{\gamma} (e^{-\gamma t} - 1) - \frac{1}{\gamma} (1 - e^{-\gamma t}) \right]\end{aligned}$$

Temporal correlation of the momentum

$$\begin{aligned}\overline{x^2(t)} &= \frac{kT}{m} \int_0^t d\xi \left[\int_0^\xi d\xi' e^{\gamma(\xi'-\xi)} + \int_\xi^t d\xi' e^{\gamma(\xi-\xi')} \right] \\&= \frac{kT}{m} \int_0^t d\xi \left[\frac{1}{\gamma} \int_0^\xi d\xi' e^{\gamma(\xi'-\xi)} - \frac{1}{\gamma} \int_\xi^t d\xi' e^{\gamma(\xi-\xi')} \right] \\&= \frac{kT}{m} \int_0^t d\xi \left[\frac{1}{\gamma} (1 - e^{-\gamma\xi}) - \frac{1}{\gamma} (e^{\gamma(\xi-t)} - 1) \right] \\&= \frac{kT}{\gamma m} \int_0^t d\xi [2 - e^{-\gamma\xi} - e^{\gamma(\xi-t)}] \\&= \frac{kT}{\gamma m} \left[2t + \frac{1}{\gamma} \int_0^t d\xi e^{-\gamma\xi} - \frac{1}{\gamma} \int_0^t d\xi e^{\gamma(\xi-t)} \right] \\&= \frac{kT}{\gamma m} \left[2t + \frac{1}{\gamma} (e^{-\gamma t} - 1) - \frac{1}{\gamma} (1 - e^{-\gamma t}) \right] \\&= \frac{2kT}{\gamma m} t \quad (e^{-\gamma t} \rightarrow 0)\end{aligned}$$

Temporal correlation of the momentum

$$\begin{aligned}\overline{x^2(t)} &= \frac{kT}{m} \int_0^t d\xi \left[\int_0^\xi d\xi' e^{\gamma(\xi'-\xi)} + \int_\xi^t d\xi' e^{\gamma(\xi-\xi')} \right] \\&= \frac{kT}{m} \int_0^t d\xi \left[\frac{1}{\gamma} \int_0^\xi d\xi' e^{\gamma(\xi'-\xi)} - \frac{1}{\gamma} \int_\xi^t d\xi' e^{\gamma(\xi-\xi')} \right] \\&= \frac{kT}{m} \int_0^t d\xi \left[\frac{1}{\gamma} (1 - e^{-\gamma\xi}) - \frac{1}{\gamma} (e^{\gamma(\xi-t)} - 1) \right] \\&= \frac{kT}{\gamma m} \int_0^t d\xi [2 - e^{-\gamma\xi} - e^{\gamma(\xi-t)}] \\&= \frac{kT}{\gamma m} \left[2t + \frac{1}{\gamma} \int_0^t d\xi e^{-\gamma\xi} - \frac{1}{\gamma} \int_0^t d\xi e^{\gamma(\xi-t)} \right] \\&= \frac{kT}{\gamma m} \left[2t + \frac{1}{\gamma} (e^{-\gamma t} - 1) - \frac{1}{\gamma} (1 - e^{-\gamma t}) \right] \\&= \frac{2kT}{\gamma m} t \quad (e^{-\gamma t} \rightarrow 0) \\&= \frac{2kT}{\alpha} t. \quad (\text{same as (10.5.8)}).\end{aligned}$$

Table of contents

1 §10. Fluctuation

- 10.1 Quasi-thermodynamics of fluctuation
- 10.5 Brownian motion
- 10.6 Diffusion and temporal correlation of Brownian particle's momentum
- 10.7 Examples of Brownian motion

§10.7 Examples of Brownian motion

§10.7 Examples of Brownian motion

- (i) Thermal noise in the circuit.

§10.7 Examples of Brownian motion

- (i) Thermal noise in the circuit.
- In the LC circuit, $L \frac{di}{dt} = -\frac{q(t)}{C}$.

§10.7 Examples of Brownian motion

- (i) Thermal noise in the circuit.
- In the LC circuit, $L \frac{di}{dt} = -\frac{q(t)}{C}$.
- Considering the resistance: $L \frac{di}{dt} = -\frac{q(t)}{C} - Ri$.

§10.7 Examples of Brownian motion

- (i) Thermal noise in the circuit.
- In the LC circuit, $L \frac{di}{dt} = -\frac{q(t)}{C}$.
- Considering the resistance: $L \frac{di}{dt} = -\frac{q(t)}{C} - Ri$.
- Reason for the resistance: vibrations of the ions scatter the moving electrons.

§10.7 Examples of Brownian motion

- (i) Thermal noise in the circuit.
- In the LC circuit, $L \frac{di}{dt} = -\frac{q(t)}{C}$.
- Considering the resistance: $L \frac{di}{dt} = -\frac{q(t)}{C} - Ri$.
- Reason for the resistance: vibrations of the ions scatter the moving electrons. Two effect: macroscopic effective resistance;

§10.7 Examples of Brownian motion

- (i) Thermal noise in the circuit.
- In the LC circuit, $L \frac{di}{dt} = -\frac{q(t)}{C}$.
- Considering the resistance: $L \frac{di}{dt} = -\frac{q(t)}{C} - Ri$.
- Reason for the resistance: vibrations of the ions scatter the moving electrons. Two effect: macroscopic effective resistance; fluctuating potential $V(t)$.

§10.7 Examples of Brownian motion

- (i) Thermal noise in the circuit.
- In the LC circuit, $L \frac{di}{dt} = -\frac{q(t)}{C}$.
- Considering the resistance: $L \frac{di}{dt} = -\frac{q(t)}{C} - Ri$.
- Reason for the resistance: vibrations of the ions scatter the moving electrons. Two effect: macroscopic effective resistance; fluctuating potential $V(t)$.
- $\therefore L \frac{di}{dt} = -\frac{q(t)}{C} - Ri + V(t)$.

§10.7 Examples of Brownian motion

- (i) Thermal noise in the circuit.
- In the LC circuit, $L \frac{di}{dt} = -\frac{q(t)}{C}$.
- Considering the resistance: $L \frac{di}{dt} = -\frac{q(t)}{C} - Ri$.
- Reason for the resistance: vibrations of the ions scatter the moving electrons. Two effect: macroscopic effective resistance; fluctuating potential $V(t)$.
- $\therefore L \frac{di}{dt} = -\frac{q(t)}{C} - Ri + V(t)$.
- Abstracting the external voltage $-\frac{q(t)}{C}$ to $\mathcal{V}(t)$,
 $L \frac{di}{dt} = \mathcal{V}(t) - Ri + V(t)$.

§10.7 Examples of Brownian motion

- (i) Thermal noise in the circuit.
- In the LC circuit, $L \frac{di}{dt} = -\frac{q(t)}{C}$.
- Considering the resistance: $L \frac{di}{dt} = -\frac{q(t)}{C} - Ri$.
- Reason for the resistance: vibrations of the ions scatter the moving electrons. Two effect: macroscopic effective resistance; fluctuating potential $V(t)$.
- $\therefore L \frac{di}{dt} = -\frac{q(t)}{C} - Ri + V(t)$.
- Abstracting the external voltage $-\frac{q(t)}{C}$ to $\mathcal{V}(t)$,
 $L \frac{di}{dt} = \mathcal{V}(t) - Ri + V(t)$.
- $m \frac{dv}{dt} = \mathcal{F} - \alpha v + F(t)$, Langevin's equation.
 $L \leftrightarrow m, i \leftrightarrow v, \mathcal{V} \leftrightarrow \mathcal{F}, R \leftrightarrow \alpha, \text{ and } V \leftrightarrow F$.

Thermal noise in the circuit

- Similar to (10.6.3), $\overline{F(t)F(t+\tau)} = 2D_p\delta(\tau)$,
 $D_p = \alpha kT$;

Thermal noise in the circuit

- Similar to (10.6.3), $\overline{F(t)F(t+\tau)} = 2D_p\delta(\tau)$,
 $D_p = \alpha kT$; $\rightarrow \overline{V(t)V(t+\tau)} = 2RkT\delta(\tau)$.

Thermal noise in the circuit

- Similar to (10.6.3), $\overline{F(t)F(t+\tau)} = 2D_p\delta(\tau)$,
 $D_p = \alpha kT$; $\rightarrow \overline{V(t)V(t+\tau)} = 2RkT\delta(\tau)$.
- Fourier transformation (time domain to frequency domain):

Thermal noise in the circuit

- Similar to (10.6.3), $\overline{F(t)F(t+\tau)} = 2D_p\delta(\tau)$,
 $D_p = \alpha kT$; $\rightarrow \overline{V(t)V(t+\tau)} = 2RkT\delta(\tau)$.
- Fourier transformation (time domain to frequency domain): $V(t) = \int_{-\infty}^{\infty} V(\omega)e^{i\omega t}d\omega$;

Thermal noise in the circuit

- Similar to (10.6.3), $\overline{F(t)F(t+\tau)} = 2D_p\delta(\tau)$,
 $D_p = \alpha kT$; $\rightarrow \overline{V(t)V(t+\tau)} = 2RkT\delta(\tau)$.
- Fourier transformation (time domain to frequency domain): $V(t) = \int_{-\infty}^{\infty} V(\omega)e^{i\omega t}d\omega$;
 $V(\omega) = \frac{1}{2\pi} \int_{-\infty}^{\infty} V(t)e^{-i\omega t}dt$.

Thermal noise in the circuit

- Similar to (10.6.3), $\overline{F(t)F(t+\tau)} = 2D_p\delta(\tau)$,
 $D_p = \alpha kT$; $\rightarrow \overline{V(t)V(t+\tau)} = 2RkT\delta(\tau)$.
- Fourier transformation (time domain to frequency domain): $V(t) = \int_{-\infty}^{\infty} V(\omega)e^{i\omega t}d\omega$;
 $V(\omega) = \frac{1}{2\pi} \int_{-\infty}^{\infty} V(t)e^{-i\omega t}dt$.
- $\overline{V(\omega)V^*(\omega')} = \overline{V(\omega)V(-\omega')}$

Thermal noise in the circuit

- Similar to (10.6.3), $\overline{F(t)F(t+\tau)} = 2D_p\delta(\tau)$,
 $D_p = \alpha kT$; $\rightarrow \overline{V(t)V(t+\tau)} = 2RkT\delta(\tau)$.
- Fourier transformation (time domain to frequency domain): $V(t) = \int_{-\infty}^{\infty} V(\omega)e^{i\omega t}d\omega$;
 $V(\omega) = \frac{1}{2\pi} \int_{-\infty}^{\infty} V(t)e^{-i\omega t}dt$.
- $\overline{V(\omega)V^*(\omega')} = \overline{V(\omega)V(-\omega')}$
 $= \frac{1}{(2\pi)^2} \iint_{-\infty}^{\infty} dt dt' \overline{V(t)V(t')} e^{-i\omega t} e^{i\omega' t'}$

Thermal noise in the circuit

- Similar to (10.6.3), $\overline{F(t)F(t+\tau)} = 2D_p\delta(\tau)$,
 $D_p = \alpha kT$; $\rightarrow \overline{V(t)V(t+\tau)} = 2RkT\delta(\tau)$.
- Fourier transformation (time domain to frequency domain): $V(t) = \int_{-\infty}^{\infty} V(\omega)e^{i\omega t}d\omega$;
 $V(\omega) = \frac{1}{2\pi} \int_{-\infty}^{\infty} V(t)e^{-i\omega t}dt$.
- $\overline{V(\omega)V^*(\omega')} = \overline{V(\omega)V(-\omega')}$
 $= \frac{1}{(2\pi)^2} \iint_{-\infty}^{\infty} dt dt' \overline{V(t)V(t')} e^{-i\omega t} e^{i\omega' t'}$
 $= \frac{1}{(2\pi)^2} \iint_{-\infty}^{\infty} dt dt' 2RkT\delta(t' - t) e^{-i\omega t} e^{i\omega' t'}$

Thermal noise in the circuit

- Similar to (10.6.3), $\overline{F(t)F(t+\tau)} = 2D_p\delta(\tau)$,
 $D_p = \alpha kT$; $\rightarrow \overline{V(t)V(t+\tau)} = 2RkT\delta(\tau)$.
- Fourier transformation (time domain to frequency domain): $V(t) = \int_{-\infty}^{\infty} V(\omega)e^{i\omega t}d\omega$;
 $V(\omega) = \frac{1}{2\pi} \int_{-\infty}^{\infty} V(t)e^{-i\omega t}dt$.
- $\overline{V(\omega)V^*(\omega')} = \overline{V(\omega)V(-\omega')}$
 $= \frac{1}{(2\pi)^2} \iint_{-\infty}^{\infty} dt dt' \overline{V(t)V(t')} e^{-i\omega t} e^{i\omega' t'}$
 $= \frac{1}{(2\pi)^2} \iint_{-\infty}^{\infty} dt dt' 2RkT\delta(t' - t) e^{-i\omega t} e^{i\omega' t'}$
 $= \frac{1}{(2\pi)^2} \int_{-\infty}^{\infty} 2RkT e^{-i(\omega - \omega')t} dt$

Thermal noise in the circuit

- Similar to (10.6.3), $\overline{F(t)F(t+\tau)} = 2D_p\delta(\tau)$,
 $D_p = \alpha kT$; $\rightarrow \overline{V(t)V(t+\tau)} = 2RkT\delta(\tau)$.
- Fourier transformation (time domain to frequency domain): $V(t) = \int_{-\infty}^{\infty} V(\omega)e^{i\omega t}d\omega$;
 $V(\omega) = \frac{1}{2\pi} \int_{-\infty}^{\infty} V(t)e^{-i\omega t}dt$.
- $\overline{V(\omega)V^*(\omega')} = \overline{V(\omega)V(-\omega')}$
 $= \frac{1}{(2\pi)^2} \iint_{-\infty}^{\infty} dt dt' \overline{V(t)V(t')} e^{-i\omega t} e^{i\omega' t'}$
 $= \frac{1}{(2\pi)^2} \iint_{-\infty}^{\infty} dt dt' 2RkT\delta(t' - t) e^{-i\omega t} e^{i\omega' t'}$
 $= \frac{1}{(2\pi)^2} \int_{-\infty}^{\infty} 2RkT e^{-i(\omega - \omega')t} dt$
 $= \frac{kTR}{\pi} \delta(\omega - \omega'), \text{ (white noise)}$
 where $\frac{1}{2\pi} \int_{-\infty}^{\infty} e^{i(\omega' - \omega)t} dt = \delta(\omega - \omega')$.

Optical adhesive and Doppler cooling

- (ii) Optical adhesive and Doppler cooling.

Optical adhesive and Doppler cooling

- (ii) Optical adhesive and Doppler cooling.
- Frequency of laser: ω_L ; of atom: ω_A . ($\omega_L \lesssim \omega_A$.)

Optical adhesive and Doppler cooling

- (ii) Optical adhesive and Doppler cooling.
- Frequency of laser: ω_L ; of atom: ω_A . ($\omega_L \lesssim \omega_A$.)
- For the atom at velocity v_z (1 direction), it feels the frequency of laser $\omega_L(1 \pm \frac{|v_z|}{c})$.

Optical adhesive and Doppler cooling

- (ii) Optical adhesive and Doppler cooling.
- Frequency of laser: ω_L ; of atom: ω_A . ($\omega_L \lesssim \omega_A$.)
- For the atom at velocity v_z (1 direction), it feels the frequency of laser $\omega_L(1 \pm \frac{|v_z|}{c})$. Higher velocity, closer to ω_A ;

Optical adhesive and Doppler cooling

- (ii) Optical adhesive and Doppler cooling.
- Frequency of laser: ω_L ; of atom: ω_A . ($\omega_L \lesssim \omega_A$.)
- For the atom at velocity v_z (1 direction), it feels the frequency of laser $\omega_L(1 \pm \frac{|v_z|}{c})$. Higher velocity, closer to ω_A ; more easier to absorb the photon and lose its momentum.

Optical adhesive and Doppler cooling

- (ii) Optical adhesive and Doppler cooling.
- Frequency of laser: ω_L ; of atom: ω_A . ($\omega_L \lesssim \omega_A$.)
- For the atom at velocity v_z (1 direction), it feels the frequency of laser $\omega_L(1 \pm \frac{|v_z|}{c})$. Higher velocity, closer to ω_A ; more easier to absorb the photon and lose its momentum. Effective viscous force $-\alpha v_z$.

Optical adhesive and Doppler cooling

- (ii) Optical adhesive and Doppler cooling.
- Frequency of laser: ω_L ; of atom: ω_A . ($\omega_L \lesssim \omega_A$.)
- For the atom at velocity v_z (1 direction), it feels the frequency of laser $\omega_L(1 \pm \frac{|v_z|}{c})$. Higher velocity, closer to ω_A ; more easier to absorb the photon and lose its momentum. Effective viscous force $-\alpha v_z$.
- When the excited atom transits to the ground state,

Optical adhesive and Doppler cooling

- (ii) Optical adhesive and Doppler cooling.
- Frequency of laser: ω_L ; of atom: ω_A . ($\omega_L \lesssim \omega_A$.)
- For the atom at velocity v_z (1 direction), it feels the frequency of laser $\omega_L(1 \pm \frac{|v_z|}{c})$. Higher velocity, closer to ω_A ; more easier to absorb the photon and lose its momentum. Effective viscous force $-\alpha v_z$.
- When the excited atom transits to the ground state, it radiates a photon and feels a repulsive force $F_z(t)$.

Optical adhesive and Doppler cooling

- (ii) Optical adhesive and Doppler cooling.
- Frequency of laser: ω_L ; of atom: ω_A . ($\omega_L \lesssim \omega_A$.)
- For the atom at velocity v_z (1 direction), it feels the frequency of laser $\omega_L(1 \pm \frac{|v_z|}{c})$. Higher velocity, closer to ω_A ; more easier to absorb the photon and lose its momentum. Effective viscous force $-\alpha v_z$.
- When the excited atom transits to the ground state, it radiates a photon and feels a repulsive force $F_z(t)$.
- Langevin's equation: $m \frac{dv_z}{dt} = -\alpha v_z + F_z$.

Optical adhesive and Doppler cooling

- (ii) Optical adhesive and Doppler cooling.
- Frequency of laser: ω_L ; of atom: ω_A . ($\omega_L \lesssim \omega_A$.)
- For the atom at velocity v_z (1 direction), it feels the frequency of laser $\omega_L(1 \pm \frac{|v_z|}{c})$. Higher velocity, closer to ω_A ; more easier to absorb the photon and lose its momentum. Effective viscous force $-\alpha v_z$.
- When the excited atom transits to the ground state, it radiates a photon and feels a repulsive force $F_z(t)$.
- Langevin's equation: $m \frac{dv_z}{dt} = -\alpha v_z + F_z$.
- Atom \leftrightarrow Brownian particle; laser photons \leftrightarrow water (optical adhesive).

Optical adhesive and Doppler cooling

- (ii) Optical adhesive and Doppler cooling.
- Frequency of laser: ω_L ; of atom: ω_A . ($\omega_L \lesssim \omega_A$.)
- For the atom at velocity v_z (1 direction), it feels the frequency of laser $\omega_L(1 \pm \frac{|v_z|}{c})$. Higher velocity, closer to ω_A ; more easier to absorb the photon and lose its momentum. Effective viscous force $-\alpha v_z$.
- When the excited atom transits to the ground state, it radiates a photon and feels a repulsive force $F_z(t)$.
- Langevin's equation: $m \frac{dv_z}{dt} = -\alpha v_z + F_z$.
- Atom \leftrightarrow Brownian particle; laser photons \leftrightarrow water (optical adhesive). (10.6.15) $\Rightarrow kT = \frac{D_p}{\alpha}$, the lowest temperature by Doppler cooling.

(iii) Magneto-optical trap

(iii) Magneto-optical trap

- However, the optical adhesive can not trap the atom, because of the dissipation.

(iii) Magneto-optical trap

- However, the optical adhesive can not trap the atom, because of the dissipation.
- Add a magnetic field (1-D) along z-axis, with strength $B_z = \lambda z$.

(iii) Magneto-optical trap

- However, the optical adhesive can not trap the atom, because of the dissipation.
- Add a magnetic field (1-D) along z-axis, with strength $B_z = \lambda z$.
- Suppose angular momentum quantum number $j = 0$ for ground state, $j = 1$ for excited state.

(iii) Magneto-optical trap

- However, the optical adhesive can not trap the atom, because of the dissipation.
- Add a magnetic field (1-D) along z-axis, with strength $B_z = \lambda z$.
- Suppose angular momentum quantum number $j = 0$ for ground state, $j = 1$ for excited state. Zeeman splitting: $j = 1 \rightarrow m_j = \pm 1, 0$, with energy level: $\varepsilon_{2,m_j} = \varepsilon_2 + \frac{e\hbar}{2m_e} B_z m_j$.

(iii) Magneto-optical trap

- However, the optical adhesive can not trap the atom, because of the dissipation.
- Add a magnetic field (1-D) along z-axis, with strength $B_z = \lambda z$.
- Suppose angular momentum quantum number $j = 0$ for ground state, $j = 1$ for excited state. Zeeman splitting: $j = 1 \rightarrow m_j = \pm 1, 0$, with energy level:
$$\varepsilon_{2,m_j} = \varepsilon_2 + \frac{e\hbar}{2m_e} B_z m_j.$$
- The lowest energy changes to $\varepsilon_{2,\text{low}} = \varepsilon_2 - \frac{e\hbar}{2m} B_z$

(iii) Magneto-optical trap

- However, the optical adhesive can not trap the atom, because of the dissipation.
- Add a magnetic field (1-D) along z-axis, with strength $B_z = \lambda z$.
- Suppose angular momentum quantum number $j = 0$ for ground state, $j = 1$ for excited state. Zeeman splitting: $j = 1 \rightarrow m_j = \pm 1, 0$, with energy level: $\varepsilon_{2,m_j} = \varepsilon_2 + \frac{e\hbar}{2m_e} B_z m_j$.
- The lowest energy changes to $\varepsilon_{2,\text{low}} = \varepsilon_2 - \frac{e\hbar}{2m} B_z = \varepsilon_2 - \frac{e\hbar}{2m} \lambda z$.

(iii) Magneto-optical trap

- However, the optical adhesive can not trap the atom, because of the dissipation.
- Add a magnetic field (1-D) along z-axis, with strength $B_z = \lambda z$.
- Suppose angular momentum quantum number $j = 0$ for ground state, $j = 1$ for excited state. Zeeman splitting: $j = 1 \rightarrow m_j = \pm 1, 0$, with energy level: $\varepsilon_{2,m_j} = \varepsilon_2 + \frac{e\hbar}{2m_e} B_z m_j$.
- The lowest energy changes to $\varepsilon_{2,\text{low}} = \varepsilon_2 - \frac{e\hbar}{2m} B_z = \varepsilon_2 - \frac{e\hbar}{2m} \lambda z$. The energy $\varepsilon_2 - \frac{e\hbar}{2m} \lambda z - \varepsilon_1$ approaching $\hbar\omega_L$ more closely with z increasing.

(iii) Magneto-optical trap

- However, the optical adhesive can not trap the atom, because of the dissipation.
- Add a magnetic field (1-D) along z-axis, with strength $B_z = \lambda z$.
- Suppose angular momentum quantum number $j = 0$ for ground state, $j = 1$ for excited state. Zeeman splitting: $j = 1 \rightarrow m_j = \pm 1, 0$, with energy level: $\varepsilon_{2,m_j} = \varepsilon_2 + \frac{e\hbar}{2m_e} B_z m_j$.
- The lowest energy changes to $\varepsilon_{2,\text{low}} = \varepsilon_2 - \frac{e\hbar}{2m} B_z = \varepsilon_2 - \frac{e\hbar}{2m} \lambda z$. The energy $\varepsilon_2 - \frac{e\hbar}{2m} \lambda z - \varepsilon_1$ approaching $\hbar\omega_L$ more closely with z increasing. Effective force: $-kz$, traps the atoms.

Table of contents

1 §10. Fluctuation

- 10.1 Quasi-thermodynamics of fluctuation
- 10.5 Brownian motion
- 10.6 Diffusion and temporal correlation of Brownian particle's momentum
- 10.7 Examples of Brownian motion