

Thermodynamics & Statistical Physics

Chapter 5. Thermodynamics of irreversible processes

Yuan-Chuan Zou
zouyc@hust.edu.cn

School of Physics, Huazhong University of Science and Technology

December 30, 2013

Table of contents

- ① §5. Thermodynamics of irreversible processes
 - 5.1 Local equilibrium, entropy flux density and production rate
 - 5.2 Linear and nonlinear processes, Onsager relation

5.1 Local equilibrium, entropy flux density, entropy local production rate

5.1 Local equilibrium, entropy flux density, entropy local production rate

- Irreversible processes.

5.1 Local equilibrium, entropy flux density, entropy local production rate

- Irreversible processes.
- 2nd Law: $dS \geq \frac{\delta Q}{T}$,

5.1 Local equilibrium, entropy flux density, entropy local production rate

- Irreversible processes.
- 2nd Law: $dS \geq \frac{\delta Q}{T}$,
= reversible; > irreversible.

5.1 Local equilibrium, entropy flux density, entropy local production rate

- Irreversible processes.
- 2nd Law: $dS \geq \frac{\delta Q}{T}$,
= reversible; > irreversible.
- $dS = d_e S + d_i S$ (e, external; i, internal),

5.1 Local equilibrium, entropy flux density, entropy local production rate

- Irreversible processes.
- 2nd Law: $dS \geq \frac{\delta Q}{T}$,
= reversible; > irreversible.
- $dS = d_e S + d_i S$ (e, external; i, internal),
 $d_e S$: exchange of heat and matter; $d_i S$: isolated part.

5.1 Local equilibrium, entropy flux density, entropy local production rate

- Irreversible processes.
- 2nd Law: $dS \geq \frac{\delta Q}{T}$,
= reversible; > irreversible.
- $dS = d_e S + d_i S$ (e, external; i, internal),
 $d_e S$: exchange of heat and matter; $d_i S$: isolated part.
- Rewrite basic equation:
$$TdS = dU + pdV - \sum_i \mu_i dn_i,$$

5.1 Local equilibrium, entropy flux density, entropy local production rate

- Irreversible processes.
- 2nd Law: $dS \geq \frac{\delta Q}{T}$,
= reversible; > irreversible.
- $dS = d_e S + d_i S$ (e, external; i, internal),
 $d_e S$: exchange of heat and matter; $d_i S$: isolated part.
- Rewrite basic equation:
 $TdS = dU + pdV - \sum_i \mu_i dn_i$, to
 $TdS = dU + pdV - \sum_i \mu_i dN_i$, where N_i is the
number of molecules of component i , and μ_i is the
chemical potential of one molecule of component i .

Local equilibrium

- No global state parameter (like T, p) for non-equilibrium system.

Local equilibrium

- No global state parameter (like T, p) for non-equilibrium system.
- Exists small parts in the system, are approximately in equilibrium, which exist (T, p, μ, \dots) .

Local equilibrium

- No global state parameter (like T, p) for non-equilibrium system.
- Exists small parts in the system, are approximately in equilibrium, which exist (T, p, μ, \dots) .
- Each small part (in volume) is an open system,
$$TdS = dU + pdV - \sum_i \mu_i dN_i.$$

Local equilibrium

- No global state parameter (like T, p) for non-equilibrium system.
- Exists small parts in the system, are approximately in equilibrium, which exist (T, p, μ, \dots) .
- Each small part (in volume) is an open system,
$$TdS = dU + pdV - \sum_i \mu_i dN_i.$$
- 1. Neglect volume change $dV = 0$;

Local equilibrium

- No global state parameter (like T, p) for non-equilibrium system.
- Exists small parts in the system, are approximately in equilibrium, which exist (T, p, μ, \dots) .
- Each small part (in volume) is an open system,
$$TdS = dU + pdV - \sum_i \mu_i dN_i.$$
- 1. Neglect volume change $dV = 0$; 2. Divided by V .

Local equilibrium

- No global state parameter (like T, p) for non-equilibrium system.
- Exists small parts in the system, are approximately in equilibrium, which exist (T, p, μ, \dots) .
- Each small part (in volume) is an open system,
 $TdS = dU + pdV - \sum_i \mu_i dN_i$.
- 1. Neglect volume change $dV = 0$; 2. Divided by V .
- $Tds = du - \sum_i \mu_i dn_i$, where n_i is the number density of component i .

Local equilibrium

- No global state parameter (like T, p) for non-equilibrium system.
- Exists small parts in the system, are approximately in equilibrium, which exist (T, p, μ, \dots) .
- Each small part (in volume) is an open system,
 $TdS = dU + pdV - \sum_i \mu_i dN_i$.
- 1. Neglect volume change $dV = 0$; 2. Divided by V .
- $Tds = du - \sum_i \mu_i dn_i$, where n_i is the number density of component i .
- For the whole component, $U = \int u d\tau$, $S = \int s d\tau$,
 $N_i = \int n_i d\tau$;

Local equilibrium

- No global state parameter (like T, p) for non-equilibrium system.
- Exists small parts in the system, are approximately in equilibrium, which exist (T, p, μ, \dots) .
- Each small part (in volume) is an open system,
 $TdS = dU + pdV - \sum_i \mu_i dN_i$.
- 1. Neglect volume change $dV = 0$; 2. Divided by V .
- $Tds = du - \sum_i \mu_i dn_i$, where n_i is the number density of component i .
- For the whole component, $U = \int u d\tau$, $S = \int s d\tau$,
 $N_i = \int n_i d\tau$; S is a critical variable.

Entropy flux density and entropy local production rate

Entropy flux density and entropy local production rate

- Define $\frac{\partial s}{\partial t} = -\nabla \cdot \vec{J}_s + \Theta$, where \vec{J}_s is the entropy flux density (external), and Θ is the entropy local production rate (internal).

Entropy flux density and entropy local production rate

- Define $\frac{\partial s}{\partial t} = -\nabla \cdot \vec{J}_s + \Theta$, where \vec{J}_s is the entropy flux density (external), and Θ is the entropy local production rate (internal).
- For the whole system:

Entropy flux density and entropy local production rate

- Define $\frac{\partial s}{\partial t} = -\nabla \cdot \vec{J}_s + \Theta$, where \vec{J}_s is the entropy flux density (external), and Θ is the entropy local production rate (internal).
- For the whole system:
$$\frac{dS}{dt} = \frac{d}{dt} \int s d\tau$$

Entropy flux density and entropy local production rate

- Define $\frac{\partial s}{\partial t} = -\nabla \cdot \vec{J}_s + \Theta$, where \vec{J}_s is the entropy flux density (external), and Θ is the entropy local production rate (internal).
- For the whole system:
$$\frac{dS}{dt} = \frac{d}{dt} \int s d\tau = \int \frac{\partial s}{\partial t} d\tau$$

Entropy flux density and entropy local production rate

- Define $\frac{\partial s}{\partial t} = -\nabla \cdot \vec{J}_s + \Theta$, where \vec{J}_s is the entropy flux density (external), and Θ is the entropy local production rate (internal).
- For the whole system:
$$\frac{dS}{dt} = \frac{d}{dt} \int s d\tau = \int \frac{\partial s}{\partial t} d\tau = \int [-\nabla \cdot \vec{J}_s + \Theta] d\tau$$

Entropy flux density and entropy local production rate

- Define $\frac{\partial s}{\partial t} = -\nabla \cdot \vec{J}_s + \Theta$, where \vec{J}_s is the entropy flux density (external), and Θ is the entropy local production rate (internal).

- For the whole system:

$$\begin{aligned}\frac{dS}{dt} &= \frac{d}{dt} \int s d\tau = \int \frac{\partial s}{\partial t} d\tau = \int [-\nabla \cdot \vec{J}_s + \Theta] d\tau \\ &= -\oint \vec{J}_s \cdot d\vec{\sigma} + \int \Theta d\tau.\end{aligned}$$

Entropy flux density and entropy local production rate

- Define $\frac{\partial s}{\partial t} = -\nabla \cdot \vec{J}_s + \Theta$, where \vec{J}_s is the entropy flux density (external), and Θ is the entropy local production rate (internal).
- For the whole system:
$$\begin{aligned}\frac{dS}{dt} &= \frac{d}{dt} \int s d\tau = \int \frac{\partial s}{\partial t} d\tau = \int [-\nabla \cdot \vec{J}_s + \Theta] d\tau \\ &= -\oint \vec{J}_s \cdot d\vec{\sigma} + \int \Theta d\tau.\end{aligned}$$
- $\frac{d_e S}{dt} = -\oint \vec{J}_s \cdot d\vec{\sigma}$, $\frac{d_i S}{dt} = \int \Theta d\tau$.

Entropy flux density and entropy local production rate

- Define $\frac{\partial s}{\partial t} = -\nabla \cdot \vec{J}_s + \Theta$, where \vec{J}_s is the entropy flux density (external), and Θ is the entropy local production rate (internal).
- For the whole system:
$$\frac{dS}{dt} = \frac{d}{dt} \int s d\tau = \int \frac{\partial s}{\partial t} d\tau = \int [-\nabla \cdot \vec{J}_s + \Theta] d\tau$$
$$= -\oint \vec{J}_s \cdot d\vec{\sigma} + \int \Theta d\tau.$$
- $\frac{d_e S}{dt} = -\oint \vec{J}_s \cdot d\vec{\sigma}$, $\frac{d_i S}{dt} = \int \Theta d\tau$.
- Apply to the simplest case: only heat conduction,

Entropy flux density and entropy local production rate

- Define $\frac{\partial s}{\partial t} = -\nabla \cdot \vec{J}_s + \Theta$, where \vec{J}_s is the entropy flux density (external), and Θ is the entropy local production rate (internal).
- For the whole system:
$$\frac{dS}{dt} = \frac{d}{dt} \int s d\tau = \int \frac{\partial s}{\partial t} d\tau = \int [-\nabla \cdot \vec{J}_s + \Theta] d\tau$$
$$= -\oint \vec{J}_s \cdot d\vec{\sigma} + \int \Theta d\tau.$$
- $\frac{d_e S}{dt} = -\oint \vec{J}_s \cdot d\vec{\sigma}$, $\frac{d_i S}{dt} = \int \Theta d\tau$.
- Apply to the simplest case: only heat conduction,
- internal energy density: $\frac{\partial u}{\partial t} = -\nabla \cdot \vec{J}_q$, where \vec{J}_q is the heat flux density.

Entropy flux density and entropy local production rate

- Define $\frac{\partial s}{\partial t} = -\nabla \cdot \vec{J}_s + \Theta$, where \vec{J}_s is the entropy flux density (external), and Θ is the entropy local production rate (internal).
- For the whole system:
$$\frac{dS}{dt} = \frac{d}{dt} \int s d\tau = \int \frac{\partial s}{\partial t} d\tau = \int [-\nabla \cdot \vec{J}_s + \Theta] d\tau$$
$$= -\oint \vec{J}_s \cdot d\vec{\sigma} + \int \Theta d\tau.$$
- $\frac{d_e S}{dt} = -\oint \vec{J}_s \cdot d\vec{\sigma}$, $\frac{d_i S}{dt} = \int \Theta d\tau$.
- Apply to the simplest case: only heat conduction,
- internal energy density: $\frac{\partial u}{\partial t} = -\nabla \cdot \vec{J}_q$, where \vec{J}_q is the heat flux density.
- No volume change, no matter flow, the basic equation: $T ds = du$

Entropy flux density and entropy local production rate

- Define $\frac{\partial s}{\partial t} = -\nabla \cdot \vec{J}_s + \Theta$, where \vec{J}_s is the entropy flux density (external), and Θ is the entropy local production rate (internal).

- For the whole system:

$$\begin{aligned}\frac{dS}{dt} &= \frac{d}{dt} \int s d\tau = \int \frac{\partial s}{\partial t} d\tau = \int [-\nabla \cdot \vec{J}_s + \Theta] d\tau \\ &= -\oint \vec{J}_s \cdot d\vec{\sigma} + \int \Theta d\tau.\end{aligned}$$

- $\frac{d_e S}{dt} = -\oint \vec{J}_s \cdot d\vec{\sigma}$, $\frac{d_i S}{dt} = \int \Theta d\tau$.

- Apply to the simplest case: only heat conduction,

- internal energy density: $\frac{\partial u}{\partial t} = -\nabla \cdot \vec{J}_q$, where \vec{J}_q is the heat flux density.

- No volume change, no matter flow, the basic equation: $T ds = du \Rightarrow \frac{\partial s}{\partial t} = \frac{1}{T} \frac{\partial u}{\partial t}$

Entropy flux density and entropy local production rate

- Define $\frac{\partial s}{\partial t} = -\nabla \cdot \vec{J}_s + \Theta$, where \vec{J}_s is the entropy flux density (external), and Θ is the entropy local production rate (internal).
- For the whole system:

$$\frac{dS}{dt} = \frac{d}{dt} \int s d\tau = \int \frac{\partial s}{\partial t} d\tau = \int [-\nabla \cdot \vec{J}_s + \Theta] d\tau$$

$$= -\oint \vec{J}_s \cdot d\vec{\sigma} + \int \Theta d\tau.$$
- $\frac{d_e S}{dt} = -\oint \vec{J}_s \cdot d\vec{\sigma}$, $\frac{d_i S}{dt} = \int \Theta d\tau$.
- Apply to the simplest case: only heat conduction,
- internal energy density: $\frac{\partial u}{\partial t} = -\nabla \cdot \vec{J}_q$, where \vec{J}_q is the heat flux density.
- No volume change, no matter flow, the basic equation: $T ds = du \Rightarrow \frac{\partial s}{\partial t} = \frac{1}{T} \frac{\partial u}{\partial t} = -\frac{1}{T} \nabla \cdot \vec{J}_q$.

Entropy flux density and entropy local production rate

- $$\frac{\partial s}{\partial t} = -\frac{1}{T} \nabla \cdot \vec{J}_q$$

Entropy flux density and entropy local production rate

- $$\frac{\partial s}{\partial t} = -\frac{1}{T} \nabla \cdot \vec{J}_q = -\left[\nabla \cdot \frac{\vec{J}_q}{T} - \vec{J}_q \cdot \nabla \frac{1}{T} \right]$$

Entropy flux density and entropy local production rate

- $$\begin{aligned}\frac{\partial s}{\partial t} &= -\frac{1}{T} \nabla \cdot \vec{J}_q = -\left[\nabla \cdot \frac{\vec{J}_q}{T} - \vec{J}_q \cdot \nabla \frac{1}{T}\right] \\ &= -\nabla \cdot \frac{\vec{J}_q}{T} + \vec{J}_q \cdot \nabla \frac{1}{T}.\end{aligned}$$

Entropy flux density and entropy local production rate

- $$\frac{\partial s}{\partial t} = -\frac{1}{T} \nabla \cdot \vec{J}_q = -\left[\nabla \cdot \frac{\vec{J}_q}{T} - \vec{J}_q \cdot \nabla \frac{1}{T}\right]$$
$$= -\nabla \cdot \frac{\vec{J}_q}{T} + \vec{J}_q \cdot \nabla \frac{1}{T}.$$
- Compare with $\frac{\partial s}{\partial t} = -\nabla \cdot \vec{J}_s + \Theta,$

Entropy flux density and entropy local production rate

- $$\frac{\partial s}{\partial t} = -\frac{1}{T} \nabla \cdot \vec{J}_q = -\left[\nabla \cdot \frac{\vec{J}_q}{T} - \vec{J}_q \cdot \nabla \frac{1}{T}\right]$$
$$= -\nabla \cdot \frac{\vec{J}_q}{T} + \vec{J}_q \cdot \nabla \frac{1}{T}.$$
- Compare with $\frac{\partial s}{\partial t} = -\nabla \cdot \vec{J}_s + \Theta,$
 $\vec{J}_s = \frac{\vec{J}_q}{T}, \Theta = \vec{J}_q \cdot \nabla \frac{1}{T}.$

Entropy flux density and entropy local production rate

- $\frac{\partial s}{\partial t} = -\frac{1}{T} \nabla \cdot \vec{J}_q = -[\nabla \cdot \frac{\vec{J}_q}{T} - \vec{J}_q \cdot \nabla \frac{1}{T}]$
 $= -\nabla \cdot \frac{\vec{J}_q}{T} + \vec{J}_q \cdot \nabla \frac{1}{T}.$
- Compare with $\frac{\partial s}{\partial t} = -\nabla \cdot \vec{J}_s + \Theta,$
 $\vec{J}_s = \frac{\vec{J}_q}{T}, \Theta = \vec{J}_q \cdot \nabla \frac{1}{T}.$
- Define $\vec{X}_q \equiv \nabla \frac{1}{T},$ as heat flow force,

Entropy flux density and entropy local production rate

- $\frac{\partial s}{\partial t} = -\frac{1}{T} \nabla \cdot \vec{J}_q = -[\nabla \cdot \frac{\vec{J}_q}{T} - \vec{J}_q \cdot \nabla \frac{1}{T}]$
 $= -\nabla \cdot \frac{\vec{J}_q}{T} + \vec{J}_q \cdot \nabla \frac{1}{T}.$
- Compare with $\frac{\partial s}{\partial t} = -\nabla \cdot \vec{J}_s + \Theta,$
 $\vec{J}_s = \frac{\vec{J}_q}{T}, \Theta = \vec{J}_q \cdot \nabla \frac{1}{T}.$
- Define $\vec{X}_q \equiv \nabla \frac{1}{T},$ as heat flow force, then
 $\Theta = \vec{J}_q \cdot \vec{X}_q.$

Entropy flux density and entropy local production rate

- $$\frac{\partial s}{\partial t} = -\frac{1}{T} \nabla \cdot \vec{J}_q = -[\nabla \cdot \frac{\vec{J}_q}{T} - \vec{J}_q \cdot \nabla \frac{1}{T}]$$
$$= -\nabla \cdot \frac{\vec{J}_q}{T} + \vec{J}_q \cdot \nabla \frac{1}{T}.$$
- Compare with $\frac{\partial s}{\partial t} = -\nabla \cdot \vec{J}_s + \Theta$,
 $\vec{J}_s = \frac{\vec{J}_q}{T}$, $\Theta = \vec{J}_q \cdot \nabla \frac{1}{T}$.
- Define $\vec{X}_q \equiv \nabla \frac{1}{T}$, as heat flow force, then
 $\Theta = \vec{J}_q \cdot \vec{X}_q$.
- If heat conduction obeys the Fourier's law,
 $\vec{J}_q = -\kappa \nabla T$, then

Entropy flux density and entropy local production rate

- $\frac{\partial s}{\partial t} = -\frac{1}{T} \nabla \cdot \vec{J}_q = -[\nabla \cdot \frac{\vec{J}_q}{T} - \vec{J}_q \cdot \nabla \frac{1}{T}]$
 $= -\nabla \cdot \frac{\vec{J}_q}{T} + \vec{J}_q \cdot \nabla \frac{1}{T}.$
- Compare with $\frac{\partial s}{\partial t} = -\nabla \cdot \vec{J}_s + \Theta,$
 $\vec{J}_s = \frac{\vec{J}_q}{T}, \Theta = \vec{J}_q \cdot \nabla \frac{1}{T}.$
- Define $\vec{X}_q \equiv \nabla \frac{1}{T},$ as heat flow force, then
 $\Theta = \vec{J}_q \cdot \vec{X}_q.$
- If heat conduction obeys the Fourier's law,
 $\vec{J}_q = -\kappa \nabla T,$ then $\Theta = \vec{J}_q \cdot \nabla \frac{1}{T}$

Entropy flux density and entropy local production rate

- $\frac{\partial s}{\partial t} = -\frac{1}{T} \nabla \cdot \vec{J}_q = -[\nabla \cdot \frac{\vec{J}_q}{T} - \vec{J}_q \cdot \nabla \frac{1}{T}]$
 $= -\nabla \cdot \frac{\vec{J}_q}{T} + \vec{J}_q \cdot \nabla \frac{1}{T}.$
- Compare with $\frac{\partial s}{\partial t} = -\nabla \cdot \vec{J}_s + \Theta,$
 $\vec{J}_s = \frac{\vec{J}_q}{T}, \Theta = \vec{J}_q \cdot \nabla \frac{1}{T}.$
- Define $\vec{X}_q \equiv \nabla \frac{1}{T},$ as heat flow force, then
 $\Theta = \vec{J}_q \cdot \vec{X}_q.$
- If heat conduction obeys the Fourier's law,
 $\vec{J}_q = -\kappa \nabla T,$ then $\Theta = \vec{J}_q \cdot \nabla \frac{1}{T} = -\vec{J}_q \cdot \frac{\nabla T}{T^2}$

Entropy flux density and entropy local production rate

- $\frac{\partial s}{\partial t} = -\frac{1}{T} \nabla \cdot \vec{J}_q = -[\nabla \cdot \frac{\vec{J}_q}{T} - \vec{J}_q \cdot \nabla \frac{1}{T}]$
 $= -\nabla \cdot \frac{\vec{J}_q}{T} + \vec{J}_q \cdot \nabla \frac{1}{T}.$
- Compare with $\frac{\partial s}{\partial t} = -\nabla \cdot \vec{J}_s + \Theta,$
 $\vec{J}_s = \frac{\vec{J}_q}{T}, \Theta = \vec{J}_q \cdot \nabla \frac{1}{T}.$
- Define $\vec{X}_q \equiv \nabla \frac{1}{T},$ as heat flow force, then
 $\Theta = \vec{J}_q \cdot \vec{X}_q.$
- If heat conduction obeys the Fourier's law,
 $\vec{J}_q = -\kappa \nabla T,$ then $\Theta = \vec{J}_q \cdot \nabla \frac{1}{T} = -\vec{J}_q \cdot \frac{\nabla T}{T^2}$
 $= \kappa \frac{(\nabla T)^2}{T^2}.$

Entropy flux density and entropy local production rate

- $$\frac{\partial s}{\partial t} = -\frac{1}{T} \nabla \cdot \vec{J}_q = -[\nabla \cdot \frac{\vec{J}_q}{T} - \vec{J}_q \cdot \nabla \frac{1}{T}]$$
$$= -\nabla \cdot \frac{\vec{J}_q}{T} + \vec{J}_q \cdot \nabla \frac{1}{T}.$$
- Compare with $\frac{\partial s}{\partial t} = -\nabla \cdot \vec{J}_s + \Theta$,
$$\vec{J}_s = \frac{\vec{J}_q}{T}, \Theta = \vec{J}_q \cdot \nabla \frac{1}{T}.$$
- Define $\vec{X}_q \equiv \nabla \frac{1}{T}$, as heat flow force, then
$$\Theta = \vec{J}_q \cdot \vec{X}_q.$$
- If heat conduction obeys the Fourier's law,
$$\vec{J}_q = -\kappa \nabla T, \text{ then } \Theta = \vec{J}_q \cdot \nabla \frac{1}{T} = -\vec{J}_q \cdot \frac{\nabla T}{T^2}$$
$$= \kappa \frac{(\nabla T)^2}{T^2}.$$
- $\Theta \geq 0.$

Entropy flux density and entropy local production rate

- Apply to a more realistic case: temperature and chemical potential, non-uniform

Entropy flux density and entropy local production rate

- Apply to a more realistic case: temperature and chemical potential, non-uniform
— heat conduction and matter flow.

Entropy flux density and entropy local production rate

- Apply to a more realistic case: temperature and chemical potential, non-uniform
— heat conduction and matter flow.
- Energy conservation: $\frac{\partial u}{\partial t} + \nabla \cdot \vec{J}_u = 0$,

Entropy flux density and entropy local production rate

- Apply to a more realistic case: temperature and chemical potential, non-uniform
— heat conduction and matter flow.
- Energy conservation: $\frac{\partial u}{\partial t} + \nabla \cdot \vec{J}_u = 0$,
matter conservation: $\frac{\partial n}{\partial t} + \nabla \cdot \vec{J}_n = 0$.

Entropy flux density and entropy local production rate

- Apply to a more realistic case: temperature and chemical potential, non-uniform
— heat conduction and matter flow.
- Energy conservation: $\frac{\partial u}{\partial t} + \nabla \cdot \vec{J}_u = 0$,
matter conservation: $\frac{\partial n}{\partial t} + \nabla \cdot \vec{J}_n = 0$.
- $\vec{J}_u = \vec{J}_q + \mu \vec{J}_n$.

Entropy flux density and entropy local production rate

- Apply to a more realistic case: temperature and chemical potential, non-uniform
— heat conduction and matter flow.
- Energy conservation: $\frac{\partial u}{\partial t} + \nabla \cdot \vec{J}_u = 0$,
matter conservation: $\frac{\partial n}{\partial t} + \nabla \cdot \vec{J}_n = 0$.
- $\vec{J}_u = \vec{J}_q + \mu \vec{J}_n$.
- $\therefore \frac{\partial u}{\partial t} = -\nabla \cdot \vec{J}_q - \nabla \cdot (\mu \vec{J}_n)$.

Entropy flux density and entropy local production rate

- Apply to a more realistic case: temperature and chemical potential, non-uniform
— heat conduction and matter flow.
- Energy conservation: $\frac{\partial u}{\partial t} + \nabla \cdot \vec{J}_u = 0$,
matter conservation: $\frac{\partial n}{\partial t} + \nabla \cdot \vec{J}_n = 0$.
- $\vec{J}_u = \vec{J}_q + \mu \vec{J}_n$.
- $\therefore \frac{\partial u}{\partial t} = -\nabla \cdot \vec{J}_q - \nabla \cdot (\mu \vec{J}_n)$.
- $ds = \frac{1}{T}(du - \mu dn)$,

Entropy flux density and entropy local production rate

- Apply to a more realistic case: temperature and chemical potential, non-uniform
— heat conduction and matter flow.
- Energy conservation: $\frac{\partial u}{\partial t} + \nabla \cdot \vec{J}_u = 0$,
matter conservation: $\frac{\partial n}{\partial t} + \nabla \cdot \vec{J}_n = 0$.
- $\vec{J}_u = \vec{J}_q + \mu \vec{J}_n$.
- $\therefore \frac{\partial u}{\partial t} = -\nabla \cdot \vec{J}_q - \nabla \cdot (\mu \vec{J}_n)$.
- $ds = \frac{1}{T}(du - \mu dn)$,
 $\Rightarrow \frac{\partial s}{\partial t} = \frac{1}{T} \frac{\partial u}{\partial t} - \frac{\mu}{T} \frac{\partial n}{\partial t}$

Entropy flux density and entropy local production rate

- Apply to a more realistic case: temperature and chemical potential, non-uniform
— heat conduction and matter flow.
- Energy conservation: $\frac{\partial u}{\partial t} + \nabla \cdot \vec{J}_u = 0$,
matter conservation: $\frac{\partial n}{\partial t} + \nabla \cdot \vec{J}_n = 0$.
- $\vec{J}_u = \vec{J}_q + \mu \vec{J}_n$.
- $\therefore \frac{\partial u}{\partial t} = -\nabla \cdot \vec{J}_q - \nabla \cdot (\mu \vec{J}_n)$.
- $ds = \frac{1}{T}(du - \mu dn)$,

$$\Rightarrow \frac{\partial s}{\partial t} = \frac{1}{T} \frac{\partial u}{\partial t} - \frac{\mu}{T} \frac{\partial n}{\partial t}$$

$$= -\frac{1}{T} \nabla \cdot \vec{J}_q - \frac{1}{T} \nabla \cdot (\mu \vec{J}_n) + \frac{\mu}{T} \nabla \cdot \vec{J}_n$$

Entropy flux density and entropy local production rate

- Apply to a more realistic case: temperature and chemical potential, non-uniform
— heat conduction and matter flow.
- Energy conservation: $\frac{\partial u}{\partial t} + \nabla \cdot \vec{J}_u = 0$,
matter conservation: $\frac{\partial n}{\partial t} + \nabla \cdot \vec{J}_n = 0$.
- $\vec{J}_u = \vec{J}_q + \mu \vec{J}_n$.
- $\therefore \frac{\partial u}{\partial t} = -\nabla \cdot \vec{J}_q - \nabla \cdot (\mu \vec{J}_n)$.
- $ds = \frac{1}{T}(du - \mu dn)$,

$$\Rightarrow \frac{\partial s}{\partial t} = \frac{1}{T} \frac{\partial u}{\partial t} - \frac{\mu}{T} \frac{\partial n}{\partial t}$$

$$= -\frac{1}{T} \nabla \cdot \vec{J}_q - \frac{1}{T} \nabla \cdot (\mu \vec{J}_n) + \frac{\mu}{T} \nabla \cdot \vec{J}_n$$

$$= -\nabla \cdot \frac{\vec{J}_q}{T} + \vec{J}_q \cdot \nabla \frac{1}{T} - \frac{\vec{J}_n}{T} \cdot \nabla \mu.$$

Entropy flux density and entropy local production rate

- $$\frac{\partial s}{\partial t} = -\nabla \cdot \frac{\vec{J}_q}{T} + \vec{J}_q \cdot \nabla \frac{1}{T} - \frac{\vec{J}_n}{T} \cdot \nabla \mu,$$
$$\frac{\partial s}{\partial t} = -\nabla \cdot \vec{J}_s + \Theta.$$

Entropy flux density and entropy local production rate

- $$\begin{aligned}\frac{\partial s}{\partial t} &= -\nabla \cdot \frac{\vec{J}_q}{T} + \vec{J}_q \cdot \nabla \frac{1}{T} - \frac{\vec{J}_n}{T} \cdot \nabla \mu, \\ \frac{\partial s}{\partial t} &= -\nabla \cdot \vec{J}_s + \Theta. \\ \Rightarrow \vec{J}_s &= \frac{\vec{J}_q}{T}, \quad \Theta = \vec{J}_q \cdot \nabla \frac{1}{T} - \vec{J}_n \cdot \frac{\nabla \mu}{T}.\end{aligned}$$

Entropy flux density and entropy local production rate

- $$\frac{\partial s}{\partial t} = -\nabla \cdot \frac{\vec{J}_q}{T} + \vec{J}_q \cdot \nabla \frac{1}{T} - \frac{\vec{J}_n}{T} \cdot \nabla \mu,$$
$$\frac{\partial s}{\partial t} = -\nabla \cdot \vec{J}_s + \Theta.$$
$$\Rightarrow \vec{J}_s = \frac{\vec{J}_q}{T}, \quad \Theta = \vec{J}_q \cdot \nabla \frac{1}{T} - \vec{J}_n \cdot \frac{\nabla \mu}{T}.$$
- Define matter flow “force” $\vec{X}_n = -\frac{1}{T} \nabla \mu$, and heat flow “force” $\vec{X}_q = \nabla \frac{1}{T}$,

Entropy flux density and entropy local production rate

- $$\frac{\partial s}{\partial t} = -\nabla \cdot \frac{\vec{J}_q}{T} + \vec{J}_q \cdot \nabla \frac{1}{T} - \frac{\vec{J}_n}{T} \cdot \nabla \mu,$$
$$\frac{\partial s}{\partial t} = -\nabla \cdot \vec{J}_s + \Theta.$$

$$\Rightarrow \vec{J}_s = \frac{\vec{J}_q}{T}, \quad \Theta = \vec{J}_q \cdot \nabla \frac{1}{T} - \vec{J}_n \cdot \frac{\nabla \mu}{T}.$$

- Define matter flow “force” $\vec{X}_n = -\frac{1}{T} \nabla \mu$, and heat flow “force” $\vec{X}_q = \nabla \frac{1}{T}$,
 $\therefore \Theta = \vec{J}_q \cdot \vec{X}_q + \vec{J}_n \cdot \vec{X}_n.$

Entropy flux density and entropy local production rate

- $$\frac{\partial s}{\partial t} = -\nabla \cdot \frac{\vec{J}_q}{T} + \vec{J}_q \cdot \nabla \frac{1}{T} - \frac{\vec{J}_n}{T} \cdot \nabla \mu,$$

$$\frac{\partial s}{\partial t} = -\nabla \cdot \vec{J}_s + \Theta.$$

$$\Rightarrow \vec{J}_s = \frac{\vec{J}_q}{T}, \quad \Theta = \vec{J}_q \cdot \nabla \frac{1}{T} - \vec{J}_n \cdot \frac{\nabla \mu}{T}.$$
- Define matter flow “force” $\vec{X}_n = -\frac{1}{T} \nabla \mu$, and heat flow “force” $\vec{X}_q = \nabla \frac{1}{T}$,
 $\therefore \Theta = \vec{J}_q \cdot \vec{X}_q + \vec{J}_n \cdot \vec{X}_n.$
- Generally, $\Theta = \sum_k \vec{J}_k \cdot \vec{X}_k.$

Table of contents

- ① §5. Thermodynamics of irreversible processes
 - 5.1 Local equilibrium, entropy flux density and production rate
 - 5.2 Linear and nonlinear processes, Onsager relation

5.2 Linear and nonlinear processes, Onsager relation

5.2 Linear and nonlinear processes, Onsager relation

- Non-uniform \rightarrow transport:

5.2 Linear and nonlinear processes, Onsager relation

- Non-uniform \rightarrow transport:
 1. Temperature

5.2 Linear and nonlinear processes, Onsager relation

- Non-uniform \rightarrow transport:
 1. Temperature \rightarrow heat flow,

5.2 Linear and nonlinear processes, Onsager relation

- Non-uniform \rightarrow transport:
 1. Temperature \rightarrow heat flow,
 $\vec{J}_q = -\kappa \nabla T$, Fourier's law.

5.2 Linear and nonlinear processes, Onsager relation

- Non-uniform \rightarrow transport:
 1. Temperature \rightarrow heat flow,
 $\vec{J}_q = -\kappa \nabla T$, Fourier's law.
 2. Density

5.2 Linear and nonlinear processes, Onsager relation

- Non-uniform \rightarrow transport:
 1. Temperature \rightarrow heat flow,
 $\vec{J}_q = -\kappa \nabla T$, Fourier's law.
 2. Density \rightarrow diffusion (matter flow),

5.2 Linear and nonlinear processes, Onsager relation

- Non-uniform \rightarrow transport:

1. Temperature \rightarrow heat flow,

$$\vec{J}_q = -\kappa \nabla T, \text{ Fourier's law.}$$

2. Density \rightarrow diffusion (matter flow),

$$\vec{J}_n = -D \nabla n, \text{ Fick's law.}$$

5.2 Linear and nonlinear processes, Onsager relation

- Non-uniform \rightarrow transport:

1. Temperature \rightarrow heat flow,

$$\vec{J}_q = -\kappa \nabla T, \text{ Fourier's law.}$$

2. Density \rightarrow diffusion (matter flow),

$$\vec{J}_n = -D \nabla n, \text{ Fick's law.}$$

3. Electric potential

5.2 Linear and nonlinear processes, Onsager relation

- Non-uniform \rightarrow transport:
 1. Temperature \rightarrow heat flow,
 $\vec{J}_q = -\kappa \nabla T$, Fourier's law.
 2. Density \rightarrow diffusion (matter flow),
 $\vec{J}_n = -D \nabla n$, Fick's law.
 3. Electric potential \rightarrow charge transport,

5.2 Linear and nonlinear processes, Onsager relation

- Non-uniform \rightarrow transport:

1. Temperature \rightarrow heat flow,

$$\vec{J}_q = -\kappa \nabla T, \text{ Fourier's law.}$$

2. Density \rightarrow diffusion (matter flow),

$$\vec{J}_n = -D \nabla n, \text{ Fick's law.}$$

3. Electric potential \rightarrow charge transport,

$$\vec{J}_e = \sigma \vec{E} = -\sigma \nabla V.$$

5.2 Linear and nonlinear processes, Onsager relation

- Non-uniform \rightarrow transport:

1. Temperature \rightarrow heat flow,

$$\vec{J}_q = -\kappa \nabla T, \text{ Fourier's law.}$$

2. Density \rightarrow diffusion (matter flow),

$$\vec{J}_n = -D \nabla n, \text{ Fick's law.}$$

3. Electric potential \rightarrow charge transport,

$$\vec{J}_e = \sigma \vec{E} = -\sigma \nabla V.$$

4. Velocity

5.2 Linear and nonlinear processes, Onsager relation

- Non-uniform \rightarrow transport:

1. Temperature \rightarrow heat flow,

$$\vec{J}_q = -\kappa \nabla T, \text{ Fourier's law.}$$

2. Density \rightarrow diffusion (matter flow),

$$\vec{J}_n = -D \nabla n, \text{ Fick's law.}$$

3. Electric potential \rightarrow charge transport,

$$\vec{J}_e = \sigma \vec{E} = -\sigma \nabla V.$$

4. Velocity \rightarrow viscosity (momentum transport),

5.2 Linear and nonlinear processes, Onsager relation

- Non-uniform \rightarrow transport:

1. Temperature \rightarrow heat flow,

$$\vec{J}_q = -\kappa \nabla T, \text{ Fourier's law.}$$

2. Density \rightarrow diffusion (matter flow),

$$\vec{J}_n = -D \nabla n, \text{ Fick's law.}$$

3. Electric potential \rightarrow charge transport,

$$\vec{J}_e = \sigma \vec{E} = -\sigma \nabla V.$$

4. Velocity \rightarrow viscosity (momentum transport),

$$P_{xy} = \eta \frac{dv}{dx}.$$

5.2 Linear and nonlinear processes, Onsager relation

- Non-uniform \rightarrow transport:

1. Temperature \rightarrow heat flow,

$$\vec{J}_q = -\kappa \nabla T, \text{ Fourier's law.}$$

2. Density \rightarrow diffusion (matter flow),

$$\vec{J}_n = -D \nabla n, \text{ Fick's law.}$$

3. Electric potential \rightarrow charge transport,

$$\vec{J}_e = \sigma \vec{E} = -\sigma \nabla V.$$

4. Velocity \rightarrow viscosity (momentum transport),

$$P_{xy} = \eta \frac{dv}{dx}.$$

- Summation:

5.2 Linear and nonlinear processes, Onsager relation

- Non-uniform \rightarrow transport:

1. Temperature \rightarrow heat flow,

$$\vec{J}_q = -\kappa \nabla T, \text{ Fourier's law.}$$

2. Density \rightarrow diffusion (matter flow),

$$\vec{J}_n = -D \nabla n, \text{ Fick's law.}$$

3. Electric potential \rightarrow charge transport,

$$\vec{J}_e = \sigma \vec{E} = -\sigma \nabla V.$$

4. Velocity \rightarrow viscosity (momentum transport),

$$P_{xy} = \eta \frac{dv}{dx}.$$

- Summation:

$$\vec{J} = L \vec{X}, \text{ where } \vec{J} \text{ is the flux density, } \vec{X} \text{ is the "force".}$$

Linear processes, Onsager relation

- For coexistence of multi-factors: (like T, ρ)

Linear processes, Onsager relation

- For coexistence of multi-factors: (like T, ρ)

$$J_k = \sum_l L_{kl} X_l.$$

Linear processes, Onsager relation

- For coexistence of multi-factors: (like T, ρ)
$$J_k = \sum_l L_{kl} X_l.$$
- Onsager relation: $L_{kl} = L_{lk}.$

Linear processes, Onsager relation

- For coexistence of multi-factors: (like T, ρ)
$$J_k = \sum_l L_{kl} X_l.$$
- Onsager relation: $L_{kl} = L_{lk}.$
- Entropy production $\Theta = \sum_k J_k X_k$

Linear processes, Onsager relation

- For coexistence of multi-factors: (like T, ρ)
$$J_k = \sum_l L_{kl} X_l.$$
- Onsager relation: $L_{kl} = L_{lk}.$
- Entropy production $\Theta = \sum_k J_k X_k = \sum_{kl} L_{kl} X_k X_l.$

Linear processes, Onsager relation

- For coexistence of multi-factors: (like T, ρ)
$$J_k = \sum_l L_{kl} X_l.$$
- Onsager relation: $L_{kl} = L_{lk}.$
- Entropy production $\Theta = \sum_k J_k X_k = \sum_{kl} L_{kl} X_k X_l.$
- Consider two “forces”,
$$\Theta = L_{11} X_1^2 + (L_{12} + L_{21}) X_1 X_2 + L_{22} X_2^2$$

Linear processes, Onsager relation

- For coexistence of multi-factors: (like T, ρ)
$$J_k = \sum_l L_{kl} X_l.$$
- Onsager relation: $L_{kl} = L_{lk}.$
- Entropy production $\Theta = \sum_k J_k X_k = \sum_{kl} L_{kl} X_k X_l.$
- Consider two “forces”,
$$\begin{aligned}\Theta &= L_{11}X_1^2 + (L_{12} + L_{21})X_1X_2 + L_{22}X_2^2 \\ &= L_{11}X_1^2 + 2L_{12}X_1X_2 + L_{22}X_2^2.\end{aligned}$$

Linear processes, Onsager relation

- For coexistence of multi-factors: (like T, ρ)
$$J_k = \sum_l L_{kl} X_l.$$
- Onsager relation: $L_{kl} = L_{lk}.$
- Entropy production $\Theta = \sum_k J_k X_k = \sum_{kl} L_{kl} X_k X_l.$
- Consider two “forces”,
$$\begin{aligned}\Theta &= L_{11} X_1^2 + (L_{12} + L_{21}) X_1 X_2 + L_{22} X_2^2 \\ &= L_{11} X_1^2 + 2L_{12} X_1 X_2 + L_{22} X_2^2.\end{aligned}$$
- $\Theta \geq 0$, requires, $L_{11} > 0$, $L_{11} L_{22} > L_{12}^2.$
(Properties of the coefficients L .)

Non-linear processes

Non-linear processes

- Linear processes can only describe the system near to the equilibrium state.

Non-linear processes

- Linear processes can only describe the system near to the equilibrium state.
- For the general case, $J_k = \sum_l L_{kl} X_l$ does not hold.

Non-linear processes

- Linear processes can only describe the system near to the equilibrium state.
- For the general case, $J_k = \sum_l L_{kl} X_l$ does not hold.
- $J_k(\{X_l\}) \equiv J_k(X_1, \dots, X_l, \dots)$, expand:

Non-linear processes

- Linear processes can only describe the system near to the equilibrium state.
- For the general case, $J_k = \sum_l L_{kl} X_l$ does not hold.
- $J_k(\{X_l\}) \equiv J_k(X_1, \dots, X_l, \dots)$, expand:
$$J_k(\{X_l\}) = J_k(0) + \sum_l \left(\frac{\partial J_k}{\partial X_l} \right)_0 X_l + \frac{1}{2} \sum_{l,n} \left(\frac{\partial^2 J_k}{\partial X_l \partial X_n} \right)_0 X_l X_n + \dots$$

Non-linear processes

- Linear processes can only describe the system near to the equilibrium state.
- For the general case, $J_k = \sum_l L_{kl} X_l$ does not hold.
- $J_k(\{X_l\}) \equiv J_k(X_1, \dots, X_l, \dots)$, expand:

$$J_k(\{X_l\}) = J_k(0) + \sum_l \left(\frac{\partial J_k}{\partial X_l} \right)_0 X_l + \frac{1}{2} \sum_{l,n} \left(\frac{\partial^2 J_k}{\partial X_l \partial X_n} \right)_0 X_l X_n + \dots$$
- $J_k(0)$ corresponds to $\{X_l\} = 0$, $J_k(0) = 0$.

Non-linear processes

- Linear processes can only describe the system near to the equilibrium state.
- For the general case, $J_k = \sum_l L_{kl} X_l$ does not hold.
- $J_k(\{X_l\}) \equiv J_k(X_1, \dots, X_l, \dots)$, expand:

$$J_k(\{X_l\}) = J_k(0) + \sum_l \left(\frac{\partial J_k}{\partial X_l} \right)_0 X_l + \frac{1}{2} \sum_{l,n} \left(\frac{\partial^2 J_k}{\partial X_l \partial X_n} \right)_0 X_l X_n + \dots$$
- $J_k(0)$ corresponds to $\{X_l\} = 0$, $J_k(0) = 0$.
 Define $L_{kl} \equiv \left(\frac{\partial J_k}{\partial X_l} \right)_0$, $L_{kln} \equiv \left(\frac{\partial^2 J_k}{\partial X_l \partial X_n} \right)_0$,

Non-linear processes

- Linear processes can only describe the system near to the equilibrium state.

- For the general case, $J_k = \sum_l L_{kl} X_l$ does not hold.

- $J_k(\{X_l\}) \equiv J_k(X_1, \dots, X_l, \dots)$, expand:

$$J_k(\{X_l\}) = J_k(0) + \sum_l \left(\frac{\partial J_k}{\partial X_l} \right)_0 X_l + \frac{1}{2} \sum_{l,n} \left(\frac{\partial^2 J_k}{\partial X_l \partial X_n} \right)_0 X_l X_n + \dots$$

- $J_k(0)$ corresponds to $\{X_l\} = 0$, $J_k(0) = 0$.

$$\text{Define } L_{kl} \equiv \left(\frac{\partial J_k}{\partial X_l} \right)_0, \quad L_{kln} \equiv \left(\frac{\partial^2 J_k}{\partial X_l \partial X_n} \right)_0,$$

$$J_k = \sum_l L_{kl} X_l + \frac{1}{2} \sum_{l,n} L_{kln} X_l X_n + \dots$$

Table of contents

- ① §5. Thermodynamics of irreversible processes
 - 5.1 Local equilibrium, entropy flux density and production rate
 - 5.2 Linear and nonlinear processes, Onsager relation