

# Thermodynamics & Statistical Physics

## Chapter 1. Basic Laws in Thermodynamics

Yuan-Chuan Zou  
zouyc@hust.edu.cn

School of Physics, Huazhong University of Science and Technology

December 30, 2013

## 1 Chpt 1. Basic Laws in Thermodynamics

### • 1.1 Thermal equilibrium state

- 1.2 Law of thermal equilibrium and the temperature
- 1.3 Equation of state
- 1.4 Work
- 1.5 First law of thermodynamics
- 1.6 Thermal capacity and enthalpy
- 1.7 Internal energy of the ideal gas
- 1.8 Adiabatic process of the ideal gas

- 1.9 Carnot cycle in the ideal gas
- 1.10 Second law of thermodynamics
- 1.11 Carnot's theorem
- 1.12 Thermodynamic temperature scale
- 1.13 Clausius theorem
- 1.14 Entropy
- 1.15 Entropy of ideal gas
- 1.16 Math expression of the 2nd law
- 1.17 Applications for the entropy increasing principle
- 1.18 Free energy and Gibbs function

# §1.1 Thermal equilibrium state

## §1.1 Thermal equilibrium state

- Objects: Macroscopic system constructed by microscopic particles.

## §1.1 Thermal equilibrium state

- Objects: Macroscopic system constructed by microscopic particles.
- Methods: thermodynamics (macro  $\rightarrow$  macro); statistical physics (micro  $\rightarrow$  macro).

## §1.1 Thermal equilibrium state

- Objects: Macroscopic system constructed by microscopic particles.
- Methods: thermodynamics (macro  $\rightarrow$  macro); statistical physics (micro  $\rightarrow$  macro).
- Systems: isolated, closed, open (particles and energy).

## §1.1 Thermal equilibrium state

- Objects: Macroscopic system constructed by microscopic particles.
- Methods: thermodynamics (macro  $\rightarrow$  macro); statistical physics (micro  $\rightarrow$  macro).
- Systems: isolated, closed, open (particles and energy).
- Thermal-equilibrium state.

## §1.1 Thermal equilibrium state

- Objects: Macroscopic system constructed by microscopic particles.
- Methods: thermodynamics (macro  $\rightarrow$  macro); statistical physics (micro  $\rightarrow$  macro).
- Systems: isolated, closed, open (particles and energy).
- Thermal-equilibrium state.
- To describe: state parameter (SP, state variables, or thermodynamic variables), function of state (FOS).



## §1.1 Thermal equilibrium state

- Objects: Macroscopic system constructed by microscopic particles.
- Methods: thermodynamics (macro  $\rightarrow$  macro); statistical physics (micro  $\rightarrow$  macro).
- Systems: isolated, closed, open (particles and energy).
- Thermal-equilibrium state.
- To describe: state parameter (SP, state variables, or thermodynamic variables), function of state (FOS).
- SP: geometry ( $V$ ), mechanics ( $p$ ), chemistry ( $n_i$ ), electromagnetics ( $E, P, H, M$ ).

## §1.1 Thermal equilibrium state

- Objects: Macroscopic system constructed by microscopic particles.
- Methods: thermodynamics (macro  $\rightarrow$  macro); statistical physics (micro  $\rightarrow$  macro).
- Systems: isolated, closed, open (particles and energy).
- Thermal-equilibrium state.
- To describe: state parameter (SP, state variables, or thermodynamic variables), function of state (FOS).
- SP: geometry ( $V$ ), mechanics ( $p$ ), chemistry ( $n_i$ ), electromagnetics ( $E, P, H, M$ ).
- FOS: functions with state variables.

## §1.1 Thermal equilibrium state

- Objects: Macroscopic system constructed by microscopic particles.
- Methods: thermodynamics (macro  $\rightarrow$  macro); statistical physics (micro  $\rightarrow$  macro).
- Systems: isolated, closed, open (particles and energy).
- Thermal-equilibrium state.
- To describe: state parameter (SP, state variables, or thermodynamic variables), function of state (FOS).
- SP: geometry ( $V$ ), mechanics ( $p$ ), chemistry ( $n_i$ ), electromagnetics ( $E, P, H, M$ ).
- FOS: functions with state variables.
- Thermal SP (additional): temperature ( $T$ ).

## 1 Chpt 1. Basic Laws in Thermodynamics

- 1.1 Thermal equilibrium state
- 1.2 Law of thermal equilibrium and the temperature
- 1.3 Equation of state
- 1.4 Work
- 1.5 First law of thermodynamics
- 1.6 Thermal capacity and enthalpy
- 1.7 Internal energy of the ideal gas
- 1.8 Adiabatic process of the ideal gas
- 1.9 Carnot cycle in the ideal gas
- 1.10 Second law of thermodynamics
- 1.11 Carnot's theorem
- 1.12 Thermodynamic temperature scale
- 1.13 Clausius theorem
- 1.14 Entropy
- 1.15 Entropy of ideal gas
- 1.16 Math expression of the 2nd law
- 1.17 Applications for the entropy increasing principle
- 1.18 Free energy and Gibbs function

## §1.2 Law of thermal equilibrium and the temperature

## §1.2 Law of thermal equilibrium and the temperature

- Considering two contacting closed systems, in thermal equilibrium, some state parameter should be equal:

## §1.2 Law of thermal equilibrium and the temperature

- Considering two contacting closed systems, in thermal equilibrium, some state parameter should be equal: Temperature ( $T$ ).

## §1.2 Law of thermal equilibrium and the temperature

- Considering two contacting closed systems, in thermal equilibrium, some state parameter should be equal: Temperature ( $T$ ).
- **0th law of thermodynamics:** Exists the  $T$ , that any two contacting systems with the same  $T$  stay in thermal equilibrium.



## §1.2 Law of thermal equilibrium and the temperature

- Considering two contacting closed systems, in thermal equilibrium, some state parameter should be equal: Temperature ( $T$ ).
- **0th law of thermodynamics:** Exists the  $T$ , that any two contacting systems with the same  $T$  stay in thermal equilibrium.
- Thermometric scale  $T$  (K) (Kelvin).
- Ideal gas scaling:  $T = 273.16\text{K} \times \lim_{p_t \rightarrow 0} (p/p_t)$ .
- Daily use, Celsius scale  $(T - 273.15)^\circ\text{C}$ .

## 1 Chpt 1. Basic Laws in Thermodynamics

- 1.1 Thermal equilibrium state
- 1.2 Law of thermal equilibrium and the temperature
- **1.3 Equation of state**
- 1.4 Work
- 1.5 First law of thermodynamics
- 1.6 Thermal capacity and enthalpy
- 1.7 Internal energy of the ideal gas
- 1.8 Adiabatic process of the ideal gas

- 1.9 Carnot cycle in the ideal gas
- 1.10 Second law of thermodynamics
- 1.11 Carnot's theorem
- 1.12 Thermodynamic temperature scale
- 1.13 Clausius theorem
- 1.14 Entropy
- 1.15 Entropy of ideal gas
- 1.16 Math expression of the 2nd law
- 1.17 Applications for the entropy increasing principle
- 1.18 Free energy and Gibbs function

## §1.3 Equation of state (EOS)

## §1.3 Equation of state (EOS)

- General EOS:  $f(p, V, T) = 0$ .

## §1.3 Equation of state (EOS)

- General EOS:  $f(p, V, T) = 0$ .
- Two methods to obtain: experiments; statistical physics.

## §1.3 Equation of state (EOS)

- General EOS:  $f(p, V, T) = 0$ .
- Two methods to obtain: experiments; statistical physics.
- Related physical variables:

## §1.3 Equation of state (EOS)

- General EOS:  $f(p, V, T) = 0$ .
- Two methods to obtain: experiments; statistical physics.
- Related physical variables:  
Coefficient of cubical expansion:  $\alpha = \frac{1}{V} \left( \frac{\partial V}{\partial T} \right)_p$ ;

## §1.3 Equation of state (EOS)

- General EOS:  $f(p, V, T) = 0$ .
- Two methods to obtain: experiments; statistical physics.
- Related physical variables:

Coefficient of cubical expansion:  $\alpha = \frac{1}{V} \left( \frac{\partial V}{\partial T} \right)_p$ ;

Pressure coefficient:  $\beta = \frac{1}{p} \left( \frac{\partial p}{\partial T} \right)_V$ ;



## §1.3 Equation of state (EOS)

- General EOS:  $f(p, V, T) = 0$ .
- Two methods to obtain: experiments; statistical physics.
- Related physical variables:

Coefficient of cubical expansion:  $\alpha = \frac{1}{V} \left( \frac{\partial V}{\partial T} \right)_p$ ;

Pressure coefficient:  $\beta = \frac{1}{p} \left( \frac{\partial p}{\partial T} \right)_V$ ;

Isothermal compressibility:  $\kappa_T = -\frac{1}{V} \left( \frac{\partial V}{\partial p} \right)_T$ .

## §1.3 Equation of state (EOS)

- General EOS:  $f(p, V, T) = 0$ .
- Two methods to obtain: experiments; statistical physics.
- Related physical variables:

Coefficient of cubical expansion:  $\alpha = \frac{1}{V} \left( \frac{\partial V}{\partial T} \right)_p$ ;

Pressure coefficient:  $\beta = \frac{1}{p} \left( \frac{\partial p}{\partial T} \right)_V$ ;

Isothermal compressibility:  $\kappa_T = -\frac{1}{V} \left( \frac{\partial V}{\partial p} \right)_T$ .

- Using  $\left( \frac{\partial T}{\partial V} \right)_p \left( \frac{\partial p}{\partial T} \right)_V \left( \frac{\partial V}{\partial p} \right)_T = -1$ ,  
 $\therefore \alpha = \kappa_T \beta p$ .

# Examples of EOS

# Examples of EOS

- Ideal **gas**:  $pV = nRT$ .

# Examples of EOS

- Ideal **gas**:  $pV = nRT$ .
- Real gas:
  - 1 van der Waals approximation:
$$\left(p + \frac{an^2}{V^2}\right)(V - nb) = nRT.$$

# Examples of EOS

- Ideal **gas**:  $pV = nRT$ .

- Real gas:

- 1 van der Waals approximation:

$$\left(p + \frac{an^2}{V^2}\right)(V - nb) = nRT.$$

- 2 Onnes approach:

$$p = \left(\frac{nRT}{V}\right) \left[1 + \frac{n}{V}B(T) + \left(\frac{n}{V}\right)^2 C(T) + \dots\right].$$

# Examples of EOS

- Ideal **gas**:  $pV = nRT$ .

- Real gas:

- 1 van der Waals approximation:

$$\left(p + \frac{an^2}{V^2}\right)(V - nb) = nRT.$$

- 2 Onnes approach:

$$p = \left(\frac{nRT}{V}\right) \left[1 + \frac{n}{V}B(T) + \left(\frac{n}{V}\right)^2 C(T) + \dots\right].$$

- Simple **solid and liquid**:

# Examples of EOS

- Ideal **gas**:  $pV = nRT$ .

- Real gas:

- 1 van der Waals approximation:

$$\left(p + \frac{an^2}{V^2}\right)(V - nb) = nRT.$$

- 2 Onnes approach:

$$p = \left(\frac{nRT}{V}\right) \left[1 + \frac{n}{V}B(T) + \left(\frac{n}{V}\right)^2 C(T) + \dots\right].$$

- Simple **solid and liquid**:

$$V(T, p) = V_0(T_0, 0)[1 + \alpha(T - T_0) - \kappa_T p].$$



# Examples of EOS

- Ideal **gas**:  $pV = nRT$ .

- Real gas:

- 1 van der Waals approximation:

$$\left(p + \frac{an^2}{V^2}\right)(V - nb) = nRT.$$

- 2 Onnes approach:

$$p = \left(\frac{nRT}{V}\right) \left[1 + \frac{n}{V}B(T) + \left(\frac{n}{V}\right)^2 C(T) + \dots\right].$$

- Simple **solid and liquid**:

$$V(T, p) = V_0(T_0, 0)[1 + \alpha(T - T_0) - \kappa_T p].$$

- **Paramagnet**:  $f(M, H, T) = 0$  (general).

# Examples of EOS

- Ideal **gas**:  $pV = nRT$ .

- Real gas:

1 van der Waals approximation:

$$\left(p + \frac{an^2}{V^2}\right)(V - nb) = nRT.$$

2 Onnes approach:

$$p = \left(\frac{nRT}{V}\right) \left[1 + \frac{n}{V}B(T) + \left(\frac{n}{V}\right)^2 C(T) + \dots\right].$$

- Simple **solid and liquid**:

$$V(T, p) = V_0(T_0, 0)[1 + \alpha(T - T_0) - \kappa_T p].$$

- **Paramagnet**:  $f(M, H, T) = 0$  (general).

$$\text{Curie's law} \text{ — } M = \frac{C}{T}H.$$

# Examples of EOS

- Ideal **gas**:  $pV = nRT$ .

- Real gas:

1 van der Waals approximation:

$$\left(p + \frac{an^2}{V^2}\right)(V - nb) = nRT.$$

2 Onnes approach:

$$p = \left(\frac{nRT}{V}\right) \left[1 + \frac{n}{V}B(T) + \left(\frac{n}{V}\right)^2 C(T) + \dots\right].$$

- Simple **solid and liquid**:

$$V(T, p) = V_0(T_0, 0)[1 + \alpha(T - T_0) - \kappa_T p].$$

- **Paramagnet**:  $f(M, H, T) = 0$  (general).

$$\text{Curie's law} \text{ — } M = \frac{C}{T}H.$$

- Extensive quantity:  $m, n, V, \Sigma M$ .

- Intensive quantity:  $p, T, H, \frac{\text{extensive quantity}}{n(\text{or } m, V)}$  like  $\rho = \frac{m}{V}$ .

## 1 Chpt 1. Basic Laws in Thermodynamics

- 1.1 Thermal equilibrium state
- 1.2 Law of thermal equilibrium and the temperature
- 1.3 Equation of state
- **1.4 Work**
- 1.5 First law of thermodynamics
- 1.6 Thermal capacity and enthalpy
- 1.7 Internal energy of the ideal gas
- 1.8 Adiabatic process of the ideal gas

- 1.9 Carnot cycle in the ideal gas
- 1.10 Second law of thermodynamics
- 1.11 Carnot's theorem
- 1.12 Thermodynamic temperature scale
- 1.13 Clausius theorem
- 1.14 Entropy
- 1.15 Entropy of ideal gas
- 1.16 Math expression of the 2nd law
- 1.17 Applications for the entropy increasing principle
- 1.18 Free energy and Gibbs function

# §1.4 Work

## §1.4 Work

- **Gas:**  $\delta W = -pdV$ . Notice the meaning of “ $\delta$ ”.

## §1.4 Work

- **Gas:**  $\delta W = -pdV$ . Notice the meaning of “ $\delta$ ”.  
Integration:  $W = -\int_{V_A}^{V_B} pdV$ , depends on the path.

## §1.4 Work

- **Gas:**  $\delta W = -pdV$ . Notice the meaning of “ $\delta$ ”.  
Integration:  $W = -\int_{V_A}^{V_B} pdV$ , depends on the path.
- **Liquid surface film:**  $\delta W = \sigma dA$ .



## §1.4 Work

- **Gas:**  $\delta W = -pdV$ . Notice the meaning of “ $\delta$ ”.  
Integration:  $W = -\int_{V_A}^{V_B} pdV$ , depends on the path.
- **Liquid surface film:**  $\delta W = \sigma dA$ .
- **Electric medium:**  $\delta W = Vdq$ , where  $V$  is electric potential.


## §1.4 Work

- **Gas:**  $\delta W = -pdV$ . Notice the meaning of “ $\delta$ ”.  
Integration:  $W = -\int_{V_A}^{V_B} pdV$ , depends on the path.
- **Liquid surface film:**  $\delta W = \sigma dA$ .
- **Electric medium:**  $\delta W = Vdq$ , where  $V$  is electric potential.
- **Magnetic medium:**  $\delta W = Vd(\frac{\mu_0 H^2}{2}) + \mu_0 V H dM$ .

## §1.4 Work


- **Gas:**  $\delta W = -pdV$ . Notice the meaning of “ $\delta$ ”.  
Integration:  $W = -\int_{V_A}^{V_B} pdV$ , depends on the path.
- **Liquid surface film:**  $\delta W = \sigma dA$ .
- **Electric medium:**  $\delta W = Vdq$ , where  $V$  is electric potential.
- **Magnetic medium:**  $\delta W = Vd(\frac{\mu_0 H^2}{2}) + \mu_0 V H dM$ .
- **Summary:**  $\delta W = \sum_i Y_i dy_i$ ,  
where  $Y_i$  is the generalized force, and  $y_i$  is the generalized displacement.

## §1.4 Work – Magnetic medium



## §1.4 Work – Magnetic medium


- **Magnetic medium:**  $\delta W = V I dt$ , with  $V$  the induced electric potential.



## §1.4 Work – Magnetic medium

- **Magnetic medium:**  $\delta W = V I dt$ , with  $V$  the induced electric potential.

$V = \frac{d\Phi}{dt} = N \frac{d(AB)}{dt}$ , where  $A$  is the area, and  $B$  is the magnetic field.




## §1.4 Work – Magnetic medium

- **Magnetic medium:**  $\delta W = V I dt$ , with  $V$  the induced electric potential.

$V = \frac{d\Phi}{dt} = N \frac{d(AB)}{dt}$ , where  $A$  is the area, and  $B$  is the magnetic field.


- From Ampere's law:  $\oint H dl = NI$ ;  $\Rightarrow Hl = NI$ .



## §1.4 Work – Magnetic medium

- **Magnetic medium:**  $\delta W = V I dt$ , with  $V$  the induced electric potential.

$V = \frac{d\Phi}{dt} = N \frac{d(AB)}{dt}$ , where  $A$  is the area, and  $B$  is the magnetic field.




- From Ampere's law:  $\oint H dl = NI$ ;  $\Rightarrow Hl = NI$ .  
 $\Rightarrow \delta W = (NA \frac{dB}{dt}) (\frac{l}{N} H) dt = Al H dB = V H dB$ .



## §1.4 Work – Magnetic medium

- **Magnetic medium:**  $\delta W = V I dt$ , with  $V$  the induced electric potential.

$V = \frac{d\Phi}{dt} = N \frac{d(AB)}{dt}$ , where  $A$  is the area, and  $B$  is the magnetic field.




- From Ampere's law:  $\oint H dl = NI$ ;  $\Rightarrow Hl = NI$ .  
 $\Rightarrow \delta W = (NA \frac{dB}{dt}) (\frac{l}{N} H) dt = Al H dB = V H dB$ .  
 Noticing  $B = \mu H = \mu_0 (H + M)$ ,

## §1.4 Work – Magnetic medium

- **Magnetic medium:**  $\delta W = V I dt$ , with  $V$  the induced electric potential.

$V = \frac{d\Phi}{dt} = N \frac{d(AB)}{dt}$ , where  $A$  is the area, and  $B$  is the magnetic field.



- From Ampere's law:  $\oint H dl = NI$ ;  $\Rightarrow Hl = NI$ .


$$\Rightarrow \delta W = (N A \frac{dB}{dt}) (\frac{l}{N} H) dt = A l H dB = V H dB.$$

Noticing  $B = \mu H = \mu_0(H + M)$ ,

$$\delta W = V H d[\mu_0(H + M)]$$

## §1.4 Work – Magnetic medium

- **Magnetic medium:**  $\delta W = V I dt$ , with  $V$  the induced electric potential.



$V = \frac{d\Phi}{dt} = N \frac{d(AB)}{dt}$ , where  $A$  is the area, and  $B$  is the magnetic field.

- From Ampere's law:  $\oint H dl = NI$ ;  $\Rightarrow Hl = NI$ .


$$\Rightarrow \delta W = (N A \frac{dB}{dt}) (\frac{l}{N} H) dt = A l H dB = V H dB.$$

Noticing  $B = \mu H = \mu_0 (H + M)$ ,

$$\delta W = V H d[\mu_0 (H + M)] = V d(\frac{\mu_0 H^2}{2}) + \mu_0 V H dM,$$

## §1.4 Work – Magnetic medium

- **Magnetic medium:**  $\delta W = V I dt$ , with  $V$  the induced electric potential.



$V = \frac{d\Phi}{dt} = N \frac{d(AB)}{dt}$ , where  $A$  is the area, and  $B$  is the magnetic field.

- From Ampere's law:  $\oint H dl = NI$ ;  $\Rightarrow Hl = NI$ .

$$\Rightarrow \delta W = (N A \frac{dB}{dt}) (\frac{l}{N} I) dt = A l H dB = V H dB.$$

Noticing  $B = \mu H = \mu_0(H + M)$ ,

$$\delta W = V H d[\mu_0(H + M)] = V d(\frac{\mu_0 H^2}{2}) + \mu_0 V H dM,$$

where  $\frac{\mu_0 H^2}{2}$  is the magnetic energy density (work by increasing the magnetic field);  $M$  is the magnetization (work done by magnetize the material).

## 1 Chpt 1. Basic Laws in Thermodynamics

- 1.1 Thermal equilibrium state
- 1.2 Law of thermal equilibrium and the temperature
- 1.3 Equation of state
- 1.4 Work
- **1.5 First law of thermodynamics**
- 1.6 Thermal capacity and enthalpy
- 1.7 Internal energy of the ideal gas
- 1.8 Adiabatic process of the ideal gas
- 1.9 Carnot cycle in the ideal gas
- 1.10 Second law of thermodynamics
- 1.11 Carnot's theorem
- 1.12 Thermodynamic temperature scale
- 1.13 Clausius theorem
- 1.14 Entropy
- 1.15 Entropy of ideal gas
- 1.16 Math expression of the 2nd law
- 1.17 Applications for the entropy increasing principle
- 1.18 Free energy and Gibbs function

# §1.5 First law of thermodynamics

## §1.5 First law of thermodynamics

- $dU = \delta Q + \delta W.$

## §1.5 First law of thermodynamics

- $dU = \delta Q + \delta W$ .
- Internal energy ( $U$ ): summation of random movement energy of molecules, including kinetic energy of molecules, potential energy between molecules, inner energy of molecules (atoms... iteration).



## §1.5 First law of thermodynamics

- $dU = \delta Q + \delta W$ .
- Internal energy ( $U$ ): summation of random movement energy of molecules, including kinetic energy of molecules, potential energy between molecules, inner energy of molecules (atoms... iteration).
- “Almost” obeys in all physical processes.

## 1 Chpt 1. Basic Laws in Thermodynamics

- 1.1 Thermal equilibrium state
- 1.2 Law of thermal equilibrium and the temperature
- 1.3 Equation of state
- 1.4 Work
- 1.5 First law of thermodynamics
- **1.6 Thermal capacity and enthalpy**
- 1.7 Internal energy of the ideal gas
- 1.8 Adiabatic process of the ideal gas
- 1.9 Carnot cycle in the ideal gas
- 1.10 Second law of thermodynamics
- 1.11 Carnot's theorem
- 1.12 Thermodynamic temperature scale
- 1.13 Clausius theorem
- 1.14 Entropy
- 1.15 Entropy of ideal gas
- 1.16 Math expression of the 2nd law
- 1.17 Applications for the entropy increasing principle
- 1.18 Free energy and Gibbs function

## §1.6 Thermal capacity and enthalpy

## §1.6 Thermal capacity and enthalpy

- Thermal capacity (heat capacity):  $C \equiv \lim_{\Delta T \rightarrow 0} \frac{\Delta Q}{\Delta T}$ .

## §1.6 Thermal capacity and enthalpy

- Thermal capacity (heat capacity):  $C \equiv \lim_{\Delta T \rightarrow 0} \frac{\Delta Q}{\Delta T}$ .
- Isochoric (constant volume) process:  
 $\Delta Q = \Delta U - \Delta W = \Delta U,$   
 $\therefore C_V = \lim_{\Delta T \rightarrow 0} \left( \frac{\Delta U}{\Delta T} \right)_V = \left( \frac{\partial U}{\partial T} \right)_V.$

## §1.6 Thermal capacity and enthalpy

- Thermal capacity (heat capacity):  $C \equiv \lim_{\Delta T \rightarrow 0} \frac{\Delta Q}{\Delta T}$ .
- Isochoric (constant volume) process:  
 $\Delta Q = \Delta U - \Delta W = \Delta U$ ,  
 $\therefore C_V = \lim_{\Delta T \rightarrow 0} \left( \frac{\Delta U}{\Delta T} \right)_V = \left( \frac{\partial U}{\partial T} \right)_V$ .
- Isobaric (constant pressure) process:  
 $C_p = \lim_{\Delta T \rightarrow 0} \left( \frac{\Delta U + p \Delta V}{\Delta T} \right)_p = \left( \frac{\partial U}{\partial T} \right)_p + p \left( \frac{\partial V}{\partial T} \right)_p$ .

## §1.6 Thermal capacity and enthalpy

- Thermal capacity (heat capacity):  $C \equiv \lim_{\Delta T \rightarrow 0} \frac{\Delta Q}{\Delta T}$ .
- Isochoric (constant volume) process:  
 $\Delta Q = \Delta U - \Delta W = \Delta U$ ,  
 $\therefore C_V = \lim_{\Delta T \rightarrow 0} \left( \frac{\Delta U}{\Delta T} \right)_V = \left( \frac{\partial U}{\partial T} \right)_V$ .
- Isobaric (constant pressure) process:  
 $C_p = \lim_{\Delta T \rightarrow 0} \left( \frac{\Delta U + p\Delta V}{\Delta T} \right)_p = \left( \frac{\partial U}{\partial T} \right)_p + p \left( \frac{\partial V}{\partial T} \right)_p$ .
- Enthalpy:  $H \equiv U + pV$ .

## §1.6 Thermal capacity and enthalpy

- Thermal capacity (heat capacity):  $C \equiv \lim_{\Delta T \rightarrow 0} \frac{\Delta Q}{\Delta T}$ .
- Isochoric (constant volume) process:  
 $\Delta Q = \Delta U - \Delta W = \Delta U$ ,  
 $\therefore C_V = \lim_{\Delta T \rightarrow 0} \left( \frac{\Delta U}{\Delta T} \right)_V = \left( \frac{\partial U}{\partial T} \right)_V$ .
- Isobaric (constant pressure) process:  
 $C_p = \lim_{\Delta T \rightarrow 0} \left( \frac{\Delta U + p \Delta V}{\Delta T} \right)_p = \left( \frac{\partial U}{\partial T} \right)_p + p \left( \frac{\partial V}{\partial T} \right)_p$ .
- Enthalpy:  $H \equiv U + pV$ .  
 $\therefore C_p = \left( \frac{\partial H}{\partial T} \right)_p$ .



## 1 Chpt 1. Basic Laws in Thermodynamics

- 1.1 Thermal equilibrium state
- 1.2 Law of thermal equilibrium and the temperature
- 1.3 Equation of state
- 1.4 Work
- 1.5 First law of thermodynamics
- 1.6 Thermal capacity and enthalpy
- 1.7 Internal energy of the ideal gas
- 1.8 Adiabatic process of the ideal gas
- 1.9 Carnot cycle in the ideal gas
- 1.10 Second law of thermodynamics
- 1.11 Carnot's theorem
- 1.12 Thermodynamic temperature scale
- 1.13 Clausius theorem
- 1.14 Entropy
- 1.15 Entropy of ideal gas
- 1.16 Math expression of the 2nd law
- 1.17 Applications for the entropy increasing principle
- 1.18 Free energy and Gibbs function

## §1.7 Internal energy of the ideal gas

## §1.7 Internal energy of the ideal gas

- General:  $f(U, T, V) = 0$ . Why no  $p$ ?

## §1.7 Internal energy of the ideal gas

- General:  $f(U, T, V) = 0$ . Why no  $p$ ?
- $\left(\frac{\partial U}{\partial V}\right)_T \left(\frac{\partial V}{\partial T}\right)_U \left(\frac{\partial T}{\partial U}\right)_V = -1$

## §1.7 Internal energy of the ideal gas

- General:  $f(U, T, V) = 0$ . Why no  $p$ ?
- $\left(\frac{\partial U}{\partial V}\right)_T \left(\frac{\partial V}{\partial T}\right)_U \left(\frac{\partial T}{\partial U}\right)_V = -1$   
 $\Rightarrow \left(\frac{\partial U}{\partial V}\right)_T = - \left(\frac{\partial U}{\partial T}\right)_V \left(\frac{\partial T}{\partial V}\right)_U$  .

## §1.7 Internal energy of the ideal gas

- General:  $f(U, T, V) = 0$ . Why no  $p$ ?
- $\left(\frac{\partial U}{\partial V}\right)_T \left(\frac{\partial V}{\partial T}\right)_U \left(\frac{\partial T}{\partial U}\right)_V = -1$   
 $\Rightarrow \left(\frac{\partial U}{\partial V}\right)_T = - \left(\frac{\partial U}{\partial T}\right)_V \left(\frac{\partial T}{\partial V}\right)_U$  .
- Joule's experiment...

## §1.7 Internal energy of the ideal gas

- General:  $f(U, T, V) = 0$ . Why no  $p$ ?
- $\left(\frac{\partial U}{\partial V}\right)_T \left(\frac{\partial V}{\partial T}\right)_U \left(\frac{\partial T}{\partial U}\right)_V = -1$   
 $\Rightarrow \left(\frac{\partial U}{\partial V}\right)_T = - \left(\frac{\partial U}{\partial T}\right)_V \left(\frac{\partial T}{\partial V}\right)_U$ .
- Joule's experiment...  
1°.  $\Delta U = Q + W = 0 + 0 = 0$ ,

## §1.7 Internal energy of the ideal gas

- General:  $f(U, T, V) = 0$ . Why no  $p$ ?
- $\left(\frac{\partial U}{\partial V}\right)_T \left(\frac{\partial V}{\partial T}\right)_U \left(\frac{\partial T}{\partial U}\right)_V = -1$   
 $\Rightarrow \left(\frac{\partial U}{\partial V}\right)_T = - \left(\frac{\partial U}{\partial T}\right)_V \left(\frac{\partial T}{\partial V}\right)_U$ .
- Joule's experiment...  
1°.  $\Delta U = Q + W = 0 + 0 = 0$ , 2°.  $\left(\frac{\partial T}{\partial V}\right)_U = 0$ ,



## §1.7 Internal energy of the ideal gas

- General:  $f(U, T, V) = 0$ . Why no  $p$ ?

- $\left(\frac{\partial U}{\partial V}\right)_T \left(\frac{\partial V}{\partial T}\right)_U \left(\frac{\partial T}{\partial U}\right)_V = -1$   
 $\Rightarrow \left(\frac{\partial U}{\partial V}\right)_T = - \left(\frac{\partial U}{\partial T}\right)_V \left(\frac{\partial T}{\partial V}\right)_U$ .

- Joule's experiment...

$$1^\circ. \Delta U = Q + W = 0 + 0 = 0, \quad 2^\circ. \left(\frac{\partial T}{\partial V}\right)_U = 0,$$

$$\therefore \left(\frac{\partial U}{\partial V}\right)_T = 0$$

## §1.7 Internal energy of the ideal gas

- General:  $f(U, T, V) = 0$ . Why no  $p$ ?

- $\left(\frac{\partial U}{\partial V}\right)_T \left(\frac{\partial V}{\partial T}\right)_U \left(\frac{\partial T}{\partial U}\right)_V = -1$   
 $\Rightarrow \left(\frac{\partial U}{\partial V}\right)_T = - \left(\frac{\partial U}{\partial T}\right)_V \left(\frac{\partial T}{\partial V}\right)_U$ .

- Joule's experiment...

$$1^\circ. \Delta U = Q + W = 0 + 0 = 0, \quad 2^\circ. \left(\frac{\partial T}{\partial V}\right)_U = 0,$$

$$\therefore \left(\frac{\partial U}{\partial V}\right)_T = 0 \Rightarrow U(T, V) \rightarrow U(T).$$

## §1.7 Internal energy of the ideal gas

- General:  $f(U, T, V) = 0$ . Why no  $p$ ?
- $\left(\frac{\partial U}{\partial V}\right)_T \left(\frac{\partial V}{\partial T}\right)_U \left(\frac{\partial T}{\partial U}\right)_V = -1$   
 $\Rightarrow \left(\frac{\partial U}{\partial V}\right)_T = - \left(\frac{\partial U}{\partial T}\right)_V \left(\frac{\partial T}{\partial V}\right)_U$ .
- Joule's experiment...  
1°.  $\Delta U = Q + W = 0 + 0 = 0$ , 2°.  $\left(\frac{\partial T}{\partial V}\right)_U = 0$ ,  
 $\therefore \left(\frac{\partial U}{\partial V}\right)_T = 0 \Rightarrow U(T, V) \rightarrow U(T)$ .
- $C_V \equiv \left(\frac{\partial U}{\partial T}\right)_V = \frac{dU}{dT}$ .

## §1.7 Internal energy of the ideal gas

- General:  $f(U, T, V) = 0$ . Why no  $p$ ?
- $\left(\frac{\partial U}{\partial V}\right)_T \left(\frac{\partial V}{\partial T}\right)_U \left(\frac{\partial T}{\partial U}\right)_V = -1$   
 $\Rightarrow \left(\frac{\partial U}{\partial V}\right)_T = - \left(\frac{\partial U}{\partial T}\right)_V \left(\frac{\partial T}{\partial V}\right)_U$ .
- Joule's experiment...  
1°.  $\Delta U = Q + W = 0 + 0 = 0$ , 2°.  $\left(\frac{\partial T}{\partial V}\right)_U = 0$ ,  
 $\therefore \left(\frac{\partial U}{\partial V}\right)_T = 0 \Rightarrow U(T, V) \rightarrow U(T)$ .
- $C_V \equiv \left(\frac{\partial U}{\partial T}\right)_V = \frac{dU}{dT}$ .
- $H = U + pV = U(T) + nRT \rightarrow H(T)$ .

## §1.7 Internal energy of the ideal gas

- General:  $f(U, T, V) = 0$ . Why no  $p$ ?

- $\left(\frac{\partial U}{\partial V}\right)_T \left(\frac{\partial V}{\partial T}\right)_U \left(\frac{\partial T}{\partial U}\right)_V = -1$   
 $\Rightarrow \left(\frac{\partial U}{\partial V}\right)_T = - \left(\frac{\partial U}{\partial T}\right)_V \left(\frac{\partial T}{\partial V}\right)_U$ .

- Joule's experiment...

$$1^\circ. \Delta U = Q + W = 0 + 0 = 0, \quad 2^\circ. \left(\frac{\partial T}{\partial V}\right)_U = 0,$$

$$\therefore \left(\frac{\partial U}{\partial V}\right)_T = 0 \Rightarrow U(T, V) \rightarrow U(T).$$

- $C_V \equiv \left(\frac{\partial U}{\partial T}\right)_V = \frac{dU}{dT}$ .
- $H = U + pV = U(T) + nRT \rightarrow H(T)$ .
- $C_p = \frac{dH}{dT} = \frac{dU}{dT} + nR$ .

# §1.7 Internal energy of the ideal gas

- General:  $f(U, T, V) = 0$ . Why no  $p$ ?

- $\left(\frac{\partial U}{\partial V}\right)_T \left(\frac{\partial V}{\partial T}\right)_U \left(\frac{\partial T}{\partial U}\right)_V = -1$   
 $\Rightarrow \left(\frac{\partial U}{\partial V}\right)_T = - \left(\frac{\partial U}{\partial T}\right)_V \left(\frac{\partial T}{\partial V}\right)_U$ .

- Joule's experiment...

$$1^\circ. \Delta U = Q + W = 0 + 0 = 0, \quad 2^\circ. \left(\frac{\partial T}{\partial V}\right)_U = 0,$$

$$\therefore \left(\frac{\partial U}{\partial V}\right)_T = 0 \Rightarrow U(T, V) \rightarrow U(T).$$

- $C_V \equiv \left(\frac{\partial U}{\partial T}\right)_V = \frac{dU}{dT}$ .
- $H = U + pV = U(T) + nRT \rightarrow H(T)$ .
- $C_p = \frac{dH}{dT} = \frac{dU}{dT} + nR$ .
- $\therefore \boxed{C_p - C_V = nR}$ ,

# §1.7 Internal energy of the ideal gas

- General:  $f(U, T, V) = 0$ . Why no  $p$ ?

- $\left(\frac{\partial U}{\partial V}\right)_T \left(\frac{\partial V}{\partial T}\right)_U \left(\frac{\partial T}{\partial U}\right)_V = -1$   
 $\Rightarrow \left(\frac{\partial U}{\partial V}\right)_T = - \left(\frac{\partial U}{\partial T}\right)_V \left(\frac{\partial T}{\partial V}\right)_U$ .

- Joule's experiment...

$$1^\circ. \Delta U = Q + W = 0 + 0 = 0, \quad 2^\circ. \left(\frac{\partial T}{\partial V}\right)_U = 0,$$

$$\therefore \left(\frac{\partial U}{\partial V}\right)_T = 0 \Rightarrow U(T, V) \rightarrow U(T).$$

- $C_V \equiv \left(\frac{\partial U}{\partial T}\right)_V = \frac{dU}{dT}$ .

- $H = U + pV = U(T) + nRT \rightarrow H(T)$ .

- $C_p = \frac{dH}{dT} = \frac{dU}{dT} + nR$ .

- $\therefore \boxed{C_p - C_V = nR}$ , define  $\gamma \equiv \frac{C_p}{C_V}$ ,

# §1.7 Internal energy of the ideal gas

- General:  $f(U, T, V) = 0$ . Why no  $p$ ?

- $\left(\frac{\partial U}{\partial V}\right)_T \left(\frac{\partial V}{\partial T}\right)_U \left(\frac{\partial T}{\partial U}\right)_V = -1$   
 $\Rightarrow \left(\frac{\partial U}{\partial V}\right)_T = - \left(\frac{\partial U}{\partial T}\right)_V \left(\frac{\partial T}{\partial V}\right)_U$ .

- Joule's experiment...

$$1^\circ. \Delta U = Q + W = 0 + 0 = 0, \quad 2^\circ. \left(\frac{\partial T}{\partial V}\right)_U = 0,$$

$$\therefore \left(\frac{\partial U}{\partial V}\right)_T = 0 \Rightarrow U(T, V) \rightarrow U(T).$$

- $C_V \equiv \left(\frac{\partial U}{\partial T}\right)_V = \frac{dU}{dT}$ .
- $H = U + pV = U(T) + nRT \rightarrow H(T)$ .
- $C_p = \frac{dH}{dT} = \frac{dU}{dT} + nR$ .
- $\therefore \boxed{C_p - C_V = nR}$ , define  $\gamma \equiv \frac{C_p}{C_V}$ ,  
 $\Rightarrow C_V = \frac{nR}{\gamma-1}, C_p = \gamma \frac{nR}{\gamma-1} = \gamma C_V$ .



## 1 Chpt 1. Basic Laws in Thermodynamics

- 1.1 Thermal equilibrium state
- 1.2 Law of thermal equilibrium and the temperature
- 1.3 Equation of state
- 1.4 Work
- 1.5 First law of thermodynamics
- 1.6 Thermal capacity and enthalpy
- 1.7 Internal energy of the ideal gas
- 1.8 Adiabatic process of the ideal gas
- 1.9 Carnot cycle in the ideal gas
- 1.10 Second law of thermodynamics
- 1.11 Carnot's theorem
- 1.12 Thermodynamic temperature scale
- 1.13 Clausius theorem
- 1.14 Entropy
- 1.15 Entropy of ideal gas
- 1.16 Math expression of the 2nd law
- 1.17 Applications for the entropy increasing principle
- 1.18 Free energy and Gibbs function

## §1.8 Adiabatic process of the ideal gas

## §1.8 Adiabatic process of the ideal gas

- Adiabatic process :  $\delta Q = 0$ ,  $\delta W = -pdV$ ,

## §1.8 Adiabatic process of the ideal gas

- Adiabatic process :  $\delta Q = 0$ ,  $\delta W = -pdV$ ,  
 $\Rightarrow C_V dT = -pdV$ .

## §1.8 Adiabatic process of the ideal gas

- Adiabatic process :  $\delta Q = 0$ ,  $\delta W = -pdV$ ,  
 $\Rightarrow C_V dT = -pdV$ .
- $pV = nRT \rightarrow pdV + Vdp = nRdT = C_V(\gamma - 1)dT$ ,

## §1.8 Adiabatic process of the ideal gas

- Adiabatic process :  $\delta Q = 0$ ,  $\delta W = -pdV$ ,  
 $\Rightarrow C_V dT = -pdV$ .
- $pV = nRT \rightarrow pdV + Vdp = nRdT = C_V(\gamma - 1)dT$ ,  
 $\Rightarrow pdV + Vdp = -(\gamma - 1)pdV$ .

## §1.8 Adiabatic process of the ideal gas

- Adiabatic process :  $\delta Q = 0$ ,  $\delta W = -pdV$ ,  
 $\Rightarrow C_V dT = -pdV$ .
- $pV = nRT \rightarrow pdV + Vdp = nRdT = C_V(\gamma - 1)dT$ ,  
 $\Rightarrow pdV + Vdp = -(\gamma - 1)pdV$ .  
 $\Rightarrow \gamma pdV + Vdp = 0$ ,

## §1.8 Adiabatic process of the ideal gas

- Adiabatic process :  $\delta Q = 0$ ,  $\delta W = -pdV$ ,  
 $\Rightarrow C_V dT = -pdV$ .
- $pV = nRT \rightarrow pdV + Vdp = nRdT = C_V(\gamma - 1)dT$ ,  
 $\Rightarrow pdV + Vdp = -(\gamma - 1)pdV$ .  
 $\Rightarrow \gamma pdV + Vdp = 0$ ,  
 $\Rightarrow \frac{dp}{p} + \gamma \frac{dV}{V} = 0$ ,



## §1.8 Adiabatic process of the ideal gas

- Adiabatic process :  $\delta Q = 0$ ,  $\delta W = -pdV$ ,  
 $\Rightarrow C_V dT = -pdV$ .
- $pV = nRT \rightarrow pdV + Vdp = nRdT = C_V(\gamma - 1)dT$ ,  
 $\Rightarrow pdV + Vdp = -(\gamma - 1)pdV$ .  
 $\Rightarrow \gamma pdV + Vdp = 0$ ,  
 $\Rightarrow \frac{dp}{p} + \gamma \frac{dV}{V} = 0$ ,  
 $\Rightarrow pV^\gamma = \text{Const.}$

# §1.8 Adiabatic process of the ideal gas

- Adiabatic process :  $\delta Q = 0$ ,  $\delta W = -pdV$ ,  
 $\Rightarrow C_V dT = -pdV$ .
- $pV = nRT \rightarrow pdV + Vdp = nRdT = C_V(\gamma - 1)dT$ ,  
 $\Rightarrow pdV + Vdp = -(\gamma - 1)pdV$ .

$$\Rightarrow \gamma pdV + Vdp = 0,$$

$$\Rightarrow \frac{dp}{p} + \gamma \frac{dV}{V} = 0,$$

$$\Rightarrow pV^\gamma = \text{Const.}$$




Figure 3-6  
 Atkins Physical Chemistry, Eighth Edition  
 © 2006 Peter Atkins and Julio de Paula

# §1.8 Adiabatic process of the ideal gas

- Adiabatic process :  $\delta Q = 0$ ,  $\delta W = -pdV$ ,  
 $\Rightarrow C_V dT = -pdV$ .
- $pV = nRT \rightarrow pdV + Vdp = nRdT = C_V(\gamma - 1)dT$ ,  
 $\Rightarrow pdV + Vdp = -(\gamma - 1)pdV$ .

$$\Rightarrow \gamma pdV + Vdp = 0,$$

$$\Rightarrow \frac{dp}{p} + \gamma \frac{dV}{V} = 0,$$

$$\Rightarrow pV^\gamma = \text{Const.}$$

$$TV^{\gamma-1} = \text{Const.},$$

$$T^\gamma p^{1-\gamma} = \text{Const.}$$




Figure 3-6  
 Atkins Physical Chemistry, Eighth Edition  
 © 2006 Peter Atkins and Julio de Paula

# §1.8 Adiabatic process of the ideal gas

- Adiabatic process :  $\delta Q = 0$ ,  $\delta W = -pdV$ ,  
 $\Rightarrow C_V dT = -pdV$ .
- $pV = nRT \rightarrow pdV + Vdp = nRdT = C_V(\gamma - 1)dT$ ,  
 $\Rightarrow pdV + Vdp = -(\gamma - 1)pdV$ .

$$\Rightarrow \gamma pdV + Vdp = 0,$$

$$\Rightarrow \frac{dp}{p} + \gamma \frac{dV}{V} = 0,$$

$$\Rightarrow pV^\gamma = \text{Const.}$$

$$TV^{\gamma-1} = \text{Const.},$$

$$T^\gamma p^{1-\gamma} = \text{Const.}$$

- Velocity of sound:  $a^2 = \gamma \frac{p}{\rho}$ .




Figure 3-6  
 Atkins Physical Chemistry, Eighth Edition  
 © 2006 Peter Atkins and Julio de Paula


## 1 Chpt 1. Basic Laws in Thermodynamics

- 1.1 Thermal equilibrium state
- 1.2 Law of thermal equilibrium and the temperature
- 1.3 Equation of state
- 1.4 Work
- 1.5 First law of thermodynamics
- 1.6 Thermal capacity and enthalpy
- 1.7 Internal energy of the ideal gas
- 1.8 Adiabatic process of the ideal gas

## • 1.9 Carnot cycle in the ideal gas


- 1.10 Second law of thermodynamics
- 1.11 Carnot's theorem
- 1.12 Thermodynamic temperature scale
- 1.13 Clausius theorem
- 1.14 Entropy
- 1.15 Entropy of ideal gas
- 1.16 Math expression of the 2nd law
- 1.17 Applications for the entropy increasing principle
- 1.18 Free energy and Gibbs function

## §1.9 Carnot cycle in the ideal gas




## §1.9 Carnot cycle in the ideal gas

- Setting  $n = 1$ ,  $pV = RT$ .



## §1.9 Carnot cycle in the ideal gas


- Setting  $n = 1$ ,  $pV = RT$ .
- Isothermal process:  $Q = -W$ .





## §1.9 Carnot cycle in the ideal gas


- Setting  $n = 1$ ,  $pV = RT$ .
- Isothermal process:  $Q = -W$ .
- $W = - \int_{V_A}^{V_B} p dV$



# §1.9 Carnot cycle in the ideal gas


- Setting  $n = 1$ ,  $pV = RT$ .
- Isothermal process:  $Q = -W$ .
- $$W = - \int_{V_A}^{V_B} p dV$$
  

$$= - \int_{V_A}^{V_B} \frac{RT}{V} dV$$




# §1.9 Carnot cycle in the ideal gas

- Setting  $n = 1$ ,  $pV = RT$ .
- Isothermal process:  $Q = -W$ .
- $W = - \int_{V_A}^{V_B} p dV$   
 $= - \int_{V_A}^{V_B} \frac{RT}{V} dV = -RT \ln \frac{V_B}{V_A}.$




## §1.9 Carnot cycle in the ideal gas

- Setting  $n = 1$ ,  $pV = RT$ .
- Isothermal process:  $Q = -W$ .
- $W = -\int_{V_A}^{V_B} p dV$   
 $= -\int_{V_A}^{V_B} \frac{RT}{V} dV = -RT \ln \frac{V_B}{V_A}$ .
- Adiabatic process:  $Q = 0$ .



# §1.9 Carnot cycle in the ideal gas


- Setting  $n = 1$ ,  $pV = RT$ .
- Isothermal process:  $Q = -W$ .
- $W = -\int_{V_A}^{V_B} p dV$   
 $= -\int_{V_A}^{V_B} \frac{RT}{V} dV = -RT \ln \frac{V_B}{V_A}$ .
- Adiabatic process:  $Q = 0$ .



1 Isothermal expansion: absorb heat  $Q_1 = RT_1 \ln \frac{V_2}{V_1}$ ,

# §1.9 Carnot cycle in the ideal gas


- Setting  $n = 1$ ,  $pV = RT$ .
- Isothermal process:  $Q = -W$ .
- $W = - \int_{V_A}^{V_B} p dV$   
 $= - \int_{V_A}^{V_B} \frac{RT}{V} dV = -RT \ln \frac{V_B}{V_A}$ .
- Adiabatic process:  $Q = 0$ .



- 1 Isothermal expansion: absorb heat  $Q_1 = RT_1 \ln \frac{V_2}{V_1}$ ,
- 2 Adiabatic expansion:  $Q = 0$ ,

# §1.9 Carnot cycle in the ideal gas


- Setting  $n = 1$ ,  $pV = RT$ .
- Isothermal process:  $Q = -W$ .
- $W = -\int_{V_A}^{V_B} p dV$   
 $= -\int_{V_A}^{V_B} \frac{RT}{V} dV = -RT \ln \frac{V_B}{V_A}$ .
- Adiabatic process:  $Q = 0$ .



- 1 Isothermal expansion: absorb heat  $Q_1 = RT_1 \ln \frac{V_2}{V_1}$ ,
- 2 Adiabatic expansion:  $Q = 0$ ,
- 3 Isothermal compression: release heat  $Q_2 = RT_2 \ln \frac{V_3}{V_4}$ ,

# §1.9 Carnot cycle in the ideal gas

- Setting  $n = 1$ ,  $pV = RT$ .
- Isothermal process:  $Q = -W$ .
- $W = - \int_{V_A}^{V_B} p dV$   
 $= - \int_{V_A}^{V_B} \frac{RT}{V} dV = -RT \ln \frac{V_B}{V_A}$ .
- Adiabatic process:  $Q = 0$ .




- 1 Isothermal expansion: absorb heat  $Q_1 = RT_1 \ln \frac{V_2}{V_1}$ ,
- 2 Adiabatic expansion:  $Q = 0$ ,
- 3 Isothermal compression: release heat  $Q_2 = RT_2 \ln \frac{V_3}{V_4}$ ,
- 4 Adiabatic compression:  $Q = 0$ .



# §1.9 Carnot cycle in the ideal gas


- Setting  $n = 1$ ,  $pV = RT$ .
- Isothermal process:  $Q = -W$ .
- $W = - \int_{V_A}^{V_B} p dV$   
 $= - \int_{V_A}^{V_B} \frac{RT}{V} dV = -RT \ln \frac{V_B}{V_A}$ .
- Adiabatic process:  $Q = 0$ .



- 1 Isothermal expansion: absorb heat  $Q_1 = RT_1 \ln \frac{V_2}{V_1}$ ,
  - 2 Adiabatic expansion:  $Q = 0$ ,
  - 3 Isothermal compression: release heat  $Q_2 = RT_2 \ln \frac{V_3}{V_4}$ ,
  - 4 Adiabatic compression:  $Q = 0$ .
- Total work:  $W = Q_1 - Q_2 = RT_1 \ln \frac{V_2}{V_1} - RT_2 \ln \frac{V_3}{V_4}$ .


# Carnot cycle in the ideal gas

- Total work of a cycle:  $W = Q_1 - Q_2 = RT_1 \ln \frac{V_2}{V_1} - RT_2 \ln \frac{V_3}{V_4}$ .



# Carnot cycle in the ideal gas


- Total work of a cycle:  $W = Q_1 - Q_2 = RT_1 \ln \frac{V_2}{V_1} - RT_2 \ln \frac{V_3}{V_4}$ .
- In adiabatic process:  $TV^{\gamma-1} = C$ ,



# Carnot cycle in the ideal gas

- Total work of a cycle:  $W = Q_1 - Q_2 = RT_1 \ln \frac{V_2}{V_1} - RT_2 \ln \frac{V_3}{V_4}$ .
- In adiabatic process:  $TV^{\gamma-1} = C$ ,

$$\left. \begin{array}{l} T_1 V_2^{\gamma-1} = T_2 V_3^{\gamma-1} \\ T_1 V_1^{\gamma-1} = T_2 V_4^{\gamma-1} \end{array} \right\} \Rightarrow \frac{V_2}{V_1} = \frac{V_3}{V_4}.$$




# Carnot cycle in the ideal gas

- Total work of a cycle:  $W = Q_1 - Q_2 = RT_1 \ln \frac{V_2}{V_1} - RT_2 \ln \frac{V_3}{V_4}$ .
- In adiabatic process:  $TV^{\gamma-1} = C$ ,

$$\left. \begin{array}{l} T_1 V_2^{\gamma-1} = T_2 V_3^{\gamma-1} \\ T_1 V_1^{\gamma-1} = T_2 V_4^{\gamma-1} \end{array} \right\} \Rightarrow \frac{V_2}{V_1} = \frac{V_3}{V_4}.$$


$$\therefore W = R(T_1 - T_2) \ln \frac{V_2}{V_1}.$$



# Carnot cycle in the ideal gas

- Total work of a cycle:  $W = Q_1 - Q_2 = RT_1 \ln \frac{V_2}{V_1} - RT_2 \ln \frac{V_3}{V_4}$ .
- In adiabatic process:  $TV^{\gamma-1} = C$ ,

$$\left. \begin{array}{l} T_1 V_2^{\gamma-1} = T_2 V_3^{\gamma-1} \\ T_1 V_1^{\gamma-1} = T_2 V_4^{\gamma-1} \end{array} \right\} \Rightarrow \frac{V_2}{V_1} = \frac{V_3}{V_4}.$$




$$\therefore W = R(T_1 - T_2) \ln \frac{V_2}{V_1}.$$

- Efficiency:  $\eta = \frac{W}{Q_1} = \frac{R(T_1 - T_2) \ln \frac{V_2}{V_1}}{RT_1 \ln \frac{V_2}{V_1}} = 1 - \frac{T_2}{T_1}.$

# Carnot cycle in the ideal gas

- Total work of a cycle:  $W = Q_1 - Q_2 = RT_1 \ln \frac{V_2}{V_1} - RT_2 \ln \frac{V_3}{V_4}$ .
- In adiabatic process:  $TV^{\gamma-1} = C$ ,

$$\left. \begin{array}{l} T_1 V_2^{\gamma-1} = T_2 V_3^{\gamma-1} \\ T_1 V_1^{\gamma-1} = T_2 V_4^{\gamma-1} \end{array} \right\} \Rightarrow \frac{V_2}{V_1} = \frac{V_3}{V_4}.$$




$$\therefore W = R(T_1 - T_2) \ln \frac{V_2}{V_1}.$$

- Efficiency:  $\eta = \frac{W}{Q_1} = \frac{R(T_1 - T_2) \ln \frac{V_2}{V_1}}{RT_1 \ln \frac{V_2}{V_1}} = 1 - \frac{T_2}{T_1}$ .
- Reverse cycle: refrigerator.

# Carnot cycle in the ideal gas

- Total work of a cycle:  $W = Q_1 - Q_2 = RT_1 \ln \frac{V_2}{V_1} - RT_2 \ln \frac{V_3}{V_4}$ .
- In adiabatic process:  $TV^{\gamma-1} = C$ ,

$$\left. \begin{array}{l} T_1 V_2^{\gamma-1} = T_2 V_3^{\gamma-1} \\ T_1 V_1^{\gamma-1} = T_2 V_4^{\gamma-1} \end{array} \right\} \Rightarrow \frac{V_2}{V_1} = \frac{V_3}{V_4}.$$



$$\therefore W = R(T_1 - T_2) \ln \frac{V_2}{V_1}.$$

- Efficiency:  $\eta = \frac{W}{Q_1} = \frac{R(T_1 - T_2) \ln \frac{V_2}{V_1}}{RT_1 \ln \frac{V_2}{V_1}} = 1 - \frac{T_2}{T_1}$ .
- Reverse cycle: refrigerator.
- Coefficient of performance:  $\eta' = \frac{Q_2}{W} = \frac{T_2}{T_1 - T_2}$ .



## 1 Chpt 1. Basic Laws in Thermodynamics

- 1.1 Thermal equilibrium state
- 1.2 Law of thermal equilibrium and the temperature
- 1.3 Equation of state
- 1.4 Work
- 1.5 First law of thermodynamics
- 1.6 Thermal capacity and enthalpy
- 1.7 Internal energy of the ideal gas
- 1.8 Adiabatic process of the ideal gas

- 1.9 Carnot cycle in the ideal gas
- 1.10 Second law of thermodynamics
- 1.11 Carnot's theorem
- 1.12 Thermodynamic temperature scale
- 1.13 Clausius theorem
- 1.14 Entropy
- 1.15 Entropy of ideal gas
- 1.16 Math expression of the 2nd law
- 1.17 Applications for the entropy increasing principle
- 1.18 Free energy and Gibbs function

## §1.10 Second law of thermodynamics

## §1.10 Second law of thermodynamics

- Direction of real processes? Reversible process, irreversible process.

## §1.10 Second law of thermodynamics

- Direction of real processes? Reversible process, irreversible process.
- Second law of thermodynamics:

## §1.10 Second law of thermodynamics

- Direction of real processes? Reversible process, irreversible process.
- Second law of thermodynamics:
  - i **Clausius statement:** No process is possible whose sole result is the transfer of heat from a body of lower temperature to a body of higher temperature.

## §1.10 Second law of thermodynamics

- Direction of real processes? Reversible process, irreversible process.
- Second law of thermodynamics:
  - i **Clausius statement:** No process is possible whose sole result is the transfer of heat from a body of lower temperature to a body of higher temperature.
  - ii **Kelvin statement:** No process is possible in which the sole result is the absorption of heat from a reservoir and its complete conversion into work.

## §1.10 Second law of thermodynamics

- Direction of real processes? Reversible process, irreversible process.
- Second law of thermodynamics:
  - i **Clausius statement**: No process is possible whose sole result is the transfer of heat from a body of lower temperature to a body of higher temperature.
  - ii **Kelvin statement**: No process is possible in which the sole result is the absorption of heat from a reservoir and its complete conversion into work.
- Equivalent! To prove:  $\bar{A} \Leftrightarrow \bar{B} \Rightarrow A \Leftrightarrow B$ .

## §1.10 Second law of thermodynamics

- Direction of real processes? Reversible process, irreversible process.
- Second law of thermodynamics:
  - i **Clausius statement**: No process is possible whose sole result is the transfer of heat from a body of lower temperature to a body of higher temperature.
  - ii **Kelvin statement**: No process is possible in which the sole result is the absorption of heat from a reservoir and its complete conversion into work.
- Equivalent! To prove:  $\bar{A} \Leftrightarrow \bar{B} \Rightarrow A \Leftrightarrow B$ .
- Direction (spontaneously): Clausius – heat from high T to low T; Kelvin – work to heat.



## 1 Chpt 1. Basic Laws in Thermodynamics

- 1.1 Thermal equilibrium state
- 1.2 Law of thermal equilibrium and the temperature
- 1.3 Equation of state
- 1.4 Work
- 1.5 First law of thermodynamics
- 1.6 Thermal capacity and enthalpy
- 1.7 Internal energy of the ideal gas
- 1.8 Adiabatic process of the ideal gas
- 1.9 Carnot cycle in the ideal gas
- 1.10 Second law of thermodynamics
- **1.11 Carnot's theorem**
- 1.12 Thermodynamic temperature scale
- 1.13 Clausius theorem
- 1.14 Entropy
- 1.15 Entropy of ideal gas
- 1.16 Math expression of the 2nd law
- 1.17 Applications for the entropy increasing principle
- 1.18 Free energy and Gibbs function

## §1.11 Carnot's theorem

## §1.11 Carnot's theorem

- **Carnot's theorem:** All irreversible heat engines between two heat reservoirs are less efficient than a reversible engine operating between the same reservoirs.

## §1.11 Carnot's theorem

- **Carnot's theorem:** All irreversible heat engines between two heat reservoirs are less efficient than a reversible engine operating between the same reservoirs.
- Reversible process:  $\eta_A = \frac{W}{Q_1}$ .
- Any process:  $\eta_B = \frac{W'}{Q_1}$ ,  
then  $\eta_A \geq \eta_B$ .

## §1.11 Carnot's theorem

- **Carnot's theorem:** All irreversible heat engines between two heat reservoirs are less efficient than a reversible engine operating between the same reservoirs.
- Reversible process:  $\eta_A = \frac{W}{Q_1}$ .
- Any process:  $\eta_B = \frac{W'}{Q'_1}$ ,  
then  $\eta_A \geq \eta_B$ .
- Independent of working medium.

## 1 Chpt 1. Basic Laws in Thermodynamics

- 1.1 Thermal equilibrium state
- 1.2 Law of thermal equilibrium and the temperature
- 1.3 Equation of state
- 1.4 Work
- 1.5 First law of thermodynamics
- 1.6 Thermal capacity and enthalpy
- 1.7 Internal energy of the ideal gas
- 1.8 Adiabatic process of the ideal gas
- 1.9 Carnot cycle in the ideal gas
- 1.10 Second law of thermodynamics
- 1.11 Carnot's theorem
- **1.12 Thermodynamic temperature scale**
- 1.13 Clausius theorem
- 1.14 Entropy
- 1.15 Entropy of ideal gas
- 1.16 Math expression of the 2nd law
- 1.17 Applications for the entropy increasing principle
- 1.18 Free energy and Gibbs function

## §1.12 Thermodynamic temperature scale

## §1.12 Thermodynamic temperature scale

- $\eta = 1 - \frac{Q_2}{Q_1}$  only function of temperature  $(T_1, T_2)$ ;



## §1.12 Thermodynamic temperature scale

- $\eta = 1 - \frac{Q_2}{Q_1}$  only function of temperature  $(T_1, T_2)$ ;  
then  $\frac{Q_2}{Q_1} = F(T_1, T_2) \dots (1)$ ,  $\frac{Q_3}{Q_2} = F(T_2, T_3) \dots (2)$ ,  
 $\frac{Q_3}{Q_1} = F(T_1, T_3) \dots (3)$ ;

## §1.12 Thermodynamic temperature scale

- $\eta = 1 - \frac{Q_2}{Q_1}$  only function of temperature  $(T_1, T_2)$ ;  
 then  $\frac{Q_2}{Q_1} = F(T_1, T_2) \dots (1)$ ,  $\frac{Q_3}{Q_2} = F(T_2, T_3) \dots (2)$ ,  
 $\frac{Q_3}{Q_1} = F(T_1, T_3) \dots (3)$ ;  $\frac{(3)}{(1)}$ ,  $\frac{Q_3}{Q_2} = \frac{F(T_1, T_3)}{F(T_1, T_2)} \cdot (4)$

## §1.12 Thermodynamic temperature scale

- $\eta = 1 - \frac{Q_2}{Q_1}$  only function of temperature ( $T_1, T_2$ );  
 then  $\frac{Q_2}{Q_1} = F(T_1, T_2) \dots (1)$ ,  $\frac{Q_3}{Q_2} = F(T_2, T_3) \dots (2)$ ,  
 $\frac{Q_3}{Q_1} = F(T_1, T_3) \dots (3)$ ;  $\frac{(3)}{(1)}$ ,  $\frac{Q_3}{Q_2} = \frac{F(T_1, T_3)}{F(T_1, T_2)} \dots (4)$
- Considering (2) and (4), means (not need  $T_1$ )  
 $F(T_2, T_3) = f(T_3)/f(T_2) \Rightarrow \frac{Q_3}{Q_2} = \frac{f(T_3)}{f(T_2)}$ .

## §1.12 Thermodynamic temperature scale

- $\eta = 1 - \frac{Q_2}{Q_1}$  only function of temperature  $(T_1, T_2)$ ;  
 then  $\frac{Q_2}{Q_1} = F(T_1, T_2) \dots (1)$ ,  $\frac{Q_3}{Q_2} = F(T_2, T_3) \dots (2)$ ,  
 $\frac{Q_3}{Q_1} = F(T_1, T_3) \dots (3)$ ;  $\frac{(3)}{(1)}$ ,  $\frac{Q_3}{Q_2} = \frac{F(T_1, T_3)}{F(T_1, T_2)} \dots (4)$
- Considering (2) and (4), means (not need  $T_1$ )  
 $F(T_2, T_3) = f(T_3)/f(T_2) \Rightarrow \frac{Q_3}{Q_2} = \frac{f(T_3)}{f(T_2)}$ .
- It is possible to choose  $f(T) \propto T$ , which satisfies the requirement above.

## §1.12 Thermodynamic temperature scale

- $\eta = 1 - \frac{Q_2}{Q_1}$  only function of temperature ( $T_1, T_2$ );  
 then  $\frac{Q_2}{Q_1} = F(T_1, T_2) \dots (1)$ ,  $\frac{Q_3}{Q_2} = F(T_2, T_3) \dots (2)$ ,  
 $\frac{Q_3}{Q_1} = F(T_1, T_3) \dots (3)$ ;  $\frac{(3)}{(1)}$ ,  $\frac{Q_3}{Q_2} = \frac{F(T_1, T_3)}{F(T_1, T_2)} \dots (4)$
- Considering (2) and (4), means (not need  $T_1$ )  
 $F(T_2, T_3) = f(T_3)/f(T_2) \Rightarrow \frac{Q_3}{Q_2} = \frac{f(T_3)}{f(T_2)}$ .
- It is possible to choose  $f(T) \propto T$ , which satisfies the requirement above.
- $\frac{T_2}{T_1} \equiv \frac{Q_2}{Q_1}$  (Consistent with ideal gas temperature scale).

## §1.12 Thermodynamic temperature scale

- $\eta = 1 - \frac{Q_2}{Q_1}$  only function of temperature ( $T_1, T_2$ );  
 then  $\frac{Q_2}{Q_1} = F(T_1, T_2) \dots (1)$ ,  $\frac{Q_3}{Q_2} = F(T_2, T_3) \dots (2)$ ,  
 $\frac{Q_3}{Q_1} = F(T_1, T_3) \dots (3)$ ;  $\frac{(3)}{(1)}$ ,  $\frac{Q_3}{Q_2} = \frac{F(T_1, T_3)}{F(T_1, T_2)} \dots (4)$
- Considering (2) and (4), means (not need  $T_1$ )  
 $F(T_2, T_3) = f(T_3)/f(T_2) \Rightarrow \frac{Q_3}{Q_2} = \frac{f(T_3)}{f(T_2)}$ .
- It is possible to choose  $f(T) \propto T$ , which satisfies the requirement above.
- $\frac{T_2}{T_1} \equiv \frac{Q_2}{Q_1}$  (Consistent with ideal gas temperature scale).
- Together with water's triple point to be 273.16 K  $\rightarrow$  thermodynamic temperature scale.

## §1.12 Thermodynamic temperature scale

- $\eta = 1 - \frac{Q_2}{Q_1}$  only function of temperature  $(T_1, T_2)$ ;  
 then  $\frac{Q_2}{Q_1} = F(T_1, T_2) \dots (1)$ ,  $\frac{Q_3}{Q_2} = F(T_2, T_3) \dots (2)$ ,  
 $\frac{Q_3}{Q_1} = F(T_1, T_3) \dots (3)$ ;  $\frac{(3)}{(1)}$ ,  $\frac{Q_3}{Q_2} = \frac{F(T_1, T_3)}{F(T_1, T_2)} \dots (4)$
- Considering (2) and (4), means (not need  $T_1$ )  
 $F(T_2, T_3) = f(T_3)/f(T_2) \Rightarrow \frac{Q_3}{Q_2} = \frac{f(T_3)}{f(T_2)}$ .
- It is possible to choose  $f(T) \propto T$ , which satisfies the requirement above.
- $\frac{T_2}{T_1} \equiv \frac{Q_2}{Q_1}$  (Consistent with ideal gas temperature scale).
- Together with water's triple point to be 273.16 K  $\rightarrow$  thermodynamic temperature scale.
- Property: independent of the working medium.

## 1 Chpt 1. Basic Laws in Thermodynamics

- 1.1 Thermal equilibrium state
- 1.2 Law of thermal equilibrium and the temperature
- 1.3 Equation of state
- 1.4 Work
- 1.5 First law of thermodynamics
- 1.6 Thermal capacity and enthalpy
- 1.7 Internal energy of the ideal gas
- 1.8 Adiabatic process of the ideal gas

- 1.9 Carnot cycle in the ideal gas
- 1.10 Second law of thermodynamics
- 1.11 Carnot's theorem
- 1.12 Thermodynamic temperature scale
- **1.13 Clausius theorem**
- 1.14 Entropy
- 1.15 Entropy of ideal gas
- 1.16 Math expression of the 2nd law
- 1.17 Applications for the entropy increasing principle
- 1.18 Free energy and Gibbs function



## §1.13 Clausius theorem

## §1.13 Clausius theorem

- For any heat machine (reversible or irreversible):

$$\eta = 1 - \frac{Q_2}{Q_1} \leq 1 - \frac{T_2}{T_1},$$

## §1.13 Clausius theorem

- For any heat machine (reversible or irreversible):

$$\eta = 1 - \frac{Q_2}{Q_1} \leq 1 - \frac{T_2}{T_1},$$

- $\Rightarrow \frac{Q_1}{T_1} - \frac{Q_2}{T_2} \leq 0.$

## §1.13 Clausius theorem

- For any heat machine (reversible or irreversible):

$$\eta = 1 - \frac{Q_2}{Q_1} \leq 1 - \frac{T_2}{T_1},$$

- $\Rightarrow \frac{Q_1}{T_1} - \frac{Q_2}{T_2} \leq 0.$

- Redefine the heat  $Q$ : absorbed heat from the heat source:  $\frac{Q_1}{T_1} + \frac{Q_2}{T_2} \leq 0.$

## §1.13 Clausius theorem

- For any heat machine (reversible or irreversible):

$$\eta = 1 - \frac{Q_2}{Q_1} \leq 1 - \frac{T_2}{T_1},$$

- $\Rightarrow \frac{Q_1}{T_1} - \frac{Q_2}{T_2} \leq 0.$

- Redefine the heat  $Q$ : absorbed heat from the heat source:  $\frac{Q_1}{T_1} + \frac{Q_2}{T_2} \leq 0.$

- For  $n$  heat sources:  $\sum_i \frac{Q_i}{T_i} \leq 0 \dots$

## §1.13 Clausius theorem

- For any heat machine (reversible or irreversible):  
$$\eta = 1 - \frac{Q_2}{Q_1} \leq 1 - \frac{T_2}{T_1},$$
- $\Rightarrow \frac{Q_1}{T_1} - \frac{Q_2}{T_2} \leq 0.$
- Redefine the heat  $Q$ : absorbed heat from the heat source:  $\frac{Q_1}{T_1} + \frac{Q_2}{T_2} \leq 0.$
- For  $n$  heat sources:  $\sum_i \frac{Q_i}{T_i} \leq 0 \dots$
- More general process (Clausius theorem):

$$\oint \frac{\delta Q}{T} \leq 0.$$

## 1 Chpt 1. Basic Laws in Thermodynamics

- 1.1 Thermal equilibrium state
- 1.2 Law of thermal equilibrium and the temperature
- 1.3 Equation of state
- 1.4 Work
- 1.5 First law of thermodynamics
- 1.6 Thermal capacity and enthalpy
- 1.7 Internal energy of the ideal gas
- 1.8 Adiabatic process of the ideal gas

- 1.9 Carnot cycle in the ideal gas
- 1.10 Second law of thermodynamics
- 1.11 Carnot's theorem
- 1.12 Thermodynamic temperature scale
- 1.13 Clausius theorem
- **1.14 Entropy**
- 1.15 Entropy of ideal gas
- 1.16 Math expression of the 2nd law
- 1.17 Applications for the entropy increasing principle
- 1.18 Free energy and Gibbs function


## §1.14 Entropy

- For **reversible** process:  $\oint \frac{\mathrm{d}Q}{T} = 0$ .




## §1.14 Entropy

- For **reversible** process:  $\oint \frac{\delta Q}{T} = 0$ .



## §1.14 Entropy

- For **reversible** process:  $\oint \frac{\delta Q}{T} = 0$ .
- $\int_A^B \frac{\delta Q_R}{T} + \int_B^A \frac{\delta Q_{R'}}{T} = 0$ ,




## §1.14 Entropy

- For **reversible** process:  $\oint \frac{\delta Q}{T} = 0$ .

- $\int_A^B \frac{\delta Q_R}{T} + \int_B^A \frac{\delta Q_{R'}}{T} = 0,$

$$\Rightarrow \int_A^B \frac{\delta Q_R}{T} = - \int_B^A \frac{\delta Q_{R'}}{T} = \int_A^B \frac{\delta Q_{R'}}{T}.$$



# §1.14 Entropy


- For **reversible** process:  $\oint \frac{\delta Q}{T} = 0$ .

- $\int_A^B \frac{\delta Q_R}{T} + \int_B^A \frac{\delta Q_{R'}}{T} = 0,$

$$\Rightarrow \int_A^B \frac{\delta Q_R}{T} = - \int_B^A \frac{\delta Q_{R'}}{T} = \int_A^B \frac{\delta Q_{R'}}{T}.$$

- Considering that  $R$  and  $R'$  are **arbitrarily** reversible processes, there exists a function of state (Entropy):

$$S_B - S_A \equiv \int_A^B \frac{\delta Q_R}{T} \text{ or } dS \equiv \frac{\delta Q}{T}.$$



## §1.14 Entropy

- For **reversible** process:  $\oint \frac{\delta Q}{T} = 0$ .


- $\int_A^B \frac{\delta Q_R}{T} + \int_B^A \frac{\delta Q_{R'}}{T} = 0,$

$$\Rightarrow \int_A^B \frac{\delta Q_R}{T} = - \int_B^A \frac{\delta Q_{R'}}{T} = \int_A^B \frac{\delta Q_{R'}}{T}.$$

- Considering that  $R$  and  $R'$  are **arbitrarily** reversible processes, there exists a function of state (Entropy):

$$S_B - S_A \equiv \int_A^B \frac{\delta Q_R}{T} \text{ or } dS \equiv \frac{\delta Q}{T}.$$

- 1st law:  $dU = \delta Q + \delta W = \delta Q - pdV$   
 $\Rightarrow dS = \frac{dU + pdV}{T}, \text{ or } dU = TdS - pdV$   
 (Basic equation of thermodynamics).



## §1.14 Entropy

- For **reversible** process:  $\oint \frac{\delta Q}{T} = 0$ .


- $\int_A^B \frac{\delta Q_R}{T} + \int_B^A \frac{\delta Q_{R'}}{T} = 0,$

$$\Rightarrow \int_A^B \frac{\delta Q_R}{T} = - \int_B^A \frac{\delta Q_{R'}}{T} = \int_A^B \frac{\delta Q_{R'}}{T}.$$

- Considering that  $R$  and  $R'$  are **arbitrarily** reversible processes, there exists a function of state (Entropy):

$$S_B - S_A \equiv \int_A^B \frac{\delta Q_R}{T} \text{ or } dS \equiv \frac{\delta Q}{T}.$$

- 1st law:  $dU = \delta Q + \delta W = \delta Q - pdV$   
 $\Rightarrow dS = \frac{dU + pdV}{T}, \text{ or } dU = TdS - pdV$   
 (Basic equation of thermodynamics).
- $S = \sum S_i$  for extensible system.



## 1 Chpt 1. Basic Laws in Thermodynamics

- 1.1 Thermal equilibrium state
- 1.2 Law of thermal equilibrium and the temperature
- 1.3 Equation of state
- 1.4 Work
- 1.5 First law of thermodynamics
- 1.6 Thermal capacity and enthalpy
- 1.7 Internal energy of the ideal gas
- 1.8 Adiabatic process of the ideal gas
- 1.9 Carnot cycle in the ideal gas
- 1.10 Second law of thermodynamics
- 1.11 Carnot's theorem
- 1.12 Thermodynamic temperature scale
- 1.13 Clausius theorem
- 1.14 Entropy
- **1.15 Entropy of ideal gas**
- 1.16 Math expression of the 2nd law
- 1.17 Applications for the entropy increasing principle
- 1.18 Free energy and Gibbs function

# §1.15 Entropy of ideal gass



## §1.15 Entropy of ideal gas

- Extensibility of entropy  $S = nS_m$ .

## §1.15 Entropy of ideal gas

- Extensibility of entropy  $S = nS_m$ .
- For 1 mol ideal gas,  $dU_m = C_{V,m}dT$ ,  $pV_m = RT$ ,

## §1.15 Entropy of ideal gass

- Extensibility of entropy  $S = nS_m$ .
- For 1 mol ideal gas,  $dU_m = C_{V,m}dT$ ,  $pV_m = RT$ ,  
 $\Rightarrow dS_m = \frac{dU_m + pdV_m}{T} = \frac{C_{V,m}dT}{T} + \frac{(RT/V_m)dV_m}{T}$

## §1.15 Entropy of ideal gass

- Extensibility of entropy  $S = nS_m$ .
- For 1 mol ideal gas,  $dU_m = C_{V,m}dT$ ,  $pV_m = RT$ ,  
$$\Rightarrow dS_m = \frac{dU_m + pdV_m}{T} = \frac{C_{V,m}dT}{T} + \frac{(RT/V_m)dV_m}{T}$$
$$= C_{V,m}\frac{dT}{T} + R\frac{dV_m}{V_m}.$$

## §1.15 Entropy of ideal gass

- Extensibility of entropy  $S = nS_m$ .
- For 1 mol ideal gas,  $dU_m = C_{V,m}dT$ ,  $pV_m = RT$ ,  
$$\Rightarrow dS_m = \frac{dU_m + pdV_m}{T} = \frac{C_{V,m}dT}{T} + \frac{(RT/V_m)dV_m}{T}$$
$$= C_{V,m}\frac{dT}{T} + R\frac{dV_m}{V_m}.$$
- $S_m = C_{V,m} \ln T + R \ln V_m + S_{m,0}$ ,

## §1.15 Entropy of ideal gas

- Extensibility of entropy  $S = nS_m$ .
- For 1 mol ideal gas,  $dU_m = C_{V,m}dT$ ,  $pV_m = RT$ ,  
$$\Rightarrow dS_m = \frac{dU_m + pdV_m}{T} = \frac{C_{V,m}dT}{T} + \frac{(RT/V_m)dV_m}{T}$$
$$= C_{V,m}\frac{dT}{T} + R\frac{dV_m}{V_m}.$$
- $S_m = C_{V,m} \ln T + R \ln V_m + S_{m,0}$ ,  
or  $S_m = C_{p,m} \ln T - R \ln p + S_{m,0}$ .

## §1.15 Entropy of ideal gas

- Extensibility of entropy  $S = nS_m$ .
- For 1 mol ideal gas,  $dU_m = C_{V,m}dT$ ,  $pV_m = RT$ ,  
$$\Rightarrow dS_m = \frac{dU_m + pdV_m}{T} = \frac{C_{V,m}dT}{T} + \frac{(RT/V_m)dV_m}{T}$$
$$= C_{V,m}\frac{dT}{T} + R\frac{dV_m}{V_m}.$$
- $S_m = C_{V,m} \ln T + R \ln V_m + S_{m,0}$ ,  
or  $S_m = C_{p,m} \ln T - R \ln p + S_{m,0}$ .
- *Example*: ideal gas, isothermal process,  $V_A \rightarrow V_B$ .  
Entropy difference?

# §1.15 Entropy of ideal gas

- Extensibility of entropy  $S = nS_m$ .
- For 1 mol ideal gas,  $dU_m = C_{V,m}dT$ ,  $pV_m = RT$ ,  

$$\Rightarrow dS_m = \frac{dU_m + pdV_m}{T} = \frac{C_{V,m}dT}{T} + \frac{(RT/V_m)dV_m}{T}$$

$$= C_{V,m}\frac{dT}{T} + R\frac{dV_m}{V_m}.$$
- $S_m = C_{V,m} \ln T + R \ln V_m + S_{m,0}$ ,  
 or  $S_m = C_{p,m} \ln T - R \ln p + S_{m,0}$ .
- *Example*: ideal gas, isothermal process,  $V_A \rightarrow V_B$ .  
 Entropy difference?  
 Initial state  $(T, V_A)$ :  $S_A = C_V \ln T + nR \ln V_A + S_0$ ,



## §1.15 Entropy of ideal gas

- Extensibility of entropy  $S = nS_m$ .
- For 1 mol ideal gas,  $dU_m = C_{V,m}dT$ ,  $pV_m = RT$ ,  

$$\Rightarrow dS_m = \frac{dU_m + pdV_m}{T} = \frac{C_{V,m}dT}{T} + \frac{(RT/V_m)dV_m}{T}$$

$$= C_{V,m}\frac{dT}{T} + R\frac{dV_m}{V_m}.$$
- $S_m = C_{V,m} \ln T + R \ln V_m + S_{m,0}$ ,  
 or  $S_m = C_{p,m} \ln T - R \ln p + S_{m,0}$ .
- *Example*: ideal gas, isothermal process,  $V_A \rightarrow V_B$ .  
 Entropy difference?  
 Initial state  $(T, V_A)$ :  $S_A = C_V \ln T + nR \ln V_A + S_0$ ,  
 Final state  $(T, V_B)$ :  $S_B = C_V \ln T + nR \ln V_B + S_0$ ,

# §1.15 Entropy of ideal gas

- Extensibility of entropy  $S = nS_m$ .
- For 1 mol ideal gas,  $dU_m = C_{V,m}dT$ ,  $pV_m = RT$ ,  

$$\Rightarrow dS_m = \frac{dU_m + pdV_m}{T} = \frac{C_{V,m}dT}{T} + \frac{(RT/V_m)dV_m}{T}$$

$$= C_{V,m} \frac{dT}{T} + R \frac{dV_m}{V_m}.$$
- $S_m = C_{V,m} \ln T + R \ln V_m + S_{m,0}$ ,  
 or  $S_m = C_{p,m} \ln T - R \ln p + S_{m,0}$ .
- *Example*: ideal gas, isothermal process,  $V_A \rightarrow V_B$ .

Entropy difference?

Initial state  $(T, V_A)$ :  $S_A = C_V \ln T + nR \ln V_A + S_0$ ,

Final state  $(T, V_B)$ :  $S_B = C_V \ln T + nR \ln V_B + S_0$ ,

$$\Rightarrow S_B - S_A = nR \ln \frac{V_B}{V_A}.$$

## 1 Chpt 1. Basic Laws in Thermodynamics


- 1.1 Thermal equilibrium state
- 1.2 Law of thermal equilibrium and the temperature
- 1.3 Equation of state
- 1.4 Work
- 1.5 First law of thermodynamics
- 1.6 Thermal capacity and enthalpy
- 1.7 Internal energy of the ideal gas
- 1.8 Adiabatic process of the ideal gas
- 1.9 Carnot cycle in the ideal gas
- 1.10 Second law of thermodynamics
- 1.11 Carnot's theorem
- 1.12 Thermodynamic temperature scale
- 1.13 Clausius theorem
- 1.14 Entropy
- 1.15 Entropy of ideal gas
- **1.16 Math expression of the 2nd law**
- 1.17 Applications for the entropy increasing principle
- 1.18 Free energy and Gibbs function

## §1.16 Math expression of the 2nd law

- Clausius's theorem:  $\oint \frac{\delta Q}{T} \leq 0$ ,

## §1.16 Math expression of the 2nd law


- Clausius's theorem:  $\oint \frac{\delta Q}{T} \leq 0$ ,



## §1.16 Math expression of the 2nd law

- Clausius's theorem:  $\oint \frac{\delta Q}{T} \leq 0$ ,

$$\Rightarrow \int_A^B \frac{\delta Q}{T} + \int_B^A \frac{\delta Q_r}{T} \leq 0,$$




## §1.16 Math expression of the 2nd law

- Clausius's theorem:  $\oint \frac{\delta Q}{T} \leq 0$ ,

$$\Rightarrow \int_A^B \frac{\delta Q}{T} + \int_B^A \frac{\delta Q_r}{T} \leq 0,$$

$$\Rightarrow \int_A^B \frac{\delta Q}{T} + (S_A - S_B) \leq 0,$$




## §1.16 Math expression of the 2nd law

- Clausius's theorem:  $\oint \frac{\delta Q}{T} \leq 0$ ,

$$\Rightarrow \int_A^B \frac{\delta Q}{T} + \int_B^A \frac{\delta Q_r}{T} \leq 0,$$

$$\Rightarrow \int_A^B \frac{\delta Q}{T} + (S_A - S_B) \leq 0,$$

$$\Rightarrow S_B - S_A \geq \int_A^B \frac{\delta Q}{T}.$$





# §1.16 Math expression of the 2nd law


- Clausius's theorem:  $\oint \frac{\delta Q}{T} \leq 0$ ,

$$\Rightarrow \int_A^B \frac{\delta Q}{T} + \int_B^A \frac{\delta Q_r}{T} \leq 0,$$

$$\Rightarrow \int_A^B \frac{\delta Q}{T} + (S_A - S_B) \leq 0,$$

$$\Rightarrow S_B - S_A \geq \int_A^B \frac{\delta Q}{T}.$$

$$\therefore \boxed{dS \geq \frac{\delta Q}{T}} \text{ Math express of the 2nd law.}$$



# §1.16 Math expression of the 2nd law

- Clausius's theorem:  $\oint \frac{\delta Q}{T} \leq 0$ ,


$$\Rightarrow \int_A^B \frac{\delta Q}{T} + \int_B^A \frac{\delta Q_r}{T} \leq 0,$$

$$\Rightarrow \int_A^B \frac{\delta Q}{T} + (S_A - S_B) \leq 0,$$

$$\Rightarrow S_B - S_A \geq \int_A^B \frac{\delta Q}{T}.$$

$$\therefore \boxed{dS \geq \frac{\delta Q}{T}} \text{ Math express of the 2nd law.}$$

- Taking into account the 1st law ( $dU = \delta Q + \delta W$ ),



# §1.16 Math expression of the 2nd law

- Clausius's theorem:  $\oint \frac{\delta Q}{T} \leq 0$ ,


$$\Rightarrow \int_A^B \frac{\delta Q}{T} + \int_B^A \frac{\delta Q_r}{T} \leq 0,$$

$$\Rightarrow \int_A^B \frac{\delta Q}{T} + (S_A - S_B) \leq 0,$$

$$\Rightarrow S_B - S_A \geq \int_A^B \frac{\delta Q}{T}.$$

$$\therefore \boxed{dS \geq \frac{\delta Q}{T}} \text{ Math express of the 2nd law.}$$

- Taking into account the 1st law ( $dU = \delta Q + \delta W$ ),  
 $dU \leq TdS + \delta W$ .



Math expression of the 2nd law

$$dS \geq \frac{\delta Q}{T}$$

- For non-equilibrium state (extensible system):

$$S = \sum S_k.$$

Math expression of the 2nd law

$$dS \geq \frac{\delta Q}{T}$$

- For non-equilibrium state (extensible system):  
 $S = \sum S_k.$
- Each  $k$  can be taken as equilibrium state:

Math expression of the 2nd law

$$dS \geq \frac{\delta Q}{T}$$

- For non-equilibrium state (extensible system):

$$S = \sum S_k.$$

- Each  $k$  can be taken as equilibrium state:

$$S_{k,B} - S_{k,A} = \int_{A_k}^{B_k} \frac{\delta Q_r}{T}.$$

## Math expression of the 2nd law

$$dS \geq \frac{\delta Q}{T}$$

- For non-equilibrium state (extensible system):

$$S = \sum S_k.$$

- Each  $k$  can be taken as equilibrium state:

$$S_{k,B} - S_{k,A} = \int_{A_k}^{B_k} \frac{\delta Q_r}{T}.$$

- Clausius theorem:  $\int_A^B \frac{\delta Q}{T} + \sum_{k=1}^n \int_{B_k}^{A_k} \frac{\delta Q_r}{T} < 0.$

## Math expression of the 2nd law

$$dS \geq \frac{\delta Q}{T}$$

- For non-equilibrium state (extensible system):

$$S = \sum S_k.$$

- Each  $k$  can be taken as equilibrium state:

$$S_{k,B} - S_{k,A} = \int_{A_k}^{B_k} \frac{\delta Q_r}{T}.$$

- Clausius theorem:  $\int_A^B \frac{\delta Q}{T} + \sum_{k=1}^n \int_{B_k}^{A_k} \frac{\delta Q_r}{T} < 0.$

$$\Rightarrow S_B - S_A > \int_A^B \frac{\delta Q}{T}.$$

- For an adiabatic process:  $\delta Q = 0;$



## Math expression of the 2nd law

$$dS \geq \frac{\delta Q}{T}$$

- For non-equilibrium state (extensible system):

$$S = \sum S_k.$$

- Each  $k$  can be taken as equilibrium state:

$$S_{k,B} - S_{k,A} = \int_{A_k}^{B_k} \frac{\delta Q_r}{T}.$$

- Clausius theorem:  $\int_A^B \frac{\delta Q}{T} + \sum_{k=1}^n \int_{B_k}^{A_k} \frac{\delta Q_r}{T} < 0.$

$$\Rightarrow S_B - S_A > \int_A^B \frac{\delta Q}{T}.$$

- For an adiabatic process:  $\delta Q = 0$ ;  $dS \geq 0.$

## Math expression of the 2nd law

$$dS \geq \frac{\delta Q}{T}$$

- For non-equilibrium state (extensible system):

$$S = \sum S_k.$$

- Each  $k$  can be taken as equilibrium state:

$$S_{k,B} - S_{k,A} = \int_{A_k}^{B_k} \frac{\delta Q_r}{T}.$$

- Clausius theorem:  $\int_A^B \frac{\delta Q}{T} + \sum_{k=1}^n \int_{B_k}^{A_k} \frac{\delta Q_r}{T} < 0.$

$$\Rightarrow S_B - S_A > \int_A^B \frac{\delta Q}{T}.$$

- For an adiabatic process:  $\delta Q = 0$ ;  $dS \geq 0.$
- Theory of heat death!

## 1 Chpt 1. Basic Laws in Thermodynamics

- 1.1 Thermal equilibrium state
- 1.2 Law of thermal equilibrium and the temperature
- 1.3 Equation of state
- 1.4 Work
- 1.5 First law of thermodynamics
- 1.6 Thermal capacity and enthalpy
- 1.7 Internal energy of the ideal gas
- 1.8 Adiabatic process of the ideal gas
- 1.9 Carnot cycle in the ideal gas
- 1.10 Second law of thermodynamics
- 1.11 Carnot's theorem
- 1.12 Thermodynamic temperature scale
- 1.13 Clausius theorem
- 1.14 Entropy
- 1.15 Entropy of ideal gas
- 1.16 Math expression of the 2nd law
- **1.17 Applications for the entropy increasing principle**
- 1.18 Free energy and Gibbs function

## §1.17 Applications for the entropy increasing principle

- 1 Calculate the entropy difference before and after the reservoir A with temperature  $T_1$  transfer energy  $Q$  to reservoir B with temperature  $T_2$  ( $T_1 > T_2$ ).

## §1.17 Applications for the entropy increasing principle

- 1 Calculate the entropy difference before and after the reservoir A with temperature  $T_1$  transfer energy  $Q$  to reservoir B with temperature  $T_2$  ( $T_1 > T_2$ ).

- $dS = \frac{\delta Q}{T}$ ?

## §1.17 Applications for the entropy increasing principle

- 1 Calculate the entropy difference before and after the reservoir A with temperature  $T_1$  transfer energy  $Q$  to reservoir B with temperature  $T_2$  ( $T_1 > T_2$ ).
  - $dS = \frac{\delta Q}{T}$ ? Irreversible!

## §1.17 Applications for the entropy increasing principle

- 1 Calculate the entropy difference before and after the reservoir A with temperature  $T_1$  transfer energy  $Q$  to reservoir B with temperature  $T_2$  ( $T_1 > T_2$ ).
  - $dS = \frac{\delta Q}{T}$ ? Irreversible!
  - Assume 2 additional reservoirs with  $T_1$  (C) and  $T_2$  (D).

## §1.17 Applications for the entropy increasing principle

- 1 Calculate the entropy difference before and after the reservoir A with temperature  $T_1$  transfer energy  $Q$  to reservoir B with temperature  $T_2$  ( $T_1 > T_2$ ).
  - $dS = \frac{\delta Q}{T}$ ? Irreversible!
  - Assume 2 additional reservoirs with  $T_1$  (C) and  $T_2$  (D). i. Let  $Q$  transferred from A to C (reversible);



## §1.17 Applications for the entropy increasing principle

- 1 Calculate the entropy difference before and after the reservoir A with temperature  $T_1$  transfer energy  $Q$  to reservoir B with temperature  $T_2$  ( $T_1 > T_2$ ).
  - $dS = \frac{\delta Q}{T}$ ? Irreversible!
  - Assume 2 additional reservoirs with  $T_1$  (C) and  $T_2$  (D). i. Let  $Q$  transferred from A to C (reversible); ii. Let  $Q$  transferred from D to B (reversible).

## §1.17 Applications for the entropy increasing principle

- 1 Calculate the entropy difference before and after the reservoir A with temperature  $T_1$  transfer energy  $Q$  to reservoir B with temperature  $T_2$  ( $T_1 > T_2$ ).
  - $dS = \frac{\delta Q}{T}$ ? Irreversible!
  - Assume 2 additional reservoirs with  $T_1$  (C) and  $T_2$  (D). i. Let  $Q$  transferred from A to C (reversible); ii. Let  $Q$  transferred from D to B (reversible).
  - For A+B,

## §1.17 Applications for the entropy increasing principle

- 1 Calculate the entropy difference before and after the reservoir A with temperature  $T_1$  transfer energy  $Q$  to reservoir B with temperature  $T_2$  ( $T_1 > T_2$ ).
  - $dS = \frac{\delta Q}{T}$ ? Irreversible!
  - Assume 2 additional reservoirs with  $T_1$  (C) and  $T_2$  (D). i. Let  $Q$  transferred from A to C (reversible); ii. Let  $Q$  transferred from D to B (reversible).
  - For A+B, step i,  $\Delta S_1 = -\frac{Q}{T_1}$ ;

## §1.17 Applications for the entropy increasing principle

- 1 Calculate the entropy difference before and after the reservoir A with temperature  $T_1$  transfer energy  $Q$  to reservoir B with temperature  $T_2$  ( $T_1 > T_2$ ).
  - $dS = \frac{\delta Q}{T}$ ? Irreversible!
  - Assume 2 additional reservoirs with  $T_1$  (C) and  $T_2$  (D). i. Let  $Q$  transferred from A to C (reversible); ii. Let  $Q$  transferred from D to B (reversible).
  - For A+B, step i,  $\Delta S_1 = -\frac{Q}{T_1}$ ; step ii,  $\Delta S_2 = \frac{Q}{T_2}$ .

## §1.17 Applications for the entropy increasing principle

- 1 Calculate the entropy difference before and after the reservoir A with temperature  $T_1$  transfer energy  $Q$  to reservoir B with temperature  $T_2$  ( $T_1 > T_2$ ).
  - $dS = \frac{\delta Q}{T}$ ? Irreversible!
  - Assume 2 additional reservoirs with  $T_1$  (C) and  $T_2$  (D). i. Let  $Q$  transferred from A to C (reversible); ii. Let  $Q$  transferred from D to B (reversible).
  - For A+B, step i,  $\Delta S_1 = -\frac{Q}{T_1}$ ; step ii,  $\Delta S_2 = \frac{Q}{T_2}$ .
  - $\therefore \Delta S = \Delta S_1 + \Delta S_2 = Q(\frac{1}{T_2} - \frac{1}{T_1})$ .

## §1.17 Applications for the entropy increasing principle

1 Calculate the entropy difference before and after the reservoir A with temperature  $T_1$  transfer energy  $Q$  to reservoir B with temperature  $T_2$  ( $T_1 > T_2$ ).

- $dS = \frac{\delta Q}{T}$ ? Irreversible!
- Assume 2 additional reservoirs with  $T_1$  (C) and  $T_2$  (D). i. Let  $Q$  transferred from A to C (reversible); ii. Let  $Q$  transferred from D to B (reversible).
- For A+B, step i,  $\Delta S_1 = -\frac{Q}{T_1}$ ; step ii,  $\Delta S_2 = \frac{Q}{T_2}$ .
- $\therefore \Delta S = \Delta S_1 + \Delta S_2 = Q(\frac{1}{T_2} - \frac{1}{T_1})$ .
- As  $S$  a state function, which doesn't depend on the process, the directly contacting A and B also derives  $\Delta S = Q(\frac{1}{T_2} - \frac{1}{T_1}) > 0$ .

## §1.17 Applications for the entropy increasing principle

- 2 Mix two cups of water with same mass, different temperature  $T_1$  and  $T_2$  in the same pressure. Calculate the entropy difference  $\Delta S$ .

## §1.17 Applications for the entropy increasing principle

2 Mix two cups of water with same mass, different temperature  $T_1$  and  $T_2$  in the same pressure.

Calculate the entropy difference  $\Delta S$ .

- Initial state:  $(T_1, p)$ ,  $(T_2, p)$ ; final state:  $(\frac{T_1+T_2}{2}, p)$ .



## §1.17 Applications for the entropy increasing principle

- 2 Mix two cups of water with same mass, different temperature  $T_1$  and  $T_2$  in the same pressure. Calculate the entropy difference  $\Delta S$ .
- Initial state:  $(T_1, p)$ ,  $(T_2, p)$ ; final state:  $(\frac{T_1+T_2}{2}, p)$ . Still irreversible process.

## §1.17 Applications for the entropy increasing principle

2 Mix two cups of water with same mass, different temperature  $T_1$  and  $T_2$  in the same pressure.

Calculate the entropy difference  $\Delta S$ .

- Initial state:  $(T_1, p)$ ,  $(T_2, p)$ ; final state:  $(\frac{T_1+T_2}{2}, p)$ .  
Still irreversible process.

- Suppose existing a reversible process, then

$$dS = \frac{dU + pdV}{T} = \frac{dH}{T} = \frac{C_p dT}{T}.$$

## §1.17 Applications for the entropy increasing principle

2 Mix two cups of water with same mass, different temperature  $T_1$  and  $T_2$  in the same pressure.

Calculate the entropy difference  $\Delta S$ .

- Initial state:  $(T_1, p)$ ,  $(T_2, p)$ ; final state:  $(\frac{T_1+T_2}{2}, p)$ .  
Still irreversible process.

- Suppose existing a reversible process, then

$$dS = \frac{dU + pdV}{T} = \frac{dH}{T} = \frac{C_p dT}{T}.$$

- Cup 1,  $\Delta S_1 = \int_{T_1}^{\frac{T_1+T_2}{2}} C_p \frac{dT}{T}$

## §1.17 Applications for the entropy increasing principle

2 Mix two cups of water with same mass, different temperature  $T_1$  and  $T_2$  in the same pressure.

Calculate the entropy difference  $\Delta S$ .

- Initial state:  $(T_1, p)$ ,  $(T_2, p)$ ; final state:  $(\frac{T_1+T_2}{2}, p)$ .  
Still irreversible process.

- Suppose existing a reversible process, then

$$dS = \frac{dU + pdV}{T} = \frac{dH}{T} = \frac{C_p dT}{T}.$$

- Cup 1,  $\Delta S_1 = \int_{T_1}^{\frac{T_1+T_2}{2}} C_p \frac{dT}{T} = C_p \ln \frac{T_1+T_2}{2T_1};$

## §1.17 Applications for the entropy increasing principle

2 Mix two cups of water with same mass, different temperature  $T_1$  and  $T_2$  in the same pressure.

Calculate the entropy difference  $\Delta S$ .

- Initial state:  $(T_1, p)$ ,  $(T_2, p)$ ; final state:  $(\frac{T_1+T_2}{2}, p)$ .  
Still irreversible process.

- Suppose existing a reversible process, then

$$dS = \frac{dU + pdV}{T} = \frac{dH}{T} = \frac{C_p dT}{T}.$$

- Cup 1,  $\Delta S_1 = \int_{T_1}^{\frac{T_1+T_2}{2}} C_p \frac{dT}{T} = C_p \ln \frac{T_1+T_2}{2T_1};$

$$\text{Cup 2, } \Delta S_2 = \int_{T_2}^{\frac{T_1+T_2}{2}} C_p \frac{dT}{T}$$

## §1.17 Applications for the entropy increasing principle

2 Mix two cups of water with same mass, different temperature  $T_1$  and  $T_2$  in the same pressure.

Calculate the entropy difference  $\Delta S$ .

- Initial state:  $(T_1, p)$ ,  $(T_2, p)$ ; final state:  $(\frac{T_1+T_2}{2}, p)$ .  
Still irreversible process.

- Suppose existing a reversible process, then

$$dS = \frac{dU + pdV}{T} = \frac{dH}{T} = \frac{C_p dT}{T}.$$

- Cup 1,  $\Delta S_1 = \int_{T_1}^{\frac{T_1+T_2}{2}} C_p \frac{dT}{T} = C_p \ln \frac{T_1+T_2}{2T_1};$

$$\text{Cup 2, } \Delta S_2 = \int_{T_2}^{\frac{T_1+T_2}{2}} C_p \frac{dT}{T} = C_p \ln \frac{T_1+T_2}{2T_2};$$

## §1.17 Applications for the entropy increasing principle

2 Mix two cups of water with same mass, different temperature  $T_1$  and  $T_2$  in the same pressure.

Calculate the entropy difference  $\Delta S$ .

- Initial state:  $(T_1, p)$ ,  $(T_2, p)$ ; final state:  $(\frac{T_1+T_2}{2}, p)$ .  
Still irreversible process.

- Suppose existing a reversible process, then

$$dS = \frac{dU+pdV}{T} = \frac{dH}{T} = \frac{C_p dT}{T}.$$

- Cup 1,  $\Delta S_1 = \int_{T_1}^{\frac{T_1+T_2}{2}} C_p \frac{dT}{T} = C_p \ln \frac{T_1+T_2}{2T_1};$

$$\text{Cup 2, } \Delta S_2 = \int_{T_2}^{\frac{T_1+T_2}{2}} C_p \frac{dT}{T} = C_p \ln \frac{T_1+T_2}{2T_2};$$

$$\therefore \Delta S = \Delta S_1 + \Delta S_2$$

## §1.17 Applications for the entropy increasing principle

2 Mix two cups of water with same mass, different temperature  $T_1$  and  $T_2$  in the same pressure.

Calculate the entropy difference  $\Delta S$ .

- Initial state:  $(T_1, p)$ ,  $(T_2, p)$ ; final state:  $(\frac{T_1+T_2}{2}, p)$ .  
Still irreversible process.

- Suppose existing a reversible process, then

$$dS = \frac{dU + pdV}{T} = \frac{dH}{T} = \frac{C_p dT}{T}.$$

- Cup 1,  $\Delta S_1 = \int_{T_1}^{\frac{T_1+T_2}{2}} C_p \frac{dT}{T} = C_p \ln \frac{T_1+T_2}{2T_1};$

$$\text{Cup 2, } \Delta S_2 = \int_{T_2}^{\frac{T_1+T_2}{2}} C_p \frac{dT}{T} = C_p \ln \frac{T_1+T_2}{2T_2};$$

$$\therefore \Delta S = \Delta S_1 + \Delta S_2 = C_p \ln \frac{(T_1+T_2)^2}{4T_1T_2}.$$



## §1.17 Applications for the entropy increasing principle

2 Mix two cups of water with same mass, different temperature  $T_1$  and  $T_2$  in the same pressure.

Calculate the entropy difference  $\Delta S$ .

- Initial state:  $(T_1, p)$ ,  $(T_2, p)$ ; final state:  $(\frac{T_1+T_2}{2}, p)$ .  
Still irreversible process.

- Suppose existing a reversible process, then

$$dS = \frac{dU+pdV}{T} = \frac{dH}{T} = \frac{C_p dT}{T}.$$

- Cup 1,  $\Delta S_1 = \int_{T_1}^{\frac{T_1+T_2}{2}} C_p \frac{dT}{T} = C_p \ln \frac{T_1+T_2}{2T_1}$ ;

$$\text{Cup 2, } \Delta S_2 = \int_{T_2}^{\frac{T_1+T_2}{2}} C_p \frac{dT}{T} = C_p \ln \frac{T_1+T_2}{2T_2};$$

$$\therefore \Delta S = \Delta S_1 + \Delta S_2 = C_p \ln \frac{(T_1+T_2)^2}{4T_1T_2}.$$

- If  $T_1 \neq T_2$ ,  $\Delta S > 0$ ;

## §1.17 Applications for the entropy increasing principle

2 Mix two cups of water with same mass, different temperature  $T_1$  and  $T_2$  in the same pressure.

Calculate the entropy difference  $\Delta S$ .

- Initial state:  $(T_1, p)$ ,  $(T_2, p)$ ; final state:  $(\frac{T_1+T_2}{2}, p)$ .  
Still irreversible process.

- Suppose existing a reversible process, then

$$dS = \frac{dU+pdV}{T} = \frac{dH}{T} = \frac{C_p dT}{T}.$$

- Cup 1,  $\Delta S_1 = \int_{T_1}^{\frac{T_1+T_2}{2}} C_p \frac{dT}{T} = C_p \ln \frac{T_1+T_2}{2T_1}$ ;

$$\text{Cup 2, } \Delta S_2 = \int_{T_2}^{\frac{T_1+T_2}{2}} C_p \frac{dT}{T} = C_p \ln \frac{T_1+T_2}{2T_2};$$

$$\therefore \Delta S = \Delta S_1 + \Delta S_2 = C_p \ln \frac{(T_1+T_2)^2}{4T_1T_2}.$$

- If  $T_1 \neq T_2$ ,  $\Delta S > 0$ ; if  $T_1 = T_2$ ,  $\Delta S = 0$ .

## §1.17 Applications for the entropy increasing principle

- 3 Ideal gas with initial temperature  $T$ , volume  $V_A$ , adiabatic and free expansion to  $V_B$ . Calculate  $\Delta S$ .

## §1.17 Applications for the entropy increasing principle

- 3 Ideal gas with initial temperature  $T$ , volume  $V_A$ ,  
adiabatic and free expansion to  $V_B$ . Calculate  $\Delta S$ .
- Free expansion,  $W = 0$ ; adiabatic,  $Q = 0$ .

## §1.17 Applications for the entropy increasing principle

- 3 Ideal gas with initial temperature  $T$ , volume  $V_A$ , adiabatic and free expansion to  $V_B$ . Calculate  $\Delta S$ .
- Free expansion,  $W = 0$ ; adiabatic,  $Q = 0$ .  
 $\rightarrow \Delta U = W + Q = 0$ ,

## §1.17 Applications for the entropy increasing principle

- 3 Ideal gas with initial temperature  $T$ , volume  $V_A$ , adiabatic and free expansion to  $V_B$ . Calculate  $\Delta S$ .
- Free expansion,  $W = 0$ ; adiabatic,  $Q = 0$ .  
 $\rightarrow \Delta U = W + Q = 0$ ,  $\rightarrow T = \text{Const.}$

## §1.17 Applications for the entropy increasing principle

- 3 Ideal gas with initial temperature  $T$ , volume  $V_A$ , adiabatic and free expansion to  $V_B$ . Calculate  $\Delta S$ .
- Free expansion,  $W = 0$ ; adiabatic,  $Q = 0$ .  
 $\rightarrow \Delta U = W + Q = 0, \rightarrow T = \text{Const.}$
  - According to (1.15.4) –  
$$S = nC_{V,m} \ln T + nR \ln V + S_0,$$

## §1.17 Applications for the entropy increasing principle

3 Ideal gas with initial temperature  $T$ , volume  $V_A$ , adiabatic and free expansion to  $V_B$ . Calculate  $\Delta S$ .

- Free expansion,  $W = 0$ ; adiabatic,  $Q = 0$ .  
 $\rightarrow \Delta U = W + Q = 0$ ,  $\rightarrow T = \text{Const.}$

- According to (1.15.4) –

$$S = nC_{V,m} \ln T + nR \ln V + S_0, \text{ so,}$$

$$S_A = C_V \ln T + nR \ln V_A + S_0,$$

$$S_B = C_V \ln T + nR \ln V_B + S_0.$$



## §1.17 Applications for the entropy increasing principle

- 3 Ideal gas with initial temperature  $T$ , volume  $V_A$ , adiabatic and free expansion to  $V_B$ . Calculate  $\Delta S$ .
- Free expansion,  $W = 0$ ; adiabatic,  $Q = 0$ .  
 $\rightarrow \Delta U = W + Q = 0, \rightarrow T = \text{Const.}$
  - According to (1.15.4) –  
 $S = nC_{V,m} \ln T + nR \ln V + S_0$ , so,  
 $S_A = C_V \ln T + nR \ln V_A + S_0$ ,  
 $S_B = C_V \ln T + nR \ln V_B + S_0$ .
  - $\Delta S = S_B - S_A = nR \ln \frac{V_B}{V_A}$ .

## §1.17 Applications for the entropy increasing principle

- 3 Ideal gas with initial temperature  $T$ , volume  $V_A$ , adiabatic and free expansion to  $V_B$ . Calculate  $\Delta S$ .
- Free expansion,  $W = 0$ ; adiabatic,  $Q = 0$ .  
 $\rightarrow \Delta U = W + Q = 0$ ,  $\rightarrow T = \text{Const.}$
  - According to (1.15.4) –  
 $S = nC_{V,m} \ln T + nR \ln V + S_0$ , so,  
 $S_A = C_V \ln T + nR \ln V_A + S_0$ ,  
 $S_B = C_V \ln T + nR \ln V_B + S_0$ .
  - $\Delta S = S_B - S_A = nR \ln \frac{V_B}{V_A}$ .
  - How about if not “free”?

## §1.17 Applications for the entropy increasing principle

$$S = nC_{V,m} \ln T + nR \ln V + S_0,$$

$$S_A = C_V \ln T + nR \ln V_A + S_0$$

- How about if not “free”? In case: Adiabatic expansion.

## §1.17 Applications for the entropy increasing principle

$$S = nC_{V,m} \ln T + nR \ln V + S_0,$$

$$S_A = C_V \ln T + nR \ln V_A + S_0$$

- How about if not “free”? In case: Adiabatic expansion.  
 $T$  changes.

## §1.17 Applications for the entropy increasing principle

$$S = nC_{V,m} \ln T + nR \ln V + S_0,$$

$$S_A = C_V \ln T + nR \ln V_A + S_0$$

- How about if not “free”? In case: Adiabatic expansion.  
 $T$  changes.
- For adiabatic process,  $T_A V_A^{\gamma-1} = T_B V_B^{\gamma-1}$

## §1.17 Applications for the entropy increasing principle

$$S = nC_{V,m} \ln T + nR \ln V + S_0,$$

$$S_A = C_V \ln T + nR \ln V_A + S_0$$

- How about if not “free”? In case: Adiabatic expansion.  
 $T$  changes.
- For adiabatic process,  $T_A V_A^{\gamma-1} = T_B V_B^{\gamma-1}$   
 $\Rightarrow T_B = (V_A/V_B)^{\gamma-1} T.$

## §1.17 Applications for the entropy increasing principle

$$S = nC_{V,m} \ln T + nR \ln V + S_0,$$

$$S_A = C_V \ln T + nR \ln V_A + S_0$$

- How about if not “free”? In case: Adiabatic expansion.

$T$  changes.

- For adiabatic process,  $T_A V_A^{\gamma-1} = T_B V_B^{\gamma-1}$

$$\Rightarrow T_B = (V_A/V_B)^{\gamma-1} T.$$

$$\therefore S_B = C_V \ln[(V_A/V_B)^{\gamma-1} T] + nR \ln V_B + S_0$$

## §1.17 Applications for the entropy increasing principle

$$S = nC_{V,m} \ln T + nR \ln V + S_0,$$

$$S_A = C_V \ln T + nR \ln V_A + S_0$$

- How about if not “free”? In case: Adiabatic expansion.

$T$  changes.

- For adiabatic process,  $T_A V_A^{\gamma-1} = T_B V_B^{\gamma-1}$

$$\Rightarrow T_B = (V_A/V_B)^{\gamma-1} T.$$

$$\therefore S_B = C_V \ln[(V_A/V_B)^{\gamma-1} T] + nR \ln V_B + S_0$$

- $\Delta S = S_B - S_A$



## §1.17 Applications for the entropy increasing principle

$$S = nC_{V,m} \ln T + nR \ln V + S_0,$$

$$S_A = \cancel{C_V \ln T} + nR \ln V_A + \cancel{S_0}$$

- How about if not “free”? In case: Adiabatic expansion.  
 $T$  changes.
- For adiabatic process,  $T_A V_A^{\gamma-1} = T_B V_B^{\gamma-1}$   
 $\Rightarrow T_B = (V_A/V_B)^{\gamma-1} T$ .  
 $\therefore S_B = \cancel{C_V \ln [(V_A/V_B)^{\gamma-1} T]} + nR \ln V_B + \cancel{S_0}$
- $\Delta S = S_B - S_A$   
 $= C_V(\gamma - 1) \ln(V_A/V_B) + nR \ln(V_B/V_A)$

## §1.17 Applications for the entropy increasing principle

$$S = nC_{V,m} \ln T + nR \ln V + S_0,$$

$$S_A = C_V \ln T + nR \ln V_A + S_0$$

- How about if not “free”? In case: Adiabatic expansion.

$T$  changes.

- For adiabatic process,  $T_A V_A^{\gamma-1} = T_B V_B^{\gamma-1}$

$$\Rightarrow T_B = (V_A/V_B)^{\gamma-1} T.$$

$$\therefore S_B = C_V \ln[(V_A/V_B)^{\gamma-1} T] + nR \ln V_B + S_0$$

- $\Delta S = S_B - S_A$   
 $= C_V(\gamma - 1) \ln(V_A/V_B) + nR \ln(V_B/V_A)$   
 $= (C_p - C_V) \ln(V_A/V_B) - nR \ln(V_A/V_B)$

## §1.17 Applications for the entropy increasing principle

$$S = nC_{V,m} \ln T + nR \ln V + S_0,$$

$$S_A = C_V \ln T + nR \ln V_A + S_0$$

- How about if not “free”? In case: Adiabatic expansion.

$T$  changes.

- For adiabatic process,  $T_A V_A^{\gamma-1} = T_B V_B^{\gamma-1}$

$$\Rightarrow T_B = (V_A/V_B)^{\gamma-1} T.$$

$$\therefore S_B = C_V \ln[(V_A/V_B)^{\gamma-1} T] + nR \ln V_B + S_0$$

- $\Delta S = S_B - S_A$   
 $= C_V(\gamma - 1) \ln(V_A/V_B) + nR \ln(V_B/V_A)$   
 $= (C_p - C_V) \ln(V_A/V_B) - nR \ln(V_A/V_B)$   
 $= 0.$

## 1 Chpt 1. Basic Laws in Thermodynamics

- 1.1 Thermal equilibrium state
- 1.2 Law of thermal equilibrium and the temperature
- 1.3 Equation of state
- 1.4 Work
- 1.5 First law of thermodynamics
- 1.6 Thermal capacity and enthalpy
- 1.7 Internal energy of the ideal gas
- 1.8 Adiabatic process of the ideal gas
- 1.9 Carnot cycle in the ideal gas
- 1.10 Second law of thermodynamics
- 1.11 Carnot's theorem
- 1.12 Thermodynamic temperature scale
- 1.13 Clausius theorem
- 1.14 Entropy
- 1.15 Entropy of ideal gas
- 1.16 Math expression of the 2nd law
- 1.17 Applications for the entropy increasing principle
- **1.18 Free energy and Gibbs function**

# §1.18 Free energy and Gibbs function

## §1.18 Free energy and Gibbs function

- Isothermal process:  $(dS \geq \delta Q/T)$

## §1.18 Free energy and Gibbs function

- Isothermal process:  $(dS \geq \delta Q/T)$
- $S_B - S_A \geq \int_A^B \frac{\delta Q}{T} = \frac{Q}{T}, \Rightarrow Q \leq T(S_B - S_A),$

## §1.18 Free energy and Gibbs function

- Isothermal process:  $(dS \geq \delta Q/T)$
- $S_B - S_A \geq \int_A^B \frac{\delta Q}{T} = \frac{Q}{T}, \Rightarrow Q \leq T(S_B - S_A),$
- $\xrightarrow{U_B - U_A = Q + W} -W \leq (U_A - U_B) - T(S_A - S_B)$



## §1.18 Free energy and Gibbs function

- Isothermal process:  $(dS \geq \delta Q/T)$
- $S_B - S_A \geq \int_A^B \frac{\delta Q}{T} = \frac{Q}{T}, \Rightarrow Q \leq T(S_B - S_A),$
- $\xrightarrow{U_B - U_A = Q + W} -W \leq (U_A - U_B) - T(S_A - S_B)$   
upper limit for the output work.

## §1.18 Free energy and Gibbs function

- Isothermal process:  $(dS \geq \delta Q/T)$
- $S_B - S_A \geq \int_A^B \frac{\delta Q}{T} = \frac{Q}{T}, \Rightarrow Q \leq T(S_B - S_A),$
- $\xrightarrow{U_B - U_A = Q + W} -W \leq (U_A - U_B) - T(S_A - S_B)$   
upper limit for the output work.
- Free energy:  $F \equiv U - TS,$

# §1.18 Free energy and Gibbs function

- Isothermal process:  $(dS \geq \delta Q/T)$
- $S_B - S_A \geq \int_A^B \frac{\delta Q}{T} = \frac{Q}{T}, \Rightarrow Q \leq T(S_B - S_A),$
- $\xrightarrow{U_B - U_A = Q + W} -W \leq (U_A - U_B) - T(S_A - S_B)$   
upper limit for the output work.
- Free energy:  $F \equiv U - TS$ , then  $-W \leq F_A - F_B$ .

## §1.18 Free energy and Gibbs function

- Isothermal process: ( $dS \geq \delta Q/T$ )
- $S_B - S_A \geq \int_A^B \frac{\delta Q}{T} = \frac{Q}{T}, \Rightarrow Q \leq T(S_B - S_A),$
- $\xrightarrow{U_B - U_A = Q + W} -W \leq (U_A - U_B) - T(S_A - S_B)$   
upper limit for the output work.
- Free energy:  $F \equiv U - TS$ , then  $-W \leq F_A - F_B$ .
- If also isochoric ( $dV = 0, W = 0$ ), i.e.,  $F_B - F_A \leq 0$ .

## §1.18 Free energy and Gibbs function

- Isothermal process: ( $dS \geq \delta Q/T$ )
- $S_B - S_A \geq \int_A^B \frac{\delta Q}{T} = \frac{Q}{T}, \Rightarrow Q \leq T(S_B - S_A),$
- $\xrightarrow{U_B - U_A = Q + W} -W \leq (U_A - U_B) - T(S_A - S_B)$   
upper limit for the output work.
- Free energy:  $F \equiv U - TS$ , then  $-W \leq F_A - F_B$ .
- If also isochoric ( $dV = 0, W = 0$ ), i.e.,  $F_B - F_A \leq 0$ .
- Or if also isobaric ( $p = \text{const.}, W = -p(V_B - V_A)$ ),

## §1.18 Free energy and Gibbs function

- Isothermal process:  $(dS \geq \delta Q/T)$
- $S_B - S_A \geq \int_A^B \frac{\delta Q}{T} = \frac{Q}{T}, \Rightarrow Q \leq T(S_B - S_A),$
- $\xrightarrow{U_B - U_A = Q + W} -W \leq (U_A - U_B) - T(S_A - S_B)$   
upper limit for the output work.
- Free energy:  $F \equiv U - TS$ , then  $-W \leq F_A - F_B$ .
- If also isochoric ( $dV = 0, W = 0$ ), i.e.,  $F_B - F_A \leq 0$ .
- Or if also isobaric ( $p = \text{const.}, W = -p(V_B - V_A)$ ),  
then  $p(V_B - V_A) \leq F_A - F_B$ ,

# §1.18 Free energy and Gibbs function

- Isothermal process: ( $dS \geq \delta Q/T$ )
- $S_B - S_A \geq \int_A^B \frac{\delta Q}{T} = \frac{Q}{T}, \Rightarrow Q \leq T(S_B - S_A),$
- $\xrightarrow{U_B - U_A = Q + W} -W \leq (U_A - U_B) - T(S_A - S_B)$   
upper limit for the output work.
- Free energy:  $F \equiv U - TS$ , then  $-W \leq F_A - F_B$ .
- If also isochoric ( $dV = 0, W = 0$ ), i.e.,  $F_B - F_A \leq 0$ .
- Or if also isobaric ( $p = \text{const.}, W = -p(V_B - V_A)$ ),  
then  $p(V_B - V_A) \leq F_A - F_B$ ,
- define Gibbs function:  
 $G \equiv F + pV (= U - TS + pV),$

# §1.18 Free energy and Gibbs function

- Isothermal process:  $(dS \geq \delta Q/T)$
- $S_B - S_A \geq \int_A^B \frac{\delta Q}{T} = \frac{Q}{T}, \Rightarrow Q \leq T(S_B - S_A),$
- $\xrightarrow{U_B - U_A = Q + W} -W \leq (U_A - U_B) - T(S_A - S_B)$   
upper limit for the output work.
- Free energy:  $F \equiv U - TS$ , then  $-W \leq F_A - F_B$ .
- If also isochoric ( $dV = 0, W = 0$ ), i.e.,  $F_B - F_A \leq 0$ .
- Or if also isobaric ( $p = \text{const.}, W = -p(V_B - V_A)$ ),  
then  $p(V_B - V_A) \leq F_A - F_B$ ,
- define Gibbs function:  
 $G \equiv F + pV (= U - TS + pV)$ , then  $G_B - G_A \leq 0$ .



# §1.18 Free energy and Gibbs function

- Isothermal process:  $(dS \geq \delta Q/T)$
- $S_B - S_A \geq \int_A^B \frac{\delta Q}{T} = \frac{Q}{T}, \Rightarrow Q \leq T(S_B - S_A),$
- $\xrightarrow{U_B - U_A = Q + W} -W \leq (U_A - U_B) - T(S_A - S_B)$   
upper limit for the output work.
- Free energy:  $F \equiv U - TS$ , then  $-W \leq F_A - F_B$ .
- If also isochoric ( $dV = 0, W = 0$ ), i.e.,  $F_B - F_A \leq 0$ .
- Or if also isobaric ( $p = \text{const.}, W = -p(V_B - V_A)$ ),  
then  $p(V_B - V_A) \leq F_A - F_B$ ,
- define Gibbs function:  
 $G \equiv F + pV (= U - TS + pV)$ , then  $G_B - G_A \leq 0$ .
- Summary: adiabatic  $dS \geq 0$ ; isothermal + isochoric  $dF \leq 0$ ; isothermal + isobaric  $dG \leq 0$ .

# §1.18 Free energy and Gibbs function

- Isothermal process: ( $dS \geq \delta Q/T$ )
- $S_B - S_A \geq \int_A^B \frac{\delta Q}{T} = \frac{Q}{T}, \Rightarrow Q \leq T(S_B - S_A),$
- $\xrightarrow{U_B - U_A = Q + W} -W \leq (U_A - U_B) - T(S_A - S_B)$   
upper limit for the output work.
- Free energy:  $F \equiv U - TS$ , then  $-W \leq F_A - F_B$ .
- If also isochoric ( $dV = 0, W = 0$ ), i.e.,  $F_B - F_A \leq 0$ .
- Or if also isobaric ( $p = \text{const.}, W = -p(V_B - V_A)$ ),  
then  $p(V_B - V_A) \leq F_A - F_B$ ,
- define Gibbs function:  
 $G \equiv F + pV (= U - TS + pV)$ , then  $G_B - G_A \leq 0$ .
- Summary: adiabatic  $dS \geq 0$ ; isothermal + isochoric  $dF \leq 0$ ; isothermal + isobaric  $dG \leq 0$ .

# Table of contents I

## ① Chpt 1. Basic Laws in Thermodynamics

- 1.1 Thermal equilibrium state
- 1.2 Law of thermal equilibrium and the temperature
- 1.3 Equation of state
- 1.4 Work
- 1.5 First law of thermodynamics
- 1.6 Thermal capacity and enthalpy
- 1.7 Internal energy of the ideal gas
- 1.8 Adiabatic process of the ideal gas
- 1.9 Carnot cycle in the ideal gas
- 1.10 Second law of thermodynamics
- 1.11 Carnot's theorem

## Table of contents II

- 1.12 Thermodynamic temperature scale
- 1.13 Clausius theorem
- 1.14 Entropy
- 1.15 Entropy of ideal gas
- 1.16 Math expression of the 2nd law
- 1.17 Applications for the entropy increasing principle
- 1.18 Free energy and Gibbs function