

Thermodynamics & Statistical Physics

Chapter 8. Bose statistics and Fermi statistics

Yuan-Chuan Zou
zouyc@hust.edu.cn

School of Physics, Huazhong University of Science and Technology

December 30, 2013

Table of contents

1 §8. Bose statistics and Fermi statistics

- 8.1 Thermal parameters in statistics
- 8.2 U of weak degenerated ideal Fermion(Boson) gas
- 8.3 Bose-Einstein Condensate (BEC)
- 8.4 Photon gas
- 8.5 Free electron gas in metal

8.1 Thermal parameters in statistics for Boson system

8.1 Thermal parameters in statistics for Boson system

- For Boson system, $a_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1}$.

8.1 Thermal parameters in statistics for Boson system

- For Boson system, $a_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1}$.
Define Grand partition function:

8.1 Thermal parameters in statistics for Boson system

- For Boson system, $a_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1}$.

Define Grand partition function:

$$\mathcal{Z} = \prod \mathcal{Z}_l = \prod (1 - e^{-\alpha-\beta\varepsilon_l})^{-\omega_l}.$$

8.1 Thermal parameters in statistics for Boson system

- For Boson system, $a_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1}$.

Define Grand partition function:

$$\mathcal{Z} = \prod \mathcal{Z}_l = \prod (1 - e^{-\alpha-\beta\varepsilon_l})^{-\omega_l}.$$

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha-\beta\varepsilon_l}).$$

8.1 Thermal parameters in statistics for Boson system

- For Boson system, $a_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1}$.

Define Grand partition function:

$$\mathcal{Z} = \prod \mathcal{Z}_l = \prod (1 - e^{-\alpha-\beta\varepsilon_l})^{-\omega_l}.$$

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha-\beta\varepsilon_l}).$$

- $\frac{\partial}{\partial \alpha} \ln \mathcal{Z}$

8.1 Thermal parameters in statistics for Boson system

- For Boson system, $a_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1}$.

Define Grand partition function:

$$\mathcal{Z} = \prod \mathcal{Z}_l = \prod (1 - e^{-\alpha-\beta\varepsilon_l})^{-\omega_l}.$$

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha-\beta\varepsilon_l}).$$

- $\frac{\partial}{\partial \alpha} \ln \mathcal{Z} = - \sum \frac{\omega_l}{1 - e^{-\alpha-\beta\varepsilon_l}} \cdot e^{-\alpha-\beta\varepsilon_l}$

8.1 Thermal parameters in statistics for Boson system

- For Boson system, $a_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1}$.

Define Grand partition function:

$$\mathcal{Z} = \prod \mathcal{Z}_l = \prod (1 - e^{-\alpha-\beta\varepsilon_l})^{-\omega_l}.$$

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha-\beta\varepsilon_l}).$$

- $\frac{\partial}{\partial \alpha} \ln \mathcal{Z} = - \sum \frac{\omega_l}{1 - e^{-\alpha-\beta\varepsilon_l}} \cdot e^{-\alpha-\beta\varepsilon_l} = - \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1}$

8.1 Thermal parameters in statistics for Boson system

- For Boson system, $a_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1}$.

Define Grand partition function:

$$\mathcal{Z} = \prod \mathcal{Z}_l = \prod (1 - e^{-\alpha-\beta\varepsilon_l})^{-\omega_l}.$$

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha-\beta\varepsilon_l}).$$

- $\frac{\partial}{\partial \alpha} \ln \mathcal{Z} = - \sum \frac{\omega_l}{1-e^{-\alpha-\beta\varepsilon_l}} \cdot e^{-\alpha-\beta\varepsilon_l} = - \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1}$
 $= - \sum a_l.$

8.1 Thermal parameters in statistics for Boson system

- For Boson system, $a_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1}$.

Define Grand partition function:

$$\mathcal{Z} = \prod \mathcal{Z}_l = \prod (1 - e^{-\alpha-\beta\varepsilon_l})^{-\omega_l}.$$

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha-\beta\varepsilon_l}).$$

- $\frac{\partial}{\partial \alpha} \ln \mathcal{Z} = - \sum \frac{\omega_l}{1 - e^{-\alpha-\beta\varepsilon_l}} \cdot e^{-\alpha-\beta\varepsilon_l} = - \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1}$
 $= - \sum a_l.$

$$\Rightarrow \boxed{N = - \frac{\partial}{\partial \alpha} \ln \mathcal{Z}}.$$

8.1 Thermal parameters in statistics for Boson system

- For Boson system, $a_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1}$.

Define Grand partition function:

$$\mathcal{Z} = \prod \mathcal{Z}_l = \prod (1 - e^{-\alpha-\beta\varepsilon_l})^{-\omega_l}.$$

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha-\beta\varepsilon_l}).$$

- $\frac{\partial}{\partial \alpha} \ln \mathcal{Z} = - \sum \frac{\omega_l}{1-e^{-\alpha-\beta\varepsilon_l}} \cdot e^{-\alpha-\beta\varepsilon_l} = - \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1}$
 $= - \sum a_l.$

$$\Rightarrow \boxed{N = -\frac{\partial}{\partial \alpha} \ln \mathcal{Z}}.$$

- $\frac{\partial}{\partial \beta} \ln \mathcal{Z}$

8.1 Thermal parameters in statistics for Boson system

- For Boson system, $a_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1}$.

Define Grand partition function:

$$\mathcal{Z} = \prod \mathcal{Z}_l = \prod (1 - e^{-\alpha-\beta\varepsilon_l})^{-\omega_l}.$$

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha-\beta\varepsilon_l}).$$

- $\frac{\partial}{\partial \alpha} \ln \mathcal{Z} = - \sum \frac{\omega_l}{1-e^{-\alpha-\beta\varepsilon_l}} \cdot e^{-\alpha-\beta\varepsilon_l} = - \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1}$
 $= - \sum a_l.$

$$\Rightarrow \boxed{N = -\frac{\partial}{\partial \alpha} \ln \mathcal{Z}}.$$

- $\frac{\partial}{\partial \beta} \ln \mathcal{Z} = - \sum \frac{\omega_l}{1-e^{-\alpha-\beta\varepsilon_l}} \cdot e^{-\alpha-\beta\varepsilon_l} \cdot \varepsilon_l$

8.1 Thermal parameters in statistics for Boson system

- For Boson system, $a_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1}$.

Define Grand partition function:

$$\mathcal{Z} = \prod \mathcal{Z}_l = \prod (1 - e^{-\alpha-\beta\varepsilon_l})^{-\omega_l}.$$

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha-\beta\varepsilon_l}).$$

- $\frac{\partial}{\partial \alpha} \ln \mathcal{Z} = - \sum \frac{\omega_l}{1-e^{-\alpha-\beta\varepsilon_l}} \cdot e^{-\alpha-\beta\varepsilon_l} = - \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1}$
 $= - \sum a_l.$

$$\Rightarrow \boxed{N = -\frac{\partial}{\partial \alpha} \ln \mathcal{Z}}.$$

- $\frac{\partial}{\partial \beta} \ln \mathcal{Z} = - \sum \frac{\omega_l}{1-e^{-\alpha-\beta\varepsilon_l}} \cdot e^{-\alpha-\beta\varepsilon_l} \cdot \varepsilon_l = - \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1} \varepsilon_l$

8.1 Thermal parameters in statistics for Boson system

- For Boson system, $a_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1}$.

Define Grand partition function:

$$\mathcal{Z} = \prod \mathcal{Z}_l = \prod (1 - e^{-\alpha-\beta\varepsilon_l})^{-\omega_l}.$$

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha-\beta\varepsilon_l}).$$

- $\frac{\partial}{\partial \alpha} \ln \mathcal{Z} = - \sum \frac{\omega_l}{1-e^{-\alpha-\beta\varepsilon_l}} \cdot e^{-\alpha-\beta\varepsilon_l} = - \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1}$
 $= - \sum a_l.$

$$\Rightarrow \boxed{N = -\frac{\partial}{\partial \alpha} \ln \mathcal{Z}}.$$

- $\frac{\partial}{\partial \beta} \ln \mathcal{Z} = - \sum \frac{\omega_l}{1-e^{-\alpha-\beta\varepsilon_l}} \cdot e^{-\alpha-\beta\varepsilon_l} \cdot \varepsilon_l = - \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1} \varepsilon_l$
 $= - \sum a_l \varepsilon_l.$

8.1 Thermal parameters in statistics for Boson system

- For Boson system, $a_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1}$.

Define Grand partition function:

$$\mathcal{Z} = \prod \mathcal{Z}_l = \prod (1 - e^{-\alpha-\beta\varepsilon_l})^{-\omega_l}.$$

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha-\beta\varepsilon_l}).$$

- $\frac{\partial}{\partial \alpha} \ln \mathcal{Z} = - \sum \frac{\omega_l}{1-e^{-\alpha-\beta\varepsilon_l}} \cdot e^{-\alpha-\beta\varepsilon_l} = - \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1}$
 $= - \sum a_l.$

$$\Rightarrow \boxed{N = -\frac{\partial}{\partial \alpha} \ln \mathcal{Z}}.$$

- $\frac{\partial}{\partial \beta} \ln \mathcal{Z} = - \sum \frac{\omega_l}{1-e^{-\alpha-\beta\varepsilon_l}} \cdot e^{-\alpha-\beta\varepsilon_l} \cdot \varepsilon_l = - \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1} \varepsilon_l$
 $= - \sum a_l \varepsilon_l.$

$$\Rightarrow \boxed{U = -\frac{\partial}{\partial \beta} \ln \mathcal{Z}}.$$

Thermal parameters in statistics for Boson system

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l})$$

- Notice $Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} - 1} \frac{\partial \varepsilon_l}{\partial y}$.

Thermal parameters in statistics for Boson system

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l})$$

- Notice $Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} - 1} \frac{\partial \varepsilon_l}{\partial y}$.

$$\frac{\partial}{\partial y} \ln \mathcal{Z}$$

Thermal parameters in statistics for Boson system

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l})$$

- Notice $Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} - 1} \frac{\partial \varepsilon_l}{\partial y}$.

$$\frac{\partial}{\partial y} \ln \mathcal{Z} = - \sum \frac{\omega_l}{1 - e^{-\alpha - \beta \varepsilon_l}} e^{-\alpha - \beta \varepsilon_l} \cdot \beta \frac{\partial \varepsilon_l}{\partial y}$$

Thermal parameters in statistics for Boson system

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l})$$

- Notice $Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} - 1} \frac{\partial \varepsilon_l}{\partial y}$.

$$\frac{\partial}{\partial y} \ln \mathcal{Z} = - \sum \frac{\omega_l}{1 - e^{-\alpha - \beta \varepsilon_l}} e^{-\alpha - \beta \varepsilon_l} \cdot \beta \frac{\partial \varepsilon_l}{\partial y} = -\beta \sum \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} - 1} \frac{\partial \varepsilon_l}{\partial y}.$$

Thermal parameters in statistics for Boson system

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l})$$

- Notice $Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} - 1} \frac{\partial \varepsilon_l}{\partial y}$.

$$\frac{\partial}{\partial y} \ln \mathcal{Z} = - \sum \frac{\omega_l}{1 - e^{-\alpha - \beta \varepsilon_l}} e^{-\alpha - \beta \varepsilon_l} \cdot \beta \frac{\partial \varepsilon_l}{\partial y} = -\beta \sum \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} - 1} \frac{\partial \varepsilon_l}{\partial y}.$$

$$\Rightarrow \boxed{Y = -\frac{1}{\beta} \frac{\partial}{\partial y} \ln \mathcal{Z}}.$$

Thermal parameters in statistics for Boson system

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l})$$

• Notice $Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} - 1} \frac{\partial \varepsilon_l}{\partial y}$.

$$\frac{\partial}{\partial y} \ln \mathcal{Z} = - \sum \frac{\omega_l}{1 - e^{-\alpha - \beta \varepsilon_l}} e^{-\alpha - \beta \varepsilon_l} \cdot \beta \frac{\partial \varepsilon_l}{\partial y} = -\beta \sum \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} - 1} \frac{\partial \varepsilon_l}{\partial y}.$$

$$\Rightarrow \boxed{Y = -\frac{1}{\beta} \frac{\partial}{\partial y} \ln \mathcal{Z}}. \quad \text{Pressure } p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln \mathcal{Z}.$$

Thermal parameters in statistics for Boson system

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l})$$

- Notice $Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} - 1} \frac{\partial \varepsilon_l}{\partial y}$.

$$\frac{\partial}{\partial y} \ln \mathcal{Z} = - \sum \frac{\omega_l}{1 - e^{-\alpha - \beta \varepsilon_l}} e^{-\alpha - \beta \varepsilon_l} \cdot \beta \frac{\partial \varepsilon_l}{\partial y} = -\beta \sum \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} - 1} \frac{\partial \varepsilon_l}{\partial y}.$$

$$\Rightarrow \boxed{Y = -\frac{1}{\beta} \frac{\partial}{\partial y} \ln \mathcal{Z}}. \quad \text{Pressure } p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln \mathcal{Z}.$$

- Together with expresses of $N, U, Y,$

Thermal parameters in statistics for Boson system

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l})$$

- Notice $Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} - 1} \frac{\partial \varepsilon_l}{\partial y}$.

$$\frac{\partial}{\partial y} \ln \mathcal{Z} = - \sum \frac{\omega_l}{1 - e^{-\alpha - \beta \varepsilon_l}} e^{-\alpha - \beta \varepsilon_l} \cdot \beta \frac{\partial \varepsilon_l}{\partial y} = -\beta \sum \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} - 1} \frac{\partial \varepsilon_l}{\partial y}.$$

$$\Rightarrow \boxed{Y = -\frac{1}{\beta} \frac{\partial}{\partial y} \ln \mathcal{Z}}. \quad \text{Pressure } p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln \mathcal{Z}.$$

- Together with expresses of $N, U, Y,$

$$\beta(dU - Ydy + \frac{\alpha}{\beta}dN)$$

Thermal parameters in statistics for Boson system

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l})$$

- Notice $Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} - 1} \frac{\partial \varepsilon_l}{\partial y}$.

$$\frac{\partial}{\partial y} \ln \mathcal{Z} = - \sum \frac{\omega_l}{1 - e^{-\alpha - \beta \varepsilon_l}} e^{-\alpha - \beta \varepsilon_l} \cdot \beta \frac{\partial \varepsilon_l}{\partial y} = -\beta \sum \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} - 1} \frac{\partial \varepsilon_l}{\partial y}.$$

$$\Rightarrow \boxed{Y = -\frac{1}{\beta} \frac{\partial}{\partial y} \ln \mathcal{Z}}. \quad \text{Pressure } p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln \mathcal{Z}.$$

- Together with expresses of N, U, Y ,

$$\beta(dU - Ydy + \frac{\alpha}{\beta}dN) = -\beta d\frac{\ln \mathcal{Z}}{\partial \beta} + \frac{\partial \ln \mathcal{Z}}{\partial y}dy - \alpha d\frac{\partial \ln \mathcal{Z}}{\partial \alpha}$$

Thermal parameters in statistics for Boson system

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l})$$

- Notice $Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} - 1} \frac{\partial \varepsilon_l}{\partial y}$.

$$\frac{\partial}{\partial y} \ln \mathcal{Z} = - \sum \frac{\omega_l}{1 - e^{-\alpha - \beta \varepsilon_l}} e^{-\alpha - \beta \varepsilon_l} \cdot \beta \frac{\partial \varepsilon_l}{\partial y} = -\beta \sum \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} - 1} \frac{\partial \varepsilon_l}{\partial y}.$$

$$\Rightarrow \boxed{Y = -\frac{1}{\beta} \frac{\partial}{\partial y} \ln \mathcal{Z}}. \quad \text{Pressure } p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln \mathcal{Z}.$$

- Together with expresses of N, U, Y ,

$$\begin{aligned} \beta(dU - Ydy + \frac{\alpha}{\beta}dN) &= -\beta d\frac{\partial \ln \mathcal{Z}}{\partial \beta} + \frac{\partial \ln \mathcal{Z}}{\partial y} dy - \alpha d\frac{\partial \ln \mathcal{Z}}{\partial \alpha} \\ &= -d(\beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}) + \frac{\partial \ln \mathcal{Z}}{\partial \beta} d\beta + \frac{\partial \ln \mathcal{Z}}{\partial y} dy - d(\alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha}) + \frac{\partial \ln \mathcal{Z}}{\partial \alpha} d\alpha \end{aligned}$$

Thermal parameters in statistics for Boson system

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l})$$

- Notice $Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} - 1} \frac{\partial \varepsilon_l}{\partial y}$.

$$\frac{\partial}{\partial y} \ln \mathcal{Z} = - \sum \frac{\omega_l}{1 - e^{-\alpha - \beta \varepsilon_l}} e^{-\alpha - \beta \varepsilon_l} \cdot \beta \frac{\partial \varepsilon_l}{\partial y} = -\beta \sum \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} - 1} \frac{\partial \varepsilon_l}{\partial y}.$$

$$\Rightarrow \boxed{Y = -\frac{1}{\beta} \frac{\partial}{\partial y} \ln \mathcal{Z}}. \quad \text{Pressure } p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln \mathcal{Z}.$$

- Together with expresses of N, U, Y ,

$$\begin{aligned} \beta(dU - Ydy + \frac{\alpha}{\beta}dN) &= -\beta d\frac{\partial \ln \mathcal{Z}}{\partial \beta} + \frac{\partial \ln \mathcal{Z}}{\partial y}dy - \alpha d\frac{\partial \ln \mathcal{Z}}{\partial \alpha} \\ &= -d(\beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}) + \frac{\partial \ln \mathcal{Z}}{\partial \beta}d\beta + \frac{\partial \ln \mathcal{Z}}{\partial y}dy - d(\alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha}) + \frac{\partial \ln \mathcal{Z}}{\partial \alpha}d\alpha \\ &= -d(\beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}) - d(\alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha}) + (\frac{\partial \ln \mathcal{Z}}{\partial \beta}d\beta + \frac{\partial \ln \mathcal{Z}}{\partial y}dy + \frac{\partial \ln \mathcal{Z}}{\partial \alpha}d\alpha) \end{aligned}$$

Thermal parameters in statistics for Boson system

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l})$$

- Notice $Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} - 1} \frac{\partial \varepsilon_l}{\partial y}$.

$$\frac{\partial}{\partial y} \ln \mathcal{Z} = - \sum \frac{\omega_l}{1 - e^{-\alpha - \beta \varepsilon_l}} e^{-\alpha - \beta \varepsilon_l} \cdot \beta \frac{\partial \varepsilon_l}{\partial y} = -\beta \sum \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} - 1} \frac{\partial \varepsilon_l}{\partial y}.$$

$$\Rightarrow \boxed{Y = -\frac{1}{\beta} \frac{\partial}{\partial y} \ln \mathcal{Z}}. \quad \text{Pressure } p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln \mathcal{Z}.$$

- Together with expresses of N, U, Y ,

$$\begin{aligned} \beta(dU - Ydy + \frac{\alpha}{\beta}dN) &= -\beta d\frac{\partial \ln \mathcal{Z}}{\partial \beta} + \frac{\partial \ln \mathcal{Z}}{\partial y}dy - \alpha d\frac{\partial \ln \mathcal{Z}}{\partial \alpha} \\ &= -d(\beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}) + \frac{\partial \ln \mathcal{Z}}{\partial \beta}d\beta + \frac{\partial \ln \mathcal{Z}}{\partial y}dy - d(\alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha}) + \frac{\partial \ln \mathcal{Z}}{\partial \alpha}d\alpha \\ &= -d(\beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}) - d(\alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha}) + (\frac{\partial \ln \mathcal{Z}}{\partial \beta}d\beta + \frac{\partial \ln \mathcal{Z}}{\partial y}dy + \frac{\partial \ln \mathcal{Z}}{\partial \alpha}d\alpha) \\ &= -d(\beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}) - d(\alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha}) + d \ln \mathcal{Z} \end{aligned}$$

Thermal parameters in statistics for Boson system

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l})$$

- Notice $Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} - 1} \frac{\partial \varepsilon_l}{\partial y}$.

$$\frac{\partial}{\partial y} \ln \mathcal{Z} = - \sum \frac{\omega_l}{1 - e^{-\alpha - \beta \varepsilon_l}} e^{-\alpha - \beta \varepsilon_l} \cdot \beta \frac{\partial \varepsilon_l}{\partial y} = -\beta \sum \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} - 1} \frac{\partial \varepsilon_l}{\partial y}.$$

$$\Rightarrow \boxed{Y = -\frac{1}{\beta} \frac{\partial}{\partial y} \ln \mathcal{Z}}. \quad \text{Pressure } p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln \mathcal{Z}.$$

- Together with expresses of N, U, Y ,

$$\begin{aligned} \beta(dU - Ydy + \frac{\alpha}{\beta}dN) &= -\beta d\frac{\partial \ln \mathcal{Z}}{\partial \beta} + \frac{\partial \ln \mathcal{Z}}{\partial y}dy - \alpha d\frac{\partial \ln \mathcal{Z}}{\partial \alpha} \\ &= -d(\beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}) + \frac{\partial \ln \mathcal{Z}}{\partial \beta}d\beta + \frac{\partial \ln \mathcal{Z}}{\partial y}dy - d(\alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha}) + \frac{\partial \ln \mathcal{Z}}{\partial \alpha}d\alpha \\ &= -d(\beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}) - d(\alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha}) + (\frac{\partial \ln \mathcal{Z}}{\partial \beta}d\beta + \frac{\partial \ln \mathcal{Z}}{\partial y}dy + \frac{\partial \ln \mathcal{Z}}{\partial \alpha}d\alpha) \\ &= -d(\beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}) - d(\alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha}) + d \ln \mathcal{Z} \\ &= d(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}). \end{aligned}$$

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l})$$

- Eq. (4.1.12), $dU = TdS - pdV + \sum \mu_i dn_i$.

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l})$$

- Eq. (4.1.12), $dU = TdS - pdV + \sum \mu_i dn_i$.
 $-p \rightarrow Y$, $dV \rightarrow dy$, $n_i \rightarrow N$,

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l})$$

- Eq. (4.1.12), $dU = TdS - pdV + \sum \mu_i dn_i$.
 $-p \rightarrow Y$, $dV \rightarrow dy$, $n_i \rightarrow N$,
 $\Rightarrow dS = \frac{1}{T}(dU - Ydy - \mu dN)$;

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l})$$

- Eq. (4.1.12), $dU = TdS - pdV + \sum \mu_i dn_i$.
 $-p \rightarrow Y$, $dV \rightarrow dy$, $n_i \rightarrow N$,
 $\Rightarrow dS = \frac{1}{T}(dU - Ydy - \mu dN)$;
 $d(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}) = \beta(dU - Ydy + \frac{\alpha}{\beta}dN)$.

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l})$$

- Eq. (4.1.12), $dU = TdS - pdV + \sum \mu_i dn_i$.
 $-p \rightarrow Y$, $dV \rightarrow dy$, $n_i \rightarrow N$,
 $\Rightarrow dS = \frac{1}{T}(dU - Ydy - \mu dN)$;
 $d(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}) = \beta(dU - Ydy + \frac{\alpha}{\beta} dN)$.
 $\Rightarrow \beta = \frac{1}{kT}$, $\alpha = -\beta\mu = -\frac{\mu}{kT}$.

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l})$$

- Eq. (4.1.12), $dU = TdS - pdV + \sum \mu_i dn_i$.
 $-p \rightarrow Y$, $dV \rightarrow dy$, $n_i \rightarrow N$,
 $\Rightarrow dS = \frac{1}{T}(dU - Ydy - \mu dN)$;
 $d(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}) = \beta(dU - Ydy + \frac{\alpha}{\beta} dN)$.
 $\Rightarrow \beta = \frac{1}{kT}$, $\alpha = -\beta\mu = -\frac{\mu}{kT}$.
 β —temperature, α —chemical potential.

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l})$$

- Eq. (4.1.12), $dU = TdS - pdV + \sum \mu_i dn_i$.
 $-p \rightarrow Y$, $dV \rightarrow dy$, $n_i \rightarrow N$,
 $\Rightarrow dS = \frac{1}{T}(dU - Ydy - \mu dN)$;
 $d(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}) = \beta(dU - Ydy + \frac{\alpha}{\beta} dN)$.
 $\Rightarrow \beta = \frac{1}{kT}$, $\alpha = -\beta\mu = -\frac{\mu}{kT}$.
 β —temperature, α —chemical potential.
- $S = k(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta})$

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l})$$

- Eq. (4.1.12), $dU = TdS - pdV + \sum \mu_i dn_i$.
 $-p \rightarrow Y$, $dV \rightarrow dy$, $n_i \rightarrow N$,
 $\Rightarrow dS = \frac{1}{T}(dU - Ydy - \mu dN)$;
 $d(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}) = \beta(dU - Ydy + \frac{\alpha}{\beta} dN)$.
 $\Rightarrow \beta = \frac{1}{kT}$, $\alpha = -\beta\mu = -\frac{\mu}{kT}$.
 β —temperature, α —chemical potential.
- $S = k(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta})$
 $= k[- \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l}) + \alpha \sum \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} - 1}$
 $+ \beta \sum \frac{\omega_l \varepsilon_l}{e^{\alpha + \beta \varepsilon_l} - 1}]$;

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l})$$

- Eq. (4.1.12), $dU = TdS - pdV + \sum \mu_i dn_i$.
 $-p \rightarrow Y$, $dV \rightarrow dy$, $n_i \rightarrow N$,
 $\Rightarrow dS = \frac{1}{T}(dU - Ydy - \mu dN)$;
 $d(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}) = \beta(dU - Ydy + \frac{\alpha}{\beta} dN)$.
 $\Rightarrow \beta = \frac{1}{kT}$, $\alpha = -\beta\mu = -\frac{\mu}{kT}$.
 β —temperature, α —chemical potential.

$$\begin{aligned} S &= k(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}) \\ &= k[- \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l}) + \alpha \sum \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} - 1} \\ &\quad + \beta \sum \frac{\omega_l \varepsilon_l}{e^{\alpha + \beta \varepsilon_l} - 1}]; \end{aligned}$$

Notice: $a_l = \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} - 1} \Rightarrow e^{\alpha + \beta \varepsilon_l} - 1 = \frac{\omega_l}{a_l}$, then

$$\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l})$$

- Eq. (4.1.12), $dU = TdS - pdV + \sum \mu_i dn_i$.
 $-p \rightarrow Y$, $dV \rightarrow dy$, $n_i \rightarrow N$,
 $\Rightarrow dS = \frac{1}{T}(dU - Ydy - \mu dN)$;
 $d(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}) = \beta(dU - Ydy + \frac{\alpha}{\beta} dN)$.
 $\Rightarrow \beta = \frac{1}{kT}$, $\alpha = -\beta\mu = -\frac{\mu}{kT}$.
 β —temperature, α —chemical potential.

$$\begin{aligned} S &= k(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}) \\ &= k[-\sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l}) + \alpha \sum \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} - 1} \\ &\quad + \beta \sum \frac{\omega_l \varepsilon_l}{e^{\alpha + \beta \varepsilon_l} - 1}]; \end{aligned}$$

Notice: $a_l = \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} - 1} \Rightarrow e^{\alpha + \beta \varepsilon_l} - 1 = \frac{\omega_l}{a_l}$, then

$$1 - e^{-\alpha - \beta \varepsilon_l} = 1 - \left(\frac{\omega_l}{a_l} + 1\right)^{-1} = 1 - \frac{a_l}{\omega_l + a_l} = \frac{\omega_l}{\omega_l + a_l}.$$

$$e^{\alpha+\beta\varepsilon_l} - 1 = \frac{\omega_l}{a_l}, \quad 1 - e^{-\alpha-\beta\varepsilon_l} = \frac{\omega_l}{\omega_l + a_l}$$

- $$S = k \left[- \sum \omega_l \ln(1 - e^{-\alpha-\beta\varepsilon_l}) + \alpha \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l} - 1} + \beta \sum \frac{\omega_l \varepsilon_l}{e^{\alpha+\beta\varepsilon_l} - 1} \right]$$

$$e^{\alpha+\beta\varepsilon_l} - 1 = \frac{\omega_l}{a_l}, \quad 1 - e^{-\alpha-\beta\varepsilon_l} = \frac{\omega_l}{\omega_l+a_l}$$

- $$\begin{aligned} S &= k \left[- \sum \omega_l \ln(1 - e^{-\alpha-\beta\varepsilon_l}) + \alpha \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l} - 1} \right. \\ &\quad \left. + \beta \sum \frac{\omega_l \varepsilon_l}{e^{\alpha+\beta\varepsilon_l} - 1} \right] \\ &= k \left[- \sum \omega_l \ln \frac{\omega_l}{a_l + \omega_l} + \sum \omega_l \frac{\alpha + \beta\varepsilon_l}{e^{\alpha+\beta\varepsilon_l} - 1} \right] \end{aligned}$$

$$e^{\alpha+\beta\varepsilon_l} - 1 = \frac{\omega_l}{a_l}, \quad 1 - e^{-\alpha-\beta\varepsilon_l} = \frac{\omega_l}{\omega_l+a_l}$$

$$\begin{aligned} \bullet S &= k \left[- \sum \omega_l \ln(1 - e^{-\alpha-\beta\varepsilon_l}) + \alpha \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1} \right. \\ &\quad \left. + \beta \sum \frac{\omega_l \varepsilon_l}{e^{\alpha+\beta\varepsilon_l}-1} \right] \\ &= k \left[- \sum \omega_l \ln \frac{\omega_l}{a_l+\omega_l} + \sum \omega_l \frac{\alpha+\beta\varepsilon_l}{e^{\alpha+\beta\varepsilon_l}-1} \right] \\ &= k \left[- \sum \omega_l \ln \frac{\omega_l}{a_l+\omega_l} + \sum \omega_l \frac{\ln(\omega_l/a_l+1)}{\omega_l/a_l} \right] \end{aligned}$$

$$e^{\alpha+\beta\varepsilon_l} - 1 = \frac{\omega_l}{a_l}, \quad 1 - e^{-\alpha-\beta\varepsilon_l} = \frac{\omega_l}{\omega_l+a_l}$$

$$\begin{aligned}
 \bullet \quad S &= k \left[- \sum \omega_l \ln(1 - e^{-\alpha-\beta\varepsilon_l}) + \alpha \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1} \right. \\
 &\quad \left. + \beta \sum \frac{\omega_l \varepsilon_l}{e^{\alpha+\beta\varepsilon_l}-1} \right] \\
 &= k \left[- \sum \omega_l \ln \frac{\omega_l}{a_l + \omega_l} + \sum \omega_l \frac{\alpha + \beta \varepsilon_l}{e^{\alpha+\beta\varepsilon_l}-1} \right] \\
 &= k \left[- \sum \omega_l \ln \frac{\omega_l}{a_l + \omega_l} + \sum \omega_l \frac{\ln(\omega_l/a_l + 1)}{\omega_l/a_l} \right] \\
 &= k \left[- \sum \omega_l \ln \frac{\omega_l}{a_l + \omega_l} + \sum a_l \ln \frac{\omega_l + a_l}{a_l} \right]
 \end{aligned}$$

$$e^{\alpha+\beta\varepsilon_l} - 1 = \frac{\omega_l}{a_l}, \quad 1 - e^{-\alpha-\beta\varepsilon_l} = \frac{\omega_l}{\omega_l+a_l}$$

$$\begin{aligned}
 \bullet S &= k \left[- \sum \omega_l \ln(1 - e^{-\alpha-\beta\varepsilon_l}) + \alpha \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1} \right. \\
 &\quad \left. + \beta \sum \frac{\omega_l \varepsilon_l}{e^{\alpha+\beta\varepsilon_l}-1} \right] \\
 &= k \left[- \sum \omega_l \ln \frac{\omega_l}{a_l+\omega_l} + \sum \omega_l \frac{\alpha+\beta\varepsilon_l}{e^{\alpha+\beta\varepsilon_l}-1} \right] \\
 &= k \left[- \sum \omega_l \ln \frac{\omega_l}{a_l+\omega_l} + \sum \omega_l \frac{\ln(\omega_l/a_l+1)}{\omega_l/a_l} \right] \\
 &= k \left[- \sum \omega_l \ln \frac{\omega_l}{a_l+\omega_l} + \sum a_l \ln \frac{\omega_l+a_l}{a_l} \right] \\
 &= k \left[\sum \omega_l \ln(a_l + \omega_l) - \sum \omega_l \ln \omega_l + \sum a_l \ln(\omega_l + a_l) \right. \\
 &\quad \left. - \sum a_l \ln a_l \right]
 \end{aligned}$$

$$e^{\alpha+\beta\varepsilon_l} - 1 = \frac{\omega_l}{a_l}, \quad 1 - e^{-\alpha-\beta\varepsilon_l} = \frac{\omega_l}{\omega_l+a_l}$$

$$\begin{aligned}
 \bullet S &= k \left[- \sum \omega_l \ln(1 - e^{-\alpha-\beta\varepsilon_l}) + \alpha \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1} \right. \\
 &\quad \left. + \beta \sum \frac{\omega_l \varepsilon_l}{e^{\alpha+\beta\varepsilon_l}-1} \right] \\
 &= k \left[- \sum \omega_l \ln \frac{\omega_l}{a_l + \omega_l} + \sum \omega_l \frac{\alpha + \beta \varepsilon_l}{e^{\alpha+\beta\varepsilon_l}-1} \right] \\
 &= k \left[- \sum \omega_l \ln \frac{\omega_l}{a_l + \omega_l} + \sum \omega_l \frac{\ln(\omega_l/a_l + 1)}{\omega_l/a_l} \right] \\
 &= k \left[- \sum \omega_l \ln \frac{\omega_l}{a_l + \omega_l} + \sum a_l \ln \frac{\omega_l + a_l}{a_l} \right] \\
 &= k \left[\sum \omega_l \ln(a_l + \omega_l) - \sum \omega_l \ln \omega_l + \sum a_l \ln(\omega_l + a_l) \right. \\
 &\quad \left. - \sum a_l \ln a_l \right] \\
 &= k \left[\sum (a_l + \omega_l) \ln(a_l + \omega_l) - \sum \omega_l \ln \omega_l - \sum a_l \ln a_l \right].
 \end{aligned}$$

$$e^{\alpha+\beta\varepsilon_l} - 1 = \frac{\omega_l}{a_l}, \quad 1 - e^{-\alpha-\beta\varepsilon_l} = \frac{\omega_l}{\omega_l+a_l}$$

$$\begin{aligned}
 \bullet S &= k \left[- \sum \omega_l \ln(1 - e^{-\alpha-\beta\varepsilon_l}) + \alpha \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1} \right. \\
 &\quad \left. + \beta \sum \frac{\omega_l \varepsilon_l}{e^{\alpha+\beta\varepsilon_l}-1} \right] \\
 &= k \left[- \sum \omega_l \ln \frac{\omega_l}{a_l + \omega_l} + \sum \omega_l \frac{\alpha + \beta \varepsilon_l}{e^{\alpha+\beta\varepsilon_l}-1} \right] \\
 &= k \left[- \sum \omega_l \ln \frac{\omega_l}{a_l + \omega_l} + \sum \omega_l \frac{\ln(\omega_l/a_l + 1)}{\omega_l/a_l} \right] \\
 &= k \left[- \sum \omega_l \ln \frac{\omega_l}{a_l + \omega_l} + \sum a_l \ln \frac{\omega_l + a_l}{a_l} \right] \\
 &= k \left[\sum \omega_l \ln(a_l + \omega_l) - \sum \omega_l \ln \omega_l + \sum a_l \ln(\omega_l + a_l) \right. \\
 &\quad \left. - \sum a_l \ln a_l \right] \\
 &= k \left[\sum (a_l + \omega_l) \ln(a_l + \omega_l) - \sum \omega_l \ln \omega_l - \sum a_l \ln a_l \right].
 \end{aligned}$$

Comparing with eq.(6.7.4),

$$e^{\alpha+\beta\varepsilon_l} - 1 = \frac{\omega_l}{a_l}, \quad 1 - e^{-\alpha-\beta\varepsilon_l} = \frac{\omega_l}{\omega_l+a_l}$$

- $$\begin{aligned}
 S &= k \left[- \sum \omega_l \ln(1 - e^{-\alpha-\beta\varepsilon_l}) + \alpha \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1} \right. \\
 &\quad \left. + \beta \sum \frac{\omega_l \varepsilon_l}{e^{\alpha+\beta\varepsilon_l}-1} \right] \\
 &= k \left[- \sum \omega_l \ln \frac{\omega_l}{a_l + \omega_l} + \sum \omega_l \frac{\alpha + \beta \varepsilon_l}{e^{\alpha+\beta\varepsilon_l}-1} \right] \\
 &= k \left[- \sum \omega_l \ln \frac{\omega_l}{a_l + \omega_l} + \sum \omega_l \frac{\ln(\omega_l/a_l + 1)}{\omega_l/a_l} \right] \\
 &= k \left[- \sum \omega_l \ln \frac{\omega_l}{a_l + \omega_l} + \sum a_l \ln \frac{\omega_l + a_l}{a_l} \right] \\
 &= k \left[\sum \omega_l \ln(a_l + \omega_l) - \sum \omega_l \ln \omega_l + \sum a_l \ln(\omega_l + a_l) \right. \\
 &\quad \left. - \sum a_l \ln a_l \right] \\
 &= k \left[\sum (a_l + \omega_l) \ln(a_l + \omega_l) - \sum \omega_l \ln \omega_l - \sum a_l \ln a_l \right].
 \end{aligned}$$

Comparing with eq.(6.7.4),

- $$\Rightarrow \boxed{S = k \ln \Omega}.$$

Thermal parameters in statistics for Fermion system

Thermal parameters in statistics for Fermion system

- Grand partition function:

$$\mathcal{Z} = \prod \mathcal{Z}_l = \prod (1 + e^{-\alpha - \beta \epsilon_l})^{\omega_l}.$$

$$\ln \mathcal{Z} = \sum \omega_l \ln(1 + e^{-\alpha - \beta \epsilon_l}).$$

- $$N = -\frac{\partial}{\partial \alpha} \ln \mathcal{Z}.$$

- $$U = -\frac{\partial}{\partial \beta} \ln \mathcal{Z}.$$

- $$Y = -\frac{1}{\beta} \frac{\partial}{\partial y} \ln \mathcal{Z}.$$

- $$S = k \ln \Omega.$$

Grand potential in thermodynamics

- Method for Bose (Fermi) statistics:

Grand potential in thermodynamics

- Method for Bose (Fermi) statistics:
 $\{\varepsilon_l\}, \{\omega_l\}$

Grand potential in thermodynamics

- Method for Bose (Fermi) statistics:
 $\{\varepsilon_l\}, \{\omega_l\} \rightarrow \ln \mathcal{Z}(\alpha, \beta, y)$

Grand potential in thermodynamics

- Method for Bose (Fermi) statistics:
 $\{\varepsilon_l\}, \{\omega_l\} \rightarrow \ln \mathcal{Z}(\alpha, \beta, y)$
 \rightarrow Other thermal variables ($N, U, S...$).

Grand potential in thermodynamics

- Method for Bose (Fermi) statistics:
 $\{\varepsilon_l\}, \{\omega_l\} \rightarrow \ln \mathcal{Z}(\alpha, \beta, y)$
 \rightarrow Other thermal variables $(N, U, S\dots)$.
- For simple system, $(\alpha, \beta, y) \leftrightarrow (\mu, T, V)$.

Grand potential in thermodynamics

- Method for Bose (Fermi) statistics:
 $\{\varepsilon_l\}, \{\omega_l\} \rightarrow \ln \mathcal{Z}(\alpha, \beta, y)$
 \rightarrow Other thermal variables $(N, U, S\dots)$.
- For simple system, $(\alpha, \beta, y) \leftrightarrow (\mu, T, V)$.
The characteristic function is grand potential:
 $J = U - TS - \mu N$ (eq. (3.2.10)).

Grand potential in thermodynamics

- Method for Bose (Fermi) statistics:
 $\{\varepsilon_l\}, \{\omega_l\} \rightarrow \ln \mathcal{Z}(\alpha, \beta, y)$
 \rightarrow Other thermal variables $(N, U, S\dots)$.
- For simple system, $(\alpha, \beta, y) \leftrightarrow (\mu, T, V)$.
The characteristic function is grand potential:
 $J = U - TS - \mu N$ (eq. (3.2.10)).
- Take (8.1.10) in,
$$J = U - \frac{1}{k\beta} \cdot k(\ln \mathcal{Z} + \alpha N + \beta U) - \mu N$$

Grand potential in thermodynamics

- Method for Bose (Fermi) statistics:
 $\{\varepsilon_l\}, \{\omega_l\} \rightarrow \ln \mathcal{Z}(\alpha, \beta, y)$
 \rightarrow Other thermal variables ($N, U, S \dots$).
- For simple system, $(\alpha, \beta, y) \leftrightarrow (\mu, T, V)$.
 The characteristic function is grand potential:
 $J = U - TS - \mu N$ (eq. (3.2.10)).
- Take (8.1.10) in,

$$J = U - \frac{1}{k\beta} \cdot k(\ln \mathcal{Z} + \alpha N + \beta U) - \mu N$$

$$= U - \frac{1}{\beta}(\ln \mathcal{Z} - \mu\beta N + \beta U) - \mu N$$

Grand potential in thermodynamics

- Method for Bose (Fermi) statistics:
 $\{\varepsilon_l\}, \{\omega_l\} \rightarrow \ln \mathcal{Z}(\alpha, \beta, y)$
 \rightarrow Other thermal variables ($N, U, S \dots$).
- For simple system, $(\alpha, \beta, y) \leftrightarrow (\mu, T, V)$.
The characteristic function is grand potential:
 $J = U - TS - \mu N$ (eq. (3.2.10)).
- Take (8.1.10) in,
$$J = U - \frac{1}{k\beta} \cdot k(\ln \mathcal{Z} + \alpha N + \beta U) - \mu N$$
$$= U - \frac{1}{\beta}(\ln \mathcal{Z} - \mu\beta N + \beta U) - \mu N = -\frac{1}{\beta} \ln \mathcal{Z}$$

Grand potential in thermodynamics

- Method for Bose (Fermi) statistics:

$$\{\varepsilon_l\}, \{\omega_l\} \rightarrow \ln \mathcal{Z}(\alpha, \beta, y)$$

→ Other thermal variables ($N, U, S...$).

- For simple system, $(\alpha, \beta, y) \leftrightarrow (\mu, T, V)$.

The characteristic function is grand potential:

$$J = U - TS - \mu N \quad (\text{eq. (3.2.10)}).$$

- Take (8.1.10) in,

$$\begin{aligned} J &= U - \frac{1}{k\beta} \cdot k(\ln \mathcal{Z} + \alpha N + \beta U) - \mu N \\ &= U - \frac{1}{\beta}(\ln \mathcal{Z} - \mu\beta N + \beta U) - \mu N = -\frac{1}{\beta} \ln \mathcal{Z} \\ &= -kT \ln \mathcal{Z}. \end{aligned}$$

Table of contents

1 §8. Bose statistics and Fermi statistics

- 8.1 Thermal parameters in statistics
- 8.2 U of weak degenerated ideal Fermion(Boson) gas
- 8.3 Bose-Einstein Condensate (BEC)
- 8.4 Photon gas
- 8.5 Free electron gas in metal

8.2 U of weak degenerated ideal Fermion(Boson) gas

8.2 U of weak degenerated ideal Fermion(Boson) gas

- Non-degeneration condition: $e^{-\alpha} \ll 1$ or $n\lambda^3 \ll 1$.
(7.2.6, 7.2.7).

8.2 U of weak degenerated ideal Fermion(Boson) gas

- Non-degeneration condition: $e^{-\alpha} \ll 1$ or $n\lambda^3 \ll 1$.
(7.2.6, 7.2.7).
- If not satisfied, but still < 1 , called weak degeneration.

8.2 U of weak degenerated ideal Fermion(Boson) gas

- Non-degeneration condition: $e^{-\alpha} \ll 1$ or $n\lambda^3 \ll 1$. (7.2.6, 7.2.7).
- If not satisfied, but still < 1 , called weak degeneration.
- Simple example: monatomic molecule (Fermion or Boson).

8.2 U of weak degenerated ideal Fermion(Boson) gas

- Non-degeneration condition: $e^{-\alpha} \ll 1$ or $n\lambda^3 \ll 1$. (7.2.6, 7.2.7).
- If not satisfied, but still < 1 , called weak degeneration.
- Simple example: monatomic molecule (Fermion or Boson).
- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$.

8.2 U of weak degenerated ideal Fermion(Boson) gas

- Non-degeneration condition: $e^{-\alpha} \ll 1$ or $n\lambda^3 \ll 1$. (7.2.6, 7.2.7).
- If not satisfied, but still < 1 , called weak degeneration.
- Simple example: monatomic molecule (Fermion or Boson).
- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$.
- Number of micro-states: $\omega = g \cdot \frac{dx dy dz dp_x dp_y dp_z}{h^3}$, where g is the degeneracy by spin.

8.2 U of weak degenerated ideal Fermion(Boson) gas

- Non-degeneration condition: $e^{-\alpha} \ll 1$ or $n\lambda^3 \ll 1$. (7.2.6, 7.2.7).
- If not satisfied, but still < 1 , called weak degeneration.
- Simple example: monatomic molecule (Fermion or Boson).
- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$.
- Number of micro-states: $\omega = g \cdot \frac{dx dy dz dp_x dp_y dp_z}{h^3}$, where g is the degeneracy by spin.
For isotropic, given volume V , $\omega' = g \cdot \frac{V}{h^3} \cdot 4\pi p^2 dp$.

8.2 U of weak degenerated ideal Fermion(Boson) gas

- Non-degeneration condition: $e^{-\alpha} \ll 1$ or $n\lambda^3 \ll 1$. (7.2.6, 7.2.7).
- If not satisfied, but still < 1 , called weak degeneration.
- Simple example: monatomic molecule (Fermion or Boson).
- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$.
- Number of micro-states: $\omega = g \cdot \frac{dx dy dz dp_x dp_y dp_z}{h^3}$, where g is the degeneracy by spin.

For isotropic, given volume V , $\omega' = g \cdot \frac{V}{h^3} \cdot 4\pi p^2 dp$.

Convert to energy: $D(\varepsilon)d\varepsilon = g \cdot \frac{V}{h^3} \cdot 4\pi \cdot 2m\varepsilon d\sqrt{2m\varepsilon}$

8.2 U of weak degenerated ideal Fermion(Boson) gas

- Non-degeneration condition: $e^{-\alpha} \ll 1$ or $n\lambda^3 \ll 1$. (7.2.6, 7.2.7).
- If not satisfied, but still < 1 , called weak degeneration.
- Simple example: monatomic molecule (Fermion or Boson).
- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$.
- Number of micro-states: $\omega = g \cdot \frac{dx dy dz dp_x dp_y dp_z}{h^3}$, where g is the degeneracy by spin.

For isotropic, given volume V , $\omega' = g \cdot \frac{V}{h^3} \cdot 4\pi p^2 dp$.

Convert to energy: $D(\varepsilon)d\varepsilon = g \cdot \frac{V}{h^3} \cdot 4\pi \cdot 2m\varepsilon d\sqrt{2m\varepsilon}$
 $= g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \varepsilon^{\frac{1}{2}} d\varepsilon$.

$$D(\varepsilon)d\varepsilon = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \varepsilon^{\frac{1}{2}} d\varepsilon$$

- Number of particles:

$$D(\varepsilon)d\varepsilon = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \varepsilon^{\frac{1}{2}} d\varepsilon$$

- Number of particles:

$$N = \sum a_l$$

$$D(\varepsilon)d\varepsilon = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \varepsilon^{\frac{1}{2}} d\varepsilon$$

- Number of particles:

$$N = \sum a_l = \sum \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} \pm 1}$$

$$D(\varepsilon)d\varepsilon = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \varepsilon^{\frac{1}{2}} d\varepsilon$$

- Number of particles:

$$N = \sum a_l = \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l} \pm 1} = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\alpha+\beta\varepsilon} \pm 1}.$$

$$D(\varepsilon)d\varepsilon = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \varepsilon^{\frac{1}{2}} d\varepsilon$$

- Number of particles:

$$N = \sum a_l = \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l} \pm 1} = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\alpha+\beta\varepsilon} \pm 1}.$$

- Internal energy:

$$D(\varepsilon)d\varepsilon = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \varepsilon^{\frac{1}{2}} d\varepsilon$$

- Number of particles:

$$N = \sum a_l = \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l} \pm 1} = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\alpha+\beta\varepsilon} \pm 1}.$$

- Internal energy:

$$U = \sum a_l \varepsilon_l$$

$$D(\varepsilon)d\varepsilon = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \varepsilon^{\frac{1}{2}} d\varepsilon$$

- Number of particles:

$$N = \sum a_l = \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l} \pm 1} = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\alpha+\beta\varepsilon} \pm 1}.$$

- Internal energy:

$$U = \sum a_l \varepsilon_l = \sum \frac{\omega_l \varepsilon_l}{e^{\alpha+\beta\varepsilon_l} \pm 1}$$

$$D(\varepsilon)d\varepsilon = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \varepsilon^{\frac{1}{2}} d\varepsilon$$

- Number of particles:

$$N = \sum a_l = \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l} \pm 1} = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\alpha+\beta\varepsilon} \pm 1}.$$

- Internal energy:

$$U = \sum a_l \varepsilon_l = \sum \frac{\omega_l \varepsilon_l}{e^{\alpha+\beta\varepsilon_l} \pm 1} = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{3}{2}} d\varepsilon}{e^{\alpha+\beta\varepsilon} \pm 1}.$$

$$D(\varepsilon)d\varepsilon = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \varepsilon^{\frac{1}{2}} d\varepsilon$$

- Number of particles:

$$N = \sum a_l = \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l} \pm 1} = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\alpha+\beta\varepsilon} \pm 1}.$$

- Internal energy:

$$U = \sum a_l \varepsilon_l = \sum \frac{\omega_l \varepsilon_l}{e^{\alpha+\beta\varepsilon_l} \pm 1} = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{3}{2}} d\varepsilon}{e^{\alpha+\beta\varepsilon} \pm 1}.$$

- Set $x = \beta\varepsilon$,

$$D(\varepsilon)d\varepsilon = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \varepsilon^{\frac{1}{2}} d\varepsilon$$

- Number of particles:

$$N = \sum a_l = \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l} \pm 1} = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\alpha+\beta\varepsilon} \pm 1}.$$

- Internal energy:

$$U = \sum a_l \varepsilon_l = \sum \frac{\omega_l \varepsilon_l}{e^{\alpha+\beta\varepsilon_l} \pm 1} = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{3}{2}} d\varepsilon}{e^{\alpha+\beta\varepsilon} \pm 1}.$$

- Set $x = \beta\varepsilon$,

$$N = g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} \int_0^\infty \frac{x^{\frac{1}{2}} dx}{e^{\alpha+x} \pm 1},$$

$$D(\varepsilon)d\varepsilon = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \varepsilon^{\frac{1}{2}} d\varepsilon$$

- Number of particles:

$$N = \sum a_l = \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l} \pm 1} = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\alpha+\beta\varepsilon} \pm 1}.$$

- Internal energy:

$$U = \sum a_l \varepsilon_l = \sum \frac{\omega_l \varepsilon_l}{e^{\alpha+\beta\varepsilon_l} \pm 1} = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{3}{2}} d\varepsilon}{e^{\alpha+\beta\varepsilon} \pm 1}.$$

- Set $x = \beta\varepsilon$,

$$N = g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} \int_0^\infty \frac{x^{\frac{1}{2}} dx}{e^{\alpha+x} \pm 1},$$

$$U = g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} kT \int_0^\infty \frac{x^{\frac{3}{2}} dx}{e^{\alpha+x} \pm 1}.$$

$$D(\varepsilon)d\varepsilon = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \varepsilon^{\frac{1}{2}} d\varepsilon$$

- Number of particles:

$$N = \sum a_l = \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l} \pm 1} = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\alpha+\beta\varepsilon} \pm 1}.$$

- Internal energy:

$$U = \sum a_l \varepsilon_l = \sum \frac{\omega_l \varepsilon_l}{e^{\alpha+\beta\varepsilon_l} \pm 1} = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{3}{2}} d\varepsilon}{e^{\alpha+\beta\varepsilon} \pm 1}.$$

- Set $x = \beta\varepsilon$,

$$N = g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} \int_0^\infty \frac{x^{\frac{1}{2}} dx}{e^{\alpha+x} \pm 1},$$

$$U = g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} kT \int_0^\infty \frac{x^{\frac{3}{2}} dx}{e^{\alpha+x} \pm 1}.$$

- As $e^{-\alpha} < 1$, $x > 0$, $\Rightarrow e^{-\alpha-x} < 1$.

$$D(\varepsilon)d\varepsilon = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \varepsilon^{\frac{1}{2}} d\varepsilon$$

- Number of particles:

$$N = \sum a_l = \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l} \pm 1} = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\alpha+\beta\varepsilon} \pm 1}.$$

- Internal energy:

$$U = \sum a_l \varepsilon_l = \sum \frac{\omega_l \varepsilon_l}{e^{\alpha+\beta\varepsilon_l} \pm 1} = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{3}{2}} d\varepsilon}{e^{\alpha+\beta\varepsilon} \pm 1}.$$

- Set $x = \beta\varepsilon$,

$$N = g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} \int_0^\infty \frac{x^{\frac{1}{2}} dx}{e^{\alpha+x} \pm 1},$$

$$U = g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} kT \int_0^\infty \frac{x^{\frac{3}{2}} dx}{e^{\alpha+x} \pm 1}.$$

- As $e^{-\alpha} < 1$, $x > 0$, $\Rightarrow e^{-\alpha-x} < 1$.

$$\frac{1}{1 \pm e^{-\alpha-x}} \simeq 1 \mp e^{-\alpha-x};$$

$$D(\varepsilon)d\varepsilon = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \varepsilon^{\frac{1}{2}} d\varepsilon$$

- Number of particles:

$$N = \sum a_l = \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l} \pm 1} = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\alpha+\beta\varepsilon} \pm 1}.$$

- Internal energy:

$$U = \sum a_l \varepsilon_l = \sum \frac{\omega_l \varepsilon_l}{e^{\alpha+\beta\varepsilon_l} \pm 1} = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{3}{2}} d\varepsilon}{e^{\alpha+\beta\varepsilon} \pm 1}.$$

- Set $x = \beta\varepsilon$,

$$N = g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} \int_0^\infty \frac{x^{\frac{1}{2}} dx}{e^{\alpha+x} \pm 1},$$

$$U = g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} kT \int_0^\infty \frac{x^{\frac{3}{2}} dx}{e^{\alpha+x} \pm 1}.$$

- As $e^{-\alpha} < 1$, $x > 0$, $\Rightarrow e^{-\alpha-x} < 1$.

$$\frac{1}{1 \pm e^{-\alpha-x}} \simeq 1 \mp e^{-\alpha-x};$$

$$\frac{1}{e^{\alpha+x} \pm 1} = \frac{1}{e^{\alpha+x} (1 \pm e^{-\alpha-x})}$$

$$D(\varepsilon)d\varepsilon = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \varepsilon^{\frac{1}{2}} d\varepsilon$$

- Number of particles:

$$N = \sum a_l = \sum \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l} \pm 1} = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\alpha+\beta\varepsilon} \pm 1}.$$

- Internal energy:

$$U = \sum a_l \varepsilon_l = \sum \frac{\omega_l \varepsilon_l}{e^{\alpha+\beta\varepsilon_l} \pm 1} = g \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{3}{2}} d\varepsilon}{e^{\alpha+\beta\varepsilon} \pm 1}.$$

- Set $x = \beta\varepsilon$,

$$N = g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} \int_0^\infty \frac{x^{\frac{1}{2}} dx}{e^{\alpha+x} \pm 1},$$

$$U = g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} kT \int_0^\infty \frac{x^{\frac{3}{2}} dx}{e^{\alpha+x} \pm 1}.$$

- As $e^{-\alpha} < 1$, $x > 0$, $\Rightarrow e^{-\alpha-x} < 1$.

$$\frac{1}{1 \pm e^{-\alpha-x}} \simeq 1 \mp e^{-\alpha-x};$$

$$\frac{1}{e^{\alpha+x} \pm 1} = \frac{1}{e^{\alpha+x}(1 \pm e^{-\alpha-x})} \simeq e^{-\alpha-x} (1 \mp e^{-\alpha-x}).$$

$$\frac{1}{e^{\alpha+x} \pm 1} \simeq e^{-\alpha-x} (1 \mp e^{-\alpha-x})$$

- Math: $\Gamma(n) \equiv \int_0^\infty e^{-x} x^{n-1} dx;$
 $\Gamma(n + \frac{1}{2}) = (n - \frac{1}{2}) \cdot (n - \frac{3}{2}) \dots \frac{1}{2} \sqrt{\pi}.$

$$\frac{1}{e^{\alpha+x} \pm 1} \simeq e^{-\alpha-x} (1 \mp e^{-\alpha-x})$$

- Math: $\Gamma(n) \equiv \int_0^\infty e^{-x} x^{n-1} dx;$
 $\Gamma(n + \frac{1}{2}) = (n - \frac{1}{2}) \cdot (n - \frac{3}{2}) \dots \frac{1}{2} \sqrt{\pi}.$
- $N = g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} \int_0^\infty \frac{x^{\frac{1}{2}} dx}{e^{\alpha+x} \pm 1}$

$$\frac{1}{e^{\alpha+x} \pm 1} \simeq e^{-\alpha-x} (1 \mp e^{-\alpha-x})$$

- Math: $\Gamma(n) \equiv \int_0^\infty e^{-x} x^{n-1} dx;$
 $\Gamma(n + \frac{1}{2}) = (n - \frac{1}{2}) \cdot (n - \frac{3}{2}) \dots \frac{1}{2} \sqrt{\pi}.$
- $N = g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} \int_0^\infty \frac{x^{\frac{1}{2}} dx}{e^{\alpha+x} \pm 1}$
 $\simeq g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} \int_0^\infty e^{-\alpha-x} (1 \mp e^{-\alpha-x}) x^{\frac{1}{2}} dx$

$$\frac{1}{e^{\alpha+x} \pm 1} \simeq e^{-\alpha-x} (1 \mp e^{-\alpha-x})$$

- Math: $\Gamma(n) \equiv \int_0^\infty e^{-x} x^{n-1} dx$;
 $\Gamma(n + \frac{1}{2}) = (n - \frac{1}{2}) \cdot (n - \frac{3}{2}) \dots \frac{1}{2} \sqrt{\pi}$.
- $N = g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} \int_0^\infty \frac{x^{\frac{1}{2}} dx}{e^{\alpha+x} \pm 1}$
 $\simeq g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} \int_0^\infty e^{-\alpha-x} (1 \mp e^{-\alpha-x}) x^{\frac{1}{2}} dx$
 $= g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} e^{-\alpha}$
 $\cdot \left[\int_0^\infty e^{-x} x^{\frac{1}{2}} dx \mp \int_0^\infty e^{-\alpha} e^{-2x} x^{\frac{1}{2}} dx \right]$

$$\frac{1}{e^{\alpha+x} \pm 1} \simeq e^{-\alpha-x} (1 \mp e^{-\alpha-x})$$

- Math: $\Gamma(n) \equiv \int_0^\infty e^{-x} x^{n-1} dx$;
 $\Gamma(n + \frac{1}{2}) = (n - \frac{1}{2}) \cdot (n - \frac{3}{2}) \dots \frac{1}{2} \sqrt{\pi}$.
- $N = g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} \int_0^\infty \frac{x^{\frac{1}{2}} dx}{e^{\alpha+x} \pm 1}$
 $\simeq g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} \int_0^\infty e^{-\alpha-x} (1 \mp e^{-\alpha-x}) x^{\frac{1}{2}} dx$
 $= g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} e^{-\alpha}$
 $\cdot \left[\int_0^\infty e^{-x} x^{\frac{1}{2}} dx \mp \int_0^\infty e^{-\alpha} e^{-2x} x^{\frac{1}{2}} dx \right]$
 $= g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} e^{-\alpha}$
 $\cdot \left[\int_0^\infty e^{-x} x^{\frac{1}{2}} dx \mp e^{-\alpha} 2^{-\frac{3}{2}} \int_0^\infty e^{-2x} (2x)^{\frac{1}{2}} d(2x) \right]$

$$\frac{1}{e^{\alpha+x} \pm 1} \simeq e^{-\alpha-x} (1 \mp e^{-\alpha-x})$$

- Math: $\Gamma(n) \equiv \int_0^\infty e^{-x} x^{n-1} dx$;
 $\Gamma(n + \frac{1}{2}) = (n - \frac{1}{2}) \cdot (n - \frac{3}{2}) \dots \frac{1}{2} \sqrt{\pi}$.
- $N = g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} \int_0^\infty \frac{x^{\frac{1}{2}} dx}{e^{\alpha+x} \pm 1}$
 $\simeq g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} \int_0^\infty e^{-\alpha-x} (1 \mp e^{-\alpha-x}) x^{\frac{1}{2}} dx$
 $= g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} e^{-\alpha}$
 $\cdot \left[\int_0^\infty e^{-x} x^{\frac{1}{2}} dx \mp \int_0^\infty e^{-\alpha} e^{-2x} x^{\frac{1}{2}} dx \right]$
 $= g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} e^{-\alpha}$
 $\cdot \left[\int_0^\infty e^{-x} x^{\frac{1}{2}} dx \mp e^{-\alpha} 2^{-\frac{3}{2}} \int_0^\infty e^{-2x} (2x)^{\frac{1}{2}} d(2x) \right]$
 $= g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} e^{-\alpha} \left[\Gamma(\frac{3}{2}) \mp e^{-\alpha} 2^{-\frac{3}{2}} \Gamma(\frac{3}{2}) \right]$

$$\frac{1}{e^{\alpha+x} \pm 1} \simeq e^{-\alpha-x} (1 \mp e^{-\alpha-x})$$

- Math: $\Gamma(n) \equiv \int_0^\infty e^{-x} x^{n-1} dx$;
 $\Gamma(n + \frac{1}{2}) = (n - \frac{1}{2}) \cdot (n - \frac{3}{2}) \dots \frac{1}{2} \sqrt{\pi}$.
- $$\begin{aligned}
 N &= g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} \int_0^\infty \frac{x^{\frac{1}{2}} dx}{e^{\alpha+x} \pm 1} \\
 &\simeq g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} \int_0^\infty e^{-\alpha-x} (1 \mp e^{-\alpha-x}) x^{\frac{1}{2}} dx \\
 &= g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} e^{-\alpha} \\
 &\quad \cdot \left[\int_0^\infty e^{-x} x^{\frac{1}{2}} dx \mp \int_0^\infty e^{-\alpha} e^{-2x} x^{\frac{1}{2}} dx \right] \\
 &= g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} e^{-\alpha} \\
 &\quad \cdot \left[\int_0^\infty e^{-x} x^{\frac{1}{2}} dx \mp e^{-\alpha} 2^{-\frac{3}{2}} \int_0^\infty e^{-2x} (2x)^{\frac{1}{2}} d(2x) \right] \\
 &= g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} e^{-\alpha} \left[\Gamma(\frac{3}{2}) \mp e^{-\alpha} 2^{-\frac{3}{2}} \Gamma(\frac{3}{2}) \right] \\
 &= g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} e^{-\alpha} \frac{\sqrt{\pi}}{2} (1 \mp e^{-\alpha} 2^{-\frac{3}{2}}) \dots
 \end{aligned}$$

$$\frac{1}{e^{\alpha+x} \pm 1} = e^{-\alpha-x} (1 \mp e^{-\alpha-x}), \quad \Gamma(n + \frac{1}{2}) = (n - \frac{1}{2}) \cdots \frac{1}{2} \sqrt{\pi}$$

- $N = g\left(\frac{2\pi mkT}{h^2}\right)^{\frac{3}{2}} V e^{-\alpha} (1 \mp 2^{-\frac{3}{2}} e^{-\alpha}).$

$$\frac{1}{e^{\alpha+x} \pm 1} = e^{-\alpha-x} (1 \mp e^{-\alpha-x}), \quad \Gamma(n + \frac{1}{2}) = (n - \frac{1}{2}) \cdots \frac{1}{2} \sqrt{\pi}$$

- $N = g\left(\frac{2\pi mkT}{h^2}\right)^{\frac{3}{2}} V e^{-\alpha} (1 \mp 2^{-\frac{3}{2}} e^{-\alpha}).$
- $U = g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} kT \int_0^\infty \frac{x^{\frac{3}{2}} dx}{e^{\alpha+x} \pm 1}$

$$\frac{1}{e^{\alpha+x}\pm 1} = e^{-\alpha-x}(1 \mp e^{-\alpha-x}), \quad \Gamma(n + \frac{1}{2}) = (n - \frac{1}{2}) \cdots \frac{1}{2} \sqrt{\pi}$$

- $N = g\left(\frac{2\pi mkT}{h^2}\right)^{\frac{3}{2}} V e^{-\alpha} (1 \mp 2^{-\frac{3}{2}} e^{-\alpha}).$
- $U = g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} kT \int_0^\infty \frac{x^{\frac{3}{2}} dx}{e^{\alpha+x}\pm 1}$
 $\simeq g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} kT \int_0^\infty e^{-\alpha-x} (1 \mp e^{-\alpha-x}) x^{\frac{3}{2}} dx$

$$\frac{1}{e^{\alpha+x}\pm 1} = e^{-\alpha-x}(1 \mp e^{-\alpha-x}), \quad \Gamma(n + \frac{1}{2}) = (n - \frac{1}{2}) \cdots \frac{1}{2} \sqrt{\pi}$$

- $N = g\left(\frac{2\pi mkT}{h^2}\right)^{\frac{3}{2}} V e^{-\alpha} (1 \mp 2^{-\frac{3}{2}} e^{-\alpha}).$
- $$\begin{aligned} U &= g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} kT \int_0^\infty \frac{x^{\frac{3}{2}} dx}{e^{\alpha+x}\pm 1} \\ &\simeq g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} kT \int_0^\infty e^{-\alpha-x} (1 \mp e^{-\alpha-x}) x^{\frac{3}{2}} dx \\ &= g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} kT e^{-\alpha} [\Gamma(\frac{5}{2}) \mp e^{-\alpha} 2^{-\frac{5}{2}} \Gamma(\frac{5}{2})] \end{aligned}$$

$$\frac{1}{e^{\alpha+x} \pm 1} = e^{-\alpha-x} (1 \mp e^{-\alpha-x}), \quad \Gamma(n + \frac{1}{2}) = (n - \frac{1}{2}) \cdots \frac{1}{2} \sqrt{\pi}$$

- $N = g \left(\frac{2\pi m k T}{h^2} \right)^{\frac{3}{2}} V e^{-\alpha} (1 \mp 2^{-\frac{3}{2}} e^{-\alpha}).$
- $$\begin{aligned} U &= g \frac{2\pi V}{h^3} (2m k T)^{\frac{3}{2}} k T \int_0^\infty \frac{x^{\frac{3}{2}} dx}{e^{\alpha+x} \pm 1} \\ &\simeq g \frac{2\pi V}{h^3} (2m k T)^{\frac{3}{2}} k T \int_0^\infty e^{-\alpha-x} (1 \mp e^{-\alpha-x}) x^{\frac{3}{2}} dx \\ &= g \frac{2\pi V}{h^3} (2m k T)^{\frac{3}{2}} k T e^{-\alpha} [\Gamma(\frac{5}{2}) \mp e^{-\alpha} 2^{-\frac{5}{2}} \Gamma(\frac{5}{2})] \\ &= g \frac{2\pi V}{h^3} (2m k T)^{\frac{3}{2}} k T e^{-\alpha} \frac{3}{2} \frac{\sqrt{\pi}}{2} (1 \mp e^{-\alpha} 2^{-\frac{5}{2}}) \end{aligned}$$

$$\frac{1}{e^{\alpha+x} \pm 1} = e^{-\alpha-x} (1 \mp e^{-\alpha-x}), \quad \Gamma(n + \frac{1}{2}) = (n - \frac{1}{2}) \cdots \frac{1}{2} \sqrt{\pi}$$

- $N = g \left(\frac{2\pi m k T}{h^2} \right)^{\frac{3}{2}} V e^{-\alpha} (1 \mp 2^{-\frac{3}{2}} e^{-\alpha}).$
- $$\begin{aligned} U &= g \frac{2\pi V}{h^3} (2m k T)^{\frac{3}{2}} k T \int_0^\infty \frac{x^{\frac{3}{2}} dx}{e^{\alpha+x} \pm 1} \\ &\simeq g \frac{2\pi V}{h^3} (2m k T)^{\frac{3}{2}} k T \int_0^\infty e^{-\alpha-x} (1 \mp e^{-\alpha-x}) x^{\frac{3}{2}} dx \\ &= g \frac{2\pi V}{h^3} (2m k T)^{\frac{3}{2}} k T e^{-\alpha} [\Gamma(\frac{5}{2}) \mp e^{-\alpha} 2^{-\frac{5}{2}} \Gamma(\frac{5}{2})] \\ &= g \frac{2\pi V}{h^3} (2m k T)^{\frac{3}{2}} k T e^{-\alpha} \frac{3}{2} \frac{\sqrt{\pi}}{2} (1 \mp e^{-\alpha} 2^{-\frac{5}{2}}) \\ &= g \left(\frac{2\pi m k T}{h^2} \right)^{\frac{3}{2}} V \frac{3}{2} k T e^{-\alpha} (1 \mp 2^{-\frac{5}{2}} e^{-\alpha}). \end{aligned}$$

$$\frac{1}{e^{\alpha+x} \pm 1} = e^{-\alpha-x} (1 \mp e^{-\alpha-x}), \quad \Gamma(n + \frac{1}{2}) = (n - \frac{1}{2}) \cdots \frac{1}{2} \sqrt{\pi}$$

- $N = g \left(\frac{2\pi mkT}{h^2} \right)^{\frac{3}{2}} V e^{-\alpha} (1 \mp 2^{-\frac{3}{2}} e^{-\alpha}).$
- $$\begin{aligned} U &= g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} kT \int_0^\infty \frac{x^{\frac{3}{2}} dx}{e^{\alpha+x} \pm 1} \\ &\simeq g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} kT \int_0^\infty e^{-\alpha-x} (1 \mp e^{-\alpha-x}) x^{\frac{3}{2}} dx \\ &= g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} kT e^{-\alpha} [\Gamma(\frac{5}{2}) \mp e^{-\alpha} 2^{-\frac{5}{2}} \Gamma(\frac{5}{2})] \\ &= g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} kT e^{-\alpha} \frac{3}{2} \frac{\sqrt{\pi}}{2} (1 \mp e^{-\alpha} 2^{-\frac{5}{2}}) \\ &= g \left(\frac{2\pi mkT}{h^2} \right)^{\frac{3}{2}} V \frac{3}{2} kT e^{-\alpha} (1 \mp 2^{-\frac{5}{2}} e^{-\alpha}). \end{aligned}$$
- $$\frac{U}{N} = \frac{g \left(\frac{2\pi mkT}{h^2} \right)^{\frac{3}{2}} V \frac{3}{2} kT e^{-\alpha} (1 \mp 2^{-\frac{5}{2}} e^{-\alpha})}{g \left(\frac{2\pi mkT}{h^2} \right)^{\frac{3}{2}} V e^{-\alpha} (1 \mp 2^{-\frac{3}{2}} e^{-\alpha})}$$

$$\frac{1}{e^{\alpha+x} \pm 1} = e^{-\alpha-x} (1 \mp e^{-\alpha-x}), \quad \Gamma(n + \frac{1}{2}) = (n - \frac{1}{2}) \cdots \frac{1}{2} \sqrt{\pi}$$

- $N = g \left(\frac{2\pi m k T}{h^2} \right)^{\frac{3}{2}} V e^{-\alpha} (1 \mp 2^{-\frac{5}{2}} e^{-\alpha}).$
- $$\begin{aligned} U &= g \frac{2\pi V}{h^3} (2m k T)^{\frac{3}{2}} k T \int_0^\infty \frac{x^{\frac{3}{2}} dx}{e^{\alpha+x} \pm 1} \\ &\simeq g \frac{2\pi V}{h^3} (2m k T)^{\frac{3}{2}} k T \int_0^\infty e^{-\alpha-x} (1 \mp e^{-\alpha-x}) x^{\frac{3}{2}} dx \\ &= g \frac{2\pi V}{h^3} (2m k T)^{\frac{3}{2}} k T e^{-\alpha} [\Gamma(\frac{5}{2}) \mp e^{-\alpha} 2^{-\frac{5}{2}} \Gamma(\frac{5}{2})] \\ &= g \frac{2\pi V}{h^3} (2m k T)^{\frac{3}{2}} k T e^{-\alpha} \frac{3}{2} \frac{\sqrt{\pi}}{2} (1 \mp e^{-\alpha} 2^{-\frac{5}{2}}) \\ &= g \left(\frac{2\pi m k T}{h^2} \right)^{\frac{3}{2}} V \frac{3}{2} k T e^{-\alpha} (1 \mp 2^{-\frac{5}{2}} e^{-\alpha}). \end{aligned}$$
- $$\frac{U}{N} = \frac{g \left(\frac{2\pi m k T}{h^2} \right)^{\frac{3}{2}} V \frac{3}{2} k T e^{-\alpha} (1 \mp 2^{-\frac{5}{2}} e^{-\alpha})}{g \left(\frac{2\pi m k T}{h^2} \right)^{\frac{3}{2}} V e^{-\alpha} (1 \mp 2^{-\frac{3}{2}} e^{-\alpha})} = \frac{3}{2} k T \frac{1 \mp 2^{-\frac{5}{2}} e^{-\alpha}}{1 \mp 2^{-\frac{3}{2}} e^{-\alpha}}$$

$$\frac{1}{e^{\alpha+x} \pm 1} = e^{-\alpha-x} (1 \mp e^{-\alpha-x}), \quad \Gamma(n + \frac{1}{2}) = (n - \frac{1}{2}) \cdots \frac{1}{2} \sqrt{\pi}$$

- $N = g\left(\frac{2\pi mkT}{h^2}\right)^{\frac{3}{2}} V e^{-\alpha} (1 \mp 2^{-\frac{3}{2}} e^{-\alpha}).$
- $$\begin{aligned} U &= g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} kT \int_0^\infty \frac{x^{\frac{3}{2}} dx}{e^{\alpha+x} \pm 1} \\ &\simeq g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} kT \int_0^\infty e^{-\alpha-x} (1 \mp e^{-\alpha-x}) x^{\frac{3}{2}} dx \\ &= g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} kT e^{-\alpha} [\Gamma(\frac{5}{2}) \mp e^{-\alpha} 2^{-\frac{5}{2}} \Gamma(\frac{5}{2})] \\ &= g \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} kT e^{-\alpha} \frac{3}{2} \frac{\sqrt{\pi}}{2} (1 \mp e^{-\alpha} 2^{-\frac{5}{2}}) \\ &= g\left(\frac{2\pi mkT}{h^2}\right)^{\frac{3}{2}} V \frac{3}{2} kT e^{-\alpha} (1 \mp 2^{-\frac{5}{2}} e^{-\alpha}). \end{aligned}$$
- $$\begin{aligned} \frac{U}{N} &= \frac{g\left(\frac{2\pi mkT}{h^2}\right)^{\frac{3}{2}} V \frac{3}{2} kT e^{-\alpha} (1 \mp 2^{-\frac{5}{2}} e^{-\alpha})}{g\left(\frac{2\pi mkT}{h^2}\right)^{\frac{3}{2}} V e^{-\alpha} (1 \mp 2^{-\frac{3}{2}} e^{-\alpha})} = \frac{3}{2} kT \frac{1 \mp 2^{-\frac{5}{2}} e^{-\alpha}}{1 \mp 2^{-\frac{3}{2}} e^{-\alpha}} \\ &\simeq \frac{3}{2} kT (1 \mp 2^{-\frac{5}{2}} e^{-\alpha}) (1 \pm 2^{-\frac{3}{2}} e^{-\alpha}) \end{aligned}$$

$$\frac{1}{e^{\alpha+x} \pm 1} = e^{-\alpha-x} (1 \mp e^{-\alpha-x}), \quad \Gamma(n + \frac{1}{2}) = (n - \frac{1}{2}) \cdots \frac{1}{2} \sqrt{\pi}$$

- $N = g \left(\frac{2\pi m k T}{h^2} \right)^{\frac{3}{2}} V e^{-\alpha} (1 \mp 2^{-\frac{3}{2}} e^{-\alpha}).$
- $$\begin{aligned} U &= g \frac{2\pi V}{h^3} (2m k T)^{\frac{3}{2}} k T \int_0^\infty \frac{x^{\frac{3}{2}} dx}{e^{\alpha+x} \pm 1} \\ &\simeq g \frac{2\pi V}{h^3} (2m k T)^{\frac{3}{2}} k T \int_0^\infty e^{-\alpha-x} (1 \mp e^{-\alpha-x}) x^{\frac{3}{2}} dx \\ &= g \frac{2\pi V}{h^3} (2m k T)^{\frac{3}{2}} k T e^{-\alpha} [\Gamma(\frac{5}{2}) \mp e^{-\alpha} 2^{-\frac{5}{2}} \Gamma(\frac{5}{2})] \\ &= g \frac{2\pi V}{h^3} (2m k T)^{\frac{3}{2}} k T e^{-\alpha} \frac{3}{2} \frac{\sqrt{\pi}}{2} (1 \mp e^{-\alpha} 2^{-\frac{5}{2}}) \\ &= g \left(\frac{2\pi m k T}{h^2} \right)^{\frac{3}{2}} V \frac{3}{2} k T e^{-\alpha} (1 \mp 2^{-\frac{5}{2}} e^{-\alpha}). \end{aligned}$$
- $$\begin{aligned} \frac{U}{N} &= \frac{g \left(\frac{2\pi m k T}{h^2} \right)^{\frac{3}{2}} V \frac{3}{2} k T e^{-\alpha} (1 \mp 2^{-\frac{5}{2}} e^{-\alpha})}{g \left(\frac{2\pi m k T}{h^2} \right)^{\frac{3}{2}} V e^{-\alpha} (1 \mp 2^{-\frac{3}{2}} e^{-\alpha})} = \frac{3}{2} k T \frac{1 \mp 2^{-\frac{5}{2}} e^{-\alpha}}{1 \mp 2^{-\frac{3}{2}} e^{-\alpha}} \\ &\simeq \frac{3}{2} k T (1 \mp 2^{-\frac{5}{2}} e^{-\alpha}) (1 \pm 2^{-\frac{3}{2}} e^{-\alpha}) \\ &= \frac{3}{2} k T [1 + (\pm 2^{-\frac{3}{2}} \mp 2^{-\frac{5}{2}}) e^{-\alpha} - 2^{-4} e^{-2\alpha}] \end{aligned}$$

$$\frac{1}{e^{\alpha+x} \pm 1} = e^{-\alpha-x} (1 \mp e^{-\alpha-x}), \quad \Gamma(n + \frac{1}{2}) = (n - \frac{1}{2}) \cdots \frac{1}{2} \sqrt{\pi}$$

- $N = g \left(\frac{2\pi m k T}{h^2} \right)^{\frac{3}{2}} V e^{-\alpha} (1 \mp 2^{-\frac{3}{2}} e^{-\alpha}).$
- $$\begin{aligned} U &= g \frac{2\pi V}{h^3} (2m k T)^{\frac{3}{2}} k T \int_0^\infty \frac{x^{\frac{3}{2}} dx}{e^{\alpha+x} \pm 1} \\ &\simeq g \frac{2\pi V}{h^3} (2m k T)^{\frac{3}{2}} k T \int_0^\infty e^{-\alpha-x} (1 \mp e^{-\alpha-x}) x^{\frac{3}{2}} dx \\ &= g \frac{2\pi V}{h^3} (2m k T)^{\frac{3}{2}} k T e^{-\alpha} [\Gamma(\frac{5}{2}) \mp e^{-\alpha} 2^{-\frac{5}{2}} \Gamma(\frac{5}{2})] \\ &= g \frac{2\pi V}{h^3} (2m k T)^{\frac{3}{2}} k T e^{-\alpha} \frac{3}{2} \frac{\sqrt{\pi}}{2} (1 \mp e^{-\alpha} 2^{-\frac{5}{2}}) \\ &= g \left(\frac{2\pi m k T}{h^2} \right)^{\frac{3}{2}} V \frac{3}{2} k T e^{-\alpha} (1 \mp 2^{-\frac{5}{2}} e^{-\alpha}). \end{aligned}$$
- $$\begin{aligned} \frac{U}{N} &= \frac{g \left(\frac{2\pi m k T}{h^2} \right)^{\frac{3}{2}} V \frac{3}{2} k T e^{-\alpha} (1 \mp 2^{-\frac{5}{2}} e^{-\alpha})}{g \left(\frac{2\pi m k T}{h^2} \right)^{\frac{3}{2}} V e^{-\alpha} (1 \mp 2^{-\frac{3}{2}} e^{-\alpha})} = \frac{3}{2} k T \frac{1 \mp 2^{-\frac{5}{2}} e^{-\alpha}}{1 \mp 2^{-\frac{3}{2}} e^{-\alpha}} \\ &\simeq \frac{3}{2} k T (1 \mp 2^{-\frac{5}{2}} e^{-\alpha}) (1 \pm 2^{-\frac{3}{2}} e^{-\alpha}) \\ &= \frac{3}{2} k T [1 + (\pm 2^{-\frac{3}{2}} \mp 2^{-\frac{5}{2}}) e^{-\alpha} - 2^{-4} e^{-2\alpha}] \\ &\simeq \frac{3}{2} k T (1 \pm 2^{-\frac{5}{2}} e^{-\alpha}). \end{aligned}$$

U of weak degenerated ideal Fermion(Boson) gas

- $\frac{U}{N} \simeq \frac{3}{2}kT(1 \pm 2^{-\frac{5}{2}}e^{-\alpha}).$

U of weak degenerated ideal Fermion(Boson) gas

- $\frac{U}{N} \simeq \frac{3}{2}kT(1 \pm 2^{-\frac{5}{2}}e^{-\alpha}).$
- For small $e^{-\alpha}$, $N = g(\frac{2\pi mkT}{h^2})^{\frac{3}{2}}Ve^{-\alpha}(1 \mp 2^{-\frac{3}{2}}e^{-\alpha})$

U of weak degenerated ideal Fermion(Boson) gas

- $\frac{U}{N} \simeq \frac{3}{2}kT(1 \pm 2^{-\frac{5}{2}}e^{-\alpha}).$
- For small $e^{-\alpha}$, $N = g(\frac{2\pi mkT}{h^2})^{\frac{3}{2}}Ve^{-\alpha}(1 \mp 2^{-\frac{3}{2}}e^{-\alpha})$
 $\simeq g(\frac{2\pi mkT}{h^2})^{\frac{3}{2}}Ve^{-\alpha},$

U of weak degenerated ideal Fermion(Boson) gas

- $\frac{U}{N} \simeq \frac{3}{2}kT(1 \pm 2^{-\frac{5}{2}}e^{-\alpha})$.
- For small $e^{-\alpha}$, $N = g\left(\frac{2\pi mkT}{h^2}\right)^{\frac{3}{2}}Ve^{-\alpha}(1 \mp 2^{-\frac{3}{2}}e^{-\alpha})$
 $\simeq g\left(\frac{2\pi mkT}{h^2}\right)^{\frac{3}{2}}Ve^{-\alpha}$,
 $\Rightarrow e^{-\alpha} \simeq \frac{N}{V}\left(\frac{h^2}{2\pi mkT}\right)^{\frac{3}{2}}g^{-1}$.

U of weak degenerated ideal Fermion(Boson) gas

- $\frac{U}{N} \simeq \frac{3}{2}kT(1 \pm 2^{-\frac{5}{2}}e^{-\alpha})$.
- For small $e^{-\alpha}$, $N = g\left(\frac{2\pi mkT}{h^2}\right)^{\frac{3}{2}}Ve^{-\alpha}(1 \mp 2^{-\frac{3}{2}}e^{-\alpha})$
 $\simeq g\left(\frac{2\pi mkT}{h^2}\right)^{\frac{3}{2}}Ve^{-\alpha}$,
 $\Rightarrow e^{-\alpha} \simeq \frac{N}{V}\left(\frac{h^2}{2\pi mkT}\right)^{\frac{3}{2}}g^{-1}$.
- $U \simeq N\frac{3}{2}kT(1 \pm 2^{-\frac{5}{2}}e^{-\alpha})$

U of weak degenerated ideal Fermion(Boson) gas

- $\frac{U}{N} \simeq \frac{3}{2}kT(1 \pm 2^{-\frac{5}{2}}e^{-\alpha}).$
- For small $e^{-\alpha}$, $N = g\left(\frac{2\pi mkT}{h^2}\right)^{\frac{3}{2}}Ve^{-\alpha}(1 \mp 2^{-\frac{3}{2}}e^{-\alpha})$
 $\simeq g\left(\frac{2\pi mkT}{h^2}\right)^{\frac{3}{2}}Ve^{-\alpha},$
 $\Rightarrow e^{-\alpha} \simeq \frac{N}{V}\left(\frac{h^2}{2\pi mkT}\right)^{\frac{3}{2}}g^{-1}.$
- $U \simeq N\frac{3}{2}kT(1 \pm 2^{-\frac{5}{2}}e^{-\alpha})$
 $\simeq \frac{3}{2}NkT[1 \pm 2^{-\frac{5}{2}}\frac{N}{V}\left(\frac{h^2}{2\pi mkT}\right)^{\frac{3}{2}}g^{-1}]$

U of weak degenerated ideal Fermion(Boson) gas

- $\frac{U}{N} \simeq \frac{3}{2}kT(1 \pm 2^{-\frac{5}{2}}e^{-\alpha}).$
- For small $e^{-\alpha}$, $N = g\left(\frac{2\pi mkT}{h^2}\right)^{\frac{3}{2}}Ve^{-\alpha}(1 \mp 2^{-\frac{3}{2}}e^{-\alpha})$
 $\simeq g\left(\frac{2\pi mkT}{h^2}\right)^{\frac{3}{2}}Ve^{-\alpha},$
 $\Rightarrow e^{-\alpha} \simeq \frac{N}{V}\left(\frac{h^2}{2\pi mkT}\right)^{\frac{3}{2}}g^{-1}.$
- $U \simeq N\frac{3}{2}kT(1 \pm 2^{-\frac{5}{2}}e^{-\alpha})$
 $\simeq \frac{3}{2}NkT\left[1 \pm 2^{-\frac{5}{2}}\frac{N}{V}\left(\frac{h^2}{2\pi mkT}\right)^{\frac{3}{2}}g^{-1}\right]$
 $= \frac{3}{2}NkT(1 \pm 2^{-\frac{5}{2}}g^{-1}n\lambda^3).$

U of weak degenerated ideal Fermion(Boson) gas

- $\frac{U}{N} \simeq \frac{3}{2}kT(1 \pm 2^{-\frac{5}{2}}e^{-\alpha}).$
- For small $e^{-\alpha}$, $N = g\left(\frac{2\pi mkT}{h^2}\right)^{\frac{3}{2}}Ve^{-\alpha}(1 \mp 2^{-\frac{3}{2}}e^{-\alpha})$
 $\simeq g\left(\frac{2\pi mkT}{h^2}\right)^{\frac{3}{2}}Ve^{-\alpha},$
 $\Rightarrow e^{-\alpha} \simeq \frac{N}{V}\left(\frac{h^2}{2\pi mkT}\right)^{\frac{3}{2}}g^{-1}.$
- $U \simeq N\frac{3}{2}kT(1 \pm 2^{-\frac{5}{2}}e^{-\alpha})$
 $\simeq \frac{3}{2}NkT\left[1 \pm 2^{-\frac{5}{2}}\frac{N}{V}\left(\frac{h^2}{2\pi mkT}\right)^{\frac{3}{2}}g^{-1}\right]$
 $= \frac{3}{2}NkT(1 \pm 2^{-\frac{5}{2}}g^{-1}n\lambda^3).$

1st term, same as the Boltzmann distribution; 2nd term, modification by the identical principle of particles (Fermions, repel; Bosons, attract).

Table of contents

1 §8. Bose statistics and Fermi statistics

- 8.1 Thermal parameters in statistics
- 8.2 U of weak degenerated ideal Fermion(Boson) gas
- 8.3 Bose-Einstein Condensate (BEC)
- 8.4 Photon gas
- 8.5 Free electron gas in metal

8.3 Bose-Einstein condensate (BEC)

8.3 Bose-Einstein condensate (BEC)

- If even $n\lambda^3 < 1$ does not satisfy, for Bosons, Bose-Einstein Condensate.

8.3 Bose-Einstein condensate (BEC)

- If even $n\lambda^3 < 1$ does not satisfy, for Bosons, Bose-Einstein Condensate.
- Object: N identical Bosons, ideal gas, nearly independent, no spin degeneracy, given (T, V) .

8.3 Bose-Einstein condensate (BEC)

- If even $n\lambda^3 < 1$ does not satisfy, for Bosons, Bose-Einstein Condensate.
- Object: N identical Bosons, ideal gas, nearly independent, no spin degeneracy, given (T, V) .
- $$a_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1} = \frac{\omega_l}{e^{\frac{\varepsilon_l-\mu}{kT}}-1}.$$

8.3 Bose-Einstein condensate (BEC)

- If even $n\lambda^3 < 1$ does not satisfy, for Bosons, Bose-Einstein Condensate.
- Object: N identical Bosons, ideal gas, nearly independent, no spin degeneracy, given (T, V) .
- $a_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1} = \frac{\omega_l}{e^{\frac{\varepsilon_l-\mu}{kT}}-1}$.
- $a_l > 0$ requires: $e^{\frac{\varepsilon_l-\mu}{kT}} - 1 > 0$

8.3 Bose-Einstein condensate (BEC)

- If even $n\lambda^3 < 1$ does not satisfy, for Bosons, Bose-Einstein Condensate.
- Object: N identical Bosons, ideal gas, nearly independent, no spin degeneracy, given (T, V) .
- $a_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1} = \frac{\omega_l}{e^{\frac{\varepsilon_l-\mu}{kT}}-1}$.
- $a_l > 0$ requires: $e^{\frac{\varepsilon_l-\mu}{kT}} - 1 > 0 \Rightarrow \varepsilon_l > \mu$, so $\varepsilon_0 > \mu$.

8.3 Bose-Einstein condensate (BEC)

- If even $n\lambda^3 < 1$ does not satisfy, for Bosons, Bose-Einstein Condensate.
- Object: N identical Bosons, ideal gas, nearly independent, no spin degeneracy, given (T, V) .
- $a_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1} = \frac{\omega_l}{e^{\frac{\varepsilon_l-\mu}{kT}}-1}$.
- $a_l > 0$ requires: $e^{\frac{\varepsilon_l-\mu}{kT}} - 1 > 0 \Rightarrow \varepsilon_l > \mu$, so $\varepsilon_0 > \mu$.
- For ideal gas, $\varepsilon_0 = 0$, $\Rightarrow \mu < 0$.

8.3 Bose-Einstein condensate (BEC)

- If even $n\lambda^3 < 1$ does not satisfy, for Bosons, Bose-Einstein Condensate.
- Object: N identical Bosons, ideal gas, nearly independent, no spin degeneracy, given (T, V) .
- $a_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1} = \frac{\omega_l}{e^{\frac{\varepsilon_l-\mu}{kT}}-1}$.
- $a_l > 0$ requires: $e^{\frac{\varepsilon_l-\mu}{kT}} - 1 > 0 \Rightarrow \varepsilon_l > \mu$, so $\varepsilon_0 > \mu$.
- For ideal gas, $\varepsilon_0 = 0$, $\Rightarrow \mu < 0$.
- $n = \frac{1}{V} \sum a_l = \frac{1}{V} \sum \frac{\omega_l}{e^{\frac{\varepsilon_l-\mu}{kT}}-1}$.

8.3 Bose-Einstein condensate (BEC)

- If even $n\lambda^3 < 1$ does not satisfy, for Bosons, Bose-Einstein Condensate.
- Object: N identical Bosons, ideal gas, nearly independent, no spin degeneracy, given (T, V) .
- $a_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}-1} = \frac{\omega_l}{e^{\frac{\varepsilon_l-\mu}{kT}}-1}$.
- $a_l > 0$ requires: $e^{\frac{\varepsilon_l-\mu}{kT}} - 1 > 0 \Rightarrow \varepsilon_l > \mu$, so $\varepsilon_0 > \mu$.
- For ideal gas, $\varepsilon_0 = 0$, $\Rightarrow \mu < 0$.
- $n = \frac{1}{V} \sum a_l = \frac{1}{V} \sum \frac{\omega_l}{e^{\frac{\varepsilon_l-\mu}{kT}}-1}$.
 $T \downarrow$, $\mu \uparrow$, until 0. Corresponds to a critical temperature: T_c .

BEC – idea gas

- $$n = \frac{1}{V} \sum \frac{\omega_l}{e^{\frac{\varepsilon_l - 0}{kT_c}} - 1}$$

BEC – idea gas

$$\bullet n = \frac{1}{V} \sum_l \frac{\omega_l}{e^{\frac{\varepsilon_l - 0}{kT_c}} - 1} = \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT_c}} - 1}$$

BEC – idea gas

$$\begin{aligned} \bullet \quad n &= \frac{1}{V} \sum_l \frac{\omega_l}{e^{\frac{\varepsilon_l - 0}{kT_c}} - 1} = \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT_c}} - 1} \\ &= \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} (kT_c)^{\frac{3}{2}} \int_0^\infty \frac{x^{\frac{1}{2}} dx}{e^x - 1} \end{aligned}$$

BEC – idea gas

$$\begin{aligned}
 \bullet \quad n &= \frac{1}{V} \sum_l \frac{\omega_l}{e^{\frac{\varepsilon_l - 0}{kT_c}} - 1} = \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT_c}} - 1} \\
 &= \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} (kT_c)^{\frac{3}{2}} \int_0^\infty \frac{x^{\frac{1}{2}} dx}{e^x - 1} \\
 &= \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} (kT_c)^{\frac{3}{2}} \frac{\sqrt{\pi}}{2} \cdot 2.612
 \end{aligned}$$

BEC – idea gas

$$\begin{aligned}
 \bullet \quad n &= \frac{1}{V} \sum_l \frac{\omega_l}{e^{\frac{\varepsilon_l - 0}{kT_c}} - 1} = \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT_c}} - 1} \\
 &= \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} (kT_c)^{\frac{3}{2}} \int_0^\infty \frac{x^{\frac{1}{2}} dx}{e^x - 1} \\
 &= \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} (kT_c)^{\frac{3}{2}} \frac{\sqrt{\pi}}{2} \cdot 2.612 \\
 \Rightarrow T_c &= \frac{2\pi}{2.612^{\frac{2}{3}}} \frac{\hbar^2}{mk} n^{\frac{2}{3}}.
 \end{aligned}$$

BEC – idea gas

- $$n = \frac{1}{V} \sum_l \frac{\omega_l}{e^{\frac{\varepsilon_l - 0}{kT_c}} - 1} = \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT_c}} - 1}$$

$$= \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} (kT_c)^{\frac{3}{2}} \int_0^\infty \frac{x^{\frac{1}{2}} dx}{e^x - 1}$$

$$= \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} (kT_c)^{\frac{3}{2}} \frac{\sqrt{\pi}}{2} \cdot 2.612$$

$$\Rightarrow T_c = \frac{2\pi}{2.612^{\frac{2}{3}}} \frac{\hbar^2}{mk} n^{\frac{2}{3}}.$$
- Notice $n\lambda^3 = n\left(\frac{h^2}{2\pi mkT}\right)^{\frac{3}{2}},$

BEC – idea gas

- $$\begin{aligned}
 n &= \frac{1}{V} \sum_l \frac{\omega_l}{e^{\frac{\varepsilon_l - 0}{kT_c}} - 1} = \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT_c}} - 1} \\
 &= \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} (kT_c)^{\frac{3}{2}} \int_0^\infty \frac{x^{\frac{1}{2}} dx}{e^x - 1} \\
 &= \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} (kT_c)^{\frac{3}{2}} \frac{\sqrt{\pi}}{2} \cdot 2.612 \\
 &\Rightarrow T_c = \frac{2\pi}{2.612^{\frac{2}{3}}} \frac{\hbar^2}{mk} n^{\frac{2}{3}}.
 \end{aligned}$$
- Notice $n\lambda^3 = n\left(\frac{h^2}{2\pi mkT}\right)^{\frac{3}{2}},$

$$n\lambda_c^3 = n\left(\frac{h^2}{2\pi mkT_c}\right)^{\frac{3}{2}}$$

BEC – idea gas

- $$n = \frac{1}{V} \sum_l \frac{\omega_l}{e^{\frac{\varepsilon_l - 0}{kT_c}} - 1} = \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT_c}} - 1}$$

$$= \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} (kT_c)^{\frac{3}{2}} \int_0^\infty \frac{x^{\frac{1}{2}} dx}{e^x - 1}$$

$$= \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} (kT_c)^{\frac{3}{2}} \frac{\sqrt{\pi}}{2} \cdot 2.612$$

$$\Rightarrow T_c = \frac{2\pi}{2.612^{\frac{2}{3}}} \frac{\hbar^2}{mk} n^{\frac{2}{3}}.$$
- Notice $n\lambda^3 = n\left(\frac{h^2}{2\pi mkT}\right)^{\frac{3}{2}},$

$$n\lambda_c^3 = n\left(\frac{h^2}{2\pi mkT_c}\right)^{\frac{3}{2}} = n\left(\frac{h^2}{2\pi mk}\right)^{\frac{3}{2}} \frac{2.612(mk)^{\frac{3}{2}}}{(2\pi)^{\frac{3}{2}} \hbar^3 n}$$

BEC – idea gas

- $$n = \frac{1}{V} \sum_l \frac{\omega_l}{e^{\frac{\varepsilon_l - 0}{kT_c}} - 1} = \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT_c}} - 1}$$

$$= \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} (kT_c)^{\frac{3}{2}} \int_0^\infty \frac{x^{\frac{1}{2}} dx}{e^x - 1}$$

$$= \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} (kT_c)^{\frac{3}{2}} \frac{\sqrt{\pi}}{2} \cdot 2.612$$

$$\Rightarrow T_c = \frac{2\pi}{2.612^{\frac{2}{3}}} \frac{\hbar^2}{mk} n^{\frac{2}{3}}.$$
- Notice $n\lambda^3 = n\left(\frac{h^2}{2\pi mkT}\right)^{\frac{3}{2}},$

$$n\lambda_c^3 = n\left(\frac{h^2}{2\pi mkT_c}\right)^{\frac{3}{2}} = n\left(\frac{h^2}{2\pi mk}\right)^{\frac{3}{2}} \frac{2.612(mk)^{\frac{3}{2}}}{(2\pi)^{\frac{3}{2}} \hbar^3 n} = 2.612.$$

BEC – idea gas

- $$n = \frac{1}{V} \sum_l \frac{\omega_l}{e^{\frac{\varepsilon_l - 0}{kT_c}} - 1} = \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT_c}} - 1}$$

$$= \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} (kT_c)^{\frac{3}{2}} \int_0^\infty \frac{x^{\frac{1}{2}} dx}{e^x - 1}$$

$$= \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} (kT_c)^{\frac{3}{2}} \frac{\sqrt{\pi}}{2} \cdot 2.612$$

$$\Rightarrow T_c = \frac{2\pi}{2.612^{\frac{2}{3}}} \frac{\hbar^2}{mk} n^{\frac{2}{3}}.$$
- Notice $n\lambda^3 = n\left(\frac{h^2}{2\pi mkT}\right)^{\frac{3}{2}},$

$$n\lambda_c^3 = n\left(\frac{h^2}{2\pi mkT_c}\right)^{\frac{3}{2}} = n\left(\frac{h^2}{2\pi mk}\right)^{\frac{3}{2}} \frac{2.612(mk)^{\frac{3}{2}}}{(2\pi)^{\frac{3}{2}} \hbar^3 n} = 2.612.$$
- What's happening if $T < T_c$?

BEC – idea gas

- $$n = \frac{1}{V} \sum_l \frac{\omega_l}{e^{\frac{\varepsilon_l - 0}{kT_c}} - 1} = \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT_c}} - 1}$$

$$= \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} (kT_c)^{\frac{3}{2}} \int_0^\infty \frac{x^{\frac{1}{2}} dx}{e^x - 1}$$

$$= \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} (kT_c)^{\frac{3}{2}} \frac{\sqrt{\pi}}{2} \cdot 2.612$$

$$\Rightarrow T_c = \frac{2\pi}{2.612^{\frac{2}{3}}} \frac{\hbar^2}{mk} n^{\frac{2}{3}}.$$
- Notice $n\lambda^3 = n\left(\frac{h^2}{2\pi mkT}\right)^{\frac{3}{2}},$

$$n\lambda_c^3 = n\left(\frac{h^2}{2\pi mkT_c}\right)^{\frac{3}{2}} = n\left(\frac{h^2}{2\pi mk}\right)^{\frac{3}{2}} \frac{2.612(mk)^{\frac{3}{2}}}{(2\pi)^{\frac{3}{2}} \hbar^3 n} = 2.612.$$
- What's happening if $T < T_c$?
- More and more particles stay at ground state.

BEC – idea gas

- $$n = \frac{1}{V} \sum_l \frac{\omega_l}{e^{\frac{\varepsilon_l - 0}{kT_c}} - 1} = \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT_c}} - 1}$$

$$= \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} (kT_c)^{\frac{3}{2}} \int_0^\infty \frac{x^{\frac{1}{2}} dx}{e^x - 1}$$

$$= \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} (kT_c)^{\frac{3}{2}} \frac{\sqrt{\pi}}{2} \cdot 2.612$$

$$\Rightarrow T_c = \frac{2\pi}{2.612^{\frac{2}{3}}} \frac{\hbar^2}{mk} n^{\frac{2}{3}}.$$
 - Notice $n\lambda^3 = n\left(\frac{h^2}{2\pi mkT}\right)^{\frac{3}{2}},$

$$n\lambda_c^3 = n\left(\frac{h^2}{2\pi mkT_c}\right)^{\frac{3}{2}} = n\left(\frac{h^2}{2\pi mk}\right)^{\frac{3}{2}} \frac{2.612(mk)^{\frac{3}{2}}}{(2\pi)^{\frac{3}{2}} \hbar^3 n} = 2.612.$$
 - What's happening if $T < T_c$?
 - More and more particles stay at ground state.
- $\mu < 0$ still obeys, $\mu \rightarrow 0$;

BEC – idea gas

- $$n = \frac{1}{V} \sum_l \frac{\omega_l}{e^{\frac{\varepsilon_l - 0}{kT_c}} - 1} = \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT_c}} - 1}$$

$$= \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} (kT_c)^{\frac{3}{2}} \int_0^\infty \frac{x^{\frac{1}{2}} dx}{e^x - 1}$$

$$= \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} (kT_c)^{\frac{3}{2}} \frac{\sqrt{\pi}}{2} \cdot 2.612$$

$$\Rightarrow T_c = \frac{2\pi}{2.612^{\frac{2}{3}}} \frac{\hbar^2}{mk} n^{\frac{2}{3}}.$$
 - Notice $n\lambda^3 = n\left(\frac{h^2}{2\pi mkT}\right)^{\frac{3}{2}},$

$$n\lambda_c^3 = n\left(\frac{h^2}{2\pi mkT_c}\right)^{\frac{3}{2}} = n\left(\frac{h^2}{2\pi mk}\right)^{\frac{3}{2}} \frac{2.612(mk)^{\frac{3}{2}}}{(2\pi)^{\frac{3}{2}} \hbar^3 n} = 2.612.$$
 - What's happening if $T < T_c$?
 - More and more particles stay at ground state.
- $\mu < 0$ still obeys, $\mu \rightarrow 0$; $\frac{2\pi}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT}} - 1} < n.$

BEC – idea gas

- $n_0(T)$ number of particles at ground state.

BEC – idea gas

- $n_0(T)$ number of particles at ground state.
- $$n_0(T) + \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT}} - 1} = n,$$

BEC – idea gas

- $n_0(T)$ number of particles at ground state.

- $$n_0(T) + \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT}} - 1} = n,$$

where
$$n_{\varepsilon>0} = \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT}} - 1}$$

BEC – idea gas

- $n_0(T)$ number of particles at ground state.

- $$n_0(T) + \frac{2\pi}{h^3}(2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT}} - 1} = n,$$

where $n_{\varepsilon>0} = \frac{2\pi}{h^3}(2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT}} - 1} = n\left(\frac{T}{T_c}\right)^{\frac{3}{2}}$ (Compared with eq.(8.3.7)).

BEC – idea gas

- $n_0(T)$ number of particles at ground state.
- $n_0(T) + \frac{2\pi}{h^3}(2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT}} - 1} = n$,
 where $n_{\varepsilon>0} = \frac{2\pi}{h^3}(2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT}} - 1} = n\left(\frac{T}{T_c}\right)^{\frac{3}{2}}$ (Compared with eq.(8.3.7)).
- $n_0(T) = n\left[1 - \left(\frac{T}{T_c}\right)^{\frac{3}{2}}\right]$.

BEC – idea gas

- $n_0(T)$ number of particles at ground state.

- $$n_0(T) + \frac{2\pi}{h^3}(2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT}} - 1} = n,$$

where $n_{\varepsilon>0} = \frac{2\pi}{h^3}(2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT}} - 1} = n\left(\frac{T}{T_c}\right)^{\frac{3}{2}}$ (Compared with eq.(8.3.7)).

- $$n_0(T) = n\left[1 - \left(\frac{T}{T_c}\right)^{\frac{3}{2}}\right].$$

The condensed part: $E = 0, U = 0, S = 0, p = 0$.

BEC – idea gas

- $n_0(T)$ number of particles at ground state.

- $$n_0(T) + \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT}} - 1} = n,$$

where $n_{\varepsilon>0} = \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT}} - 1} = n \left(\frac{T}{T_c}\right)^{\frac{3}{2}}$ (Compared with eq.(8.3.7)).

- $$n_0(T) = n \left[1 - \left(\frac{T}{T_c}\right)^{\frac{3}{2}}\right].$$

The condensed part: $E = 0, U = 0, S = 0, p = 0$.

- $$U = U_{\varepsilon>0}$$

BEC – idea gas

- $n_0(T)$ number of particles at ground state.

- $$n_0(T) + \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT}} - 1} = n,$$

where $n_{\varepsilon>0} = \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT}} - 1} = n \left(\frac{T}{T_c}\right)^{\frac{3}{2}}$ (Compared with eq.(8.3.7)).

- $$n_0(T) = n \left[1 - \left(\frac{T}{T_c}\right)^{\frac{3}{2}}\right].$$

The condensed part: $E = 0, U = 0, S = 0, p = 0$.

- $$U = U_{\varepsilon>0} = \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon \cdot \varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT}} - 1}$$

BEC – idea gas

- $n_0(T)$ number of particles at ground state.

- $$n_0(T) + \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT}} - 1} = n,$$

where $n_{\varepsilon>0} = \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT}} - 1} = n \left(\frac{T}{T_c}\right)^{\frac{3}{2}}$ (Compared with eq.(8.3.7)).

- $$n_0(T) = n \left[1 - \left(\frac{T}{T_c}\right)^{\frac{3}{2}}\right].$$

The condensed part: $E = 0, U = 0, S = 0, p = 0$.

- $$\begin{aligned} U &= U_{\varepsilon>0} = \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon \cdot \varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT}} - 1} \\ &= \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} (kT)^{\frac{5}{2}} \int_0^\infty \frac{x^{\frac{3}{2}} dx}{e^x - 1} \end{aligned}$$

BEC – idea gas

- $n_0(T)$ number of particles at ground state.

- $$n_0(T) + \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT}} - 1} = n,$$

where $n_{\varepsilon>0} = \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT}} - 1} = n \left(\frac{T}{T_c}\right)^{\frac{3}{2}}$ (Compared with eq.(8.3.7)).

- $$n_0(T) = n \left[1 - \left(\frac{T}{T_c}\right)^{\frac{3}{2}}\right].$$

The condensed part: $E = 0, U = 0, S = 0, p = 0$.

- $$\begin{aligned} U &= U_{\varepsilon>0} = \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon \cdot \varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT}} - 1} \\ &= \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} (kT)^{\frac{5}{2}} \int_0^\infty \frac{x^{\frac{3}{2}} dx}{e^x - 1} \\ &= \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} (kT) \cdot \frac{3}{4} \sqrt{\pi} \cdot 1.341 \end{aligned}$$

BEC – idea gas

- $n_0(T)$ number of particles at ground state.

- $$n_0(T) + \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT}} - 1} = n,$$

where $n_{\varepsilon>0} = \frac{2\pi}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT}} - 1} = n \left(\frac{T}{T_c}\right)^{\frac{3}{2}}$ (Compared with eq.(8.3.7)).

- $$n_0(T) = n \left[1 - \left(\frac{T}{T_c}\right)^{\frac{3}{2}}\right].$$

The condensed part: $E = 0, U = 0, S = 0, p = 0$.

- $$\begin{aligned} U &= U_{\varepsilon>0} = \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon \cdot \varepsilon^{\frac{1}{2}} d\varepsilon}{e^{\frac{\varepsilon}{kT}} - 1} \\ &= \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} (kT)^{\frac{5}{2}} \int_0^\infty \frac{x^{\frac{3}{2}} dx}{e^x - 1} \\ &= \frac{2\pi V}{h^3} (2mkT)^{\frac{3}{2}} (kT) \cdot \frac{3}{4} \sqrt{\pi} \cdot 1.341 = 0.77 N kT \left(\frac{T}{T_c}\right)^{\frac{3}{2}}. \end{aligned}$$

BEC – idea gas

- $U = 0.77NkT\left(\frac{T}{T_c}\right)^{\frac{3}{2}}.$

BEC – idea gas

- $U = 0.77NkT\left(\frac{T}{T_c}\right)^{\frac{3}{2}}.$
- $C_V = \left(\frac{\partial U}{\partial T}\right)_V$

BEC – idea gas

- $U = 0.77 N k T \left(\frac{T}{T_c} \right)^{\frac{3}{2}}.$
- $C_V = \left(\frac{\partial U}{\partial T} \right)_V = \frac{5}{2} \frac{U}{T}$

BEC – idea gas

- $U = 0.77NkT\left(\frac{T}{T_c}\right)^{\frac{3}{2}}.$
- $C_V = \left(\frac{\partial U}{\partial T}\right)_V = \frac{5}{2}\frac{U}{T} = 1.925Nk\left(\frac{T}{T_c}\right)^{\frac{3}{2}}.$

BEC – idea gas

- $U = 0.77NkT\left(\frac{T}{T_c}\right)^{\frac{3}{2}}.$
- $C_V = \left(\frac{\partial U}{\partial T}\right)_V = \frac{5}{2}\frac{U}{T} = 1.925Nk\left(\frac{T}{T_c}\right)^{\frac{3}{2}}.$
- In high temperature, U , C_V go back to the ideal gas $\frac{3}{2}NkT$ and $\frac{3}{2}Nk$.

BEC – idea gas

- $U = 0.77NkT\left(\frac{T}{T_c}\right)^{\frac{3}{2}}.$
- $C_V = \left(\frac{\partial U}{\partial T}\right)_V = \frac{5}{2}\frac{U}{T} = 1.925Nk\left(\frac{T}{T_c}\right)^{\frac{3}{2}}.$
- In high temperature, U , C_V go back to the ideal gas $\frac{3}{2}NkT$ and $\frac{3}{2}Nk$.

There is a break for C_V at $T = T_c$. (Jump on derivative of C_V , continuous phase transition.)

BEC – idea gas

- $U = 0.77NkT\left(\frac{T}{T_c}\right)^{\frac{3}{2}}.$
- $C_V = \left(\frac{\partial U}{\partial T}\right)_V = \frac{5}{2}\frac{U}{T} = 1.925Nk\left(\frac{T}{T_c}\right)^{\frac{3}{2}}.$
- In high temperature, U , C_V go back to the ideal gas $\frac{3}{2}NkT$ and $\frac{3}{2}Nk$.
There is a break for C_V at $T = T_c$. (Jump on derivative of C_V , continuous phase transition.)
- ^4He at 3.13K, transfer to superfluid.

BEC – idea gas

- $U = 0.77NkT\left(\frac{T}{T_c}\right)^{\frac{3}{2}}.$
- $C_V = \left(\frac{\partial U}{\partial T}\right)_V = \frac{5}{2}\frac{U}{T} = 1.925Nk\left(\frac{T}{T_c}\right)^{\frac{3}{2}}.$
- In high temperature, U , C_V go back to the ideal gas $\frac{3}{2}NkT$ and $\frac{3}{2}Nk$.

There is a break for C_V at $T = T_c$. (Jump on derivative of C_V , continuous phase transition.)

- ^4He at 3.13K, transfer to superfluid.
- Methods to get BEC: $n\lambda^3 > 2.612$, mainly reduce temperature.

BEC – idea gas

- $U = 0.77NkT\left(\frac{T}{T_c}\right)^{\frac{3}{2}}.$
- $C_V = \left(\frac{\partial U}{\partial T}\right)_V = \frac{5}{2}\frac{U}{T} = 1.925Nk\left(\frac{T}{T_c}\right)^{\frac{3}{2}}.$
- In high temperature, U , C_V go back to the ideal gas $\frac{3}{2}NkT$ and $\frac{3}{2}Nk$.

There is a break for C_V at $T = T_c$. (Jump on derivative of C_V , continuous phase transition.)

- ^4He at 3.13K, transfer to superfluid.
- Methods to get BEC: $n\lambda^3 > 2.612$, mainly reduce temperature.
- Magneto optical trap – 3-D harmonic oscillator.

BEC – realization

- $\varepsilon_{n_x, n_y, n_z} = \hbar\omega_x\left(n_x + \frac{1}{2}\right) + \hbar\omega_y\left(n_y + \frac{1}{2}\right) + \hbar\omega_z\left(n_z + \frac{1}{2}\right).$

BEC – realization

- $\varepsilon_{n_x, n_y, n_z} = \hbar\omega_x(n_x + \frac{1}{2}) + \hbar\omega_y(n_y + \frac{1}{2}) + \hbar\omega_z(n_z + \frac{1}{2})$.
- Large number of particles at the ground state:
$$\varepsilon_0 = \frac{\hbar}{2}(\omega_x + \omega_y + \omega_z).$$

BEC – realization

- $\varepsilon_{n_x, n_y, n_z} = \hbar\omega_x(n_x + \frac{1}{2}) + \hbar\omega_y(n_y + \frac{1}{2}) + \hbar\omega_z(n_z + \frac{1}{2})$.
- Large number of particles at the ground state:
 $\varepsilon_0 = \frac{\hbar}{2}(\omega_x + \omega_y + \omega_z)$.
- Particle number at energy level (n_x, n_y, n_z) :

BEC – realization

- $\varepsilon_{n_x, n_y, n_z} = \hbar\omega_x(n_x + \frac{1}{2}) + \hbar\omega_y(n_y + \frac{1}{2}) + \hbar\omega_z(n_z + \frac{1}{2})$.

- Large number of particles at the ground state:

$$\varepsilon_0 = \frac{\hbar}{2}(\omega_x + \omega_y + \omega_z).$$

- Particle number at energy level (n_x, n_y, n_z) :

$$a_l = a_{n_x, n_y, n_z} = \frac{1}{e^{\frac{1}{kT} [\hbar\omega_x(n_x + \frac{1}{2}) + \hbar\omega_y(n_y + \frac{1}{2}) + \hbar\omega_z(n_z + \frac{1}{2}) - \mu]} - 1}.$$

BEC – realization

- $\varepsilon_{n_x, n_y, n_z} = \hbar\omega_x(n_x + \frac{1}{2}) + \hbar\omega_y(n_y + \frac{1}{2}) + \hbar\omega_z(n_z + \frac{1}{2})$.
- Large number of particles at the ground state:
 $\varepsilon_0 = \frac{\hbar}{2}(\omega_x + \omega_y + \omega_z)$.
- Particle number at energy level (n_x, n_y, n_z) :

$$a_l = a_{n_x, n_y, n_z} = \frac{1}{e^{\frac{1}{kT}[\hbar\omega_x(n_x + \frac{1}{2}) + \hbar\omega_y(n_y + \frac{1}{2}) + \hbar\omega_z(n_z + \frac{1}{2}) - \mu]} - 1}.$$
- $a_l \geq 0$ requires $\mu < \varepsilon_0 = \frac{\hbar}{2}(\omega_x + \omega_y + \omega_z)$.

BEC – realization

- $\varepsilon_{n_x, n_y, n_z} = \hbar\omega_x(n_x + \frac{1}{2}) + \hbar\omega_y(n_y + \frac{1}{2}) + \hbar\omega_z(n_z + \frac{1}{2})$.
- Large number of particles at the ground state:
 $\varepsilon_0 = \frac{\hbar}{2}(\omega_x + \omega_y + \omega_z)$.
- Particle number at energy level (n_x, n_y, n_z) :

$$a_l = a_{n_x, n_y, n_z} = \frac{1}{e^{\frac{1}{kT}[\hbar\omega_x(n_x + \frac{1}{2}) + \hbar\omega_y(n_y + \frac{1}{2}) + \hbar\omega_z(n_z + \frac{1}{2}) - \mu]} - 1}.$$
- $a_l \geq 0$ requires $\mu < \varepsilon_0 = \frac{\hbar}{2}(\omega_x + \omega_y + \omega_z)$.
- T_c satisfies: at $T = T_c$, $\mu \rightarrow \varepsilon_0$,

BEC – realization

- $\varepsilon_{n_x, n_y, n_z} = \hbar\omega_x(n_x + \frac{1}{2}) + \hbar\omega_y(n_y + \frac{1}{2}) + \hbar\omega_z(n_z + \frac{1}{2})$.

- Large number of particles at the ground state:

$$\varepsilon_0 = \frac{\hbar}{2}(\omega_x + \omega_y + \omega_z).$$

- Particle number at energy level (n_x, n_y, n_z) :

$$a_l = a_{n_x, n_y, n_z} = \frac{1}{e^{\frac{1}{kT}[\hbar\omega_x(n_x + \frac{1}{2}) + \hbar\omega_y(n_y + \frac{1}{2}) + \hbar\omega_z(n_z + \frac{1}{2}) - \mu]} - 1}.$$

- $a_l \geq 0$ requires $\mu < \varepsilon_0 = \frac{\hbar}{2}(\omega_x + \omega_y + \omega_z)$.

- T_c satisfies: at $T = T_c$, $\mu \rightarrow \varepsilon_0$, i.e.

$$N = \sum a_l = \sum_{n_x, n_y, n_z} \frac{1}{e^{\frac{\hbar}{kT}(n_x\omega_x + n_y\omega_y + n_z\omega_z)} - 1}.$$

BEC – realization

- $\varepsilon_{n_x, n_y, n_z} = \hbar\omega_x(n_x + \frac{1}{2}) + \hbar\omega_y(n_y + \frac{1}{2}) + \hbar\omega_z(n_z + \frac{1}{2})$.

- Large number of particles at the ground state:

$$\varepsilon_0 = \frac{\hbar}{2}(\omega_x + \omega_y + \omega_z).$$

- Particle number at energy level (n_x, n_y, n_z) :

$$a_l = a_{n_x, n_y, n_z} = \frac{1}{e^{\frac{1}{kT}[\hbar\omega_x(n_x + \frac{1}{2}) + \hbar\omega_y(n_y + \frac{1}{2}) + \hbar\omega_z(n_z + \frac{1}{2}) - \mu]} - 1}.$$

- $a_l \geq 0$ requires $\mu < \varepsilon_0 = \frac{\hbar}{2}(\omega_x + \omega_y + \omega_z)$.

- T_c satisfies: at $T = T_c$, $\mu \rightarrow \varepsilon_0$, i.e.

$$N = \sum a_l = \sum_{n_x, n_y, n_z} \frac{1}{e^{\frac{\hbar}{kT}(n_x\omega_x + n_y\omega_y + n_z\omega_z)} - 1}.$$

In case of $\frac{\hbar\omega}{kT} \ll 1$, summation to integration:

BEC – realization

- $\varepsilon_{n_x, n_y, n_z} = \hbar\omega_x(n_x + \frac{1}{2}) + \hbar\omega_y(n_y + \frac{1}{2}) + \hbar\omega_z(n_z + \frac{1}{2})$.
- Large number of particles at the ground state:

$$\varepsilon_0 = \frac{\hbar}{2}(\omega_x + \omega_y + \omega_z).$$

- Particle number at energy level (n_x, n_y, n_z) :

$$a_l = a_{n_x, n_y, n_z} = \frac{1}{e^{\frac{1}{kT}[\hbar\omega_x(n_x + \frac{1}{2}) + \hbar\omega_y(n_y + \frac{1}{2}) + \hbar\omega_z(n_z + \frac{1}{2}) - \mu]} - 1}.$$

- $a_l \geq 0$ requires $\mu < \varepsilon_0 = \frac{\hbar}{2}(\omega_x + \omega_y + \omega_z)$.
- T_c satisfies: at $T = T_c$, $\mu \rightarrow \varepsilon_0$, i.e.

$$N = \sum a_l = \sum_{n_x, n_y, n_z} \frac{1}{e^{\frac{\hbar}{kT}(n_x\omega_x + n_y\omega_y + n_z\omega_z)} - 1}.$$

In case of $\frac{\hbar\omega}{kT} \ll 1$, summation to integration:

$$N = \iiint \frac{1}{e^{\frac{\hbar}{kT_c}(n_x\omega_x + n_y\omega_y + n_z\omega_z)} - 1} dn_x dn_y dn_z.$$

BEC – realization

- $$N = \iiint \frac{1}{e^{\frac{\hbar}{kT_c}(n_x\omega_x + n_y\omega_y + n_z\omega_z)} - 1} dn_x dn_y dn_z$$

BEC – realization

$$\begin{aligned}
 \bullet \quad N &= \iiint \frac{1}{e^{\frac{\hbar}{kT_c}(n_x\omega_x+n_y\omega_y+n_z\omega_z)} - 1} dn_x dn_y dn_z \\
 &= \iiint \frac{\left[\left(\frac{\hbar}{kT_c}\right)^3 \omega_x \omega_y \omega_z\right]^{-1}}{e^{\frac{\hbar}{kT_c}(n_x\omega_x+n_y\omega_y+n_z\omega_z)} - 1} d\frac{\hbar\omega_x}{kT_c} n_x d\frac{\hbar\omega_y}{kT_c} n_y d\frac{\hbar\omega_z}{kT_c} n_z
 \end{aligned}$$

BEC – realization

$$\begin{aligned}
 \bullet \quad N &= \iiint \frac{1}{e^{\frac{\hbar}{kT_c}(n_x\omega_x+n_y\omega_y+n_z\omega_z)} - 1} dn_x dn_y dn_z \\
 &= \iiint \frac{[(\frac{\hbar}{kT_c})^3 \omega_x \omega_y \omega_z]^{-1}}{e^{\frac{\hbar}{kT_c}(n_x\omega_x+n_y\omega_y+n_z\omega_z)} - 1} d\frac{\hbar\omega_x}{kT_c} n_x d\frac{\hbar\omega_y}{kT_c} n_y d\frac{\hbar\omega_z}{kT_c} n_z \\
 &= \left(\frac{kT_c}{\hbar}\right)^3 (\omega_x \omega_y \omega_z)^{-1} \iiint \frac{1}{e^{x+y+z} - 1} dx dy dz
 \end{aligned}$$

BEC – realization

$$\begin{aligned}
 \bullet \quad N &= \iiint \frac{1}{e^{\frac{\hbar}{kT_c}(n_x\omega_x+n_y\omega_y+n_z\omega_z)} - 1} dn_x dn_y dn_z \\
 &= \iiint \frac{[(\frac{\hbar}{kT_c})^3 \omega_x \omega_y \omega_z]^{-1}}{e^{\frac{\hbar}{kT_c}(n_x\omega_x+n_y\omega_y+n_z\omega_z)} - 1} d\frac{\hbar\omega_x}{kT_c} n_x d\frac{\hbar\omega_y}{kT_c} n_y d\frac{\hbar\omega_z}{kT_c} n_z \\
 &= (\frac{kT_c}{\hbar})^3 (\omega_x \omega_y \omega_z)^{-1} \iiint \frac{1}{e^{x+y+z} - 1} dx dy dz \\
 &\quad (\text{define } \bar{\omega} = (\omega_x \omega_y \omega_z)^{\frac{1}{3}})
 \end{aligned}$$

BEC – realization

$$\begin{aligned}
\bullet \quad N &= \iiint \frac{1}{e^{\frac{\hbar}{kT_c}(n_x\omega_x+n_y\omega_y+n_z\omega_z)} - 1} dn_x dn_y dn_z \\
&= \iiint \frac{\left[\left(\frac{\hbar}{kT_c}\right)^3 \omega_x \omega_y \omega_z\right]^{-1}}{e^{\frac{\hbar}{kT_c}(n_x\omega_x+n_y\omega_y+n_z\omega_z)} - 1} d\frac{\hbar\omega_x}{kT_c} n_x d\frac{\hbar\omega_y}{kT_c} n_y d\frac{\hbar\omega_z}{kT_c} n_z \\
&= \left(\frac{kT_c}{\hbar}\right)^3 (\omega_x \omega_y \omega_z)^{-1} \iiint \frac{1}{e^{x+y+z} - 1} dx dy dz \\
&\quad \text{(define } \bar{\omega} = (\omega_x \omega_y \omega_z)^{\frac{1}{3}} \text{)} \\
&= \left(\frac{kT_c}{\hbar\bar{\omega}}\right)^3 \iiint \frac{1}{e^{x+y+z} - 1} dx dy dz
\end{aligned}$$

BEC – realization

$$\begin{aligned}
\bullet N &= \iiint \frac{1}{e^{\frac{\hbar}{kT_c}(n_x\omega_x+n_y\omega_y+n_z\omega_z)} - 1} dn_x dn_y dn_z \\
&= \iiint \frac{[(\frac{\hbar}{kT_c})^3 \omega_x \omega_y \omega_z]^{-1}}{e^{\frac{\hbar}{kT_c}(n_x\omega_x+n_y\omega_y+n_z\omega_z)} - 1} d\frac{\hbar\omega_x}{kT_c} n_x d\frac{\hbar\omega_y}{kT_c} n_y d\frac{\hbar\omega_z}{kT_c} n_z \\
&= (\frac{kT_c}{\hbar})^3 (\omega_x \omega_y \omega_z)^{-1} \iiint \frac{1}{e^{x+y+z} - 1} dx dy dz \\
&\quad (\text{define } \bar{\omega} = (\omega_x \omega_y \omega_z)^{\frac{1}{3}}) \\
&= (\frac{kT_c}{\hbar\bar{\omega}})^3 \iiint \frac{1}{e^{x+y+z} - 1} dx dy dz \\
&= (\frac{kT_c}{\hbar\bar{\omega}})^3 \iiint \frac{1}{e^{x+y+z} [1 - e^{-(x+y+z)}]} dx dy dz
\end{aligned}$$

BEC – realization

$$\begin{aligned}
\bullet N &= \iiint \frac{1}{e^{\frac{\hbar}{kT_c}(n_x\omega_x+n_y\omega_y+n_z\omega_z)} - 1} dn_x dn_y dn_z \\
&= \iiint \frac{[(\frac{\hbar}{kT_c})^3 \omega_x \omega_y \omega_z]^{-1}}{e^{\frac{\hbar}{kT}(n_x\omega_x+n_y\omega_y+n_z\omega_z)} - 1} d\frac{\hbar\omega_x}{kT_c} n_x d\frac{\hbar\omega_y}{kT_c} n_y d\frac{\hbar\omega_z}{kT_c} n_z \\
&= (\frac{kT_c}{\hbar})^3 (\omega_x \omega_y \omega_z)^{-1} \iiint \frac{1}{e^{x+y+z} - 1} dx dy dz \\
&\quad (\text{define } \bar{\omega} = (\omega_x \omega_y \omega_z)^{\frac{1}{3}}) \\
&= (\frac{kT_c}{\hbar\bar{\omega}})^3 \iiint \frac{1}{e^{x+y+z} - 1} dx dy dz \\
&= (\frac{kT_c}{\hbar\bar{\omega}})^3 \iiint \frac{1}{e^{x+y+z} [1 - e^{-(x+y+z)}]} dx dy dz \\
&= (\frac{kT_c}{\hbar\bar{\omega}})^3 \iiint e^{-(x+y+z)} \sum_{l=0}^{\infty} e^{-l(x+y+z)} dx dy dz
\end{aligned}$$

BEC – realization

$$\begin{aligned}
\bullet N &= \iiint \frac{1}{e^{\frac{\hbar}{kT_c}(n_x\omega_x+n_y\omega_y+n_z\omega_z)} - 1} dn_x dn_y dn_z \\
&= \iiint \frac{[(\frac{\hbar}{kT_c})^3 \omega_x \omega_y \omega_z]^{-1}}{e^{\frac{\hbar}{kT_c}(n_x\omega_x+n_y\omega_y+n_z\omega_z)} - 1} d\frac{\hbar\omega_x}{kT_c} n_x d\frac{\hbar\omega_y}{kT_c} n_y d\frac{\hbar\omega_z}{kT_c} n_z \\
&= (\frac{kT_c}{\hbar})^3 (\omega_x \omega_y \omega_z)^{-1} \iiint \frac{1}{e^{x+y+z} - 1} dx dy dz \\
&\quad (\text{define } \bar{\omega} = (\omega_x \omega_y \omega_z)^{\frac{1}{3}}) \\
&= (\frac{kT_c}{\hbar\bar{\omega}})^3 \iiint \frac{1}{e^{x+y+z} - 1} dx dy dz \\
&= (\frac{kT_c}{\hbar\bar{\omega}})^3 \iiint \frac{1}{e^{x+y+z} [1 - e^{-(x+y+z)}]} dx dy dz \\
&= (\frac{kT_c}{\hbar\bar{\omega}})^3 \iiint e^{-(x+y+z)} \sum_{l=0}^{\infty} e^{-l(x+y+z)} dx dy dz \\
&= (\frac{kT_c}{\hbar\bar{\omega}})^3 \iiint \sum_{l=1}^{\infty} e^{-l(x+y+z)} dx dy dz \dots
\end{aligned}$$

BEC – realization

- $$N = \left(\frac{kT_c}{\hbar\bar{\omega}}\right)^3 \iiint \sum_{l=1}^{\infty} e^{-l(x+y+z)} dx dy dz$$

BEC – realization

- $$N = \left(\frac{kT_c}{\hbar\bar{\omega}}\right)^3 \iiint \sum_{l=1}^{\infty} e^{-l(x+y+z)} dx dy dz$$
$$= \left(\frac{kT_c}{\hbar\bar{\omega}}\right)^3 \sum_{l=1}^{\infty} \int_0^{\infty} e^{-lx} dx \int_0^{\infty} e^{-ly} dy \int_0^{\infty} e^{-lz} dz$$

BEC – realization

- $$\begin{aligned} N &= \left(\frac{kT_c}{\hbar\bar{\omega}}\right)^3 \iiint \sum_{l=1}^{\infty} e^{-l(x+y+z)} dx dy dz \\ &= \left(\frac{kT_c}{\hbar\bar{\omega}}\right)^3 \sum_{l=1}^{\infty} \int_0^{\infty} e^{-lx} dx \int_0^{\infty} e^{-ly} dy \int_0^{\infty} e^{-lz} dz \\ &= \left(\frac{kT_c}{\hbar\bar{\omega}}\right)^3 \sum_{l=1}^{\infty} \frac{1}{l^3} \end{aligned}$$

BEC – realization

- $$\begin{aligned} N &= \left(\frac{kT_c}{\hbar\bar{\omega}}\right)^3 \iiint \sum_{l=1}^{\infty} e^{-l(x+y+z)} dx dy dz \\ &= \left(\frac{kT_c}{\hbar\bar{\omega}}\right)^3 \sum_{l=1}^{\infty} \int_0^{\infty} e^{-lx} dx \int_0^{\infty} e^{-ly} dy \int_0^{\infty} e^{-lz} dz \\ &= \left(\frac{kT_c}{\hbar\bar{\omega}}\right)^3 \sum_{l=1}^{\infty} \frac{1}{l^3} \\ &= 1.202 \left(\frac{kT_c}{\hbar\bar{\omega}}\right)^3, \end{aligned}$$

BEC – realization

- $$\begin{aligned} N &= \left(\frac{kT_c}{\hbar\bar{\omega}}\right)^3 \iiint \sum_{l=1}^{\infty} e^{-l(x+y+z)} dx dy dz \\ &= \left(\frac{kT_c}{\hbar\bar{\omega}}\right)^3 \sum_{l=1}^{\infty} \int_0^{\infty} e^{-lx} dx \int_0^{\infty} e^{-ly} dy \int_0^{\infty} e^{-lz} dz \\ &= \left(\frac{kT_c}{\hbar\bar{\omega}}\right)^3 \sum_{l=1}^{\infty} \frac{1}{l^3} \\ &= 1.202 \left(\frac{kT_c}{\hbar\bar{\omega}}\right)^3, \\ &\Rightarrow kT_c = \hbar\bar{\omega} \left(\frac{N}{1.202}\right)^{\frac{1}{3}}. \end{aligned}$$

BEC – realization

- $$N = \left(\frac{kT_c}{\hbar\bar{\omega}}\right)^3 \iiint \sum_{l=1}^{\infty} e^{-l(x+y+z)} dx dy dz$$

$$= \left(\frac{kT_c}{\hbar\bar{\omega}}\right)^3 \sum_{l=1}^{\infty} \int_0^{\infty} e^{-lx} dx \int_0^{\infty} e^{-ly} dy \int_0^{\infty} e^{-lz} dz$$

$$= \left(\frac{kT_c}{\hbar\bar{\omega}}\right)^3 \sum_{l=1}^{\infty} \frac{1}{l^3}$$

$$= 1.202 \left(\frac{kT_c}{\hbar\bar{\omega}}\right)^3,$$
- $$\Rightarrow kT_c = \hbar\bar{\omega} \left(\frac{N}{1.202}\right)^{\frac{1}{3}}.$$
- For $T < T_c$, N_0 particles stay at the ground state (BEC).

BEC – realization

- $$N = \left(\frac{kT_c}{\hbar\bar{\omega}}\right)^3 \iiint \sum_{l=1}^{\infty} e^{-l(x+y+z)} dx dy dz$$

$$= \left(\frac{kT_c}{\hbar\bar{\omega}}\right)^3 \sum_{l=1}^{\infty} \int_0^{\infty} e^{-lx} dx \int_0^{\infty} e^{-ly} dy \int_0^{\infty} e^{-lz} dz$$

$$= \left(\frac{kT_c}{\hbar\bar{\omega}}\right)^3 \sum_{l=1}^{\infty} \frac{1}{l^3}$$

$$= 1.202 \left(\frac{kT_c}{\hbar\bar{\omega}}\right)^3,$$
- $$\Rightarrow kT_c = \hbar\bar{\omega} \left(\frac{N}{1.202}\right)^{\frac{1}{3}}.$$
- For $T < T_c$, N_0 particles stay at the ground state (BEC).

$$N - N_0 = 1.202 \left(\frac{kT}{\hbar\bar{\omega}}\right)^3$$

BEC – realization

- $$N = \left(\frac{kT_c}{\hbar\bar{\omega}}\right)^3 \iiint \sum_{l=1}^{\infty} e^{-l(x+y+z)} dx dy dz$$

$$= \left(\frac{kT_c}{\hbar\bar{\omega}}\right)^3 \sum_{l=1}^{\infty} \int_0^{\infty} e^{-lx} dx \int_0^{\infty} e^{-ly} dy \int_0^{\infty} e^{-lz} dz$$

$$= \left(\frac{kT_c}{\hbar\bar{\omega}}\right)^3 \sum_{l=1}^{\infty} \frac{1}{l^3}$$

$$= 1.202 \left(\frac{kT_c}{\hbar\bar{\omega}}\right)^3,$$
- $$\Rightarrow kT_c = \hbar\bar{\omega} \left(\frac{N}{1.202}\right)^{\frac{1}{3}}.$$
- For $T < T_c$, N_0 particles stay at the ground state (BEC).

$$N - N_0 = 1.202 \left(\frac{kT}{\hbar\bar{\omega}}\right)^3 = N \left(\frac{T}{T_c}\right)^3.$$

Table of contents

1 §8. Bose statistics and Fermi statistics

- 8.1 Thermal parameters in statistics
- 8.2 U of weak degenerated ideal Fermion(Boson) gas
- 8.3 Bose-Einstein Condensate (BEC)
- 8.4 Photon gas
- 8.5 Free electron gas in metal

8.4 Photon gas

8.4 Photon gas

- Thermal radiation in the cavity: waves or photons.

8.4 Photon gas

- Thermal radiation in the cavity: waves or photons.
- Number of photons is not conserved.

8.4 Photon gas

- Thermal radiation in the cavity: waves or photons.
- Number of photons is not conserved.
- Photons are Bosons.

8.4 Photon gas

- Thermal radiation in the cavity: waves or photons.
- Number of photons is not conserved.
- Photons are Bosons.
- Bose statistics without conserved N , lack of α :

8.4 Photon gas

- Thermal radiation in the cavity: waves or photons.
- Number of photons is not conserved.
- Photons are Bosons.
- Bose statistics without conserved N , lack of α :

$$a_l = \frac{\omega_l}{e^{\beta\varepsilon_l} - 1}.$$

8.4 Photon gas

- Thermal radiation in the cavity: waves or photons.
- Number of photons is not conserved.
- Photons are Bosons.
- Bose statistics without conserved N , lack of α :

$$a_l = \frac{\omega_l}{e^{\beta \varepsilon_l} - 1}.$$

- I.e., $\alpha = 0$ or $\mu = 0$:

8.4 Photon gas

- Thermal radiation in the cavity: waves or photons.
- Number of photons is not conserved.
- Photons are Bosons.
- Bose statistics without conserved N , lack of α :
$$a_l = \frac{\omega_l}{e^{\beta\varepsilon_l} - 1}.$$
- I.e., $\alpha = 0$ or $\mu = 0$:
 N is not conserved, open system.

8.4 Photon gas

- Thermal radiation in the cavity: waves or photons.
- Number of photons is not conserved.
- Photons are Bosons.
- Bose statistics without conserved N , lack of α :
$$a_l = \frac{\omega_l}{e^{\beta\epsilon_l} - 1}.$$
- I.e., $\alpha = 0$ or $\mu = 0$:
 N is not conserved, open system. For a given (T, V) ,
in thermal equilibrium,

8.4 Photon gas

- Thermal radiation in the cavity: waves or photons.
- Number of photons is not conserved.
- Photons are Bosons.
- Bose statistics without conserved N , lack of α :
$$a_l = \frac{\omega_l}{e^{\beta\epsilon_l} - 1}.$$
- I.e., $\alpha = 0$ or $\mu = 0$:
 N is not conserved, open system. For a given (T, V) , in thermal equilibrium, the condition for the equilibrium:

8.4 Photon gas

- Thermal radiation in the cavity: waves or photons.
- Number of photons is not conserved.
- Photons are Bosons.
- Bose statistics without conserved N , lack of α :
$$a_l = \frac{\omega_l}{e^{\beta\varepsilon_l} - 1}.$$
- I.e., $\alpha = 0$ or $\mu = 0$:
 N is not conserved, open system. For a given (T, V) , in thermal equilibrium, the condition for the equilibrium: $\Delta F = 0$ (§3.1).

8.4 Photon gas

- Thermal radiation in the cavity: waves or photons.
- Number of photons is not conserved.
- Photons are Bosons.
- Bose statistics without conserved N , lack of α :

$$a_l = \frac{\omega_l}{e^{\beta\epsilon_l} - 1}.$$

- I.e., $\alpha = 0$ or $\mu = 0$:

N is not conserved, open system. For a given (T, V) , in thermal equilibrium, the condition for the equilibrium: $\Delta F = 0$ (§3.1).

I.e., $(\frac{\partial F}{\partial N})_{T,V} = 0$.

8.4 Photon gas

- Thermal radiation in the cavity: waves or photons.
- Number of photons is not conserved.
- Photons are Bosons.
- Bose statistics without conserved N , lack of α :

$$a_l = \frac{\omega_l}{e^{\beta\epsilon_l} - 1}.$$

- I.e., $\alpha = 0$ or $\mu = 0$:

N is not conserved, open system. For a given (T, V) , in thermal equilibrium, the condition for the equilibrium: $\Delta F = 0$ (§3.1).

I.e., $(\frac{\partial F}{\partial N})_{T,V} = 0$.

Notice $dF = -SdT - pdV + \mu dN$,

8.4 Photon gas

- Thermal radiation in the cavity: waves or photons.
- Number of photons is not conserved.
- Photons are Bosons.
- Bose statistics without conserved N , lack of α :

$$a_l = \frac{\omega_l}{e^{\beta\epsilon_l} - 1}.$$

- I.e., $\alpha = 0$ or $\mu = 0$:

N is not conserved, open system. For a given (T, V) , in thermal equilibrium, the condition for the equilibrium: $\Delta F = 0$ (§3.1).

$$\text{I.e., } \left(\frac{\partial F}{\partial N}\right)_{T,V} = 0.$$

$$\text{Notice } dF = -SdT - pdV + \mu dN,$$

$$\therefore \mu = \left(\frac{\partial F}{\partial N}\right)_{T,V} = 0.$$

Photon gas

- Number of quantum states: $2 \cdot \frac{V 4\pi p^2 dp}{h^3} = \frac{8\pi V p^2 dp}{h^3}$,
where 2 is because of the polarization.

Photon gas

- Number of quantum states: $2 \cdot \frac{V 4\pi p^2 dp}{h^3} = \frac{8\pi V p^2 dp}{h^3}$,
where 2 is because of the polarization.
- Convert to frequency $\hbar\omega = \varepsilon = pc, \Rightarrow p = \frac{\hbar\omega}{c}$,

Photon gas

- Number of quantum states: $2 \cdot \frac{V 4\pi p^2 dp}{h^3} = \frac{8\pi V p^2 dp}{h^3}$, where 2 is because of the polarization.
- Convert to frequency $\hbar\omega = \varepsilon = pc, \Rightarrow p = \frac{\hbar\omega}{c}$,
$$D(\omega)d\omega = \frac{8\pi V (\frac{\hbar\omega}{c})^2 d(\frac{\hbar\omega}{c})}{h^3}$$

Photon gas

- Number of quantum states: $2 \cdot \frac{V 4\pi p^2 dp}{h^3} = \frac{8\pi V p^2 dp}{h^3}$, where 2 is because of the polarization.
- Convert to frequency $\hbar\omega = \varepsilon = pc, \Rightarrow p = \frac{\hbar\omega}{c}$,
$$D(\omega)d\omega = \frac{8\pi V (\frac{\hbar\omega}{c})^2 d(\frac{\hbar\omega}{c})}{h^3} = \frac{V}{\pi^2 c^3} \omega^2 d\omega.$$

Photon gas

- Number of quantum states: $2 \cdot \frac{V 4\pi p^2 dp}{h^3} = \frac{8\pi V p^2 dp}{h^3}$,
where 2 is because of the polarization.
- Convert to frequency $\hbar\omega = \varepsilon = pc, \Rightarrow p = \frac{\hbar\omega}{c}$,
$$D(\omega)d\omega = \frac{8\pi V (\frac{\hbar\omega}{c})^2 d(\frac{\hbar\omega}{c})}{h^3} = \frac{V}{\pi^2 c^3} \omega^2 d\omega.$$
- Number of photons at $(\omega, \omega + d\omega)$:

Photon gas

- Number of quantum states: $2 \cdot \frac{V4\pi p^2 dp}{h^3} = \frac{8\pi V p^2 dp}{h^3}$,
where 2 is because of the polarization.
- Convert to frequency $\hbar\omega = \varepsilon = pc, \Rightarrow p = \frac{\hbar\omega}{c}$,

$$D(\omega)d\omega = \frac{8\pi V(\frac{\hbar\omega}{c})^2 d(\frac{\hbar\omega}{c})}{h^3} = \frac{V}{\pi^2 c^3} \omega^2 d\omega.$$
- Number of photons at $(\omega, \omega + d\omega)$:

$$a(\omega)d\omega = \frac{V}{\pi^2 c^3} \frac{\omega^2 d\omega}{e^{\frac{\hbar\omega}{kT}} - 1}.$$

Photon gas

- Number of quantum states: $2 \cdot \frac{V4\pi p^2 dp}{h^3} = \frac{8\pi V p^2 dp}{h^3}$,
where 2 is because of the polarization.
- Convert to frequency $\hbar\omega = \varepsilon = pc, \Rightarrow p = \frac{\hbar\omega}{c}$,

$$D(\omega)d\omega = \frac{8\pi V(\frac{\hbar\omega}{c})^2 d(\frac{\hbar\omega}{c})}{h^3} = \frac{V}{\pi^2 c^3} \omega^2 d\omega.$$
- Number of photons at $(\omega, \omega + d\omega)$:

$$a(\omega)d\omega = \frac{V}{\pi^2 c^3} \frac{\omega^2 d\omega}{e^{\frac{\hbar\omega}{kT}} - 1}.$$
- Internal energy: $U(\omega, T)d\omega = \frac{V}{\pi^2 c^3} \frac{\hbar\omega^3 d\omega}{e^{\frac{\hbar\omega}{kT}} - 1}.$

Photon gas

- Number of quantum states: $2 \cdot \frac{V4\pi p^2 dp}{h^3} = \frac{8\pi V p^2 dp}{h^3}$,
where 2 is because of the polarization.
- Convert to frequency $\hbar\omega = \varepsilon = pc, \Rightarrow p = \frac{\hbar\omega}{c}$,

$$D(\omega)d\omega = \frac{8\pi V(\frac{\hbar\omega}{c})^2 d(\frac{\hbar\omega}{c})}{h^3} = \frac{V}{\pi^2 c^3} \omega^2 d\omega.$$
- Number of photons at $(\omega, \omega + d\omega)$:

$$a(\omega)d\omega = \frac{V}{\pi^2 c^3} \frac{\omega^2 d\omega}{e^{\frac{\hbar\omega}{kT}} - 1}.$$
- Internal energy: $U(\omega, T)d\omega = \frac{V}{\pi^2 c^3} \frac{\hbar\omega^3 d\omega}{e^{\frac{\hbar\omega}{kT}} - 1}.$
- Total internal energy: $U = \int_0^\infty U(\omega, T)d\omega$

Photon gas

- Number of quantum states: $2 \cdot \frac{V4\pi p^2 dp}{h^3} = \frac{8\pi V p^2 dp}{h^3}$,
where 2 is because of the polarization.
- Convert to frequency $\hbar\omega = \varepsilon = pc, \Rightarrow p = \frac{\hbar\omega}{c}$,

$$D(\omega)d\omega = \frac{8\pi V (\frac{\hbar\omega}{c})^2 d(\frac{\hbar\omega}{c})}{h^3} = \frac{V}{\pi^2 c^3} \omega^2 d\omega.$$
- Number of photons at $(\omega, \omega + d\omega)$:

$$a(\omega)d\omega = \frac{V}{\pi^2 c^3} \frac{\omega^2 d\omega}{e^{\frac{\hbar\omega}{kT}} - 1}.$$
- Internal energy: $U(\omega, T)d\omega = \frac{V}{\pi^2 c^3} \frac{\hbar\omega^3 d\omega}{e^{\frac{\hbar\omega}{kT}} - 1}.$
- Total internal energy: $U = \int_0^\infty U(\omega, T)d\omega$

$$= \int_0^\infty \frac{V}{\pi^2 c^3} \frac{\hbar\omega^3 d\omega}{e^{\frac{\hbar\omega}{kT}} - 1}$$

Photon gas

- Number of quantum states: $2 \cdot \frac{V4\pi p^2 dp}{h^3} = \frac{8\pi V p^2 dp}{h^3}$,
where 2 is because of the polarization.
- Convert to frequency $\hbar\omega = \varepsilon = pc, \Rightarrow p = \frac{\hbar\omega}{c}$,

$$D(\omega)d\omega = \frac{8\pi V (\frac{\hbar\omega}{c})^2 d(\frac{\hbar\omega}{c})}{h^3} = \frac{V}{\pi^2 c^3} \omega^2 d\omega.$$
- Number of photons at $(\omega, \omega + d\omega)$:

$$a(\omega)d\omega = \frac{V}{\pi^2 c^3} \frac{\omega^2 d\omega}{e^{\frac{\hbar\omega}{kT}} - 1}.$$
- Internal energy: $U(\omega, T)d\omega = \frac{V}{\pi^2 c^3} \frac{\hbar\omega^3 d\omega}{e^{\frac{\hbar\omega}{kT}} - 1}.$
- Total internal energy: $U = \int_0^\infty U(\omega, T)d\omega$

$$= \int_0^\infty \frac{V}{\pi^2 c^3} \frac{\hbar\omega^3 d\omega}{e^{\frac{\hbar\omega}{kT}} - 1} = \int_0^\infty \frac{V\hbar}{\pi^2 c^3} \left(\frac{\hbar}{kT}\right)^{-4} \frac{(\frac{\hbar\omega}{kT})^3 d(\frac{\hbar\omega}{kT})}{e^{\frac{\hbar\omega}{kT}} - 1}$$

Photon gas

- Number of quantum states: $2 \cdot \frac{V4\pi p^2 dp}{h^3} = \frac{8\pi V p^2 dp}{h^3}$,
where 2 is because of the polarization.
- Convert to frequency $\hbar\omega = \varepsilon = pc, \Rightarrow p = \frac{\hbar\omega}{c}$,

$$D(\omega)d\omega = \frac{8\pi V (\frac{\hbar\omega}{c})^2 d(\frac{\hbar\omega}{c})}{h^3} = \frac{V}{\pi^2 c^3} \omega^2 d\omega.$$
- Number of photons at $(\omega, \omega + d\omega)$:

$$a(\omega)d\omega = \frac{V}{\pi^2 c^3} \frac{\omega^2 d\omega}{e^{\frac{\hbar\omega}{kT}} - 1}.$$
- Internal energy: $U(\omega, T)d\omega = \frac{V}{\pi^2 c^3} \frac{\hbar\omega^3 d\omega}{e^{\frac{\hbar\omega}{kT}} - 1}.$
- Total internal energy: $U = \int_0^\infty U(\omega, T)d\omega$

$$= \int_0^\infty \frac{V}{\pi^2 c^3} \frac{\hbar\omega^3 d\omega}{e^{\frac{\hbar\omega}{kT}} - 1} = \int_0^\infty \frac{V\hbar}{\pi^2 c^3} \left(\frac{\hbar}{kT}\right)^{-4} \frac{(\frac{\hbar\omega}{kT})^3 d(\frac{\hbar\omega}{kT})}{e^{\frac{\hbar\omega}{kT}} - 1}$$

$$= \frac{V(kT)^4}{\pi^2 c^3 \hbar^3} \int_0^\infty \frac{x^3 dx}{e^x - 1}$$

Photon gas

- Number of quantum states: $2 \cdot \frac{V4\pi p^2 dp}{h^3} = \frac{8\pi V p^2 dp}{h^3}$,
where 2 is because of the polarization.
- Convert to frequency $\hbar\omega = \varepsilon = pc, \Rightarrow p = \frac{\hbar\omega}{c}$,

$$D(\omega)d\omega = \frac{8\pi V (\frac{\hbar\omega}{c})^2 d(\frac{\hbar\omega}{c})}{h^3} = \frac{V}{\pi^2 c^3} \omega^2 d\omega.$$
- Number of photons at $(\omega, \omega + d\omega)$:

$$a(\omega)d\omega = \frac{V}{\pi^2 c^3} \frac{\omega^2 d\omega}{e^{\frac{\hbar\omega}{kT}} - 1}.$$
- Internal energy: $U(\omega, T)d\omega = \frac{V}{\pi^2 c^3} \frac{\hbar\omega^3 d\omega}{e^{\frac{\hbar\omega}{kT}} - 1}.$
- Total internal energy: $U = \int_0^\infty U(\omega, T)d\omega$

$$= \int_0^\infty \frac{V}{\pi^2 c^3} \frac{\hbar\omega^3 d\omega}{e^{\frac{\hbar\omega}{kT}} - 1} = \int_0^\infty \frac{V\hbar}{\pi^2 c^3} \left(\frac{\hbar}{kT}\right)^{-4} \frac{(\frac{\hbar\omega}{kT})^3 d(\frac{\hbar\omega}{kT})}{e^{\frac{\hbar\omega}{kT}} - 1}$$

$$= \frac{V(kT)^4}{\pi^2 c^3 \hbar^3} \int_0^\infty \frac{x^3 dx}{e^x - 1} = \frac{\pi^2 k^4}{15 c^3 \hbar^3} T^4 V.$$

Photon gas

- Number of quantum states: $2 \cdot \frac{V4\pi p^2 dp}{h^3} = \frac{8\pi V p^2 dp}{h^3}$,
where 2 is because of the polarization.
- Convert to frequency $\hbar\omega = \varepsilon = pc, \Rightarrow p = \frac{\hbar\omega}{c}$,

$$D(\omega)d\omega = \frac{8\pi V(\frac{\hbar\omega}{c})^2 d(\frac{\hbar\omega}{c})}{h^3} = \frac{V}{\pi^2 c^3} \omega^2 d\omega.$$
- Number of photons at $(\omega, \omega + d\omega)$:

$$a(\omega)d\omega = \frac{V}{\pi^2 c^3} \frac{\omega^2 d\omega}{e^{\frac{\hbar\omega}{kT}} - 1}.$$
- Internal energy: $U(\omega, T)d\omega = \frac{V}{\pi^2 c^3} \frac{\hbar\omega^3 d\omega}{e^{\frac{\hbar\omega}{kT}} - 1}.$
- Total internal energy: $U = \int_0^\infty U(\omega, T)d\omega$

$$= \int_0^\infty \frac{V}{\pi^2 c^3} \frac{\hbar\omega^3 d\omega}{e^{\frac{\hbar\omega}{kT}} - 1} = \int_0^\infty \frac{V\hbar}{\pi^2 c^3} \left(\frac{\hbar}{kT}\right)^{-4} \frac{(\frac{\hbar\omega}{kT})^3 d(\frac{\hbar\omega}{kT})}{e^{\frac{\hbar\omega}{kT}} - 1}$$

$$= \frac{V(kT)^4}{\pi^2 c^3 \hbar^3} \int_0^\infty \frac{x^3 dx}{e^x - 1} = \frac{\pi^2 k^4}{15 c^3 \hbar^3} T^4 V. \quad (a = \frac{\pi^2 k^4}{15 c^3 \hbar^3}).$$

Photon gas $D(\omega)d\omega = \frac{V}{\pi^2 c^3} \omega^2 d\omega$

- Maximum for $U(\omega, T)d\omega = \frac{V}{\pi^2 c^3} \frac{\hbar \omega^3 d\omega}{e^{\frac{\hbar \omega}{kT}} - 1}$.

Photon gas $D(\omega)d\omega = \frac{V}{\pi^2 c^3} \omega^2 d\omega$

- Maximum for $U(\omega, T)d\omega = \frac{V}{\pi^2 c^3} \frac{\hbar \omega^3 d\omega}{e^{\frac{\hbar \omega}{kT}} - 1}$:

$$\frac{d}{dx} \left(\frac{x^3}{e^x - 1} \right) = 0$$

Photon gas $D(\omega)d\omega = \frac{V}{\pi^2 c^3} \omega^2 d\omega$

- Maximum for $U(\omega, T)d\omega = \frac{V}{\pi^2 c^3} \frac{\hbar \omega^3 d\omega}{e^{\frac{\hbar \omega}{kT}} - 1}$:

$$\frac{d}{dx} \left(\frac{x^3}{e^x - 1} \right) = 0 \Rightarrow x_m = \frac{\hbar \omega_m}{kT} \simeq 2.822, \text{ Wien's displacement law.}$$

Photon gas $D(\omega)d\omega = \frac{V}{\pi^2 c^3} \omega^2 d\omega$

- Maximum for $U(\omega, T)d\omega = \frac{V}{\pi^2 c^3} \frac{\hbar \omega^3 d\omega}{e^{\frac{\hbar \omega}{kT}} - 1}$:
 $\frac{d}{dx} \left(\frac{x^3}{e^x - 1} \right) = 0 \Rightarrow x_m = \frac{\hbar \omega_m}{kT} \simeq 2.822$, Wien's displacement law.
- Grand partition function for the photons:

Photon gas $D(\omega)d\omega = \frac{V}{\pi^2 c^3} \omega^2 d\omega$

- Maximum for $U(\omega, T)d\omega = \frac{V}{\pi^2 c^3} \frac{\hbar \omega^3 d\omega}{e^{\frac{\hbar \omega}{kT}} - 1}$:
 $\frac{d}{dx} \left(\frac{x^3}{e^x - 1} \right) = 0 \Rightarrow x_m = \frac{\hbar \omega_m}{kT} \simeq 2.822$, Wien's displacement law.
- Grand partition function for the photons:
 $\ln \mathcal{Z} = - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l})$

Photon gas $D(\omega)d\omega = \frac{V}{\pi^2 c^3} \omega^2 d\omega$

- Maximum for $U(\omega, T)d\omega = \frac{V}{\pi^2 c^3} \frac{\hbar \omega^3 d\omega}{e^{\frac{\hbar \omega}{kT}} - 1}$:

$$\frac{d}{dx} \left(\frac{x^3}{e^x - 1} \right) = 0 \Rightarrow x_m = \frac{\hbar \omega_m}{kT} \simeq 2.822, \text{ Wien's displacement law.}$$

- Grand partition function for the photons:

$$\begin{aligned} \ln \mathcal{Z} &= - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l}) \\ &= - \frac{V}{\pi^2 c^3} \int_0^\infty \omega^2 \ln(1 - e^{-\beta \hbar \omega}) d\omega \end{aligned}$$

Photon gas $D(\omega)d\omega = \frac{V}{\pi^2 c^3} \omega^2 d\omega$

- Maximum for $U(\omega, T)d\omega = \frac{V}{\pi^2 c^3} \frac{\hbar \omega^3 d\omega}{e^{\frac{\hbar \omega}{kT}} - 1}$:

$$\frac{d}{dx} \left(\frac{x^3}{e^x - 1} \right) = 0 \Rightarrow x_m = \frac{\hbar \omega_m}{kT} \simeq 2.822, \text{ Wien's displacement law.}$$

- Grand partition function for the photons:

$$\begin{aligned} \ln \mathcal{Z} &= - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l}) \\ &= - \frac{V}{\pi^2 c^3} \int_0^\infty \omega^2 \ln(1 - e^{-\beta \hbar \omega}) d\omega \\ &= - \frac{V}{\pi^2 c^3} (\beta \hbar)^{-3} \int_0^\infty (\beta \hbar \omega)^2 \ln(1 - e^{-\beta \hbar \omega}) d(\beta \hbar \omega) \end{aligned}$$

Photon gas $D(\omega)d\omega = \frac{V}{\pi^2 c^3} \omega^2 d\omega$

- Maximum for $U(\omega, T)d\omega = \frac{V}{\pi^2 c^3} \frac{\hbar \omega^3 d\omega}{e^{\frac{\hbar \omega}{kT}} - 1}$:

$$\frac{d}{dx} \left(\frac{x^3}{e^x - 1} \right) = 0 \Rightarrow x_m = \frac{\hbar \omega_m}{kT} \simeq 2.822, \text{ Wien's displacement law.}$$

- Grand partition function for the photons:

$$\begin{aligned} \ln \mathcal{Z} &= - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l}) \\ &= - \frac{V}{\pi^2 c^3} \int_0^\infty \omega^2 \ln(1 - e^{-\beta \hbar \omega}) d\omega \\ &= - \frac{V}{\pi^2 c^3} (\beta \hbar)^{-3} \int_0^\infty (\beta \hbar \omega)^2 \ln(1 - e^{-\beta \hbar \omega}) d(\beta \hbar \omega) \\ &= - \frac{V}{\pi^2 c^3} (\beta \hbar)^{-3} \int_0^\infty x^2 \ln(1 - e^{-x}) dx \end{aligned}$$

Photon gas $D(\omega)d\omega = \frac{V}{\pi^2 c^3} \omega^2 d\omega$

- Maximum for $U(\omega, T)d\omega = \frac{V}{\pi^2 c^3} \frac{\hbar \omega^3 d\omega}{e^{\frac{\hbar \omega}{kT}} - 1}$:

$$\frac{d}{dx} \left(\frac{x^3}{e^x - 1} \right) = 0 \Rightarrow x_m = \frac{\hbar \omega_m}{kT} \simeq 2.822, \text{ Wien's displacement law.}$$

- Grand partition function for the photons:

$$\begin{aligned} \ln \mathcal{Z} &= - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l}) \\ &= - \frac{V}{\pi^2 c^3} \int_0^\infty \omega^2 \ln(1 - e^{-\beta \hbar \omega}) d\omega \\ &= - \frac{V}{\pi^2 c^3} (\beta \hbar)^{-3} \int_0^\infty (\beta \hbar \omega)^2 \ln(1 - e^{-\beta \hbar \omega}) d(\beta \hbar \omega) \\ &= - \frac{V}{\pi^2 c^3} (\beta \hbar)^{-3} \int_0^\infty x^2 \ln(1 - e^{-x}) dx \\ &= - \frac{V}{\pi^2 c^3} \frac{1}{(\beta \hbar)^3} \int_0^\infty \ln(1 - e^{-x}) d\frac{x^3}{3} \end{aligned}$$

Photon gas $D(\omega)d\omega = \frac{V}{\pi^2 c^3} \omega^2 d\omega$

- Maximum for $U(\omega, T)d\omega = \frac{V}{\pi^2 c^3} \frac{\hbar \omega^3 d\omega}{e^{\frac{\hbar \omega}{kT}} - 1}$:

$$\frac{d}{dx} \left(\frac{x^3}{e^x - 1} \right) = 0 \Rightarrow x_m = \frac{\hbar \omega_m}{kT} \simeq 2.822, \text{ Wien's displacement law.}$$

- Grand partition function for the photons:

$$\begin{aligned} \ln \mathcal{Z} &= - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l}) \\ &= - \frac{V}{\pi^2 c^3} \int_0^\infty \omega^2 \ln(1 - e^{-\beta \hbar \omega}) d\omega \\ &= - \frac{V}{\pi^2 c^3} (\beta \hbar)^{-3} \int_0^\infty (\beta \hbar \omega)^2 \ln(1 - e^{-\beta \hbar \omega}) d(\beta \hbar \omega) \\ &= - \frac{V}{\pi^2 c^3} (\beta \hbar)^{-3} \int_0^\infty x^2 \ln(1 - e^{-x}) dx \\ &= - \frac{V}{\pi^2 c^3} \frac{1}{(\beta \hbar)^3} \int_0^\infty \ln(1 - e^{-x}) d\frac{x^3}{3} \\ &= - \frac{V}{\pi^2 c^3} \frac{1}{(\beta \hbar)^3} \left[\frac{x^3}{3} \ln(1 - e^{-x}) \Big|_0^\infty - \int_0^\infty \frac{x^3}{3} \cdot \frac{e^{-x}}{1 - e^{-x}} dx \right] \end{aligned}$$

Photon gas $D(\omega)d\omega = \frac{V}{\pi^2 c^3} \omega^2 d\omega$

- Maximum for $U(\omega, T)d\omega = \frac{V}{\pi^2 c^3} \frac{\hbar \omega^3 d\omega}{e^{\frac{\hbar \omega}{kT}} - 1}$:

$$\frac{d}{dx} \left(\frac{x^3}{e^x - 1} \right) = 0 \Rightarrow x_m = \frac{\hbar \omega_m}{kT} \simeq 2.822, \text{ Wien's displacement law.}$$

- Grand partition function for the photons:

$$\begin{aligned} \ln \mathcal{Z} &= - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l}) \\ &= - \frac{V}{\pi^2 c^3} \int_0^\infty \omega^2 \ln(1 - e^{-\beta \hbar \omega}) d\omega \\ &= - \frac{V}{\pi^2 c^3} (\beta \hbar)^{-3} \int_0^\infty (\beta \hbar \omega)^2 \ln(1 - e^{-\beta \hbar \omega}) d(\beta \hbar \omega) \\ &= - \frac{V}{\pi^2 c^3} (\beta \hbar)^{-3} \int_0^\infty x^2 \ln(1 - e^{-x}) dx \\ &= - \frac{V}{\pi^2 c^3} \frac{1}{(\beta \hbar)^3} \int_0^\infty \ln(1 - e^{-x}) d\frac{x^3}{3} \\ &= - \frac{V}{\pi^2 c^3} \frac{1}{(\beta \hbar)^3} \left[\frac{x^3}{3} \ln(1 - e^{-x}) \Big|_0^\infty - \int_0^\infty \frac{x^3}{3} \cdot \frac{e^{-x}}{1 - e^{-x}} dx \right] \\ &= \frac{V}{\pi^2 c^3} \frac{1}{(\beta \hbar)^3} \frac{1}{3} \int_0^\infty \frac{x^3}{e^x - 1} dx \end{aligned}$$

Photon gas $D(\omega)d\omega = \frac{V}{\pi^2 c^3} \omega^2 d\omega$

- Maximum for $U(\omega, T)d\omega = \frac{V}{\pi^2 c^3} \frac{\hbar \omega^3 d\omega}{e^{\frac{\hbar \omega}{kT}} - 1}$:

$$\frac{d}{dx} \left(\frac{x^3}{e^x - 1} \right) = 0 \Rightarrow x_m = \frac{\hbar \omega_m}{kT} \simeq 2.822, \text{ Wien's displacement law.}$$

- Grand partition function for the photons:

$$\begin{aligned} \ln \mathcal{Z} &= - \sum \omega_l \ln(1 - e^{-\alpha - \beta \varepsilon_l}) \\ &= - \frac{V}{\pi^2 c^3} \int_0^\infty \omega^2 \ln(1 - e^{-\beta \hbar \omega}) d\omega \\ &= - \frac{V}{\pi^2 c^3} (\beta \hbar)^{-3} \int_0^\infty (\beta \hbar \omega)^2 \ln(1 - e^{-\beta \hbar \omega}) d(\beta \hbar \omega) \\ &= - \frac{V}{\pi^2 c^3} (\beta \hbar)^{-3} \int_0^\infty x^2 \ln(1 - e^{-x}) dx \\ &= - \frac{V}{\pi^2 c^3} \frac{1}{(\beta \hbar)^3} \int_0^\infty \ln(1 - e^{-x}) d \frac{x^3}{3} \\ &= - \frac{V}{\pi^2 c^3} \frac{1}{(\beta \hbar)^3} \left[\frac{x^3}{3} \ln(1 - e^{-x}) \Big|_0^\infty - \int_0^\infty \frac{x^3}{3} \cdot \frac{e^{-x}}{1 - e^{-x}} dx \right] \\ &= \frac{V}{\pi^2 c^3} \frac{1}{(\beta \hbar)^3} \frac{1}{3} \int_0^\infty \frac{x^3}{e^x - 1} dx = \frac{\pi^2 V}{45 c^3} \frac{1}{(\beta \hbar)^3}. \end{aligned}$$

Photon gas

$$U = \frac{\pi^2 k^4}{15c^3 \hbar^3} T^4 V,$$

$$\ln \mathcal{Z} = \frac{\pi^2 V}{45c^3} \frac{1}{(\beta \hbar)^3}.$$

- Internal energy:

$$U = -\frac{\partial}{\partial \beta} \ln \mathcal{Z}$$

Photon gas

$$U = \frac{\pi^2 k^4}{15c^3 \hbar^3} T^4 V,$$

$$\ln \mathcal{Z} = \frac{\pi^2 V}{45c^3} \frac{1}{(\beta \hbar)^3}.$$

- Internal energy:

$$U = -\frac{\partial}{\partial \beta} \ln \mathcal{Z} = -\frac{\pi^2 V}{45c^3} \frac{1}{\hbar^3} \frac{-3}{\beta^4}$$

Photon gas

$$U = \frac{\pi^2 k^4}{15c^3 \hbar^3} T^4 V, \quad \ln \mathcal{Z} = \frac{\pi^2 V}{45c^3} \frac{1}{(\beta \hbar)^3}.$$

- Internal energy:

$$U = -\frac{\partial}{\partial \beta} \ln \mathcal{Z} = -\frac{\pi^2 V}{45c^3} \frac{1}{\hbar^3} \frac{-3}{\beta^4} = \frac{\pi^2 k^4}{15c^3 \hbar^3} T^4 V.$$

Photon gas

$$U = \frac{\pi^2 k^4}{15c^3 \hbar^3} T^4 V, \quad \ln \mathcal{Z} = \frac{\pi^2 V}{45c^3} \frac{1}{(\beta \hbar)^3}.$$

- Internal energy:

$$U = -\frac{\partial}{\partial \beta} \ln \mathcal{Z} = -\frac{\pi^2 V}{45c^3} \frac{1}{\hbar^3} \frac{-3}{\beta^4} = \frac{\pi^2 k^4}{15c^3 \hbar^3} T^4 V.$$

- Pressure:

$$p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln \mathcal{Z}$$

Photon gas

$$U = \frac{\pi^2 k^4}{15c^3 \hbar^3} T^4 V, \quad \ln \mathcal{Z} = \frac{\pi^2 V}{45c^3} \frac{1}{(\beta \hbar)^3}.$$

- Internal energy:

$$U = -\frac{\partial}{\partial \beta} \ln \mathcal{Z} = -\frac{\pi^2 V}{45c^3} \frac{1}{\hbar^3} \frac{-3}{\beta^4} = \frac{\pi^2 k^4}{15c^3 \hbar^3} T^4 V.$$

- Pressure:

$$p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln \mathcal{Z} = \frac{1}{\beta} \frac{\pi^2}{45c^3} \frac{1}{(\beta \hbar)^3}$$

Photon gas

$$U = \frac{\pi^2 k^4}{15c^3 \hbar^3} T^4 V, \quad \ln \mathcal{Z} = \frac{\pi^2 V}{45c^3} \frac{1}{(\beta \hbar)^3}.$$

- Internal energy:

$$U = -\frac{\partial}{\partial \beta} \ln \mathcal{Z} = -\frac{\pi^2 V}{45c^3} \frac{1}{\hbar^3} \frac{-3}{\beta^4} = \frac{\pi^2 k^4}{15c^3 \hbar^3} T^4 V.$$

- Pressure:

$$p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln \mathcal{Z} = \frac{1}{\beta} \frac{\pi^2}{45c^3} \frac{1}{(\beta \hbar)^3} = \frac{\pi^2 k^4}{45c^3 \hbar^3} T^4$$

Photon gas

$$U = \frac{\pi^2 k^4}{15c^3 \hbar^3} T^4 V, \quad \ln \mathcal{Z} = \frac{\pi^2 V}{45c^3} \frac{1}{(\beta \hbar)^3}.$$

- Internal energy:

$$U = -\frac{\partial}{\partial \beta} \ln \mathcal{Z} = -\frac{\pi^2 V}{45c^3} \frac{1}{\hbar^3} \frac{-3}{\beta^4} = \frac{\pi^2 k^4}{15c^3 \hbar^3} T^4 V.$$

- Pressure:

$$p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln \mathcal{Z} = \frac{1}{\beta} \frac{\pi^2}{45c^3} \frac{1}{(\beta \hbar)^3} = \frac{\pi^2 k^4}{45c^3 \hbar^3} T^4 = \frac{1}{3} \frac{U}{V}.$$

Photon gas

$$U = \frac{\pi^2 k^4}{15c^3 \hbar^3} T^4 V, \quad \ln \mathcal{Z} = \frac{\pi^2 V}{45c^3} \frac{1}{(\beta \hbar)^3}.$$

- Internal energy:

$$U = -\frac{\partial}{\partial \beta} \ln \mathcal{Z} = -\frac{\pi^2 V}{45c^3} \frac{1}{\hbar^3} \frac{-3}{\beta^4} = \frac{\pi^2 k^4}{15c^3 \hbar^3} T^4 V.$$

- Pressure:

$$p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln \mathcal{Z} = \frac{1}{\beta} \frac{\pi^2}{45c^3} \frac{1}{(\beta \hbar)^3} = \frac{\pi^2 k^4}{45c^3 \hbar^3} T^4 = \frac{1}{3} \frac{U}{V}.$$

- Entropy:

$$S = k(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta})$$

Photon gas

$$U = \frac{\pi^2 k^4}{15c^3 \hbar^3} T^4 V, \quad \ln \mathcal{Z} = \frac{\pi^2 V}{45c^3} \frac{1}{(\beta \hbar)^3}.$$

- Internal energy:

$$U = -\frac{\partial}{\partial \beta} \ln \mathcal{Z} = -\frac{\pi^2 V}{45c^3} \frac{1}{\hbar^3} \frac{-3}{\beta^4} = \frac{\pi^2 k^4}{15c^3 \hbar^3} T^4 V.$$

- Pressure:

$$p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln \mathcal{Z} = \frac{1}{\beta} \frac{\pi^2}{45c^3} \frac{1}{(\beta \hbar)^3} = \frac{\pi^2 k^4}{45c^3 \hbar^3} T^4 = \frac{1}{3} \frac{U}{V}.$$

- Entropy:

$$S = k(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}) = k(\ln \mathcal{Z} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta})$$

Photon gas

$$U = \frac{\pi^2 k^4}{15c^3 \hbar^3} T^4 V, \quad \ln \mathcal{Z} = \frac{\pi^2 V}{45c^3} \frac{1}{(\beta \hbar)^3}.$$

- Internal energy:

$$U = -\frac{\partial}{\partial \beta} \ln \mathcal{Z} = -\frac{\pi^2 V}{45c^3} \frac{1}{\hbar^3} \frac{-3}{\beta^4} = \frac{\pi^2 k^4}{15c^3 \hbar^3} T^4 V.$$

- Pressure:

$$p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln \mathcal{Z} = \frac{1}{\beta} \frac{\pi^2}{45c^3} \frac{1}{(\beta \hbar)^3} = \frac{\pi^2 k^4}{45c^3 \hbar^3} T^4 = \frac{1}{3} \frac{U}{V}.$$

- Entropy:

$$\begin{aligned} S &= k \left(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta} \right) = k \left(\ln \mathcal{Z} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta} \right) \\ &= k \left[\frac{\pi^2 V}{45c^3} \frac{1}{(\beta \hbar)^3} + \beta \frac{\pi^2 k^4}{15c^3 \hbar^3} T^4 V \right] \end{aligned}$$

Photon gas

$$U = \frac{\pi^2 k^4}{15c^3 \hbar^3} T^4 V,$$

$$\ln \mathcal{Z} = \frac{\pi^2 V}{45c^3} \frac{1}{(\beta \hbar)^3}.$$

- Internal energy:

$$U = -\frac{\partial}{\partial \beta} \ln \mathcal{Z} = -\frac{\pi^2 V}{45c^3} \frac{1}{\hbar^3} \frac{-3}{\beta^4} = \frac{\pi^2 k^4}{15c^3 \hbar^3} T^4 V.$$

- Pressure:

$$p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln \mathcal{Z} = \frac{1}{\beta} \frac{\pi^2}{45c^3} \frac{1}{(\beta \hbar)^3} = \frac{\pi^2 k^4}{45c^3 \hbar^3} T^4 = \frac{1}{3} \frac{U}{V}.$$

- Entropy:

$$\begin{aligned} S &= k \left(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta} \right) = k \left(\ln \mathcal{Z} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta} \right) \\ &= k \left[\frac{\pi^2 V}{45c^3} \frac{1}{(\beta \hbar)^3} + \beta \frac{\pi^2 k^4}{15c^3 \hbar^3} T^4 V \right] \\ &= k \left(\frac{\pi^2 V}{45c^3} \frac{k^3 T^3}{\hbar^3} + \frac{\pi^2 V}{15c^3} \frac{k^3 T^3}{\hbar^3} \right) \end{aligned}$$

Photon gas

$$U = \frac{\pi^2 k^4}{15c^3 \hbar^3} T^4 V, \quad \ln \mathcal{Z} = \frac{\pi^2 V}{45c^3} \frac{1}{(\beta \hbar)^3}.$$

- Internal energy:

$$U = -\frac{\partial}{\partial \beta} \ln \mathcal{Z} = -\frac{\pi^2 V}{45c^3} \frac{1}{\hbar^3} \frac{-3}{\beta^4} = \frac{\pi^2 k^4}{15c^3 \hbar^3} T^4 V.$$

- Pressure:

$$p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln \mathcal{Z} = \frac{1}{\beta} \frac{\pi^2}{45c^3} \frac{1}{(\beta \hbar)^3} = \frac{\pi^2 k^4}{45c^3 \hbar^3} T^4 = \frac{1}{3} \frac{U}{V}.$$

- Entropy:

$$\begin{aligned} S &= k \left(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta} \right) = k \left(\ln \mathcal{Z} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta} \right) \\ &= k \left[\frac{\pi^2 V}{45c^3} \frac{1}{(\beta \hbar)^3} + \beta \frac{\pi^2 k^4}{15c^3 \hbar^3} T^4 V \right] \\ &= k \left(\frac{\pi^2 V}{45c^3} \frac{k^3 T^3}{\hbar^3} + \frac{\pi^2 V}{15c^3} \frac{k^3 T^3}{\hbar^3} \right) \\ &= \frac{4}{45} \frac{\pi^2 k^4 V}{c^3 \hbar^3} T^3. \quad (\text{same as (2.6.4)}) \end{aligned}$$

Table of contents

1 §8. Bose statistics and Fermi statistics

- 8.1 Thermal parameters in statistics
- 8.2 U of weak degenerated ideal Fermion(Boson) gas
- 8.3 Bose-Einstein Condensate (BEC)
- 8.4 Photon gas
- 8.5 Free electron gas in metal

8.5 Free electron gas in metal

- Degenerated Fermi gas ($e^{\alpha} \gg 1$ or $n\lambda^3 \gg 1$).

8.5 Free electron gas in metal

- Degenerated Fermi gas ($e^{\alpha} \gg 1$ or $n\lambda^3 \gg 1$).
- Problem: in the metal, electron's contribution on C_V is negligible.

8.5 Free electron gas in metal

- Degenerated Fermi gas ($e^{\alpha} \gg 1$ or $n\lambda^3 \gg 1$).
- Problem: in the metal, electron's contribution on C_V is negligible.
- Model: the free electrons are in degeneracy, needs Fermi statistics.

8.5 Free electron gas in metal

- Degenerated Fermi gas ($e^{\alpha} \gg 1$ or $n\lambda^3 \gg 1$).
- Problem: in the metal, electron's contribution on C_V is negligible.
- Model: the free electrons are in degeneracy, needs Fermi statistics.
- Example: Copper (Cu)

8.5 Free electron gas in metal

- Degenerated Fermi gas ($e^{\alpha} \gg 1$ or $n\lambda^3 \gg 1$).
- Problem: in the metal, electron's contribution on C_V is negligible.
- Model: the free electrons are in degeneracy, needs Fermi statistics.
- Example: Copper (Cu)
 $\rho = 8.9 \times 10^3 \text{kg} \cdot \text{m}^{-3},$

8.5 Free electron gas in metal

- Degenerated Fermi gas ($e^{\alpha} \gg 1$ or $n\lambda^3 \gg 1$).
- Problem: in the metal, electron's contribution on C_V is negligible.
- Model: the free electrons are in degeneracy, needs Fermi statistics.
- Example: Copper (Cu)
 $\rho = 8.9 \times 10^3 \text{kg} \cdot \text{m}^{-3}$, atomic weight: 63,

8.5 Free electron gas in metal

- Degenerated Fermi gas ($e^{\alpha} \gg 1$ or $n\lambda^3 \gg 1$).
- Problem: in the metal, electron's contribution on C_V is negligible.
- Model: the free electrons are in degeneracy, needs Fermi statistics.
- Example: Copper (Cu)
 $\rho = 8.9 \times 10^3 \text{kg} \cdot \text{m}^{-3}$, atomic weight: 63, number density $n_e \sim n_{\text{Cu}} \simeq \frac{8.9 \times 10^3 \text{kg} \cdot \text{m}^{-3}}{63 \times 1.67 \times 10^{-27} \text{kg}}$

8.5 Free electron gas in metal

- Degenerated Fermi gas ($e^{\alpha} \gg 1$ or $n\lambda^3 \gg 1$).
- Problem: in the metal, electron's contribution on C_V is negligible.
- Model: the free electrons are in degeneracy, needs Fermi statistics.
- Example: Copper (Cu)

$\rho = 8.9 \times 10^3 \text{kg} \cdot \text{m}^{-3}$, atomic weight: 63, number density $n_e \sim n_{\text{Cu}} \simeq \frac{8.9 \times 10^3 \text{kg} \cdot \text{m}^{-3}}{63 \times 1.67 \times 10^{-27} \text{kg}} \simeq 8.5 \times 10^{28} \text{m}^{-3}$.

8.5 Free electron gas in metal

- Degenerated Fermi gas ($e^{\alpha} \gg 1$ or $n\lambda^3 \gg 1$).
- Problem: in the metal, electron's contribution on C_V is negligible.
- Model: the free electrons are in degeneracy, needs Fermi statistics.
- Example: Copper (Cu)

$$\rho = 8.9 \times 10^3 \text{kg} \cdot \text{m}^{-3}, \text{ atomic weight: } 63, \text{ number density } n_e \sim n_{\text{Cu}} \simeq \frac{8.9 \times 10^3 \text{kg} \cdot \text{m}^{-3}}{63 \times 1.67 \times 10^{-27} \text{kg}} \simeq 8.5 \times 10^{28} \text{m}^{-3}.$$

$$n\lambda^3 = n \left(\frac{h^2}{2\pi m k T} \right)^{\frac{3}{2}}$$

8.5 Free electron gas in metal

- Degenerated Fermi gas ($e^\alpha \gg 1$ or $n\lambda^3 \gg 1$).
- Problem: in the metal, electron's contribution on C_V is negligible.
- Model: the free electrons are in degeneracy, needs Fermi statistics.

- Example: Copper (Cu)

$\rho = 8.9 \times 10^3 \text{ kg} \cdot \text{m}^{-3}$, atomic weight: 63, number density $n_e \sim n_{\text{Cu}} \simeq \frac{8.9 \times 10^3 \text{ kg} \cdot \text{m}^{-3}}{63 \times 1.67 \times 10^{-27} \text{ kg}} \simeq 8.5 \times 10^{28} \text{ m}^{-3}$.

$$n\lambda^3 = n \left(\frac{h^2}{2\pi m k T} \right)^{\frac{3}{2}}$$

$$\simeq 8.5 \times 10^{28} \text{ m}^{-3} \left(\frac{(6.626 \times 10^{-34} \text{ J} \cdot \text{s})^2}{2\pi \times 9.1 \times 10^{-31} \text{ kg} \cdot 1.38 \times 10^{-23} \text{ J} \cdot \text{K}^{-1} \cdot 300 \text{ K} \frac{T}{300 \text{ K}}} \right)^{\frac{3}{2}}$$

8.5 Free electron gas in metal

- Degenerated Fermi gas ($e^\alpha \gg 1$ or $n\lambda^3 \gg 1$).
- Problem: in the metal, electron's contribution on C_V is negligible.
- Model: the free electrons are in degeneracy, needs Fermi statistics.

- Example: Copper (Cu)

$\rho = 8.9 \times 10^3 \text{ kg} \cdot \text{m}^{-3}$, atomic weight: 63, number density $n_e \sim n_{\text{Cu}} \simeq \frac{8.9 \times 10^3 \text{ kg} \cdot \text{m}^{-3}}{63 \times 1.67 \times 10^{-27} \text{ kg}} \simeq 8.5 \times 10^{28} \text{ m}^{-3}$.

$$\begin{aligned}
 n\lambda^3 &= n \left(\frac{h^2}{2\pi m k T} \right)^{\frac{3}{2}} \\
 &\simeq 8.5 \times 10^{28} \text{ m}^{-3} \left(\frac{(6.626 \times 10^{-34} \text{ J} \cdot \text{s})^2}{2\pi \times 9.1 \times 10^{-31} \text{ kg} \cdot 1.38 \times 10^{-23} \text{ J} \cdot \text{K}^{-1} \cdot 300 \text{ K} \frac{T}{300 \text{ K}}} \right)^{\frac{3}{2}} \\
 &\simeq 3400 \left(\frac{T}{300 \text{ K}} \right)^{-\frac{3}{2}}
 \end{aligned}$$

8.5 Free electron gas in metal

- Degenerated Fermi gas ($e^\alpha \gg 1$ or $n\lambda^3 \gg 1$).
- Problem: in the metal, electron's contribution on C_V is negligible.
- Model: the free electrons are in degeneracy, needs Fermi statistics.

- Example: Copper (Cu)

$\rho = 8.9 \times 10^3 \text{ kg} \cdot \text{m}^{-3}$, atomic weight: 63, number density $n_e \sim n_{\text{Cu}} \simeq \frac{8.9 \times 10^3 \text{ kg} \cdot \text{m}^{-3}}{63 \times 1.67 \times 10^{-27} \text{ kg}} \simeq 8.5 \times 10^{28} \text{ m}^{-3}$.

$$\begin{aligned}
 n\lambda^3 &= n \left(\frac{h^2}{2\pi m k T} \right)^{\frac{3}{2}} \\
 &\simeq 8.5 \times 10^{28} \text{ m}^{-3} \left(\frac{(6.626 \times 10^{-34} \text{ J} \cdot \text{s})^2}{2\pi \times 9.1 \times 10^{-31} \text{ kg} \cdot 1.38 \times 10^{-23} \text{ J} \cdot \text{K}^{-1} \cdot 300 \text{ K} \frac{T}{300 \text{ K}}} \right)^{\frac{3}{2}} \\
 &\simeq 3400 \left(\frac{T}{300 \text{ K}} \right)^{-\frac{3}{2}} \gg 1.
 \end{aligned}$$

Free electron gas in metal

- Fermi distribution: $f = \frac{a_l}{\omega_l} = \frac{1}{e^{\alpha + \beta \varepsilon_l} + 1} = \frac{1}{e^{\frac{\varepsilon_l - \mu}{kT}} + 1}.$

Free electron gas in metal

- Fermi distribution: $f = \frac{a_l}{\omega_l} = \frac{1}{e^{\alpha+\beta\varepsilon_l}+1} = \frac{1}{e^{\frac{\varepsilon_l-\mu}{kT}}+1}$.
- Number of quantum states: $2 \cdot \frac{V \cdot 4\pi p^2 dp}{h^3}$, where 2 is because of electron's spin.

Free electron gas in metal

- Fermi distribution: $f = \frac{a_l}{\omega_l} = \frac{1}{e^{\alpha + \beta \varepsilon_l} + 1} = \frac{1}{e^{\frac{\varepsilon_l - \mu}{kT}} + 1}$.
- Number of quantum states: $2 \cdot \frac{V \cdot 4\pi p^2 dp}{h^3}$, where 2 is because of electron's spin.
- Convert to kinetic energy: $D(\varepsilon)d\varepsilon = \frac{4\pi V}{h^3}(2m)^{\frac{3}{2}}\varepsilon^{\frac{1}{2}}d\varepsilon$.

Free electron gas in metal

- Fermi distribution: $f = \frac{a_l}{\omega_l} = \frac{1}{e^{\alpha+\beta\varepsilon_l}+1} = \frac{1}{e^{\frac{\varepsilon_l-\mu}{kT}}+1}$.
- Number of quantum states: $2 \cdot \frac{V \cdot 4\pi p^2 dp}{h^3}$, where 2 is because of electron's spin.
- Convert to kinetic energy: $D(\varepsilon)d\varepsilon = \frac{4\pi V}{h^3}(2m)^{\frac{3}{2}}\varepsilon^{\frac{1}{2}}d\varepsilon$.
- Number of electron at $(\varepsilon, \varepsilon + d\varepsilon)$: ($a_l = f \cdot \omega_l$)

Free electron gas in metal

- Fermi distribution: $f = \frac{a_l}{\omega_l} = \frac{1}{e^{\alpha+\beta\varepsilon_l}+1} = \frac{1}{e^{\frac{\varepsilon_l-\mu}{kT}}+1}$.
- Number of quantum states: $2 \cdot \frac{V \cdot 4\pi p^2 dp}{h^3}$, where 2 is because of electron's spin.
- Convert to kinetic energy: $D(\varepsilon)d\varepsilon = \frac{4\pi V}{h^3}(2m)^{\frac{3}{2}}\varepsilon^{\frac{1}{2}}d\varepsilon$.
- Number of electron at $(\varepsilon, \varepsilon + d\varepsilon)$: $(a_l = f \cdot \omega_l)$

$$\frac{4\pi V}{h^3}(2m)^{\frac{3}{2}}\frac{\varepsilon^{\frac{1}{2}}}{e^{\frac{\varepsilon-\mu}{kT}}+1}d\varepsilon.$$

Free electron gas in metal

- Fermi distribution: $f = \frac{a_l}{\omega_l} = \frac{1}{e^{\alpha+\beta\varepsilon_l}+1} = \frac{1}{e^{\frac{\varepsilon_l-\mu}{kT}}+1}$.
- Number of quantum states: $2 \cdot \frac{V \cdot 4\pi p^2 dp}{h^3}$, where 2 is because of electron's spin.
- Convert to kinetic energy: $D(\varepsilon)d\varepsilon = \frac{4\pi V}{h^3}(2m)^{\frac{3}{2}}\varepsilon^{\frac{1}{2}}d\varepsilon$.
- Number of electron at $(\varepsilon, \varepsilon + d\varepsilon)$: $(a_l = f \cdot \omega_l)$

$$\frac{4\pi V}{h^3}(2m)^{\frac{3}{2}} \frac{\varepsilon^{\frac{1}{2}}}{e^{\frac{\varepsilon-\mu}{kT}}+1} d\varepsilon.$$
- Total number: $N = \frac{4\pi V}{h^3}(2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}}}{e^{\frac{\varepsilon-\mu}{kT}}+1} d\varepsilon.$

Free electron gas in metal

$$f = \frac{1}{e^{\frac{\varepsilon - \mu}{kT}} + 1}$$

- Case: $T=0\text{K}$.

Free electron gas in metal

$$f = \frac{1}{e^{\frac{\varepsilon - \mu}{kT}} + 1}$$

- Case: $T=0\text{K}$.
- $f = 1$ if $\varepsilon < \mu(0)$; $f = 0$ if $\varepsilon > \mu(0)$.

Free electron gas in metal

$$f = \frac{1}{e^{\frac{\varepsilon - \mu}{kT}} + 1}$$

- Case: $T=0\text{K}$.
- $f = 1$ if $\varepsilon < \mu(0)$; $f = 0$ if $\varepsilon > \mu(0)$.
- Picture: $T=0\text{K}$, electrons stay at the lowest possible state.

Free electron gas in metal

$$f = \frac{1}{e^{\frac{\varepsilon - \mu}{kT}} + 1}$$

- Case: $T=0\text{K}$.
- $f = 1$ if $\varepsilon < \mu(0)$; $f = 0$ if $\varepsilon > \mu(0)$.
- Picture: $T=0\text{K}$, electrons stay at the lowest possible state.
- $$N = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}}}{e^{\frac{\varepsilon - \mu}{kT}} + 1} d\varepsilon$$

Free electron gas in metal

$$f = \frac{1}{e^{\frac{\varepsilon - \mu}{kT}} + 1}$$

- Case: $T=0\text{K}$.
- $f = 1$ if $\varepsilon < \mu(0)$; $f = 0$ if $\varepsilon > \mu(0)$.
- Picture: $T=0\text{K}$, electrons stay at the lowest possible state.
- $$N = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}}}{e^{\frac{\varepsilon - \mu}{kT}} + 1} d\varepsilon = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^{\mu(0)} \varepsilon^{\frac{1}{2}} d\varepsilon$$

Free electron gas in metal

$$f = \frac{1}{e^{\frac{\varepsilon - \mu}{kT}} + 1}$$

- Case: $T=0\text{K}$.
- $f = 1$ if $\varepsilon < \mu(0)$; $f = 0$ if $\varepsilon > \mu(0)$.
- Picture: $T=0\text{K}$, electrons stay at the lowest possible state.
- $$N = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}}}{e^{\frac{\varepsilon - \mu}{kT}} + 1} d\varepsilon = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^{\mu(0)} \varepsilon^{\frac{1}{2}} d\varepsilon$$

$$= \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \frac{2}{3} \mu(0)^{\frac{3}{2}}.$$

Free electron gas in metal

$$f = \frac{1}{e^{\frac{\varepsilon - \mu}{kT}} + 1}$$

- Case: $T=0\text{K}$.
- $f = 1$ if $\varepsilon < \mu(0)$; $f = 0$ if $\varepsilon > \mu(0)$.
- Picture: $T=0\text{K}$, electrons stay at the lowest possible state.
- $$N = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}}}{e^{\frac{\varepsilon - \mu}{kT}} + 1} d\varepsilon = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^{\mu(0)} \varepsilon^{\frac{1}{2}} d\varepsilon$$

$$= \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \frac{2}{3} \mu(0)^{\frac{3}{2}}.$$
- $\Rightarrow \mu(0) = \frac{\hbar^2}{2m} \left(3\pi^2 \frac{N}{V} \right)^{\frac{2}{3}}$, Fermi energy level.

Free electron gas in metal

$$f = \frac{1}{e^{\frac{\varepsilon - \mu}{kT}} + 1}$$

- Case: $T=0\text{K}$.
- $f = 1$ if $\varepsilon < \mu(0)$; $f = 0$ if $\varepsilon > \mu(0)$.
- Picture: $T=0\text{K}$, electrons stay at the lowest possible state.
- $$N = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}}}{e^{\frac{\varepsilon - \mu}{kT}} + 1} d\varepsilon = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^{\mu(0)} \varepsilon^{\frac{1}{2}} d\varepsilon$$

$$= \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \frac{2}{3} \mu(0)^{\frac{3}{2}}.$$
- $\Rightarrow \mu(0) = \frac{\hbar^2}{2m} \left(3\pi^2 \frac{N}{V} \right)^{\frac{2}{3}}$, Fermi energy level.
- Define $\frac{p_F^2}{2m} = \mu(0)$,

Free electron gas in metal

$$f = \frac{1}{e^{\frac{\varepsilon - \mu}{kT}} + 1}$$

- Case: $T=0\text{K}$.
- $f = 1$ if $\varepsilon < \mu(0)$; $f = 0$ if $\varepsilon > \mu(0)$.
- Picture: $T=0\text{K}$, electrons stay at the lowest possible state.
- $$N = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}}}{e^{\frac{\varepsilon - \mu}{kT}} + 1} d\varepsilon = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^{\mu(0)} \varepsilon^{\frac{1}{2}} d\varepsilon$$

$$= \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \frac{2}{3} \mu(0)^{\frac{3}{2}}.$$
- $\Rightarrow \mu(0) = \frac{\hbar^2}{2m} \left(3\pi^2 \frac{N}{V} \right)^{\frac{2}{3}}$, Fermi energy level.
- Define $\frac{p_F^2}{2m} = \mu(0)$, Fermi momentum: $p_F = (3\pi^2 n)^{\frac{1}{3}} \hbar$,

Free electron gas in metal

$$f = \frac{1}{e^{\frac{\varepsilon - \mu}{kT}} + 1}$$

- Case: $T=0\text{K}$.
- $f = 1$ if $\varepsilon < \mu(0)$; $f = 0$ if $\varepsilon > \mu(0)$.
- Picture: $T=0\text{K}$, electrons stay at the lowest possible state.
- $$N = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}}}{e^{\frac{\varepsilon - \mu}{kT}} + 1} d\varepsilon = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^{\mu(0)} \varepsilon^{\frac{1}{2}} d\varepsilon$$

$$= \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \frac{2}{3} \mu(0)^{\frac{3}{2}}.$$
- $\Rightarrow \mu(0) = \frac{\hbar^2}{2m} \left(3\pi^2 \frac{N}{V} \right)^{\frac{2}{3}}$, Fermi energy level.
- Define $\frac{p_F^2}{2m} = \mu(0)$, Fermi momentum: $p_F = (3\pi^2 n)^{\frac{1}{3}} \hbar$, the maximal momentum for the electrons.

Free electron gas in metal

$$f = \frac{1}{e^{\frac{\varepsilon - \mu}{kT}} + 1}$$

- Case: $T=0\text{K}$.
- $f = 1$ if $\varepsilon < \mu(0)$; $f = 0$ if $\varepsilon > \mu(0)$.
- Picture: $T=0\text{K}$, electrons stay at the lowest possible state.
- $$N = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}}}{e^{\frac{\varepsilon - \mu}{kT}} + 1} d\varepsilon = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^{\mu(0)} \varepsilon^{\frac{1}{2}} d\varepsilon$$

$$= \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \frac{2}{3} \mu(0)^{\frac{3}{2}}.$$
- $\Rightarrow \mu(0) = \frac{\hbar^2}{2m} \left(3\pi^2 \frac{N}{V} \right)^{\frac{2}{3}}$, Fermi energy level.
- Define $\frac{p_F^2}{2m} = \mu(0)$, Fermi momentum: $p_F = (3\pi^2 n)^{\frac{1}{3}} \hbar$, the maximal momentum for the electrons.
- Define $kT_F = \mu(0)$,

Free electron gas in metal

$$f = \frac{1}{e^{\frac{\varepsilon - \mu}{kT}} + 1}$$

- Case: $T=0K$.
- $f = 1$ if $\varepsilon < \mu(0)$; $f = 0$ if $\varepsilon > \mu(0)$.
- Picture: $T=0K$, electrons stay at the lowest possible state.
- $$N = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}}}{e^{\frac{\varepsilon - \mu}{kT}} + 1} d\varepsilon = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^{\mu(0)} \varepsilon^{\frac{1}{2}} d\varepsilon$$

$$= \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \frac{2}{3} \mu(0)^{\frac{3}{2}}.$$
- $\Rightarrow \mu(0) = \frac{\hbar^2}{2m} \left(3\pi^2 \frac{N}{V} \right)^{\frac{2}{3}}$, Fermi energy level.
- Define $\frac{p_F^2}{2m} = \mu(0)$, Fermi momentum: $p_F = (3\pi^2 n)^{\frac{1}{3}} \hbar$, the maximal momentum for the electrons.
- Define $kT_F = \mu(0)$, Fermi temperature: $T_F = \frac{\mu(0)}{k}$, a criterion for degeneracy ($T < T_F$).

Free electron gas in metal

- Fermi temperature for the copper:

$$T_F = \frac{\mu^{(0)}}{k} = \frac{1}{k} \frac{\hbar^2}{2m} \left(3\pi^2 \frac{N}{V} \right)^{\frac{2}{3}}$$

Free electron gas in metal

- Fermi temperature for the copper:

$$\begin{aligned} T_F &= \frac{\mu^{(0)}}{k} = \frac{1}{k} \frac{\hbar^2}{2m} \left(3\pi^2 \frac{N}{V} \right)^{\frac{2}{3}} \\ &= \frac{1}{k} \frac{\hbar^2}{2m} \left(3\pi^2 \cdot 8.5 \times 10^{28} \text{m}^{-3} \right)^{\frac{2}{3}} \end{aligned}$$

Free electron gas in metal

- Fermi temperature for the copper:

$$\begin{aligned} T_F &= \frac{\mu^{(0)}}{k} = \frac{1}{k} \frac{\hbar^2}{2m} \left(3\pi^2 \frac{N}{V} \right)^{\frac{2}{3}} \\ &= \frac{1}{k} \frac{\hbar^2}{2m} \left(3\pi^2 \cdot 8.5 \times 10^{28} \text{m}^{-3} \right)^{\frac{2}{3}} \simeq 8.2 \times 10^4 \text{K}. \end{aligned}$$

Free electron gas in metal

- Fermi temperature for the copper:

$$\begin{aligned} T_F &= \frac{\mu(0)}{k} = \frac{1}{k} \frac{\hbar^2}{2m} \left(3\pi^2 \frac{N}{V} \right)^{\frac{2}{3}} \\ &= \frac{1}{k} \frac{\hbar^2}{2m} \left(3\pi^2 \cdot 8.5 \times 10^{28} \text{m}^{-3} \right)^{\frac{2}{3}} \simeq 8.2 \times 10^4 \text{K}. \end{aligned}$$

- Internal energy: $U(0) = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{3}{2}}}{e^{\frac{\varepsilon-\mu}{kT}} + 1} d\varepsilon$

Free electron gas in metal

- Fermi temperature for the copper:

$$\begin{aligned} T_F &= \frac{\mu^{(0)}}{k} = \frac{1}{k} \frac{\hbar^2}{2m} \left(3\pi^2 \frac{N}{V} \right)^{\frac{2}{3}} \\ &= \frac{1}{k} \frac{\hbar^2}{2m} \left(3\pi^2 \cdot 8.5 \times 10^{28} \text{m}^{-3} \right)^{\frac{2}{3}} \simeq 8.2 \times 10^4 \text{K}. \end{aligned}$$

- Internal energy: $U(0) = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{3}{2}}}{e^{\frac{\varepsilon-\mu}{kT}} + 1} d\varepsilon$
 $= \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^{\mu^{(0)}} \varepsilon^{\frac{3}{2}} d\varepsilon$

Free electron gas in metal

- Fermi temperature for the copper:

$$T_F = \frac{\mu(0)}{k} = \frac{1}{k} \frac{\hbar^2}{2m} \left(3\pi^2 \frac{N}{V} \right)^{\frac{2}{3}}$$

$$= \frac{1}{k} \frac{\hbar^2}{2m} \left(3\pi^2 \cdot 8.5 \times 10^{28} \text{m}^{-3} \right)^{\frac{2}{3}} \simeq 8.2 \times 10^4 \text{K}.$$

- Internal energy: $U(0) = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{3}{2}}}{e^{\frac{\varepsilon-\mu}{kT}} + 1} d\varepsilon$
- $$= \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^{\mu(0)} \varepsilon^{\frac{3}{2}} d\varepsilon = \frac{3}{5} N \mu(0).$$

Free electron gas in metal

- Fermi temperature for the copper:

$$T_F = \frac{\mu(0)}{k} = \frac{1}{k} \frac{\hbar^2}{2m} \left(3\pi^2 \frac{N}{V} \right)^{\frac{2}{3}}$$

$$= \frac{1}{k} \frac{\hbar^2}{2m} \left(3\pi^2 \cdot 8.5 \times 10^{28} \text{m}^{-3} \right)^{\frac{2}{3}} \simeq 8.2 \times 10^4 \text{K}.$$

- Internal energy: $U(0) = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{3}{2}}}{e^{\frac{\varepsilon-\mu}{kT}} + 1} d\varepsilon$
- $= \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^{\mu(0)} \varepsilon^{\frac{3}{2}} d\varepsilon = \frac{3}{5} N \mu(0).$
- Pressure: $p = \frac{2}{3} \frac{U}{V}$

Free electron gas in metal

- Fermi temperature for the copper:

$$\begin{aligned} T_F &= \frac{\mu(0)}{k} = \frac{1}{k} \frac{\hbar^2}{2m} \left(3\pi^2 \frac{N}{V} \right)^{\frac{2}{3}} \\ &= \frac{1}{k} \frac{\hbar^2}{2m} \left(3\pi^2 \cdot 8.5 \times 10^{28} \text{m}^{-3} \right)^{\frac{2}{3}} \simeq 8.2 \times 10^4 \text{K}. \end{aligned}$$

- Internal energy: $U(0) = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{3}{2}}}{e^{\frac{\varepsilon-\mu}{kT}} + 1} d\varepsilon$
 $= \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^{\mu(0)} \varepsilon^{\frac{3}{2}} d\varepsilon = \frac{3}{5} N \mu(0).$
- Pressure: $p = \frac{2}{3} \frac{U}{V} = \frac{2}{5} n \mu(0).$

Free electron gas in metal

- Fermi temperature for the copper:

$$T_F = \frac{\mu(0)}{k} = \frac{1}{k} \frac{\hbar^2}{2m} \left(3\pi^2 \frac{N}{V} \right)^{\frac{2}{3}}$$

$$= \frac{1}{k} \frac{\hbar^2}{2m} \left(3\pi^2 \cdot 8.5 \times 10^{28} \text{m}^{-3} \right)^{\frac{2}{3}} \simeq 8.2 \times 10^4 \text{K}.$$

- Internal energy: $U(0) = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{3}{2}}}{e^{\frac{\varepsilon-\mu}{kT}} + 1} d\varepsilon$

$$= \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^{\mu(0)} \varepsilon^{\frac{3}{2}} d\varepsilon = \frac{3}{5} N \mu(0).$$

- Pressure: $p = \frac{2}{3} \frac{U}{V} = \frac{2}{5} n \mu(0).$

$$p_{\text{Cu}} = \frac{2}{5} \cdot 8.5 \times 10^{28} \text{m}^{-3} \cdot 1.38 \times 10^{-23} \text{J} \cdot \text{K}^{-1} \cdot 8.2 \times 10^4 \text{K}$$

Free electron gas in metal

- Fermi temperature for the copper:

$$T_F = \frac{\mu(0)}{k} = \frac{1}{k} \frac{\hbar^2}{2m} \left(3\pi^2 \frac{N}{V} \right)^{\frac{2}{3}}$$

$$= \frac{1}{k} \frac{\hbar^2}{2m} \left(3\pi^2 \cdot 8.5 \times 10^{28} \text{m}^{-3} \right)^{\frac{2}{3}} \simeq 8.2 \times 10^4 \text{K}.$$

- Internal energy: $U(0) = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{3}{2}}}{e^{\frac{\varepsilon-\mu}{kT}} + 1} d\varepsilon$

$$= \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^{\mu(0)} \varepsilon^{\frac{3}{2}} d\varepsilon = \frac{3}{5} N \mu(0).$$

- Pressure: $p = \frac{2}{3} \frac{U}{V} = \frac{2}{5} n \mu(0).$

$$p_{\text{Cu}} = \frac{2}{5} \cdot 8.5 \times 10^{28} \text{m}^{-3} \cdot 1.38 \times 10^{-23} \text{J} \cdot \text{K}^{-1} \cdot 8.2 \times 10^4 \text{K}$$

$$\simeq 3.8 \times 10^{10} \text{Pa}.$$

Free electron gas in metal

- Fermi temperature for the copper:

$$T_F = \frac{\mu(0)}{k} = \frac{1}{k} \frac{\hbar^2}{2m} \left(3\pi^2 \frac{N}{V} \right)^{\frac{2}{3}}$$

$$= \frac{1}{k} \frac{\hbar^2}{2m} \left(3\pi^2 \cdot 8.5 \times 10^{28} \text{m}^{-3} \right)^{\frac{2}{3}} \simeq 8.2 \times 10^4 \text{K}.$$

- Internal energy: $U(0) = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{3}{2}}}{e^{\frac{\varepsilon-\mu}{kT}} + 1} d\varepsilon$

$$= \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^{\mu(0)} \varepsilon^{\frac{3}{2}} d\varepsilon = \frac{3}{5} N \mu(0).$$

- Pressure: $p = \frac{2}{3} \frac{U}{V} = \frac{2}{5} n \mu(0).$

$$p_{\text{Cu}} = \frac{2}{5} \cdot 8.5 \times 10^{28} \text{m}^{-3} \cdot 1.38 \times 10^{-23} \text{J} \cdot \text{K}^{-1} \cdot 8.2 \times 10^4 \text{K}$$

$$\simeq 3.8 \times 10^{10} \text{Pa}.$$

- Entropy: $S = 0.$

Free electron gas in metal

$$f = \frac{1}{e^{\frac{\varepsilon - \mu}{kT}} + 1}$$

- Case: $T > 0\text{K}$.

Free electron gas in metal

$$f = \frac{1}{e^{\frac{\varepsilon - \mu}{kT}} + 1}$$

- Case: $T > 0\text{K}$.


$$f \begin{cases} > \frac{1}{2}, & \varepsilon < \mu; \\ = \frac{1}{2}, & \varepsilon = \mu; \\ < \frac{1}{2}, & \varepsilon > \mu. \end{cases}$$

- Number of states:

$$\frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \varepsilon^{\frac{1}{2}} d\varepsilon.$$

- Energy density:

$$\frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \frac{\varepsilon^{\frac{3}{2}}}{e^{\frac{\varepsilon - \mu}{kT}} + 1} d\varepsilon.$$



Free electron gas in metal

- Estimate: only the electrons at $(\mu - kT, \mu + kT)$ contributes to the C_V .

Free electron gas in metal

- Estimate: only the electrons at $(\mu - kT, \mu + kT)$ contributes to the C_V .

$$N_{\text{eff}} \simeq \frac{kT}{\mu} N.$$

Free electron gas in metal

- Estimate: only the electrons at $(\mu - kT, \mu + kT)$ contributes to the C_V .

$$N_{\text{eff}} \simeq \frac{kT}{\mu} N.$$

Acts like classical free particle, so $C_V \simeq \frac{3}{2} N_{\text{eff}} k$

Free electron gas in metal

- Estimate: only the electrons at $(\mu - kT, \mu + kT)$ contributes to the C_V .

$$N_{\text{eff}} \simeq \frac{kT}{\mu} N.$$

Acts like classical free particle, so $C_V \simeq \frac{3}{2} N_{\text{eff}} k$
 $= \frac{3}{2} \frac{kT}{\mu} N k \simeq \frac{3}{2} N k \frac{T}{T_F}.$

Free electron gas in metal

- Estimate: only the electrons at $(\mu - kT, \mu + kT)$ contributes to the C_V .

$$N_{\text{eff}} \simeq \frac{kT}{\mu} N.$$

Acts like classical free particle, so $C_V \simeq \frac{3}{2} N_{\text{eff}} k$
 $= \frac{3}{2} \frac{kT}{\mu} N k \simeq \frac{3}{2} N k \frac{T}{T_F}.$

- Quantitative:

Free electron gas in metal

- Estimate: only the electrons at $(\mu - kT, \mu + kT)$ contributes to the C_V .

$$N_{\text{eff}} \simeq \frac{kT}{\mu} N.$$

Acts like classical free particle, so $C_V \simeq \frac{3}{2} N_{\text{eff}} k$
 $= \frac{3}{2} \frac{kT}{\mu} N k \simeq \frac{3}{2} N k \frac{T}{T_F}.$

- Quantitative:

$$N = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}}}{e^{\frac{\varepsilon - \mu}{kT}} + 1} d\varepsilon,$$

Free electron gas in metal

- Estimate: only the electrons at $(\mu - kT, \mu + kT)$ contributes to the C_V .

$$N_{\text{eff}} \simeq \frac{kT}{\mu} N.$$

Acts like classical free particle, so $C_V \simeq \frac{3}{2} N_{\text{eff}} k$
 $= \frac{3}{2} \frac{kT}{\mu} N k \simeq \frac{3}{2} N k \frac{T}{T_F}.$

- Quantitative:

$$N = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}}}{e^{\frac{\varepsilon-\mu}{kT}} + 1} d\varepsilon,$$

$$E = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{3}{2}}}{e^{\frac{\varepsilon-\mu}{kT}} + 1} d\varepsilon.$$

Free electron gas in metal

- Estimate: only the electrons at $(\mu - kT, \mu + kT)$ contributes to the C_V .

$$N_{\text{eff}} \simeq \frac{kT}{\mu} N.$$

Acts like classical free particle, so $C_V \simeq \frac{3}{2} N_{\text{eff}} k$
 $= \frac{3}{2} \frac{kT}{\mu} N k \simeq \frac{3}{2} N k \frac{T}{T_F}.$

- Quantitative:

$$N = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}}}{e^{\frac{\varepsilon-\mu}{kT}} + 1} d\varepsilon,$$

$$E = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{3}{2}}}{e^{\frac{\varepsilon-\mu}{kT}} + 1} d\varepsilon.$$

- Calculable in principle:

Free electron gas in metal

- Estimate: only the electrons at $(\mu - kT, \mu + kT)$ contributes to the C_V .

$$N_{\text{eff}} \simeq \frac{kT}{\mu} N.$$

Acts like classical free particle, so $C_V \simeq \frac{3}{2} N_{\text{eff}} k$
 $= \frac{3}{2} \frac{kT}{\mu} N k \simeq \frac{3}{2} N k \frac{T}{T_F}.$

- Quantitative:

$$N = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{1}{2}}}{e^{\frac{\varepsilon - \mu}{kT}} + 1} d\varepsilon,$$

$$E = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}} \int_0^\infty \frac{\varepsilon^{\frac{3}{2}}}{e^{\frac{\varepsilon - \mu}{kT}} + 1} d\varepsilon.$$

- Calculable in principle:

$$N = \frac{2}{3} C \mu^{\frac{3}{2}} \left[1 + \frac{\pi^2}{8} \left(\frac{kT}{\mu} \right)^2 \right],$$

$$E = \frac{2}{5} C \mu^{\frac{5}{2}} \left[1 + \frac{5\pi^2}{8} \left(\frac{kT}{\mu} \right)^2 \right], \text{ where } C = \frac{4\pi V}{h^3} (2m)^{\frac{3}{2}}.$$

$$N = \frac{2}{3}C\mu^{\frac{3}{2}}\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right], \quad E = \frac{2}{5}C\mu^{\frac{5}{2}}\left[1 + \frac{5\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right]$$

- As $kT \ll \mu$, from N ,

$$N = \frac{2}{3}C\mu^{\frac{3}{2}}\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right], \quad E = \frac{2}{5}C\mu^{\frac{5}{2}}\left[1 + \frac{5\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right]$$

- As $kT \ll \mu$, from N , $\mu(0)^{\frac{3}{2}} \simeq \frac{3N}{2C}$

$$N = \frac{2}{3}C\mu^{\frac{3}{2}}\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right], \quad E = \frac{2}{5}C\mu^{\frac{5}{2}}\left[1 + \frac{5\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right]$$

- As $kT \ll \mu$, from N , $\mu(0)^{\frac{3}{2}} \simeq \frac{3N}{2C} = \mu^{\frac{3}{2}}\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right]$

$$N = \frac{2}{3}C\mu^{\frac{3}{2}}\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right], \quad E = \frac{2}{5}C\mu^{\frac{5}{2}}\left[1 + \frac{5\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right]$$

- As $kT \ll \mu$, from N , $\mu(0)^{\frac{3}{2}} \simeq \frac{3N}{2C} = \mu^{\frac{3}{2}}\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right] \simeq \mu^{\frac{3}{2}}\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu(0)}\right)^2\right]$

$$N = \frac{2}{3}C\mu^{\frac{3}{2}}\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right], \quad E = \frac{2}{5}C\mu^{\frac{5}{2}}\left[1 + \frac{5\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right]$$

- As $kT \ll \mu$, from N , $\mu(0)^{\frac{3}{2}} \simeq \frac{3N}{2C} = \mu^{\frac{3}{2}}\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right]$
 $\simeq \mu^{\frac{3}{2}}\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu(0)}\right)^2\right] \Rightarrow \mu(0) \simeq \mu\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu(0)}\right)^2\right]^{\frac{2}{3}}$

$$N = \frac{2}{3}C\mu^{\frac{3}{2}}\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right], \quad E = \frac{2}{5}C\mu^{\frac{5}{2}}\left[1 + \frac{5\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right]$$

- As $kT \ll \mu$, from N , $\mu(0)^{\frac{3}{2}} \simeq \frac{3N}{2C} = \mu^{\frac{3}{2}}\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right]$
 $\simeq \mu^{\frac{3}{2}}\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu(0)}\right)^2\right] \Rightarrow \mu(0) \simeq \mu\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu(0)}\right)^2\right]^{\frac{2}{3}}$
 $\Rightarrow \mu \simeq \mu(0)\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu(0)}\right)^2\right]^{-\frac{2}{3}}$

$$N = \frac{2}{3}C\mu^{\frac{3}{2}}[1 + \frac{\pi^2}{8}(\frac{kT}{\mu})^2], \quad E = \frac{2}{5}C\mu^{\frac{5}{2}}[1 + \frac{5\pi^2}{8}(\frac{kT}{\mu})^2]$$

- As $kT \ll \mu$, from N , $\mu(0)^{\frac{3}{2}} \simeq \frac{3N}{2C} = \mu^{\frac{3}{2}}[1 + \frac{\pi^2}{8}(\frac{kT}{\mu})^2]$
 $\simeq \mu^{\frac{3}{2}}[1 + \frac{\pi^2}{8}(\frac{kT}{\mu(0)})^2] \Rightarrow \mu(0) \simeq \mu[1 + \frac{\pi^2}{8}(\frac{kT}{\mu(0)})^2]^{\frac{2}{3}}$
 $\Rightarrow \mu \simeq \mu(0)[1 + \frac{\pi^2}{8}(\frac{kT}{\mu(0)})^2]^{-\frac{2}{3}} \simeq \mu(0)[1 - \frac{\pi^2}{12}(\frac{kT}{\mu(0)})^2].$

$$N = \frac{2}{3}C\mu^{\frac{3}{2}}\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right], \quad E = \frac{2}{5}C\mu^{\frac{5}{2}}\left[1 + \frac{5\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right]$$

- As $kT \ll \mu$, from N , $\mu(0)^{\frac{3}{2}} \simeq \frac{3N}{2C} = \mu^{\frac{3}{2}}\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right]$
 $\simeq \mu^{\frac{3}{2}}\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu(0)}\right)^2\right] \Rightarrow \mu(0) \simeq \mu\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu(0)}\right)^2\right]^{\frac{2}{3}}$
 $\Rightarrow \mu \simeq \mu(0)\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu(0)}\right)^2\right]^{-\frac{2}{3}} \simeq \mu(0)\left[1 - \frac{\pi^2}{12}\left(\frac{kT}{\mu(0)}\right)^2\right].$
- $U = \frac{3}{5}N\mu \frac{1 + \frac{5\pi^2}{8}\left(\frac{kT}{\mu}\right)^2}{1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu}\right)^2}$

$$N = \frac{2}{3}C\mu^{\frac{3}{2}}\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right], \quad E = \frac{2}{5}C\mu^{\frac{5}{2}}\left[1 + \frac{5\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right]$$

- As $kT \ll \mu$, from N , $\mu(0)^{\frac{3}{2}} \simeq \frac{3N}{2C} = \mu^{\frac{3}{2}}\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right]$
 $\simeq \mu^{\frac{3}{2}}\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu(0)}\right)^2\right] \Rightarrow \mu(0) \simeq \mu\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu(0)}\right)^2\right]^{\frac{2}{3}}$
 $\Rightarrow \mu \simeq \mu(0)\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu(0)}\right)^2\right]^{-\frac{2}{3}} \simeq \mu(0)\left[1 - \frac{\pi^2}{12}\left(\frac{kT}{\mu(0)}\right)^2\right].$
- $U = \frac{3}{5}N\mu \frac{1 + \frac{5\pi^2}{8}\left(\frac{kT}{\mu}\right)^2}{1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu}\right)^2} \simeq \frac{3}{5}N\mu\left[1 + \frac{5\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right]\left[1 - \frac{\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right]$

$$N = \frac{2}{3}C\mu^{\frac{3}{2}}\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right], \quad E = \frac{2}{5}C\mu^{\frac{5}{2}}\left[1 + \frac{5\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right]$$

- As $kT \ll \mu$, from N , $\mu(0)^{\frac{3}{2}} \simeq \frac{3N}{2C} = \mu^{\frac{3}{2}}\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right]$
 $\simeq \mu^{\frac{3}{2}}\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu(0)}\right)^2\right] \Rightarrow \mu(0) \simeq \mu\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu(0)}\right)^2\right]^{\frac{2}{3}}$
 $\Rightarrow \mu \simeq \mu(0)\left[1 + \frac{\pi^2}{8}\left(\frac{kT}{\mu(0)}\right)^2\right]^{-\frac{2}{3}} \simeq \mu(0)\left[1 - \frac{\pi^2}{12}\left(\frac{kT}{\mu(0)}\right)^2\right].$
- $U = \frac{3}{5}N\mu\frac{1+\frac{5\pi^2}{8}\left(\frac{kT}{\mu}\right)^2}{1+\frac{\pi^2}{8}\left(\frac{kT}{\mu}\right)^2} \simeq \frac{3}{5}N\mu\left[1 + \frac{5\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right]\left[1 - \frac{\pi^2}{8}\left(\frac{kT}{\mu}\right)^2\right]$
 $\simeq \frac{3}{5}N\mu\left[1 + \frac{\pi^2}{2}\left(\frac{kT}{\mu}\right)^2\right]$

$$N = \frac{2}{3}C\mu^{\frac{3}{2}}[1 + \frac{\pi^2}{8}(\frac{kT}{\mu})^2], \quad E = \frac{2}{5}C\mu^{\frac{5}{2}}[1 + \frac{5\pi^2}{8}(\frac{kT}{\mu})^2]$$

- As $kT \ll \mu$, from N , $\mu(0)^{\frac{3}{2}} \simeq \frac{3N}{2C} = \mu^{\frac{3}{2}}[1 + \frac{\pi^2}{8}(\frac{kT}{\mu})^2]$
 $\simeq \mu^{\frac{3}{2}}[1 + \frac{\pi^2}{8}(\frac{kT}{\mu(0)})^2] \Rightarrow \mu(0) \simeq \mu[1 + \frac{\pi^2}{8}(\frac{kT}{\mu(0)})^2]^{\frac{2}{3}}$
 $\Rightarrow \mu \simeq \mu(0)[1 + \frac{\pi^2}{8}(\frac{kT}{\mu(0)})^2]^{-\frac{2}{3}} \simeq \mu(0)[1 - \frac{\pi^2}{12}(\frac{kT}{\mu(0)})^2].$
- $U = \frac{3}{5}N\mu \frac{1 + \frac{5\pi^2}{8}(\frac{kT}{\mu})^2}{1 + \frac{\pi^2}{8}(\frac{kT}{\mu})^2} \simeq \frac{3}{5}N\mu[1 + \frac{5\pi^2}{8}(\frac{kT}{\mu})^2][1 - \frac{\pi^2}{8}(\frac{kT}{\mu})^2]$
 $\simeq \frac{3}{5}N\mu[1 + \frac{\pi^2}{2}(\frac{kT}{\mu})^2]$
 $\simeq \frac{3}{5}N\mu(0)[1 - \frac{\pi^2}{12}(\frac{kT}{\mu(0)})^2][1 + \frac{\pi^2}{2}(\frac{kT}{\mu(0)})^2]$

$$N = \frac{2}{3}C\mu^{\frac{3}{2}}[1 + \frac{\pi^2}{8}(\frac{kT}{\mu})^2], \quad E = \frac{2}{5}C\mu^{\frac{5}{2}}[1 + \frac{5\pi^2}{8}(\frac{kT}{\mu})^2]$$

- As $kT \ll \mu$, from N , $\mu(0)^{\frac{3}{2}} \simeq \frac{3N}{2C} = \mu^{\frac{3}{2}}[1 + \frac{\pi^2}{8}(\frac{kT}{\mu})^2]$
 $\simeq \mu^{\frac{3}{2}}[1 + \frac{\pi^2}{8}(\frac{kT}{\mu(0)})^2] \Rightarrow \mu(0) \simeq \mu[1 + \frac{\pi^2}{8}(\frac{kT}{\mu(0)})^2]^{\frac{2}{3}}$
 $\Rightarrow \mu \simeq \mu(0)[1 + \frac{\pi^2}{8}(\frac{kT}{\mu(0)})^2]^{-\frac{2}{3}} \simeq \mu(0)[1 - \frac{\pi^2}{12}(\frac{kT}{\mu(0)})^2].$
- $U = \frac{3}{5}N\mu \frac{1 + \frac{5\pi^2}{8}(\frac{kT}{\mu})^2}{1 + \frac{\pi^2}{8}(\frac{kT}{\mu})^2} \simeq \frac{3}{5}N\mu[1 + \frac{5\pi^2}{8}(\frac{kT}{\mu})^2][1 - \frac{\pi^2}{8}(\frac{kT}{\mu})^2]$
 $\simeq \frac{3}{5}N\mu[1 + \frac{\pi^2}{2}(\frac{kT}{\mu})^2]$
 $\simeq \frac{3}{5}N\mu(0)[1 - \frac{\pi^2}{12}(\frac{kT}{\mu(0)})^2][1 + \frac{\pi^2}{2}(\frac{kT}{\mu(0)})^2]$
 $\simeq \frac{3}{5}N\mu(0)[1 + \frac{5\pi^2}{12}(\frac{kT}{\mu(0)})^2].$

$$N = \frac{2}{3}C\mu^{\frac{3}{2}}[1 + \frac{\pi^2}{8}(\frac{kT}{\mu})^2], \quad E = \frac{2}{5}C\mu^{\frac{5}{2}}[1 + \frac{5\pi^2}{8}(\frac{kT}{\mu})^2]$$

- As $kT \ll \mu$, from N , $\mu(0)^{\frac{3}{2}} \simeq \frac{3N}{2C} = \mu^{\frac{3}{2}}[1 + \frac{\pi^2}{8}(\frac{kT}{\mu})^2]$
 $\simeq \mu^{\frac{3}{2}}[1 + \frac{\pi^2}{8}(\frac{kT}{\mu(0)})^2] \Rightarrow \mu(0) \simeq \mu[1 + \frac{\pi^2}{8}(\frac{kT}{\mu(0)})^2]^{\frac{2}{3}}$
 $\Rightarrow \mu \simeq \mu(0)[1 + \frac{\pi^2}{8}(\frac{kT}{\mu(0)})^2]^{-\frac{2}{3}} \simeq \mu(0)[1 - \frac{\pi^2}{12}(\frac{kT}{\mu(0)})^2].$
- $U = \frac{3}{5}N\mu \frac{1 + \frac{5\pi^2}{8}(\frac{kT}{\mu})^2}{1 + \frac{\pi^2}{8}(\frac{kT}{\mu})^2} \simeq \frac{3}{5}N\mu[1 + \frac{5\pi^2}{8}(\frac{kT}{\mu})^2][1 - \frac{\pi^2}{8}(\frac{kT}{\mu})^2]$
 $\simeq \frac{3}{5}N\mu[1 + \frac{\pi^2}{2}(\frac{kT}{\mu})^2]$
 $\simeq \frac{3}{5}N\mu(0)[1 - \frac{\pi^2}{12}(\frac{kT}{\mu(0)})^2][1 + \frac{\pi^2}{2}(\frac{kT}{\mu(0)})^2]$
 $\simeq \frac{3}{5}N\mu(0)[1 + \frac{5\pi^2}{12}(\frac{kT}{\mu(0)})^2].$
- $\therefore C_V = (\frac{\partial U}{\partial T})_V = Nk \frac{\pi^2 k}{2\mu(0)} T = \frac{\pi^2}{2} Nk \frac{T}{T_F} \propto T.$

Table of contents

1 §8. Bose statistics and Fermi statistics

- 8.1 Thermal parameters in statistics
- 8.2 U of weak degenerated ideal Fermion(Boson) gas
- 8.3 Bose-Einstein Condensate (BEC)
- 8.4 Photon gas
- 8.5 Free electron gas in metal