

# Thermodynamics & Statistical Physics

## Exercises

Yuan-Chuan Zou  
zouyc@hust.edu.cn

School of Physics, Huazhong University of Science and Technology

December 30, 2013

# Table of contents

- ① §6. Most probable distribution of nearly independent particles
- ② §7. Boltzmann statistics
- ③ §8. Bose statistics and Fermi statistics
  - Bose statistics
  - Fermi statistics
  - Fermion gas in astrophysics

A10.41 Discuss: if the nearly independent ideal gas has no degenerated levels, there is no most probable distribution.

A10.41 Discuss: if the nearly independent ideal gas has no degenerated levels, there is no most probable distribution.

- For a given distribution  $(\varepsilon_l, \omega_l, a_l)$ ,

A10.41 Discuss: if the nearly independent ideal gas has no degenerated levels, there is no most probable distribution.

- For a given distribution  $(\varepsilon_l, \omega_l, a_l)$ ,
- for Boson gas, the number of micro-states (6.5.6):

A10.41 Discuss: if the nearly independent ideal gas has no degenerated levels, there is no most probable distribution.

- For a given distribution  $(\varepsilon_l, \omega_l, a_l)$ ,
- for Boson gas, the number of micro-states (6.5.6):

$$\Omega_{\text{B.S.}} = \prod \frac{(\omega_l + a_l - 1)!}{a_l! (\omega_l - 1)!}$$

A10.41 Discuss: if the nearly independent ideal gas has no degenerated levels, there is no most probable distribution.

- For a given distribution  $(\varepsilon_l, \omega_l, a_l)$ ,
- for Boson gas, the number of micro-states (6.5.6):

$$\Omega_{\text{B.S.}} = \prod \frac{(\omega_l + a_l - 1)!}{a_l! (\omega_l - 1)!} = \prod \frac{a_l!}{a_l!}$$

A10.41 Discuss: if the nearly independent ideal gas has no degenerated levels, there is no most probable distribution.

- For a given distribution  $(\varepsilon_l, \omega_l, a_l)$ ,
- for Boson gas, the number of micro-states (6.5.6):

$$\Omega_{\text{B.S.}} = \prod \frac{(\omega_l + a_l - 1)!}{a_l! (\omega_l - 1)!} = \prod \frac{a_l!}{a_l!} = 1;$$



A10.41 Discuss: if the nearly independent ideal gas has no degenerated levels, there is no most probable distribution.

- For a given distribution  $(\varepsilon_l, \omega_l, a_l)$ ,
- for Boson gas, the number of micro-states (6.5.6):
$$\Omega_{\text{B.S.}} = \prod \frac{(\omega_l + a_l - 1)!}{a_l! (\omega_l - 1)!} = \prod \frac{a_l!}{a_l!} = 1;$$
- for Fermion gas, the number of micro-states (6.5.7):

A10.41 Discuss: if the nearly independent ideal gas has no degenerated levels, there is no most probable distribution.

- For a given distribution  $(\varepsilon_l, \omega_l, a_l)$ ,
- for Boson gas, the number of micro-states (6.5.6):  

$$\Omega_{\text{B.S.}} = \prod \frac{(\omega_l + a_l - 1)!}{a_l! (\omega_l - 1)!} = \prod \frac{a_l!}{a_l!} = 1;$$
- for Fermion gas, the number of micro-states (6.5.7):  

$$\Omega_{\text{F.D.}} = \prod \frac{\omega_l!}{a_l! (\omega_l - a_l)!}$$

A10.41 Discuss: if the nearly independent ideal gas has no degenerated levels, there is no most probable distribution.

- For a given distribution  $(\varepsilon_l, \omega_l, a_l)$ ,
- for Boson gas, the number of micro-states (6.5.6):  

$$\Omega_{\text{B.S.}} = \prod \frac{(\omega_l + a_l - 1)!}{a_l! (\omega_l - 1)!} = \prod \frac{a_l!}{a_l!} = 1;$$
- for Fermion gas, the number of micro-states (6.5.7):  

$$\Omega_{\text{F.D.}} = \prod \frac{\omega_l!}{a_l! (\omega_l - a_l)!} = \prod \frac{1}{a_l! (1 - a_l)!}$$

A10.41 Discuss: if the nearly independent ideal gas has no degenerated levels, there is no most probable distribution.

- For a given distribution  $(\varepsilon_l, \omega_l, a_l)$ ,
- for Boson gas, the number of micro-states (6.5.6):  

$$\Omega_{\text{B.S.}} = \prod \frac{(\omega_l + a_l - 1)!}{a_l! (\omega_l - 1)!} = \prod \frac{a_l!}{a_l!} = 1;$$
- for Fermion gas, the number of micro-states (6.5.7):  

$$\Omega_{\text{F.D.}} = \prod \frac{\omega_l!}{a_l! (\omega_l - a_l)!} = \prod \frac{1}{a_l! (1 - a_l)!} = 1;$$

A10.41 Discuss: if the nearly independent ideal gas has no degenerated levels, there is no most probable distribution.

- For a given distribution  $(\varepsilon_l, \omega_l, a_l)$ ,
- for Boson gas, the number of micro-states (6.5.6):  

$$\Omega_{\text{B.S.}} = \prod \frac{(\omega_l + a_l - 1)!}{a_l! (\omega_l - 1)!} = \prod \frac{a_l!}{a_l!} = 1;$$
- for Fermion gas, the number of micro-states (6.5.7):  

$$\Omega_{\text{F.D.}} = \prod \frac{\omega_l!}{a_l! (\omega_l - a_l)!} = \prod \frac{1}{a_l! (1 - a_l)!} = 1;$$
- There is no maximum for the number of micro-states.

# Table of contents

- 1 §6. Most probable distribution of nearly independent particles
- 2 §7. Boltzmann statistics
- 3 §8. Bose statistics and Fermi statistics
  - Bose statistics
  - Fermi statistics
  - Fermion gas in astrophysics

A8.2 For classical ideal gas, if the total number of molecules is  $N$ , the number with speed less than the most probable speed  $v_m$  is  $N_m$ , calculate  $\frac{N_m}{N}$ .

A8.2 For classical ideal gas, if the total number of molecules is  $N$ , the number with speed less than the most probable speed  $v_m$  is  $N_m$ , calculate  $\frac{N_m}{N}$ .

- According to the Maxwell's speed distribution, the number of molecules at  $(v, v + dv)$ :



A8.2 For classical ideal gas, if the total number of molecules is  $N$ , the number with speed less than the most probable speed  $v_m$  is  $N_m$ , calculate  $\frac{N_m}{N}$ .

- According to the Maxwell's speed distribution, the number of molecules at  $(v, v + dv)$ :

$$dN(v) = 4\pi N \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2 dv.$$

A8.2 For classical ideal gas, if the total number of molecules is  $N$ , the number with speed less than the most probable speed  $v_m$  is  $N_m$ , calculate  $\frac{N_m}{N}$ .

- According to the Maxwell's speed distribution, the number of molecules at  $(v, v + dv)$ :

$$dN(v) = 4\pi N \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2 dv.$$

The most probable speed:  $v_m = \sqrt{\frac{2kT}{m}}.$

A8.2 For classical ideal gas, if the total number of molecules is  $N$ , the number with speed less than the most probable speed  $v_m$  is  $N_m$ , calculate  $\frac{N_m}{N}$ .

- According to the Maxwell's speed distribution, the number of molecules at  $(v, v + dv)$ :

$$dN(v) = 4\pi N \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2 dv.$$

The most probable speed:  $v_m = \sqrt{\frac{2kT}{m}}$ .

$$\frac{N_m}{N} = \int_0^{v_m} 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2 dv$$

A8.2 For classical ideal gas, if the total number of molecules is  $N$ , the number with speed less than the most probable speed  $v_m$  is  $N_m$ , calculate  $\frac{N_m}{N}$ .

- According to the Maxwell's speed distribution, the number of molecules at  $(v, v + dv)$ :

$$dN(v) = 4\pi N \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2 dv.$$

The most probable speed:  $v_m = \sqrt{\frac{2kT}{m}}$ .

$$\begin{aligned} \frac{N_m}{N} &= \int_0^{v_m} 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2 dv \\ &= \int_0^{(\sqrt{\frac{m}{2kT}})} \sqrt{\frac{2kT}{m}} \frac{4}{\sqrt{\pi}} e^{-\frac{m}{2kT}v^2} \left(\sqrt{\frac{m}{2kT}}v\right)^2 d\left(\sqrt{\frac{m}{2kT}}v\right) \end{aligned}$$

A8.2 For classical ideal gas, if the total number of molecules is  $N$ , the number with speed less than the most probable speed  $v_m$  is  $N_m$ , calculate  $\frac{N_m}{N}$ .

- According to the Maxwell's speed distribution, the number of molecules at  $(v, v + dv)$ :

$$dN(v) = 4\pi N \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2 dv.$$

The most probable speed:  $v_m = \sqrt{\frac{2kT}{m}}$ .

$$\begin{aligned} \frac{N_m}{N} &= \int_0^{v_m} 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2 dv \\ &= \int_0^{(\sqrt{\frac{m}{2kT}})\sqrt{\frac{2kT}{m}}} \frac{4}{\sqrt{\pi}} e^{-\frac{m}{2kT}v^2} \left(\sqrt{\frac{m}{2kT}}v\right)^2 d\left(\sqrt{\frac{m}{2kT}}v\right) \\ &= \frac{4}{\sqrt{\pi}} \int_0^1 x^2 e^{-x^2} dx \end{aligned}$$

A8.2 For classical ideal gas, if the total number of molecules is  $N$ , the number with speed less than the most probable speed  $v_m$  is  $N_m$ , calculate  $\frac{N_m}{N}$ .

- According to the Maxwell's speed distribution, the number of molecules at  $(v, v + dv)$ :

$$dN(v) = 4\pi N \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2 dv.$$

The most probable speed:  $v_m = \sqrt{\frac{2kT}{m}}$ .

$$\begin{aligned} \frac{N_m}{N} &= \int_0^{v_m} 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2 dv \\ &= \int_0^{(\sqrt{\frac{m}{2kT}})\sqrt{\frac{2kT}{m}}} \frac{4}{\sqrt{\pi}} e^{-\frac{m}{2kT}v^2} \left(\sqrt{\frac{m}{2kT}}v\right)^2 d\left(\sqrt{\frac{m}{2kT}}v\right) \\ &= \frac{4}{\sqrt{\pi}} \int_0^1 x^2 e^{-x^2} dx \\ &\simeq 0.4276; \end{aligned}$$

A8.2 For classical ideal gas, if the total number of molecules is  $N$ , the number with speed less than the most probable speed  $v_m$  is  $N_m$ , calculate  $\frac{N_m}{N}$ .

- According to the Maxwell's speed distribution, the number of molecules at  $(v, v + dv)$ :

$$dN(v) = 4\pi N \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2 dv.$$

The most probable speed:  $v_m = \sqrt{\frac{2kT}{m}}$ .

$$\begin{aligned} \frac{N_m}{N} &= \int_0^{v_m} 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2 dv \\ &= \int_0^{(\sqrt{\frac{m}{2kT}})\sqrt{\frac{2kT}{m}}} \frac{4}{\sqrt{\pi}} e^{-\frac{m}{2kT}v^2} \left(\sqrt{\frac{m}{2kT}}v\right)^2 d\left(\sqrt{\frac{m}{2kT}}v\right) \\ &= \frac{4}{\sqrt{\pi}} \int_0^1 x^2 e^{-x^2} dx \\ &\simeq 0.4276; \end{aligned}$$

Not depending on  $T$ .

A8.21 Prove: for monatomic classical ideal gas, entropy:

$$S = \frac{5}{2}Nk - Nk \ln(n\lambda^3).$$



A8.21 Prove: for monatomic classical ideal gas, entropy:  
$$S = \frac{5}{2}Nk - Nk \ln(n\lambda^3).$$

- Partition function:  $Z_1 = V(\frac{2\pi m}{h^2\beta})^{3/2}. \quad (7.5.21)$

A8.21 Prove: for monatomic classical ideal gas, entropy:  

$$S = \frac{5}{2}Nk - Nk \ln(n\lambda^3).$$

- Partition function:  $Z_1 = V(\frac{2\pi m}{h^2\beta})^{3/2}$ . (7.5.21)

Entropy:

$$S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) - k \ln N! \quad (7.1.13')$$

A8.21 Prove: for monatomic classical ideal gas, entropy:  
 $S = \frac{5}{2}Nk - Nk \ln(n\lambda^3).$

- Partition function:  $Z_1 = V(\frac{2\pi m}{h^2\beta})^{3/2}. \quad (7.5.21)$

Entropy:

$$S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) - k \ln N! \quad (7.1.13')$$

$$\simeq Nk\{\ln[V(\frac{2\pi m}{h^2\beta})^{\frac{3}{2}}] - \beta \frac{\partial}{\partial \beta} \ln[V(\frac{2\pi m}{h^2\beta})^{\frac{3}{2}}]\} - kN(\ln N - 1)$$

A8.21 Prove: for monatomic classical ideal gas, entropy:  
 $S = \frac{5}{2}Nk - Nk \ln(n\lambda^3).$

- Partition function:  $Z_1 = V(\frac{2\pi m}{h^2\beta})^{3/2}$ . (7.5.21)

Entropy:

$$\begin{aligned} S &= Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) - k \ln N! \quad (7.1.13') \\ &\simeq Nk\{\ln[V(\frac{2\pi m}{h^2\beta})^{3/2}] - \beta \frac{\partial}{\partial \beta} \ln[V(\frac{2\pi m}{h^2\beta})^{3/2}]\} - kN(\ln N - 1) \\ &= Nk\{\ln[V(\frac{2\pi mkT}{h^2})^{3/2}] - \beta \frac{\partial}{\partial \beta} \ln \beta^{-3/2}\} - (\ln N - 1) \end{aligned}$$

A8.21 Prove: for monatomic classical ideal gas, entropy:  
 $S = \frac{5}{2}Nk - Nk \ln(n\lambda^3).$

- Partition function:  $Z_1 = V(\frac{2\pi m}{h^2\beta})^{3/2}. \quad (7.5.21)$

Entropy:

$$\begin{aligned}
 S &= Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) - k \ln N! \quad (7.1.13') \\
 &\simeq Nk\{\ln[V(\frac{2\pi m}{h^2\beta})^{\frac{3}{2}}] - \beta \frac{\partial}{\partial \beta} \ln[V(\frac{2\pi m}{h^2\beta})^{\frac{3}{2}}]\} - kN(\ln N - 1) \\
 &= Nk\{\ln[V(\frac{2\pi mkT}{h^2})^{3/2}] - \beta \frac{\partial}{\partial \beta} \ln \beta^{-3/2}] - (\ln N - 1)\} \\
 &= Nk\{\ln[V(\frac{2\pi mkT}{h^2})^{3/2}] + \frac{3}{2} - (\ln N - 1)\}
 \end{aligned}$$

A8.21 Prove: for monatomic classical ideal gas, entropy:

$$S = \frac{5}{2}Nk - Nk \ln(n\lambda^3).$$

- Partition function:  $Z_1 = V(\frac{2\pi m}{h^2\beta})^{3/2}$ . (7.5.21)

Entropy:

$$\begin{aligned} S &= Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) - k \ln N! \quad (7.1.13') \\ &\simeq Nk\{\ln[V(\frac{2\pi m}{h^2\beta})^{3/2}] - \beta \frac{\partial}{\partial \beta} \ln[V(\frac{2\pi m}{h^2\beta})^{3/2}]\} - kN(\ln N - 1) \\ &= Nk\{\ln[V(\frac{2\pi mkT}{h^2})^{3/2}] - \beta \frac{\partial}{\partial \beta} \ln \beta^{-3/2}\} - (\ln N - 1) \\ &= Nk\{\ln[V(\frac{2\pi mkT}{h^2})^{3/2}] + \frac{3}{2} - (\ln N - 1)\} \\ &= Nk\{\ln[\frac{V}{N}(\frac{2\pi mkT}{h^2})^{3/2}] + \frac{5}{2}\} \end{aligned}$$

A8.21 Prove: for monatomic classical ideal gas, entropy:

$$S = \frac{5}{2}Nk - Nk \ln(n\lambda^3).$$

- Partition function:  $Z_1 = V(\frac{2\pi m}{h^2\beta})^{3/2}$ . (7.5.21)

Entropy:

$$\begin{aligned} S &= Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) - k \ln N! \quad (7.1.13') \\ &\simeq Nk\{\ln[V(\frac{2\pi m}{h^2\beta})^{3/2}] - \beta \frac{\partial}{\partial \beta} \ln[V(\frac{2\pi m}{h^2\beta})^{3/2}]\} - kN(\ln N - 1) \\ &= Nk\{\ln[V(\frac{2\pi mkT}{h^2})^{3/2}] - \beta \frac{\partial}{\partial \beta} \ln \beta^{-3/2}\} - (\ln N - 1) \\ &= Nk\{\ln[V(\frac{2\pi mkT}{h^2})^{3/2}] + \frac{3}{2} - (\ln N - 1)\} \\ &= Nk\{\ln[\frac{V}{N}(\frac{2\pi mkT}{h^2})^{3/2}] + \frac{5}{2}\} \\ &= Nk\{-\ln(n\lambda^3) + \frac{5}{2}\} \quad (\text{before (7.2.7)}) \end{aligned}$$

A8.21 Prove: for monatomic classical ideal gas, entropy:

$$S = \frac{5}{2}Nk - Nk \ln(n\lambda^3).$$

- Partition function:  $Z_1 = V(\frac{2\pi m}{h^2\beta})^{3/2}$ . (7.5.21)

Entropy:

$$\begin{aligned} S &= Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) - k \ln N! \quad (7.1.13') \\ &\simeq Nk\{\ln[V(\frac{2\pi m}{h^2\beta})^{3/2}] - \beta \frac{\partial}{\partial \beta} \ln[V(\frac{2\pi m}{h^2\beta})^{3/2}]\} - kN(\ln N - 1) \\ &= Nk\{\ln[V(\frac{2\pi mkT}{h^2})^{3/2}] - \beta \frac{\partial}{\partial \beta} \ln \beta^{-3/2}\} - (\ln N - 1) \\ &= Nk\{\ln[V(\frac{2\pi mkT}{h^2})^{3/2}] + \frac{3}{2} - (\ln N - 1)\} \\ &= Nk\{\ln[\frac{V}{N}(\frac{2\pi mkT}{h^2})^{3/2}] + \frac{5}{2}\} \\ &= Nk\{-\ln(n\lambda^3) + \frac{5}{2}\} \quad (\text{before (7.2.7)}) \\ &= \frac{5}{2}Nk - Nk \ln(n\lambda^3); \end{aligned}$$



A8.21 Prove: for monatomic classical ideal gas, entropy:

$$S = \frac{5}{2}Nk - Nk \ln(n\lambda^3).$$

- Partition function:  $Z_1 = V(\frac{2\pi m}{h^2\beta})^{3/2}$ . (7.5.21)

Entropy:

$$S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) - k \ln N! \quad (7.1.13')$$

$$\simeq Nk\{\ln[V(\frac{2\pi m}{h^2\beta})^{3/2}] - \beta \frac{\partial}{\partial \beta} \ln[V(\frac{2\pi m}{h^2\beta})^{3/2}]\} - kN(\ln N - 1)$$

$$= Nk\{\ln[V(\frac{2\pi mkT}{h^2})^{3/2}] - \beta \frac{\partial}{\partial \beta} \ln \beta^{-3/2}\} - (\ln N - 1)\}$$

$$= Nk\{\ln[V(\frac{2\pi mkT}{h^2})^{3/2}] + \frac{3}{2} - (\ln N - 1)\}$$

$$= Nk\{\ln[\frac{V}{N}(\frac{2\pi mkT}{h^2})^{3/2}] + \frac{5}{2}\}$$

$$= Nk\{-\ln(n\lambda^3) + \frac{5}{2}\} \quad (\text{before (7.2.7)})$$

$$= \frac{5}{2}Nk - Nk \ln(n\lambda^3);$$

- $S = \frac{3}{2}Nk \ln T + Nk \ln \frac{V}{N} + \frac{3}{2}Nk[\frac{5}{2} + \ln \frac{2\pi mk}{h^2}]. \quad (7.6.2)$

A8.26 Monatomic classical ideal gas, calculate the probability for atoms at  $(\varepsilon, \varepsilon + d\varepsilon)$ ,  $\rho(\varepsilon)$ .

A8.26 Monatomic classical ideal gas, calculate the probability for atoms at  $(\varepsilon, \varepsilon + d\varepsilon)$ ,  $\rho(\varepsilon)$ .

- The Maxwell's speed distribution, the number of molecules at  $(v, v + dv)$ :

A8.26 Monatomic classical ideal gas, calculate the probability for atoms at  $(\varepsilon, \varepsilon + d\varepsilon)$ ,  $\rho(\varepsilon)$ .

- The Maxwell's speed distribution, the number of molecules at  $(v, v + dv)$ :

$$\rho(v)dv = \frac{dN(v)}{N} = 4\pi\left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2 dv.$$

A8.26 Monatomic classical ideal gas, calculate the probability for atoms at  $(\varepsilon, \varepsilon + d\varepsilon)$ ,  $\rho(\varepsilon)$ .

- The Maxwell's speed distribution, the number of molecules at  $(v, v + dv)$ :

$$\rho(v)dv = \frac{dN(v)}{N} = 4\pi\left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2 dv.$$

Convert to energy ( $v = \sqrt{\frac{2\varepsilon}{m}}$ ):

A8.26 Monatomic classical ideal gas, calculate the probability for atoms at  $(\varepsilon, \varepsilon + d\varepsilon)$ ,  $\rho(\varepsilon)$ .

- The Maxwell's speed distribution, the number of molecules at  $(v, v + dv)$ :

$$\rho(v)dv = \frac{dN(v)}{N} = 4\pi\left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2 dv.$$

Convert to energy ( $v = \sqrt{\frac{2\varepsilon}{m}}$ ):

$$\rho(\varepsilon)d\varepsilon = 4\pi\left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}\frac{2\varepsilon}{m}} \frac{2\varepsilon}{m} d\sqrt{\frac{2\varepsilon}{m}}$$

A8.26 Monatomic classical ideal gas, calculate the probability for atoms at  $(\varepsilon, \varepsilon + d\varepsilon)$ ,  $\rho(\varepsilon)$ .

- The Maxwell's speed distribution, the number of molecules at  $(v, v + dv)$ :

$$\rho(v)dv = \frac{dN(v)}{N} = 4\pi\left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2 dv.$$

Convert to energy ( $v = \sqrt{\frac{2\varepsilon}{m}}$ ):

$$\begin{aligned} \rho(\varepsilon)d\varepsilon &= 4\pi\left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}\frac{2\varepsilon}{m}} \frac{2\varepsilon}{m} d\sqrt{\frac{2\varepsilon}{m}} \\ &= \frac{2}{\sqrt{\pi}}(kT)^{-\frac{3}{2}} \varepsilon^{\frac{1}{2}} e^{-\frac{\varepsilon}{kT}} d\varepsilon; \end{aligned}$$

## A8.26 Monatomic classical ideal gas, calculate the probability for atoms at $(\varepsilon, \varepsilon + d\varepsilon)$ , $\rho(\varepsilon)$ .

- The Maxwell's speed distribution, the number of molecules at  $(v, v + dv)$ :

$$\rho(v)dv = \frac{dN(v)}{N} = 4\pi\left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2 dv.$$

Convert to energy ( $v = \sqrt{\frac{2\varepsilon}{m}}$ ):

$$\begin{aligned} \rho(\varepsilon)d\varepsilon &= 4\pi\left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}\frac{2\varepsilon}{m}} \frac{2\varepsilon}{m} d\sqrt{\frac{2\varepsilon}{m}} \\ &= \frac{2}{\sqrt{\pi}} (kT)^{-\frac{3}{2}} \varepsilon^{\frac{1}{2}} e^{-\frac{\varepsilon}{kT}} d\varepsilon; \end{aligned}$$

- Compare with the black-body:

$$\rho(\omega)d\omega = \frac{V}{\pi^2 c^3} \frac{\omega^2}{e^{\frac{\hbar\omega}{kT}} - 1} d\omega. \quad (8.4.6)$$



A8.33 Atomic hydrogen in a stellar atmosphere, with average kinetic energy 1.0 eV. Calculate, 1. Temperature of the stellar atmosphere; 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

A8.33 Atomic hydrogen in a stellar atmosphere, with average kinetic energy 1.0 eV. Calculate, 1. Temperature of the stellar atmosphere; 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

- $n\lambda^3 = n\left(\frac{h^2}{2\pi mkT}\right)^{\frac{3}{2}}. \quad n = ?$

A8.33 Atomic hydrogen in a stellar atmosphere, with average kinetic energy 1.0 eV. Calculate, 1. Temperature of the stellar atmosphere; 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

- $n\lambda^3 = n\left(\frac{h^2}{2\pi mkT}\right)^{\frac{3}{2}}. \quad n = ?$

Solar photosphere density  $2 \times 10^{-4} \text{kg m}^{-3}$  (wikipedia);

A8.33 Atomic hydrogen in a stellar atmosphere, with average kinetic energy 1.0 eV. Calculate, 1. Temperature of the stellar atmosphere; 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

- $n\lambda^3 = n\left(\frac{h^2}{2\pi mkT}\right)^{\frac{3}{2}}. \quad n = ?$

Solar photosphere density  $2 \times 10^{-4} \text{kg m}^{-3}$  (wikipedia);  
 number density  $n \simeq \frac{2 \times 10^{-4} \text{kg m}^{-3}}{1.67 \times 10^{-27} \text{kg}} \simeq 10^{23} \text{m}^{-3}.$

A8.33 Atomic hydrogen in a stellar atmosphere, with average kinetic energy 1.0 eV. Calculate, 1. Temperature of the stellar atmosphere; 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

- $n\lambda^3 = n\left(\frac{h^2}{2\pi mkT}\right)^{\frac{3}{2}}. \quad n = ?$

Solar photosphere density  $2 \times 10^{-4} \text{kg m}^{-3}$  (wikipedia);

number density  $n \simeq \frac{2 \times 10^{-4} \text{kg m}^{-3}}{1.67 \times 10^{-27} \text{kg}} \simeq 10^{23} \text{m}^{-3}.$

$$n\lambda^3 \simeq 10^{23} \text{m}^{-3} \left[ \frac{(6.626 \times 10^{-34} \text{J}\cdot\text{s})^2}{2\pi \cdot 1.67 \times 10^{-27} \text{kg} \cdot k \cdot \frac{1 \text{eV}}{k}} \right]^{3/2}$$

A8.33 Atomic hydrogen in a stellar atmosphere, with average kinetic energy 1.0 eV. Calculate, 1. Temperature of the stellar atmosphere; 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

- $n\lambda^3 = n\left(\frac{h^2}{2\pi mkT}\right)^{\frac{3}{2}} \cdot n = ?$

Solar photosphere density  $2 \times 10^{-4} \text{ kg m}^{-3}$  (wikipedia);

number density  $n \simeq \frac{2 \times 10^{-4} \text{ kg m}^{-3}}{1.67 \times 10^{-27} \text{ kg}} \simeq 10^{23} \text{ m}^{-3}$ .

$$n\lambda^3 \simeq 10^{23} \text{ m}^{-3} \left[ \frac{(6.626 \times 10^{-34} \text{ J}\cdot\text{s})^2}{2\pi \cdot 1.67 \times 10^{-27} \text{ kg} \cdot k \cdot \frac{1 \text{ eV}}{k}} \right]^{3/2}$$

$$\simeq 10^{23} \text{ m}^{-3} \left[ \frac{(6.626 \times 10^{-34} \text{ J}\cdot\text{s})^2}{2\pi \cdot 1.67 \times 10^{-27} \text{ kg} \cdot 1.6 \times 10^{-19} \text{ J}} \right]^{3/2}$$

A8.33 Atomic hydrogen in a stellar atmosphere, with average kinetic energy 1.0 eV. Calculate, 1. Temperature of the stellar atmosphere; 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

- $n\lambda^3 = n\left(\frac{h^2}{2\pi mkT}\right)^{\frac{3}{2}} \cdot n = ?$

Solar photosphere density  $2 \times 10^{-4} \text{ kg m}^{-3}$  (wikipedia);

number density  $n \simeq \frac{2 \times 10^{-4} \text{ kg m}^{-3}}{1.67 \times 10^{-27} \text{ kg}} \simeq 10^{23} \text{ m}^{-3}$ .

$$n\lambda^3 \simeq 10^{23} \text{ m}^{-3} \left[ \frac{(6.626 \times 10^{-34} \text{ J}\cdot\text{s})^2}{2\pi \cdot 1.67 \times 10^{-27} \text{ kg} \cdot k \cdot \frac{1 \text{ eV}}{k}} \right]^{3/2}$$

$$\simeq 10^{23} \text{ m}^{-3} \left[ \frac{(6.626 \times 10^{-34} \text{ J}\cdot\text{s})^2}{2\pi \cdot 1.67 \times 10^{-27} \text{ kg} \cdot 1.6 \times 10^{-19} \text{ J}} \right]^{3/2} \simeq 4.2 \times 10^{-10}$$

A8.33 Atomic hydrogen in a stellar atmosphere, with average kinetic energy 1.0 eV. Calculate, 1. Temperature of the stellar atmosphere; 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

- $n\lambda^3 = n\left(\frac{h^2}{2\pi mkT}\right)^{\frac{3}{2}} \cdot n = ?$

Solar photosphere density  $2 \times 10^{-4} \text{ kg m}^{-3}$  (wikipedia);

number density  $n \simeq \frac{2 \times 10^{-4} \text{ kg m}^{-3}}{1.67 \times 10^{-27} \text{ kg}} \simeq 10^{23} \text{ m}^{-3}$ .

$$n\lambda^3 \simeq 10^{23} \text{ m}^{-3} \left[ \frac{(6.626 \times 10^{-34} \text{ J}\cdot\text{s})^2}{2\pi \cdot 1.67 \times 10^{-27} \text{ kg} \cdot k \cdot \frac{1 \text{ eV}}{k}} \right]^{3/2}$$

$$\simeq 10^{23} \text{ m}^{-3} \left[ \frac{(6.626 \times 10^{-34} \text{ J}\cdot\text{s})^2}{2\pi \cdot 1.67 \times 10^{-27} \text{ kg} \cdot 1.6 \times 10^{-19} \text{ J}} \right]^{3/2} \simeq 4.2 \times 10^{-10}$$

$\ll 1$ , Boltzmann statistics.



A8.33 Atomic hydrogen in a stellar atmosphere, with average kinetic energy 1.0 eV. Calculate, 1. Temperature of the stellar atmosphere; 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

- $n\lambda^3 = n\left(\frac{h^2}{2\pi mkT}\right)^{\frac{3}{2}}. \quad n = ?$

Solar photosphere density  $2 \times 10^{-4} \text{ kg m}^{-3}$  (wikipedia);

number density  $n \simeq \frac{2 \times 10^{-4} \text{ kg m}^{-3}}{1.67 \times 10^{-27} \text{ kg}} \simeq 10^{23} \text{ m}^{-3}.$

$$n\lambda^3 \simeq 10^{23} \text{ m}^{-3} \left[ \frac{(6.626 \times 10^{-34} \text{ J}\cdot\text{s})^2}{2\pi \cdot 1.67 \times 10^{-27} \text{ kg} \cdot k \cdot \frac{1 \text{ eV}}{k}} \right]^{3/2}$$

$$\simeq 10^{23} \text{ m}^{-3} \left[ \frac{(6.626 \times 10^{-34} \text{ J}\cdot\text{s})^2}{2\pi \cdot 1.67 \times 10^{-27} \text{ kg} \cdot 1.6 \times 10^{-19} \text{ J}} \right]^{3/2} \simeq 4.2 \times 10^{-10}$$

$\ll 1$ , Boltzmann statistics.

- 1.  $\frac{3}{2}kT = \bar{\varepsilon} = 1 \text{ eV};$

A8.33 Atomic hydrogen in a stellar atmosphere, with average kinetic energy 1.0 eV. Calculate, 1. Temperature of the stellar atmosphere; 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

- $n\lambda^3 = n\left(\frac{h^2}{2\pi mkT}\right)^{\frac{3}{2}}. \quad n = ?$

Solar photosphere density  $2 \times 10^{-4} \text{kg m}^{-3}$  (wikipedia);

number density  $n \simeq \frac{2 \times 10^{-4} \text{kg m}^{-3}}{1.67 \times 10^{-27} \text{kg}} \simeq 10^{23} \text{m}^{-3}.$

$$n\lambda^3 \simeq 10^{23} \text{m}^{-3} \left[ \frac{(6.626 \times 10^{-34} \text{J}\cdot\text{s})^2}{2\pi \cdot 1.67 \times 10^{-27} \text{kg} \cdot k \cdot \frac{1 \text{eV}}{k}} \right]^{3/2}$$

$$\simeq 10^{23} \text{m}^{-3} \left[ \frac{(6.626 \times 10^{-34} \text{J}\cdot\text{s})^2}{2\pi \cdot 1.67 \times 10^{-27} \text{kg} \cdot 1.6 \times 10^{-19} \text{J}} \right]^{3/2} \simeq 4.2 \times 10^{-10} \\ \ll 1, \text{ Boltzmann statistics.}$$

- 1.  $\frac{3}{2}kT = \bar{\varepsilon} = 1 \text{eV}; \Rightarrow T \simeq 7.7 \times 10^3 \text{K}.$

... 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

- 2. Energy level:  $\varepsilon_n = -\frac{13.6}{n^2}\text{eV}$ ;

... 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

- 2. Energy level:  $\varepsilon_n = -\frac{13.6}{n^2}\text{eV}$ ;  
Boltzmann distribution:  $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$ ,

... 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

- 2. Energy level:  $\varepsilon_n = -\frac{13.6}{n^2}\text{eV}$ ;  
Boltzmann distribution:  $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$ , so,  
$$\frac{N_3}{N_1} = \frac{a_3}{a_1} = 3^2 \cdot e^{\frac{\varepsilon_1 - \varepsilon_3}{kT}}$$

... 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

- 2. Energy level:  $\varepsilon_n = -\frac{13.6}{n^2}\text{eV}$ ;  
 Boltzmann distribution:  $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$ , so,  

$$\frac{N_3}{N_1} = \frac{a_3}{a_1} = 3^2 \cdot e^{\frac{\varepsilon_1 - \varepsilon_3}{kT}} = 3^2 e^{\frac{-1.36\text{eV} \cdot 8/9}{2/3\text{eV}}}$$

... 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

- 2. Energy level:  $\varepsilon_n = -\frac{13.6}{n^2}\text{eV}$ ;

Boltzmann distribution:  $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$ , so,

$$\frac{N_3}{N_1} = \frac{a_3}{a_1} = 3^2 \cdot e^{\frac{\varepsilon_1 - \varepsilon_3}{kT}} = 3^2 e^{\frac{-1.36\text{eV} \cdot 8/9}{2/3\text{eV}}} \simeq 1.33 \times 10^{-8}.$$

... 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

- 2. Energy level:  $\varepsilon_n = -\frac{13.6}{n^2}\text{eV}$ ;

Boltzmann distribution:  $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$ , so,

$$\frac{N_3}{N_1} = \frac{a_3}{a_1} = 3^2 \cdot e^{\frac{\varepsilon_1 - \varepsilon_3}{kT}} = 3^2 e^{\frac{-1.36\text{eV} \cdot 8/9}{2/3\text{eV}}} \simeq 1.33 \times 10^{-8}.$$

- 3. Ionized number:



... 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

- 2. Energy level:  $\varepsilon_n = -\frac{13.6}{n^2}\text{eV}$ ;  
Boltzmann distribution:  $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$ , so,  
$$\frac{N_3}{N_1} = \frac{a_3}{a_1} = 3^2 \cdot e^{\frac{\varepsilon_1 - \varepsilon_3}{kT}} = 3^2 e^{\frac{-1.36\text{eV} \cdot 8/9}{2/3\text{eV}}} \simeq 1.33 \times 10^{-8}.$$
- 3. Ionized number:  
Ionized hydrogen, energy is the kinetic energy (continuous), electron's spin (1/2).

... 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

- 2. Energy level:  $\varepsilon_n = -\frac{13.6}{n^2}\text{eV}$ ;

Boltzmann distribution:  $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$ , so,

$$\frac{N_3}{N_1} = \frac{a_3}{a_1} = 3^2 \cdot e^{\frac{\varepsilon_1 - \varepsilon_3}{kT}} = 3^2 e^{\frac{-1.36\text{eV} \cdot 8/9}{2/3\text{eV}}} \simeq 1.33 \times 10^{-8}.$$

- 3. Ionized number:

Ionized hydrogen, energy is the kinetic energy (continuous), electron's spin ( $1/2$ ).

$$a(\varepsilon)d\varepsilon = e^{-\alpha - \beta \varepsilon} \cdot 2 \cdot \frac{2\pi V}{h^3} (2m)^{3/2} \varepsilon^{1/2} d\varepsilon$$

... 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

- 2. Energy level:  $\varepsilon_n = -\frac{13.6}{n^2}\text{eV}$ ;

Boltzmann distribution:  $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$ , so,

$$\frac{N_3}{N_1} = \frac{a_3}{a_1} = 3^2 \cdot e^{\frac{\varepsilon_1 - \varepsilon_3}{kT}} = 3^2 e^{\frac{-1.36\text{eV} \cdot 8/9}{2/3\text{eV}}} \simeq 1.33 \times 10^{-8}.$$

- 3. Ionized number:

Ionized hydrogen, energy is the kinetic energy (continuous), electron's spin (1/2).

$$a(\varepsilon)d\varepsilon = e^{-\alpha - \beta \varepsilon} \cdot 2 \cdot \frac{2\pi V}{h^3} (2m)^{3/2} \varepsilon^{1/2} d\varepsilon$$

$$N_I = \frac{4\pi V}{h^3} (2m)^{3/2} \int_0^\infty e^{-\alpha - \beta \varepsilon} \varepsilon^{1/2} d\varepsilon$$

... 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

- 2. Energy level:  $\varepsilon_n = -\frac{13.6}{n^2}\text{eV}$ ;

Boltzmann distribution:  $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$ , so,

$$\frac{N_3}{N_1} = \frac{a_3}{a_1} = 3^2 \cdot e^{\frac{\varepsilon_1 - \varepsilon_3}{kT}} = 3^2 e^{\frac{-1.36\text{eV} \cdot 8/9}{2/3\text{eV}}} \simeq 1.33 \times 10^{-8}.$$

- 3. Ionized number:

Ionized hydrogen, energy is the kinetic energy (continuous), electron's spin ( $1/2$ ).

$$a(\varepsilon)d\varepsilon = e^{-\alpha - \beta \varepsilon} \cdot 2 \cdot \frac{2\pi V}{h^3} (2m)^{3/2} \varepsilon^{1/2} d\varepsilon$$

$$N_I = \frac{4\pi V}{h^3} (2m)^{3/2} \int_0^\infty e^{-\alpha - \beta \varepsilon} \varepsilon^{1/2} d\varepsilon$$

$$= \frac{4\pi V}{h^3} (2m)^{3/2} e^{-\alpha} \beta^{-3/2} \int_0^\infty e^{-\beta \varepsilon} (\beta \varepsilon)^{1/2} d(\beta \varepsilon)$$

... 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

- 2. Energy level:  $\varepsilon_n = -\frac{13.6}{n^2}\text{eV}$ ;

Boltzmann distribution:  $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$ , so,

$$\frac{N_3}{N_1} = \frac{a_3}{a_1} = 3^2 \cdot e^{\frac{\varepsilon_1 - \varepsilon_3}{kT}} = 3^2 e^{\frac{-1.36\text{eV} \cdot 8/9}{2/3\text{eV}}} \simeq 1.33 \times 10^{-8}.$$

- 3. Ionized number:

Ionized hydrogen, energy is the kinetic energy (continuous), electron's spin ( $1/2$ ).

$$a(\varepsilon)d\varepsilon = e^{-\alpha - \beta \varepsilon} \cdot 2 \cdot \frac{2\pi V}{h^3} (2m)^{3/2} \varepsilon^{1/2} d\varepsilon$$

$$N_I = \frac{4\pi V}{h^3} (2m)^{3/2} \int_0^\infty e^{-\alpha - \beta \varepsilon} \varepsilon^{1/2} d\varepsilon$$

$$= \frac{4\pi V}{h^3} (2m)^{3/2} e^{-\alpha} \beta^{-3/2} \int_0^\infty e^{-\beta \varepsilon} (\beta \varepsilon)^{1/2} d(\beta \varepsilon)$$

$$= \frac{4\pi V}{h^3} (2m)^{3/2} e^{-\alpha} \beta^{-3/2} \frac{\sqrt{\pi}}{2},$$

... 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

- 2. Energy level:  $\varepsilon_n = -\frac{13.6}{n^2}\text{eV}$ ;

Boltzmann distribution:  $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$ , so,

$$\frac{N_3}{N_1} = \frac{a_3}{a_1} = 3^2 \cdot e^{\frac{\varepsilon_1 - \varepsilon_3}{kT}} = 3^2 e^{\frac{-1.36\text{eV} \cdot 8/9}{2/3\text{eV}}} \simeq 1.33 \times 10^{-8}.$$

- 3. Ionized number:

Ionized hydrogen, energy is the kinetic energy (continuous), electron's spin ( $1/2$ ).

$$a(\varepsilon)d\varepsilon = e^{-\alpha - \beta \varepsilon} \cdot 2 \cdot \frac{2\pi V}{h^3} (2m)^{3/2} \varepsilon^{1/2} d\varepsilon$$

$$N_I = \frac{4\pi V}{h^3} (2m)^{3/2} \int_0^\infty e^{-\alpha - \beta \varepsilon} \varepsilon^{1/2} d\varepsilon$$

$$= \frac{4\pi V}{h^3} (2m)^{3/2} e^{-\alpha} \beta^{-3/2} \int_0^\infty e^{-\beta \varepsilon} (\beta \varepsilon)^{1/2} d(\beta \varepsilon)$$

$$= \frac{4\pi V}{h^3} (2m)^{3/2} e^{-\alpha} \beta^{-3/2} \frac{\sqrt{\pi}}{2},$$

... 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

- $$N_I = \frac{4\pi V}{h^3} (2m)^{3/2} e^{-\alpha} \beta^{-3/2} \frac{\sqrt{\pi}}{2},$$

... 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

- $$N_I = \frac{4\pi V}{h^3} (2m)^{3/2} e^{-\alpha} \beta^{-3/2} \frac{\sqrt{\pi}}{2},$$

$$N_1 = N e^{-\alpha - \frac{13.6\text{eV}}{kT}}; \text{ (?? see saha.pdf)}$$



... 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

- $$N_I = \frac{4\pi V}{h^3} (2m)^{3/2} e^{-\alpha} \beta^{-3/2} \frac{\sqrt{\pi}}{2},$$

$$N_1 = N e^{-\alpha - \frac{13.6\text{eV}}{kT}}; \text{ (?? see saha.pdf)}$$

$$\Rightarrow \frac{N_I}{N_1} = \frac{4\pi}{n h^3} (2m)^{3/2} e^{-\frac{13.6\text{eV}}{kT}} \beta^{-3/2}$$

... 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

- $$N_I = \frac{4\pi V}{h^3} (2m)^{3/2} e^{-\alpha} \beta^{-3/2} \frac{\sqrt{\pi}}{2},$$

$$N_1 = N e^{-\alpha - \frac{13.6\text{eV}}{kT}}; \text{ (?? see saha.pdf)}$$

$$\Rightarrow \frac{N_I}{N_1} = \frac{4\pi}{n h^3} (2m)^{3/2} e^{-\frac{13.6\text{eV}}{kT}} \beta^{-3/2}$$

$$= \frac{2}{n h^3} (2\pi m k T)^{3/2} e^{-\frac{13.6\text{eV}}{kT}}$$

... 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

- $$N_I = \frac{4\pi V}{h^3} (2m)^{3/2} e^{-\alpha} \beta^{-3/2} \frac{\sqrt{\pi}}{2},$$

$$N_1 = N e^{-\alpha - \frac{13.6\text{eV}}{kT}}; \text{ (?? see saha.pdf)}$$

$$\Rightarrow \frac{N_I}{N_1} = \frac{4\pi}{n h^3} (2m)^{3/2} e^{-\frac{13.6\text{eV}}{kT}} \beta^{-3/2}$$

$$= \frac{2}{n h^3} (2\pi m k T)^{3/2} e^{-\frac{13.6\text{eV}}{kT}}$$

with  $n \simeq 10^{23} \text{m}^{-3}$ ,  $h = 6.626 \times 10^{-34} \text{J} \cdot \text{s}$ ,  
 $kT \simeq 2/3 \text{eV}$ ,  $m_e = 9.1 \times 10^{-31} \text{kg}$

... 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

- $$N_I = \frac{4\pi V}{h^3} (2m)^{3/2} e^{-\alpha} \beta^{-3/2} \frac{\sqrt{\pi}}{2},$$

$$N_1 = N e^{-\alpha - \frac{13.6\text{eV}}{kT}}; \text{ (?? see saha.pdf)}$$

$$\Rightarrow \frac{N_I}{N_1} = \frac{4\pi}{n h^3} (2m)^{3/2} e^{-\frac{13.6\text{eV}}{kT}} \beta^{-3/2}$$

$$= \frac{2}{n h^3} (2\pi m k T)^{3/2} e^{-\frac{13.6\text{eV}}{kT}}$$

with  $n \simeq 10^{23} \text{m}^{-3}$ ,  $h = 6.626 \times 10^{-34} \text{J} \cdot \text{s}$ ,  
 $kT \simeq 2/3 \text{eV}$ ,  $m_e = 9.1 \times 10^{-31} \text{kg}$

$$\Rightarrow \frac{N_I}{N_1} \simeq \frac{2 \cdot (2\pi \cdot 9.1 \times 10^{-31} \text{kg} \cdot 2/3 \cdot 1.6 \times 10^{-19})^{3/2}}{10^{23} \text{m}^{-3} \cdot (6.626 \times 10^{-34} \text{J} \cdot \text{s})^3} e^{-\frac{13.6\text{eV}}{2/3\text{eV}}}$$

... 2. ratio of the number of hydrogen at the 2nd excited level ( $n = 3$ ) and the ground level ( $n = 1$ ); 3. number of ionized hydrogen.

- $$N_I = \frac{4\pi V}{h^3} (2m)^{3/2} e^{-\alpha} \beta^{-3/2} \frac{\sqrt{\pi}}{2},$$

$$N_1 = N e^{-\alpha - \frac{13.6\text{eV}}{kT}}; \text{ (?? see saha.pdf)}$$

$$\Rightarrow \frac{N_I}{N_1} = \frac{4\pi}{nh^3} (2m)^{3/2} e^{-\frac{13.6\text{eV}}{kT}} \beta^{-3/2}$$

$$= \frac{2}{nh^3} (2\pi m kT)^{3/2} e^{-\frac{13.6\text{eV}}{kT}}$$

with  $n \simeq 10^{23} \text{m}^{-3}$ ,  $h = 6.626 \times 10^{-34} \text{J} \cdot \text{s}$ ,  
 $kT \simeq 2/3 \text{eV}$ ,  $m_e = 9.1 \times 10^{-31} \text{kg}$

$$\Rightarrow \frac{N_I}{N_1} \simeq \frac{2 \cdot (2\pi \cdot 9.1 \times 10^{-31} \text{kg} \cdot 2/3 \cdot 1.6 \times 10^{-19})^{3/2}}{10^{23} \text{m}^{-3} \cdot (6.626 \times 10^{-34} \text{J} \cdot \text{s})^3} e^{-\frac{13.6\text{eV}}{2/3\text{eV}}}$$

$$\simeq 4.5 \times 10^{-5} \ll 1.$$

# Table of contents

- ① §6. Most probable distribution of nearly independent particles
- ② §7. Boltzmann statistics
- ③ §8. Bose statistics and Fermi statistics
  - Bose statistics
  - Fermi statistics
  - Fermion gas in astrophysics

A10.1 Boson gas with number  $N$ , two energy levels,  $0, \varepsilon$ ; degeneracy  $\omega_1, \omega_2$ . Determine the temperature  $T$  when  $N_1 = 2N_2$ .

A10.1 Boson gas with number  $N$ , two energy levels,  $0, \varepsilon$ ; degeneracy  $\omega_1, \omega_2$ . Determine the temperature  $T$  when  $N_1 = 2N_2$ .

- $N_1 = \frac{2}{3}N, N_2 = \frac{1}{3}N.$



A10.1 Boson gas with number  $N$ , two energy levels,  $0, \varepsilon$ ; degeneracy  $\omega_1, \omega_2$ . Determine the temperature  $T$  when  $N_1 = 2N_2$ .

- $N_1 = \frac{2}{3}N, N_2 = \frac{1}{3}N$ .

Bose distribution:  $a_l = \frac{\omega_l}{e^{\frac{\varepsilon_l - \mu}{kT}} - 1}$ .

A10.1 Boson gas with number  $N$ , two energy levels,  $0, \varepsilon$ ; degeneracy  $\omega_1, \omega_2$ . Determine the temperature  $T$  when  $N_1 = 2N_2$ .

- $N_1 = \frac{2}{3}N, N_2 = \frac{1}{3}N$ .

Bose distribution:  $a_l = \frac{\omega_l}{e^{\frac{\varepsilon_l - \mu}{kT}} - 1}$ .

$$N_1 = a_1 = \frac{\omega_1}{e^{\frac{\varepsilon_1 - \mu}{kT}} - 1} = \frac{\omega_1}{e^{\frac{-\mu}{kT}} - 1} = \frac{2}{3}N$$

A10.1 Boson gas with number  $N$ , two energy levels,  $0, \varepsilon$ ; degeneracy  $\omega_1, \omega_2$ . Determine the temperature  $T$  when  $N_1 = 2N_2$ .

- $N_1 = \frac{2}{3}N, N_2 = \frac{1}{3}N$ .

Bose distribution:  $a_l = \frac{\omega_l}{e^{\frac{\varepsilon_l - \mu}{kT}} - 1}$ .

$$N_1 = a_1 = \frac{\omega_1}{e^{\frac{\varepsilon_1 - \mu}{kT}} - 1} = \frac{\omega_1}{e^{\frac{-\mu}{kT}} - 1} = \frac{2}{3}N$$

$$\Rightarrow \mu = -kT \ln\left(1 + \frac{3}{2} \frac{\omega_1}{N}\right);$$

A10.1 Boson gas with number  $N$ , two energy levels,  $0, \varepsilon$ ; degeneracy  $\omega_1, \omega_2$ . Determine the temperature  $T$  when  $N_1 = 2N_2$ .

- $N_1 = \frac{2}{3}N, N_2 = \frac{1}{3}N.$

Bose distribution:  $a_l = \frac{\omega_l}{e^{\frac{\varepsilon_l - \mu}{kT}} - 1}.$

$$N_1 = a_1 = \frac{\omega_1}{e^{\frac{\varepsilon_1 - \mu}{kT}} - 1} = \frac{\omega_1}{e^{\frac{-\mu}{kT}} - 1} = \frac{2}{3}N$$

$$\Rightarrow \mu = -kT \ln\left(1 + \frac{3}{2} \frac{\omega_1}{N}\right);$$

$$N_2 = a_2 = \frac{\omega_2}{e^{\frac{\varepsilon_2 - \mu}{kT}} - 1} = \frac{\omega_2}{e^{\frac{\varepsilon - \mu}{kT}} - 1} = \frac{1}{3}N$$

A10.1 Boson gas with number  $N$ , two energy levels,  $0, \varepsilon$ ; degeneracy  $\omega_1, \omega_2$ . Determine the temperature  $T$  when  $N_1 = 2N_2$ .

- $N_1 = \frac{2}{3}N, N_2 = \frac{1}{3}N.$

Bose distribution:  $a_l = \frac{\omega_l}{e^{\frac{\varepsilon_l - \mu}{kT}} - 1}.$

$$N_1 = a_1 = \frac{\omega_1}{e^{\frac{\varepsilon_1 - \mu}{kT}} - 1} = \frac{\omega_1}{e^{\frac{-\mu}{kT}} - 1} = \frac{2}{3}N$$

$$\Rightarrow \mu = -kT \ln\left(1 + \frac{3}{2} \frac{\omega_1}{N}\right);$$

$$N_2 = a_2 = \frac{\omega_2}{e^{\frac{\varepsilon_2 - \mu}{kT}} - 1} = \frac{\omega_2}{e^{\frac{\varepsilon - \mu}{kT}} - 1} = \frac{1}{3}N$$

$$\Rightarrow \varepsilon - \mu = kT \ln\left(1 + \frac{3\omega_2}{N}\right)$$

A10.1 Boson gas with number  $N$ , two energy levels,  $0, \varepsilon$ ; degeneracy  $\omega_1, \omega_2$ . Determine the temperature  $T$  when  $N_1 = 2N_2$ .

- $N_1 = \frac{2}{3}N, N_2 = \frac{1}{3}N.$

Bose distribution:  $a_l = \frac{\omega_l}{e^{\frac{\varepsilon_l - \mu}{kT}} - 1}.$

$$N_1 = a_1 = \frac{\omega_1}{e^{\frac{\varepsilon_1 - \mu}{kT}} - 1} = \frac{\omega_1}{e^{\frac{-\mu}{kT}} - 1} = \frac{2}{3}N$$

$$\Rightarrow \mu = -kT \ln\left(1 + \frac{3}{2} \frac{\omega_1}{N}\right);$$

$$N_2 = a_2 = \frac{\omega_2}{e^{\frac{\varepsilon_2 - \mu}{kT}} - 1} = \frac{\omega_2}{e^{\frac{\varepsilon - \mu}{kT}} - 1} = \frac{1}{3}N$$

$$\Rightarrow \varepsilon - \mu = kT \ln\left(1 + \frac{3\omega_2}{N}\right)$$

$$\Rightarrow T = \frac{\varepsilon}{k \ln\left(\frac{2N+6\omega_2}{2N+3\omega_1}\right)}.$$

# Table of contents

- 1 §6. Most probable distribution of nearly independent particles
- 2 §7. Boltzmann statistics
- 3 §8. Bose statistics and Fermi statistics
  - Bose statistics
  - Fermi statistics
  - Fermion gas in astrophysics

A9.14 For the free electrons in metal, prove: the probability to find an electron at state with energy  $\mu - \Delta$  is equal to not to find an electron at state with energy  $\mu + \Delta$ .



A9.14 For the free electrons in metal, prove: the probability to find an electron at state with energy  $\mu - \Delta$  is equal to not to find an electron at state with energy  $\mu + \Delta$ .

- Fermi distribution at a quantum state:

$$f(\varepsilon) = \frac{1}{e^{\beta(\varepsilon-\mu)} + 1}.$$

A9.14 For the free electrons in metal, prove: the probability to find an electron at state with energy  $\mu - \Delta$  is equal to not to find an electron at state with energy  $\mu + \Delta$ .

- Fermi distribution at a quantum state:

$$f(\varepsilon) = \frac{1}{e^{\beta(\varepsilon-\mu)}+1}.$$

The probability to find an electron at  $\mu - \Delta$ :

$$\frac{1}{e^{\beta[(\mu-\Delta)-\mu]}+1} = \frac{1}{e^{-\beta\Delta}+1};$$

A9.14 For the free electrons in metal, prove: the probability to find an electron at state with energy  $\mu - \Delta$  is equal to not to find an electron at state with energy  $\mu + \Delta$ .

- Fermi distribution at a quantum state:

$$f(\varepsilon) = \frac{1}{e^{\beta(\varepsilon-\mu)} + 1}.$$

The probability to find an electron at  $\mu - \Delta$ :

$$\frac{1}{e^{\beta[(\mu-\Delta)-\mu]} + 1} = \frac{1}{e^{-\beta\Delta} + 1};$$

The probability to find an electron at  $\mu + \Delta$ :

$$\frac{1}{e^{\beta[(\mu+\Delta)-\mu]} + 1} = \frac{1}{e^{\beta\Delta} + 1};$$

A9.14 For the free electrons in metal, prove: the probability to find an electron at state with energy  $\mu - \Delta$  is equal to not to find an electron at state with energy  $\mu + \Delta$ .

- Fermi distribution at a quantum state:

$$f(\varepsilon) = \frac{1}{e^{\beta(\varepsilon-\mu)} + 1}.$$

The probability to find an electron at  $\mu - \Delta$ :

$$\frac{1}{e^{\beta[(\mu-\Delta)-\mu]} + 1} = \frac{1}{e^{-\beta\Delta} + 1};$$

The probability to find an electron at  $\mu + \Delta$ :

$$\frac{1}{e^{\beta[(\mu+\Delta)-\mu]} + 1} = \frac{1}{e^{\beta\Delta} + 1};$$

The probability not to find an electron at  $\mu + \Delta$ :

$$1 - \frac{1}{e^{\beta\Delta} + 1}$$

A9.14 For the free electrons in metal, prove: the probability to find an electron at state with energy  $\mu - \Delta$  is equal to not to find an electron at state with energy  $\mu + \Delta$ .

- Fermi distribution at a quantum state:

$$f(\varepsilon) = \frac{1}{e^{\beta(\varepsilon-\mu)}+1}.$$

The probability to find an electron at  $\mu - \Delta$ :

$$\frac{1}{e^{\beta[(\mu-\Delta)-\mu]}+1} = \frac{1}{e^{-\beta\Delta}+1};$$

The probability to find an electron at  $\mu + \Delta$ :

$$\frac{1}{e^{\beta[(\mu+\Delta)-\mu]}+1} = \frac{1}{e^{\beta\Delta}+1};$$

The probability not to find an electron at  $\mu + \Delta$ :

$$1 - \frac{1}{e^{\beta\Delta}+1} = \frac{e^{\beta\Delta}}{e^{\beta\Delta}+1} = \frac{1}{e^{-\beta\Delta}+1}.$$

A9.15 Consider box with volume  $V$ , temperature  $T$ , contains photons and electron-positron pairs.  $\gamma \rightleftharpoons e^+ + e^-$  is in equilibrium. Calculate 1. chemical potentials; 2. number of electron-positron pairs in limits  $kT \gg m_e c^2$ , and  $kT \ll m_e c^2$ .

A9.15 Consider box with volume  $V$ , temperature  $T$ , contains photons and electron-positron pairs.  $\gamma \rightleftharpoons e^+ + e^-$  is in equilibrium. Calculate 1. chemical potentials; 2. number of electron-positron pairs in limits  $kT \gg m_e c^2$ , and  $kT \ll m_e c^2$ .

- 1. For chemical reaction  $A \rightleftharpoons B + C$  in equilibrium, there exists  $\mu_A = \mu_B + \mu_C$ .

A9.15 Consider box with volume  $V$ , temperature  $T$ , contains photons and electron-positron pairs.  $\gamma \rightleftharpoons e^+ + e^-$  is in equilibrium. Calculate 1. chemical potentials; 2. number of electron-positron pairs in limits  $kT \gg m_e c^2$ , and  $kT \ll m_e c^2$ .

- 1. For chemical reaction  $A \rightleftharpoons B + C$  in equilibrium, there exists  $\mu_A = \mu_B + \mu_C$ .  
And  $\mu_\gamma = 0$ ,



A9.15 Consider box with volume  $V$ , temperature  $T$ , contains photons and electron-positron pairs.  $\gamma \rightleftharpoons e^+ + e^-$  is in equilibrium. Calculate 1. chemical potentials; 2. number of electron-positron pairs in limits  $kT \gg m_e c^2$ , and  $kT \ll m_e c^2$ .

- 1. For chemical reaction  $A \rightleftharpoons B + C$  in equilibrium, there exists  $\mu_A = \mu_B + \mu_C$ .  
And  $\mu_\gamma = 0$ , so  $\mu_{e^+} + \mu_{e^-} = 0$ .

A9.15 Consider box with volume  $V$ , temperature  $T$ , contains photons and electron-positron pairs.  $\gamma \rightleftharpoons e^+ + e^-$  is in equilibrium. Calculate 1. chemical potentials; 2. number of electron-positron pairs in limits  $kT \gg m_e c^2$ , and  $kT \ll m_e c^2$ .

- 1. For chemical reaction  $A \rightleftharpoons B + C$  in equilibrium, there exists  $\mu_A = \mu_B + \mu_C$ .  
And  $\mu_\gamma = 0$ , so  $\mu_{e^+} + \mu_{e^-} = 0$ . Because of the symmetry between electron and positron,  
 $\mu_{e^+} = \mu_{e^-} = 0$ .

A9.15 Consider box with volume  $V$ , temperature  $T$ , contains photons and electron-positron pairs.  $\gamma \rightleftharpoons e^+ + e^-$  is in equilibrium. Calculate 1. chemical potentials; 2. number of electron-positron pairs in limits  $kT \gg m_e c^2$ , and  $kT \ll m_e c^2$ .

- 1. For chemical reaction  $A \rightleftharpoons B + C$  in equilibrium, there exists  $\mu_A = \mu_B + \mu_C$ .

And  $\mu_\gamma = 0$ , so  $\mu_{e^+} + \mu_{e^-} = 0$ . Because of the symmetry between electron and positron,

$$\mu_{e^+} = \mu_{e^-} = 0.$$

- 2. Fermi distribution:  $f(\varepsilon) = \frac{1}{e^{\beta(\varepsilon - \mu)} + 1}$ .

A9.15 Consider box with volume  $V$ , temperature  $T$ , contains photons and electron-positron pairs.  $\gamma \rightleftharpoons e^+ + e^-$  is in equilibrium. Calculate 1. chemical potentials; 2. number of electron-positron pairs in limits  $kT \gg m_e c^2$ , and  $kT \ll m_e c^2$ .

- 1. For chemical reaction  $A \rightleftharpoons B + C$  in equilibrium, there exists  $\mu_A = \mu_B + \mu_C$ .

And  $\mu_\gamma = 0$ , so  $\mu_{e^+} + \mu_{e^-} = 0$ . Because of the symmetry between electron and positron,

$$\mu_{e^+} = \mu_{e^-} = 0.$$

- 2. Fermi distribution:  $f(\varepsilon) = \frac{1}{e^{\beta(\varepsilon - \mu)} + 1}$ .

$$N_{e^\pm} = \int_0^\infty f(\varepsilon) 2 \cdot \frac{V 4\pi p^2 dp}{h^3}$$

A9.15 Consider box with volume  $V$ , temperature  $T$ , contains photons and electron-positron pairs.  $\gamma \rightleftharpoons e^+ + e^-$  is in equilibrium. Calculate 1. chemical potentials; 2. number of electron-positron pairs in limits  $kT \gg m_e c^2$ , and  $kT \ll m_e c^2$ .

- 1. For chemical reaction  $A \rightleftharpoons B + C$  in equilibrium, there exists  $\mu_A = \mu_B + \mu_C$ .

And  $\mu_\gamma = 0$ , so  $\mu_{e^+} + \mu_{e^-} = 0$ . Because of the symmetry between electron and positron,

$$\mu_{e^+} = \mu_{e^-} = 0.$$

- 2. Fermi distribution:  $f(\varepsilon) = \frac{1}{e^{\beta(\varepsilon - \mu)} + 1}$ .

$$N_{e^\pm} = \int_0^\infty f(\varepsilon) 2 \cdot \frac{V 4\pi p^2 dp}{h^3} = \int_0^\infty \frac{1}{e^{\frac{\varepsilon}{kT}} + 1} 2 \cdot \frac{V 4\pi p^2 dp}{h^3}$$

... 2. number of electron-positron pairs in limits  $kT \gg m_e c^2$ , and  $kT \ll m_e c^2$ .

- $$N_{e^\pm} = \int_0^\infty \frac{1}{e^{\frac{\epsilon}{kT}} + 1} 2 \cdot \frac{V 4\pi p^2 dp}{h^3}.$$

... 2. number of electron-positron pairs in limits  $kT \gg m_e c^2$ , and  $kT \ll m_e c^2$ .

- $N_{e^\pm} = \int_0^\infty \frac{1}{e^{\frac{\varepsilon}{kT}} + 1} 2 \cdot \frac{V 4\pi p^2 dp}{h^3}.$
- $kT \gg m_e c^2$ , relativistic,  $\varepsilon \simeq pc$ ,

... 2. number of electron-positron pairs in limits  $kT \gg m_e c^2$ , and  $kT \ll m_e c^2$ .

- $N_{e^\pm} = \int_0^\infty \frac{1}{e^{\frac{\varepsilon}{kT}} + 1} 2 \cdot \frac{V 4\pi p^2 dp}{h^3}.$
- $kT \gg m_e c^2$ , relativistic,  $\varepsilon \simeq pc$ ,  

$$N_{e^\pm} = \int_0^\infty \frac{1}{e^{\frac{pc}{kT}} + 1} 2 \cdot \frac{V 4\pi p^2 dp}{h^3} = \frac{V (kT)^3}{\pi^2 (\hbar c)^3} \int_0^\infty \frac{x^2 dx}{e^x + 1}.$$



... 2. number of electron-positron pairs in limits  $kT \gg m_e c^2$ , and  $kT \ll m_e c^2$ .

- $N_{e^\pm} = \int_0^\infty \frac{1}{e^{\frac{\varepsilon}{kT}} + 1} 2 \cdot \frac{V 4\pi p^2 dp}{h^3}.$
- $kT \gg m_e c^2$ , relativistic,  $\varepsilon \simeq pc$ ,  

$$N_{e^\pm} = \int_0^\infty \frac{1}{e^{\frac{pc}{kT}} + 1} 2 \cdot \frac{V 4\pi p^2 dp}{h^3} = \frac{V (kT)^3}{\pi^2 (\hbar c)^3} \int_0^\infty \frac{x^2 dx}{e^x + 1}.$$
- $kT \ll m_e c^2$ ,  $\varepsilon = \sqrt{(pc)^2 + (m_e c^2)^2} \simeq m_e c^2 + \frac{p^2}{2m_e}$   
 and  $e^{\frac{\varepsilon}{kT}} \gg 1,$

... 2. number of electron-positron pairs in limits  $kT \gg m_e c^2$ , and  $kT \ll m_e c^2$ .

- $N_{e^\pm} = \int_0^\infty \frac{1}{e^{\frac{\varepsilon}{kT}} + 1} 2 \cdot \frac{V 4\pi p^2 dp}{h^3}.$
- $kT \gg m_e c^2$ , relativistic,  $\varepsilon \simeq pc$ ,  

$$N_{e^\pm} = \int_0^\infty \frac{1}{e^{\frac{pc}{kT}} + 1} 2 \cdot \frac{V 4\pi p^2 dp}{h^3} = \frac{V (kT)^3}{\pi^2 (\hbar c)^3} \int_0^\infty \frac{x^2 dx}{e^x + 1}.$$
- $kT \ll m_e c^2$ ,  $\varepsilon = \sqrt{(pc)^2 + (m_e c^2)^2} \simeq m_e c^2 + \frac{p^2}{2m_e}$   
 and  $e^{\frac{\varepsilon}{kT}} \gg 1$ ,  $N_{e^\pm} = \int_0^\infty \frac{1}{e^{\frac{m_e c^2 + \frac{p^2}{2m_e}}{kT}}} 2 \cdot \frac{V 4\pi p^2 dp}{h^3}$

... 2. number of electron-positron pairs in limits  $kT \gg m_e c^2$ , and  $kT \ll m_e c^2$ .

- $N_{e^\pm} = \int_0^\infty \frac{1}{e^{\frac{\varepsilon}{kT}} + 1} 2 \cdot \frac{V 4\pi p^2 dp}{h^3}.$
- $kT \gg m_e c^2$ , relativistic,  $\varepsilon \simeq pc$ ,  

$$N_{e^\pm} = \int_0^\infty \frac{1}{e^{\frac{pc}{kT}} + 1} 2 \cdot \frac{V 4\pi p^2 dp}{h^3} = \frac{V (kT)^3}{\pi^2 (\hbar c)^3} \int_0^\infty \frac{x^2 dx}{e^x + 1}.$$
- $kT \ll m_e c^2$ ,  $\varepsilon = \sqrt{(pc)^2 + (m_e c^2)^2} \simeq m_e c^2 + \frac{p^2}{2m_e}$   
 and  $e^{\frac{\varepsilon}{kT}} \gg 1$ ,  $N_{e^\pm} = \int_0^\infty \frac{1}{e^{\frac{m_e c^2 + \frac{p^2}{2m_e}}{kT}}} 2 \cdot \frac{V 4\pi p^2 dp}{h^3}$   

$$= e^{-\frac{m_e c^2}{kT}} \int_0^\infty e^{-\frac{p^2}{2m_e kT}} 2 \cdot \frac{V 4\pi p^2 dp}{h^3}$$

... 2. number of electron-positron pairs in limits  $kT \gg m_e c^2$ , and  $kT \ll m_e c^2$ .

- $N_{e^\pm} = \int_0^\infty \frac{1}{e^{\frac{\varepsilon}{kT}} + 1} 2 \cdot \frac{V 4\pi p^2 dp}{h^3}.$

- $kT \gg m_e c^2$ , relativistic,  $\varepsilon \simeq pc$ ,  

$$N_{e^\pm} = \int_0^\infty \frac{1}{e^{\frac{pc}{kT}} + 1} 2 \cdot \frac{V 4\pi p^2 dp}{h^3} = \frac{V (kT)^3}{\pi^2 (\hbar c)^3} \int_0^\infty \frac{x^2 dx}{e^x + 1}.$$

- $kT \ll m_e c^2$ ,  $\varepsilon = \sqrt{(pc)^2 + (m_e c^2)^2} \simeq m_e c^2 + \frac{p^2}{2m_e}$   
 and  $e^{\frac{\varepsilon}{kT}} \gg 1$ , 
$$N_{e^\pm} = \int_0^\infty \frac{1}{e^{\frac{m_e c^2 + \frac{p^2}{2m_e}}{kT}}} 2 \cdot \frac{V 4\pi p^2 dp}{h^3}$$

$$= e^{-\frac{m_e c^2}{kT}} \int_0^\infty e^{-\frac{p^2}{2m_e kT}} 2 \cdot \frac{V 4\pi p^2 dp}{h^3} = 2V \left( \frac{2\pi m_e kT}{h^2} \right)^{\frac{3}{2}} e^{-\frac{m_e c^2}{kT}}.$$

A9.23 Fermi energy of Fermi gas is  $\varepsilon_F$ . Calculate the average energy at  $T = 0\text{K}$ .

A9.23 Fermi energy of Fermi gas is  $\varepsilon_F$ . Calculate the average energy at  $T = 0\text{K}$ .

- Two cases according to the velocity comparing with  $c$ .

A9.23 Fermi energy of Fermi gas is  $\varepsilon_F$ . Calculate the average energy at  $T = 0\text{K}$ .

- Two cases according to the velocity comparing with  $c$ .
- a. Non-relativistic case,  $\varepsilon = \frac{p^2}{2m}$ ,

A9.23 Fermi energy of Fermi gas is  $\varepsilon_F$ . Calculate the average energy at  $T = 0\text{K}$ .

- Two cases according to the velocity comparing with  $c$ .
- a. Non-relativistic case,  $\varepsilon = \frac{p^2}{2m}$ ,  
number of state:  $D(\varepsilon)d\varepsilon = g \cdot \frac{V4\pi p^2 dp}{h^3} = C\varepsilon^{1/2}d\varepsilon$ .



A9.23 Fermi energy of Fermi gas is  $\varepsilon_F$ . Calculate the average energy at  $T = 0\text{K}$ .

- Two cases according to the velocity comparing with  $c$ .
- a. Non-relativistic case,  $\varepsilon = \frac{p^2}{2m}$ ,  
number of state:  $D(\varepsilon)d\varepsilon = g \cdot \frac{V4\pi p^2 dp}{h^3} = C\varepsilon^{1/2}d\varepsilon$ .  
average energy:  $\bar{\varepsilon} = \frac{\int_0^{\varepsilon_F} \varepsilon^{3/2} d\varepsilon}{\int_0^{\varepsilon_F} \varepsilon^{1/2} d\varepsilon}$

A9.23 Fermi energy of Fermi gas is  $\varepsilon_F$ . Calculate the average energy at  $T = 0\text{K}$ .

- Two cases according to the velocity comparing with  $c$ .
- a. Non-relativistic case,  $\varepsilon = \frac{p^2}{2m}$ ,  
 number of state:  $D(\varepsilon)d\varepsilon = g \cdot \frac{V4\pi p^2 dp}{h^3} = C\varepsilon^{1/2}d\varepsilon$ .  
 average energy:  $\bar{\varepsilon} = \frac{\int_0^{\varepsilon_F} \varepsilon^{3/2} d\varepsilon}{\int_0^{\varepsilon_F} \varepsilon^{1/2} d\varepsilon} = \frac{3}{5}\varepsilon_F$ .

A9.23 Fermi energy of Fermi gas is  $\varepsilon_F$ . Calculate the average energy at  $T = 0\text{K}$ .

- Two cases according to the velocity comparing with  $c$ .
- a. Non-relativistic case,  $\varepsilon = \frac{p^2}{2m}$ ,  
number of state:  $D(\varepsilon)d\varepsilon = g \cdot \frac{V4\pi p^2 dp}{h^3} = C\varepsilon^{1/2}d\varepsilon$ .  
average energy:  $\bar{\varepsilon} = \frac{\int_0^{\varepsilon_F} \varepsilon^{3/2} d\varepsilon}{\int_0^{\varepsilon_F} \varepsilon^{1/2} d\varepsilon} = \frac{3}{5}\varepsilon_F$ .
- b. Relativistic case,  $\varepsilon = pc$ ,

A9.23 Fermi energy of Fermi gas is  $\varepsilon_F$ . Calculate the average energy at  $T = 0\text{K}$ .

- Two cases according to the velocity comparing with  $c$ .
- a. Non-relativistic case,  $\varepsilon = \frac{p^2}{2m}$ ,  
number of state:  $D(\varepsilon)d\varepsilon = g \cdot \frac{V4\pi p^2 dp}{h^3} = C\varepsilon^{1/2}d\varepsilon$ .  
average energy:  $\bar{\varepsilon} = \frac{\int_0^{\varepsilon_F} \varepsilon^{3/2} d\varepsilon}{\int_0^{\varepsilon_F} \varepsilon^{1/2} d\varepsilon} = \frac{3}{5}\varepsilon_F$ .
- b. Relativistic case,  $\varepsilon = pc$ ,  
number of state:  $D(\varepsilon)d\varepsilon = g \cdot \frac{V4\pi p^2 dp}{h^3} = C'\varepsilon^2 d\varepsilon$ .

A9.23 Fermi energy of Fermi gas is  $\varepsilon_F$ . Calculate the average energy at  $T = 0\text{K}$ .

- Two cases according to the velocity comparing with  $c$ .
- a. Non-relativistic case,  $\varepsilon = \frac{p^2}{2m}$ ,  
 number of state:  $D(\varepsilon)d\varepsilon = g \cdot \frac{V4\pi p^2 dp}{h^3} = C\varepsilon^{1/2}d\varepsilon$ .  
 average energy:  $\bar{\varepsilon} = \frac{\int_0^{\varepsilon_F} \varepsilon^{3/2} d\varepsilon}{\int_0^{\varepsilon_F} \varepsilon^{1/2} d\varepsilon} = \frac{3}{5}\varepsilon_F$ .
- b. Relativistic case,  $\varepsilon = pc$ ,  
 number of state:  $D(\varepsilon)d\varepsilon = g \cdot \frac{V4\pi p^2 dp}{h^3} = C'\varepsilon^2 d\varepsilon$ .  
 average energy:  $\bar{\varepsilon} = \frac{\int_0^{\varepsilon_F} \varepsilon^3 d\varepsilon}{\int_0^{\varepsilon_F} \varepsilon^2 d\varepsilon}$

A9.23 Fermi energy of Fermi gas is  $\varepsilon_F$ . Calculate the average energy at  $T = 0\text{K}$ .

- Two cases according to the velocity comparing with  $c$ .
- a. Non-relativistic case,  $\varepsilon = \frac{p^2}{2m}$ ,  
number of state:  $D(\varepsilon)d\varepsilon = g \cdot \frac{V4\pi p^2 dp}{h^3} = C\varepsilon^{1/2}d\varepsilon$ .  
average energy:  $\bar{\varepsilon} = \frac{\int_0^{\varepsilon_F} \varepsilon^{3/2} d\varepsilon}{\int_0^{\varepsilon_F} \varepsilon^{1/2} d\varepsilon} = \frac{3}{5}\varepsilon_F$ .
- b. Relativistic case,  $\varepsilon = pc$ ,  
number of state:  $D(\varepsilon)d\varepsilon = g \cdot \frac{V4\pi p^2 dp}{h^3} = C'\varepsilon^2 d\varepsilon$ .  
average energy:  $\bar{\varepsilon} = \frac{\int_0^{\varepsilon_F} \varepsilon^3 d\varepsilon}{\int_0^{\varepsilon_F} \varepsilon^2 d\varepsilon} = \frac{3}{4}\varepsilon_F$ .

A9.32 Prove  $\mathcal{Z}$  for strong degenerated Fermi **gas**:

$\mathcal{Z}(\alpha, \beta, V) = e^{a[\frac{4}{15}(-\alpha)^{5/2} + \frac{\pi^2}{6}(-\alpha)^{1/2} - \frac{7\pi^4}{1440}(-\alpha)^{-3/2}]} \beta^{-3/2}$ , where  
 $a = \frac{2\pi g V (2m)^{3/2}}{h^3}$ ,  $g$  is the inner degeneracy. Then calculate the grand potential  $J(T, V, \mu)$ . (non-relativistic)

A9.32 Prove  $\mathcal{Z}$  for strong degenerated Fermi **gas**:

$\mathcal{Z}(\alpha, \beta, V) = e^{a[\frac{4}{15}(-\alpha)^{5/2} + \frac{\pi^2}{6}(-\alpha)^{1/2} - \frac{7\pi^4}{1440}(-\alpha)^{-3/2}]} \beta^{-3/2}$ , where  $a = \frac{2\pi g V (2m)^{3/2}}{h^3}$ ,  $g$  is the inner degeneracy. Then calculate the grand potential  $J(T, V, \mu)$ . (non-relativistic)

$$\bullet D(\varepsilon)d\varepsilon = g \cdot \frac{V 4\pi p^2 dp}{h^3} = \frac{2\pi g V (2m)^{3/2}}{h^3} \varepsilon^{1/2} d\varepsilon = a \varepsilon^{1/2} d\varepsilon.$$



A9.32 Prove  $\mathcal{Z}$  for strong degenerated Fermi **gas**:

$\mathcal{Z}(\alpha, \beta, V) = e^{a[\frac{4}{15}(-\alpha)^{5/2} + \frac{\pi^2}{6}(-\alpha)^{1/2} - \frac{7\pi^4}{1440}(-\alpha)^{-3/2}]} \beta^{-3/2}$ , where  $a = \frac{2\pi g V (2m)^{3/2}}{h^3}$ ,  $g$  is the inner degeneracy. Then calculate the grand potential  $J(T, V, \mu)$ . (non-relativistic)

- $D(\varepsilon)d\varepsilon = g \cdot \frac{V 4\pi p^2 dp}{h^3} = \frac{2\pi g V (2m)^{3/2}}{h^3} \varepsilon^{1/2} d\varepsilon = a \varepsilon^{1/2} d\varepsilon.$   
 $\ln \mathcal{Z} = \sum \omega_l \ln(1 + e^{-\alpha - \beta \varepsilon_l})$

A9.32 Prove  $\mathcal{Z}$  for strong degenerated Fermi **gas**:

$\mathcal{Z}(\alpha, \beta, V) = e^{a[\frac{4}{15}(-\alpha)^{5/2} + \frac{\pi^2}{6}(-\alpha)^{1/2} - \frac{7\pi^4}{1440}(-\alpha)^{-3/2}]} \beta^{-3/2}$ , where  $a = \frac{2\pi g V (2m)^{3/2}}{h^3}$ ,  $g$  is the inner degeneracy. Then calculate the grand potential  $J(T, V, \mu)$ . (non-relativistic)

- $D(\varepsilon)d\varepsilon = g \cdot \frac{V 4\pi p^2 dp}{h^3} = \frac{2\pi g V (2m)^{3/2}}{h^3} \varepsilon^{1/2} d\varepsilon = a \varepsilon^{1/2} d\varepsilon.$   
 $\ln \mathcal{Z} = \sum \omega_l \ln(1 + e^{-\alpha - \beta \varepsilon_l})$   
 $= \int_0^\infty D(\varepsilon) \ln(1 + e^{-\alpha - \beta \varepsilon}) d\varepsilon$

A9.32 Prove  $\mathcal{Z}$  for strong degenerated Fermi **gas**:

$\mathcal{Z}(\alpha, \beta, V) = e^{a[\frac{4}{15}(-\alpha)^{5/2} + \frac{\pi^2}{6}(-\alpha)^{1/2} - \frac{7\pi^4}{1440}(-\alpha)^{-3/2}]\beta^{-3/2}}$ , where  $a = \frac{2\pi g V (2m)^{3/2}}{h^3}$ ,  $g$  is the inner degeneracy. Then calculate the grand potential  $J(T, V, \mu)$ . (non-relativistic)

- $$D(\varepsilon)d\varepsilon = g \cdot \frac{V 4\pi p^2 dp}{h^3} = \frac{2\pi g V (2m)^{3/2}}{h^3} \varepsilon^{1/2} d\varepsilon = a \varepsilon^{1/2} d\varepsilon.$$

$$\ln \mathcal{Z} = \sum \omega_l \ln(1 + e^{-\alpha - \beta \varepsilon_l})$$

$$= \int_0^\infty D(\varepsilon) \ln(1 + e^{-\alpha - \beta \varepsilon}) d\varepsilon$$

$$= a \int_0^\infty \varepsilon^{1/2} \ln(1 + e^{-\alpha - \beta \varepsilon}) d\varepsilon$$

A9.32 Prove  $\mathcal{Z}$  for strong degenerated Fermi **gas**:

$\mathcal{Z}(\alpha, \beta, V) = e^{a[\frac{4}{15}(-\alpha)^{5/2} + \frac{\pi^2}{6}(-\alpha)^{1/2} - \frac{7\pi^4}{1440}(-\alpha)^{-3/2}]\beta^{-3/2}}$ , where  $a = \frac{2\pi g V (2m)^{3/2}}{h^3}$ ,  $g$  is the inner degeneracy. Then calculate the grand potential  $J(T, V, \mu)$ . (non-relativistic)

$$\begin{aligned}
 \bullet \quad D(\varepsilon)d\varepsilon &= g \cdot \frac{V 4\pi p^2 dp}{h^3} = \frac{2\pi g V (2m)^{3/2}}{h^3} \varepsilon^{1/2} d\varepsilon = a \varepsilon^{1/2} d\varepsilon. \\
 \ln \mathcal{Z} &= \sum \omega_l \ln(1 + e^{-\alpha - \beta \varepsilon_l}) \\
 &= \int_0^\infty D(\varepsilon) \ln(1 + e^{-\alpha - \beta \varepsilon}) d\varepsilon \\
 &= a \int_0^\infty \varepsilon^{1/2} \ln(1 + e^{-\alpha - \beta \varepsilon}) d\varepsilon \\
 &= a \left\{ \frac{2}{3} \varepsilon^{3/2} \ln(1 + e^{-\alpha - \beta \varepsilon}) \Big|_0^\infty + \frac{2}{3} \int_0^\infty \varepsilon^{3/2} \frac{\beta e^{-\alpha - \beta \varepsilon}}{1 + e^{-\alpha - \beta \varepsilon}} d\varepsilon \right\}
 \end{aligned}$$

A9.32 Prove  $\mathcal{Z}$  for strong degenerated Fermi **gas**:

$\mathcal{Z}(\alpha, \beta, V) = e^{a[\frac{4}{15}(-\alpha)^{5/2} + \frac{\pi^2}{6}(-\alpha)^{1/2} - \frac{7\pi^4}{1440}(-\alpha)^{-3/2}]\beta^{-3/2}}$ , where  $a = \frac{2\pi g V (2m)^{3/2}}{h^3}$ ,  $g$  is the inner degeneracy. Then calculate the grand potential  $J(T, V, \mu)$ . (non-relativistic)

$$\begin{aligned}
 \bullet \quad D(\varepsilon)d\varepsilon &= g \cdot \frac{V 4\pi p^2 dp}{h^3} = \frac{2\pi g V (2m)^{3/2}}{h^3} \varepsilon^{1/2} d\varepsilon = a \varepsilon^{1/2} d\varepsilon. \\
 \ln \mathcal{Z} &= \sum \omega_l \ln(1 + e^{-\alpha - \beta \varepsilon_l}) \\
 &= \int_0^\infty D(\varepsilon) \ln(1 + e^{-\alpha - \beta \varepsilon}) d\varepsilon \\
 &= a \int_0^\infty \varepsilon^{1/2} \ln(1 + e^{-\alpha - \beta \varepsilon}) d\varepsilon \\
 &= a \left\{ \frac{2}{3} \varepsilon^{3/2} \ln(1 + e^{-\alpha - \beta \varepsilon}) \Big|_0^\infty + \frac{2}{3} \int_0^\infty \varepsilon^{3/2} \frac{\beta e^{-\alpha - \beta \varepsilon}}{1 + e^{-\alpha - \beta \varepsilon}} d\varepsilon \right\} \\
 &\dots (\text{A grand dic of phys problems \& solutions 5, P}_{322}).
 \end{aligned}$$

A9.32 Prove  $\mathcal{Z}$  for strong degenerated Fermi gas:

$\mathcal{Z}(\alpha, \beta, V) = e^{a[\frac{4}{15}(-\alpha)^{5/2} + \frac{\pi^2}{6}(-\alpha)^{1/2} - \frac{7\pi^4}{1440}(-\alpha)^{-3/2}]} \beta^{-3/2}$ , where  $a = \frac{2\pi g V (2m)^{3/2}}{h^3}$ ,  $g$  is the inner degeneracy. Then calculate the grand potential  $J(T, V, \mu)$ . (non-relativistic)

$$\begin{aligned}
 \bullet \quad D(\varepsilon) d\varepsilon &= g \cdot \frac{V 4\pi p^2 dp}{h^3} = \frac{2\pi g V (2m)^{3/2}}{h^3} \varepsilon^{1/2} d\varepsilon = a \varepsilon^{1/2} d\varepsilon. \\
 \ln \mathcal{Z} &= \sum \omega_l \ln(1 + e^{-\alpha - \beta \varepsilon_l}) \\
 &= \int_0^\infty D(\varepsilon) \ln(1 + e^{-\alpha - \beta \varepsilon}) d\varepsilon \\
 &= a \int_0^\infty \varepsilon^{1/2} \ln(1 + e^{-\alpha - \beta \varepsilon}) d\varepsilon \\
 &= a \left\{ \frac{2}{3} \varepsilon^{3/2} \ln(1 + e^{-\alpha - \beta \varepsilon}) \Big|_0^\infty + \frac{2}{3} \int_0^\infty \varepsilon^{3/2} \frac{\beta e^{-\alpha - \beta \varepsilon}}{1 + e^{-\alpha - \beta \varepsilon}} d\varepsilon \right\} \\
 &\text{..(A grand dic of phys problems \& solutions 5, P}_{322}\text{.)} \\
 &= a \left[ \frac{4}{15} (-\alpha)^{5/2} + \frac{\pi^2}{6} (-\alpha)^{1/2} - \frac{7\pi^4}{1440} (-\alpha)^{-3/2} \right] \beta^{-3/2}.
 \end{aligned}$$

A9.32 Prove  $\mathcal{Z}$  for strong degenerated Fermi gas:

$\mathcal{Z}(\alpha, \beta, V) = e^{a[\frac{4}{15}(-\alpha)^{5/2} + \frac{\pi^2}{6}(-\alpha)^{1/2} - \frac{7\pi^4}{1440}(-\alpha)^{-3/2}]} \beta^{-3/2}$ , where  $a = \frac{2\pi g V (2m)^{3/2}}{h^3}$ ,  $g$  is the inner degeneracy. Then calculate the grand potential  $J(T, V, \mu)$ . (non-relativistic)

- $D(\varepsilon)d\varepsilon = g \cdot \frac{V 4\pi p^2 dp}{h^3} = \frac{2\pi g V (2m)^{3/2}}{h^3} \varepsilon^{1/2} d\varepsilon = a \varepsilon^{1/2} d\varepsilon.$   
 $\ln \mathcal{Z} = \sum \omega_l \ln(1 + e^{-\alpha - \beta \varepsilon_l})$   
 $= \int_0^\infty D(\varepsilon) \ln(1 + e^{-\alpha - \beta \varepsilon}) d\varepsilon$   
 $= a \int_0^\infty \varepsilon^{1/2} \ln(1 + e^{-\alpha - \beta \varepsilon}) d\varepsilon$   
 $= a \left\{ \frac{2}{3} \varepsilon^{3/2} \ln(1 + e^{-\alpha - \beta \varepsilon}) \Big|_0^\infty + \frac{2}{3} \int_0^\infty \varepsilon^{3/2} \frac{\beta e^{-\alpha - \beta \varepsilon}}{1 + e^{-\alpha - \beta \varepsilon}} d\varepsilon \right\}$   
 ..(A grand dic of phys problems & solutions 5, P<sub>322</sub>.)  
 $= a \left[ \frac{4}{15} (-\alpha)^{5/2} + \frac{\pi^2}{6} (-\alpha)^{1/2} - \frac{7\pi^4}{1440} (-\alpha)^{-3/2} \right] \beta^{-3/2}.$
- $J = -kT \ln \mathcal{Z} = -a \left( \frac{4}{15} \mu^{5/2} + \frac{\pi^2}{6\beta^2} \mu^{1/2} - \frac{7\pi^2}{1440\beta^4} \mu^{-3/2} \right).$

A9.33 Using the partition function of the strong degenerated Fermi gas, calculate  $\mu$ ,  $p$ ,  $U$ ,  $F$ ,  $S$  and  $H$ .



A9.33 Using the partition function of the strong degenerated Fermi gas, calculate  $\mu$ ,  $p$ ,  $U$ ,  $F$ ,  $S$  and  $H$ .

- We have  $D(\varepsilon) = a\varepsilon^{\frac{1}{2}}$ , where  $a = \frac{2\pi gV(2m)^{3/2}}{h^3}$ ;

A9.33 Using the partition function of the strong degenerated Fermi gas, calculate  $\mu$ ,  $p$ ,  $U$ ,  $F$ ,  $S$  and  $H$ .

- We have  $D(\varepsilon) = a\varepsilon^{\frac{1}{2}}$ , where  $a = \frac{2\pi gV(2m)^{3/2}}{h^3}$ ;  
The total number  $N = \int_0^{\mu_0} D(\varepsilon)d\varepsilon = \frac{2}{3}a\mu_0^{\frac{3}{2}}$ .

A9.33 Using the partition function of the strong degenerated Fermi gas, calculate  $\mu$ ,  $p$ ,  $U$ ,  $F$ ,  $S$  and  $H$ .

- We have  $D(\varepsilon) = a\varepsilon^{\frac{1}{2}}$ , where  $a = \frac{2\pi gV(2m)^{3/2}}{h^3}$ ;

$$\text{The total number } N = \int_0^{\mu_0} D(\varepsilon) d\varepsilon = \frac{2}{3} a \mu_0^{\frac{3}{2}}.$$

$$\Rightarrow \mu_0 = \mu(T = 0\text{K}) = \left(\frac{3N}{2a}\right)^{\frac{2}{3}} \dots\dots (1)$$

A9.33 Using the partition function of the strong degenerated Fermi gas, calculate  $\mu$ ,  $p$ ,  $U$ ,  $F$ ,  $S$  and  $H$ .

- We have  $D(\varepsilon) = a\varepsilon^{\frac{1}{2}}$ , where  $a = \frac{2\pi gV(2m)^{3/2}}{h^3}$ ;

$$\text{The total number } N = \int_0^{\mu_0} D(\varepsilon)d\varepsilon = \frac{2}{3}a\mu_0^{\frac{3}{2}}.$$

$$\Rightarrow \mu_0 = \mu(T = 0\text{K}) = \left(\frac{3N}{2a}\right)^{\frac{2}{3}} \dots\dots (1)$$

- $\ln \mathcal{Z} = a\left[\frac{4}{15}(-\alpha)^{5/2} + \frac{\pi^2}{6}(-\alpha)^{1/2} - \frac{7\pi^4}{1440}(-\alpha)^{-3/2}\right]\beta^{-3/2}$

A9.33 Using the partition function of the strong degenerated Fermi gas, calculate  $\mu$ ,  $p$ ,  $U$ ,  $F$ ,  $S$  and  $H$ .

- We have  $D(\varepsilon) = a\varepsilon^{\frac{1}{2}}$ , where  $a = \frac{2\pi gV(2m)^{3/2}}{h^3}$ ;

$$\text{The total number } N = \int_0^{\mu_0} D(\varepsilon) d\varepsilon = \frac{2}{3} a \mu_0^{\frac{3}{2}}.$$

$$\Rightarrow \mu_0 = \mu(T = 0\text{K}) = \left(\frac{3N}{2a}\right)^{\frac{2}{3}} \dots\dots (1)$$

- $\ln \mathcal{Z} = a \left[ \frac{4}{15} (-\alpha)^{5/2} + \frac{\pi^2}{6} (-\alpha)^{1/2} - \frac{7\pi^4}{1440} (-\alpha)^{-3/2} \right] \beta^{-3/2}$   
 $= \frac{2}{5} N \beta \mu_0^{-\frac{3}{2}} \mu^{\frac{5}{2}} \left[ 1 + \frac{5}{12} \left( \frac{\pi}{\beta \mu} \right)^2 - \frac{7}{384} \left( \frac{\pi}{\beta \mu} \right)^4 \right] \dots\dots (2)$

# A9.33 Using the partition function of the strong degenerated Fermi gas, calculate $\mu$ , $p$ , $U$ , $F$ , $S$ and $H$ .

- We have  $D(\varepsilon) = a\varepsilon^{\frac{1}{2}}$ , where  $a = \frac{2\pi gV(2m)^{3/2}}{h^3}$ ;

$$\text{The total number } N = \int_0^{\mu_0} D(\varepsilon)d\varepsilon = \frac{2}{3}a\mu_0^{\frac{3}{2}}.$$

$$\Rightarrow \mu_0 = \mu(T = 0\text{K}) = \left(\frac{3N}{2a}\right)^{\frac{2}{3}} \dots\dots (1)$$

- $\ln \mathcal{Z} = a\left[\frac{4}{15}(-\alpha)^{5/2} + \frac{\pi^2}{6}(-\alpha)^{1/2} - \frac{7\pi^4}{1440}(-\alpha)^{-3/2}\right]\beta^{-3/2}$

$$= \frac{2}{5}N\beta\mu_0^{-\frac{3}{2}}\mu^{\frac{5}{2}}\left[1 + \frac{5}{12}\left(\frac{\pi}{\beta\mu}\right)^2 - \frac{7}{384}\left(\frac{\pi}{\beta\mu}\right)^4\right] \dots\dots (2)$$

$$N = -\left(\frac{\partial \ln \mathcal{Z}}{\partial \alpha}\right)_{\beta, V}$$

# A9.33 Using the partition function of the strong degenerated Fermi gas, calculate $\mu$ , $p$ , $U$ , $F$ , $S$ and $H$ .

- We have  $D(\varepsilon) = a\varepsilon^{\frac{1}{2}}$ , where  $a = \frac{2\pi gV(2m)^{3/2}}{h^3}$ ;

$$\text{The total number } N = \int_0^{\mu_0} D(\varepsilon)d\varepsilon = \frac{2}{3}a\mu_0^{\frac{3}{2}}.$$

$$\Rightarrow \mu_0 = \mu(T = 0\text{K}) = \left(\frac{3N}{2a}\right)^{\frac{2}{3}} \dots\dots (1)$$

- $\ln \mathcal{Z} = a\left[\frac{4}{15}(-\alpha)^{5/2} + \frac{\pi^2}{6}(-\alpha)^{1/2} - \frac{7\pi^4}{1440}(-\alpha)^{-3/2}\right]\beta^{-3/2}$

$$= \frac{2}{5}N\beta\mu_0^{-\frac{3}{2}}\mu^{\frac{5}{2}}\left[1 + \frac{5}{12}\left(\frac{\pi}{\beta\mu}\right)^2 - \frac{7}{384}\left(\frac{\pi}{\beta\mu}\right)^4\right] \dots\dots (2)$$

$$N = -\left(\frac{\partial \ln \mathcal{Z}}{\partial \alpha}\right)_{\beta, V}$$

$$= a\left[\frac{2}{3}(-\alpha)^{3/2} + \frac{\pi^2}{12}(-\alpha)^{-1/2} - \frac{7\pi^4}{960}(-\alpha)^{-5/2}\right]\beta^{-3/2}$$

# A9.33 Using the partition function of the strong degenerated Fermi gas, calculate $\mu$ , $p$ , $U$ , $F$ , $S$ and $H$ .

- We have  $D(\varepsilon) = a\varepsilon^{\frac{1}{2}}$ , where  $a = \frac{2\pi gV(2m)^{3/2}}{h^3}$ ;

$$\text{The total number } N = \int_0^{\mu_0} D(\varepsilon)d\varepsilon = \frac{2}{3}a\mu_0^{\frac{3}{2}}.$$

$$\Rightarrow \mu_0 = \mu(T = 0\text{K}) = \left(\frac{3N}{2a}\right)^{\frac{2}{3}} \dots\dots (1)$$

- $\ln \mathcal{Z} = a\left[\frac{4}{15}(-\alpha)^{5/2} + \frac{\pi^2}{6}(-\alpha)^{1/2} - \frac{7\pi^4}{1440}(-\alpha)^{-3/2}\right]\beta^{-3/2}$

$$= \frac{2}{5}N\beta\mu_0^{-\frac{3}{2}}\mu^{\frac{5}{2}}\left[1 + \frac{5}{12}\left(\frac{\pi}{\beta\mu}\right)^2 - \frac{7}{384}\left(\frac{\pi}{\beta\mu}\right)^4\right] \dots\dots (2)$$

$$N = -\left(\frac{\partial \ln \mathcal{Z}}{\partial \alpha}\right)_{\beta, V}$$

$$= a\left[\frac{2}{3}(-\alpha)^{3/2} + \frac{\pi^2}{12}(-\alpha)^{-1/2} - \frac{7\pi^4}{960}(-\alpha)^{-5/2}\right]\beta^{-3/2}$$

$$= \frac{2}{3}a\left(\mu^{\frac{3}{2}} + \frac{\pi^2}{8\beta^2}\mu^{-\frac{1}{2}} + \frac{7\pi^2}{640\beta^4}\mu^{-\frac{5}{2}}\right) \dots\dots (3)$$



# A9.33 Using the partition function of the strong degenerated Fermi gas, calculate $\mu$ , $p$ , $U$ , $F$ , $S$ and $H$ .

- We have  $D(\varepsilon) = a\varepsilon^{\frac{1}{2}}$ , where  $a = \frac{2\pi gV(2m)^{3/2}}{h^3}$ ;

$$\text{The total number } N = \int_0^{\mu_0} D(\varepsilon)d\varepsilon = \frac{2}{3}a\mu_0^{\frac{3}{2}}.$$

$$\Rightarrow \mu_0 = \mu(T = 0\text{K}) = \left(\frac{3N}{2a}\right)^{\frac{2}{3}} \dots\dots (1)$$

- $\ln \mathcal{Z} = a\left[\frac{4}{15}(-\alpha)^{5/2} + \frac{\pi^2}{6}(-\alpha)^{1/2} - \frac{7\pi^4}{1440}(-\alpha)^{-3/2}\right]\beta^{-3/2}$

$$= \frac{2}{5}N\beta\mu_0^{-\frac{3}{2}}\mu^{\frac{5}{2}}\left[1 + \frac{5}{12}\left(\frac{\pi}{\beta\mu}\right)^2 - \frac{7}{384}\left(\frac{\pi}{\beta\mu}\right)^4\right] \dots\dots (2)$$

$$N = -\left(\frac{\partial \ln \mathcal{Z}}{\partial \alpha}\right)_{\beta, V}$$

$$= a\left[\frac{2}{3}(-\alpha)^{3/2} + \frac{\pi^2}{12}(-\alpha)^{-1/2} - \frac{7\pi^4}{960}(-\alpha)^{-5/2}\right]\beta^{-3/2}$$

$$= \frac{2}{3}a\left(\mu^{\frac{3}{2}} + \frac{\pi^2}{8\beta^2}\mu^{-\frac{1}{2}} + \frac{7\pi^2}{640\beta^4}\mu^{-\frac{5}{2}}\right) \dots\dots (3)$$

- Combine (1) and (3),  $\mu = \mu_0\left[1 + \frac{1}{8}\left(\frac{\pi}{\beta\mu}\right)^2 + \frac{7}{640}\left(\frac{\pi}{\beta\mu}\right)^4\right]^{-\frac{2}{3}}$

... For the strong degenerated Fermi gas, calculate  $\mu$ ,  $p$ ,  $U$ ,  $F$ ,  $S$  and  $H$ .

$$\ln \mathcal{Z} = \frac{2}{5} N \beta \mu_0^{-\frac{3}{2}} \mu^{\frac{5}{2}} \left[ 1 + \frac{5}{12} \left( \frac{\pi}{\beta \mu} \right)^2 - \frac{7}{384} \left( \frac{\pi}{\beta \mu} \right)^4 \right] \dots (2)$$

- $\mu = \mu_0 \left[ 1 + \frac{1}{8} \left( \frac{\pi}{\beta \mu} \right)^2 + \frac{7}{640} \left( \frac{\pi}{\beta \mu} \right)^4 \right]^{-\frac{2}{3}}.$

... For the strong degenerated Fermi gas, calculate  $\mu$ ,  $p$ ,  $U$ ,  $F$ ,  $S$  and  $H$ .

$$\ln \mathcal{Z} = \frac{2}{5} N \beta \mu_0^{-\frac{3}{2}} \mu^{\frac{5}{2}} \left[ 1 + \frac{5}{12} \left( \frac{\pi}{\beta \mu} \right)^2 - \frac{7}{384} \left( \frac{\pi}{\beta \mu} \right)^4 \right] \dots (2)$$

- $\mu = \mu_0 \left[ 1 + \frac{1}{8} \left( \frac{\pi}{\beta \mu} \right)^2 + \frac{7}{640} \left( \frac{\pi}{\beta \mu} \right)^4 \right]^{-\frac{2}{3}}.$

For the strong degenerated Fermi gas,  $kT \ll \mu_0$ ,

$\mu \simeq \mu_0$ , so,  $\beta \mu \gg 1$ ; and define  $\theta = \frac{\pi}{\beta \mu_0} = \frac{\pi kT}{\mu_0} \ll 1$ .

... For the strong degenerated Fermi gas, calculate  $\mu$ ,  $p$ ,  $U$ ,  $F$ ,  $S$  and  $H$ .

$$\ln \mathcal{Z} = \frac{2}{5} N \beta \mu_0^{-\frac{3}{2}} \mu^{\frac{5}{2}} \left[ 1 + \frac{5}{12} \left( \frac{\pi}{\beta \mu} \right)^2 - \frac{7}{384} \left( \frac{\pi}{\beta \mu} \right)^4 \right] \dots (2)$$

- $\mu = \mu_0 \left[ 1 + \frac{1}{8} \left( \frac{\pi}{\beta \mu} \right)^2 + \frac{7}{640} \left( \frac{\pi}{\beta \mu} \right)^4 \right]^{-\frac{2}{3}}.$

For the strong degenerated Fermi gas,  $kT \ll \mu_0$ ,

$\mu \simeq \mu_0$ , so,  $\beta \mu \gg 1$ ; and define  $\theta = \frac{\pi}{\beta \mu_0} = \frac{\pi kT}{\mu_0} \ll 1$ .

$$\mu \simeq \mu_0 \left( 1 - \frac{1}{12} \theta^2 - \frac{1}{80} \theta^4 \right) \dots (4)$$

... For the strong degenerated Fermi gas, calculate  $\mu$ ,  $p$ ,  $U$ ,  $F$ ,  $S$  and  $H$ .

$$\ln \mathcal{Z} = \frac{2}{5} N \beta \mu_0^{-\frac{3}{2}} \mu^{\frac{5}{2}} \left[ 1 + \frac{5}{12} \left( \frac{\pi}{\beta \mu} \right)^2 - \frac{7}{384} \left( \frac{\pi}{\beta \mu} \right)^4 \right] \dots (2)$$

- $\mu = \mu_0 \left[ 1 + \frac{1}{8} \left( \frac{\pi}{\beta \mu} \right)^2 + \frac{7}{640} \left( \frac{\pi}{\beta \mu} \right)^4 \right]^{-\frac{2}{3}}.$

For the strong degenerated Fermi gas,  $kT \ll \mu_0$ ,

$\mu \simeq \mu_0$ , so,  $\beta \mu \gg 1$ ; and define  $\theta = \frac{\pi}{\beta \mu_0} = \frac{\pi kT}{\mu_0} \ll 1$ .

$$\mu \simeq \mu_0 \left( 1 - \frac{1}{12} \theta^2 - \frac{1}{80} \theta^4 \right) \dots (4)$$

- Combine (2) and (4),

$$\ln \mathcal{Z} \simeq$$

... For the strong degenerated Fermi gas, calculate  $\mu$ ,  $p$ ,  $U$ ,  $F$ ,  $S$  and  $H$ .

$$\ln \mathcal{Z} = \frac{2}{5} N \beta \mu_0^{-\frac{3}{2}} \mu^{\frac{5}{2}} \left[ 1 + \frac{5}{12} \left( \frac{\pi}{\beta \mu} \right)^2 - \frac{7}{384} \left( \frac{\pi}{\beta \mu} \right)^4 \right] \dots (2)$$

- $\mu = \mu_0 \left[ 1 + \frac{1}{8} \left( \frac{\pi}{\beta \mu} \right)^2 + \frac{7}{640} \left( \frac{\pi}{\beta \mu} \right)^4 \right]^{-\frac{2}{3}}.$

For the strong degenerated Fermi gas,  $kT \ll \mu_0$ ,

$\mu \simeq \mu_0$ , so,  $\beta \mu \gg 1$ ; and define  $\theta = \frac{\pi}{\beta \mu_0} = \frac{\pi kT}{\mu_0} \ll 1$ .

$$\mu \simeq \mu_0 \left( 1 - \frac{1}{12} \theta^2 - \frac{1}{80} \theta^4 \right) \dots (4)$$

- Combine (2) and (4),

$$\ln \mathcal{Z} \simeq \frac{2\pi}{5} N \theta^{-1} \left( 1 + \frac{5}{12} \theta^2 - \frac{1}{16} \theta^4 \right).$$

... For the strong degenerated Fermi gas, calculate  $\mu$ ,  $p$ ,  $U$ ,  $F$ ,  $S$  and  $H$ .

$$\ln \mathcal{Z} = \frac{2}{5} N \beta \mu_0^{-\frac{3}{2}} \mu^{\frac{5}{2}} \left[ 1 + \frac{5}{12} \left( \frac{\pi}{\beta \mu} \right)^2 - \frac{7}{384} \left( \frac{\pi}{\beta \mu} \right)^4 \right] \dots (2)$$

- $\mu = \mu_0 \left[ 1 + \frac{1}{8} \left( \frac{\pi}{\beta \mu} \right)^2 + \frac{7}{640} \left( \frac{\pi}{\beta \mu} \right)^4 \right]^{-\frac{2}{3}}.$

For the strong degenerated Fermi gas,  $kT \ll \mu_0$ ,

$\mu \simeq \mu_0$ , so,  $\beta \mu \gg 1$ ; and define  $\theta = \frac{\pi}{\beta \mu_0} = \frac{\pi kT}{\mu_0} \ll 1$ .

$$\mu \simeq \mu_0 \left( 1 - \frac{1}{12} \theta^2 - \frac{1}{80} \theta^4 \right) \dots (4)$$

- Combine (2) and (4),

$$\ln \mathcal{Z} \simeq \frac{2\pi}{5} N \theta^{-1} \left( 1 + \frac{5}{12} \theta^2 - \frac{1}{16} \theta^4 \right).$$

As  $J = -pV = -kT \ln \mathcal{Z}$ ,

$$p = \frac{kT}{V} \ln \mathcal{Z} \simeq \frac{2N}{5V} \mu_0 \left( 1 + \frac{5}{12} \theta^2 - \frac{1}{16} \theta^4 \right);$$

... For the strong degenerated Fermi gas, calculate  $\mu$ ,  $p$ ,  $U$ ,  $F$ ,  $S$  and  $H$ .

$$\ln \mathcal{Z} = \frac{2}{5} N \beta \mu_0^{-\frac{3}{2}} \mu^{\frac{5}{2}} \left[ 1 + \frac{5}{12} \left( \frac{\pi}{\beta \mu} \right)^2 - \frac{7}{384} \left( \frac{\pi}{\beta \mu} \right)^4 \right] \dots (2)$$

- $\mu = \mu_0 \left[ 1 + \frac{1}{8} \left( \frac{\pi}{\beta \mu} \right)^2 + \frac{7}{640} \left( \frac{\pi}{\beta \mu} \right)^4 \right]^{-\frac{2}{3}}.$

For the strong degenerated Fermi gas,  $kT \ll \mu_0$ ,

$\mu \simeq \mu_0$ , so,  $\beta \mu \gg 1$ ; and define  $\theta = \frac{\pi}{\beta \mu_0} = \frac{\pi kT}{\mu_0} \ll 1$ .

$$\mu \simeq \mu_0 \left( 1 - \frac{1}{12} \theta^2 - \frac{1}{80} \theta^4 \right) \dots (4)$$

- Combine (2) and (4),

$$\ln \mathcal{Z} \simeq \frac{2\pi}{5} N \theta^{-1} \left( 1 + \frac{5}{12} \theta^2 - \frac{1}{16} \theta^4 \right).$$

As  $J = -pV = -kT \ln \mathcal{Z}$ ,

$$p = \frac{kT}{V} \ln \mathcal{Z} \simeq \frac{2N}{5V} \mu_0 \left( 1 + \frac{5}{12} \theta^2 - \frac{1}{16} \theta^4 \right);$$

Internal energy:  $U = \frac{3V}{2} p \simeq \frac{3}{5} N \mu_0 \left( 1 + \frac{5}{12} \theta^2 - \frac{1}{16} \theta^4 \right).$



... For the strong degenerated Fermi gas, calculate  $\mu$ ,  $p$ ,  $U$ ,  $F$ ,  $S$  and  $H$ .  $\mu \simeq \mu_0(1 - \frac{1}{12}\theta^2 - \frac{1}{80}\theta^4).....(4)$

$$p \simeq \frac{2N}{5V}\mu_0(1 + \frac{5}{12}\theta^2 - \frac{1}{16}\theta^4)$$

- Free energy:  $F = N\mu + J = N\mu - pV \simeq N\mu_0(1 - \frac{1}{12}\theta^2 - \frac{1}{80}\theta^4) - \frac{2N}{5V}\mu_0(1 + \frac{5}{12}\theta^2 - \frac{1}{16}\theta^4) \cdot V$

... For the strong degenerated Fermi gas, calculate  $\mu$ ,  $p$ ,  $U$ ,  $F$ ,  $S$  and  $H$ .  $\mu \simeq \mu_0(1 - \frac{1}{12}\theta^2 - \frac{1}{80}\theta^4).....(4)$

$$p \simeq \frac{2N}{5V}\mu_0(1 + \frac{5}{12}\theta^2 - \frac{1}{16}\theta^4)$$

- Free energy:  $F = N\mu + J = N\mu - pV \simeq N\mu_0(1 - \frac{1}{12}\theta^2 - \frac{1}{80}\theta^4) - \frac{2N}{5V}\mu_0(1 + \frac{5}{12}\theta^2 - \frac{1}{16}\theta^4) \cdot V = \frac{3N}{5}\mu_0(1 - \frac{5}{12}\theta^2 + \frac{1}{48}\theta^4).$

... For the strong degenerated Fermi gas, calculate  $\mu$ ,  $p$ ,  $U$ ,  $F$ ,  $S$  and  $H$ .  $\mu \simeq \mu_0(1 - \frac{1}{12}\theta^2 - \frac{1}{80}\theta^4).....(4)$

$$p \simeq \frac{2N}{5V}\mu_0(1 + \frac{5}{12}\theta^2 - \frac{1}{16}\theta^4)$$

- Free energy:  $F = N\mu + J = N\mu - pV \simeq N\mu_0(1 - \frac{1}{12}\theta^2 - \frac{1}{80}\theta^4) - \frac{2N}{5V}\mu_0(1 + \frac{5}{12}\theta^2 - \frac{1}{16}\theta^4) \cdot V = \frac{3N}{5}\mu_0(1 - \frac{5}{12}\theta^2 + \frac{1}{48}\theta^4)$ .
- Entropy:  $S = \frac{U-F}{T} \simeq \frac{1}{2T}N\mu_0\theta^2(1 - \frac{1}{10}\theta^2) = \frac{\pi k}{2}N\theta(1 - \frac{1}{10}\theta^2)$ .

... For the strong degenerated Fermi gas, calculate  $\mu$ ,  $p$ ,  $U$ ,  $F$ ,  $S$  and  $H$ .  $\mu \simeq \mu_0(1 - \frac{1}{12}\theta^2 - \frac{1}{80}\theta^4).....(4)$

$$p \simeq \frac{2N}{5V}\mu_0(1 + \frac{5}{12}\theta^2 - \frac{1}{16}\theta^4)$$

- Free energy:  $F = N\mu + J = N\mu - pV \simeq N\mu_0(1 - \frac{1}{12}\theta^2 - \frac{1}{80}\theta^4) - \frac{2N}{5V}\mu_0(1 + \frac{5}{12}\theta^2 - \frac{1}{16}\theta^4) \cdot V = \frac{3N}{5}\mu_0(1 - \frac{5}{12}\theta^2 + \frac{1}{48}\theta^4).$
- Entropy:  $S = \frac{U-F}{T} \simeq \frac{1}{2T}N\mu_0\theta^2(1 - \frac{1}{10}\theta^2) = \frac{\pi k}{2}N\theta(1 - \frac{1}{10}\theta^2).$
- Enthalpy:  
 $H = U + pV = \frac{5}{2}pV \simeq N\mu_0(1 + \frac{5}{12}\theta^2 - \frac{1}{16}\theta^4).$

A9.35 Calculate expansion coefficient  $\alpha$ , pressure coefficient  $\beta$ , and isothermal compressibility  $\kappa_T$ , for the strong degenerated Fermi gas.

A9.35 Calculate expansion coefficient  $\alpha$ , pressure coefficient  $\beta$ , and isothermal compressibility  $\kappa_T$ , for the strong degenerated Fermi gas.

- (Convert to  $(p, V, T)$ ) We have  $p = \frac{2}{5V} N \mu_0 (1 + \frac{5}{12} \theta^2)$ , where  $\theta = \frac{\pi}{\beta \mu_0} = \frac{\pi k T}{\mu_0}$ ,  $\mu_0 = (\frac{3N}{2a})^{\frac{2}{3}}$ ,  $a = 2\pi g V (\frac{2m}{h^2})^{\frac{3}{2}}$ . Define:  $b = \frac{h^2}{2m} (\frac{3N}{4\pi g})^{\frac{2}{3}}$ .

A9.35 Calculate expansion coefficient  $\alpha$ , pressure coefficient  $\beta$ , and isothermal compressibility  $\kappa_T$ , for the strong degenerated Fermi gas.

- (Convert to  $(p, V, T)$ ) We have  $p = \frac{2}{5V} N \mu_0 (1 + \frac{5}{12} \theta^2)$ , where  $\theta = \frac{\pi}{\beta \mu_0} = \frac{\pi k T}{\mu_0}$ ,  $\mu_0 = (\frac{3N}{2a})^{\frac{2}{3}}$ ,  $a = 2\pi g V (\frac{2m}{h^2})^{\frac{3}{2}}$ .  
Define:  $b = \frac{h^2}{2m} (\frac{3N}{4\pi g})^{\frac{2}{3}}$ .  
 $\Rightarrow p = \frac{2bN}{5} V^{-\frac{5}{3}} + \frac{\pi^2 k^2}{6b} N V^{-\frac{1}{3}} T^2$ .

A9.35 Calculate expansion coefficient  $\alpha$ , pressure coefficient  $\beta$ , and isothermal compressibility  $\kappa_T$ , for the strong degenerated Fermi gas.

- (Convert to  $(p, V, T)$ ) We have  $p = \frac{2}{5V} N \mu_0 (1 + \frac{5}{12} \theta^2)$ , where  $\theta = \frac{\pi}{\beta \mu_0} = \frac{\pi k T}{\mu_0}$ ,  $\mu_0 = (\frac{3N}{2a})^{\frac{2}{3}}$ ,  $a = 2\pi g V (\frac{2m}{h^2})^{\frac{3}{2}}$ .  
Define:  $b = \frac{h^2}{2m} (\frac{3N}{4\pi g})^{\frac{2}{3}}$ .  
 $\Rightarrow p = \frac{2bN}{5} V^{-\frac{5}{3}} + \frac{\pi^2 k^2}{6b} N V^{-\frac{1}{3}} T^2$ .
- $\alpha \equiv \frac{1}{V} (\frac{\partial V}{\partial T})_p = \frac{\pi^2 k^2}{2\mu_0^2} (1 - \frac{1}{12} \theta^2) T$ .



A9.35 Calculate expansion coefficient  $\alpha$ , pressure coefficient  $\beta$ , and isothermal compressibility  $\kappa_T$ , for the strong degenerated Fermi gas.

- (Convert to  $(p, V, T)$ ) We have  $p = \frac{2}{5V} N \mu_0 (1 + \frac{5}{12} \theta^2)$ , where  $\theta = \frac{\pi}{\beta \mu_0} = \frac{\pi k T}{\mu_0}$ ,  $\mu_0 = (\frac{3N}{2a})^{\frac{2}{3}}$ ,  $a = 2\pi g V (\frac{2m}{h^2})^{\frac{3}{2}}$ .  
Define:  $b = \frac{h^2}{2m} (\frac{3N}{4\pi g})^{\frac{2}{3}}$ .  
 $\Rightarrow p = \frac{2bN}{5} V^{-\frac{5}{3}} + \frac{\pi^2 k^2}{6b} N V^{-\frac{1}{3}} T^2$ .
- $\alpha \equiv \frac{1}{V} (\frac{\partial V}{\partial T})_p = \frac{\pi^2 k^2}{2\mu_0^2} (1 - \frac{1}{12} \theta^2) T$ .  
 $\beta \equiv \frac{1}{p} (\frac{\partial p}{\partial T})_V = \frac{5\pi^2 k^2}{6\mu_0^2} (1 - \frac{5}{12} \theta^2) T$ .

A9.35 Calculate expansion coefficient  $\alpha$ , pressure coefficient  $\beta$ , and isothermal compressibility  $\kappa_T$ , for the strong degenerated Fermi gas.

- (Convert to  $(p, V, T)$ ) We have  $p = \frac{2}{5V} N \mu_0 (1 + \frac{5}{12} \theta^2)$ , where  $\theta = \frac{\pi}{\beta \mu_0} = \frac{\pi k T}{\mu_0}$ ,  $\mu_0 = (\frac{3N}{2a})^{\frac{2}{3}}$ ,  $a = 2\pi g V (\frac{2m}{h^2})^{\frac{3}{2}}$ .  
Define:  $b = \frac{h^2}{2m} (\frac{3N}{4\pi g})^{\frac{2}{3}}$ .  
 $\Rightarrow p = \frac{2bN}{5} V^{-\frac{5}{3}} + \frac{\pi^2 k^2}{6b} N V^{-\frac{1}{3}} T^2$ .
- $\alpha \equiv \frac{1}{V} (\frac{\partial V}{\partial T})_p = \frac{\pi^2 k^2}{2\mu_0^2} (1 - \frac{1}{12} \theta^2) T$ .  
 $\beta \equiv \frac{1}{p} (\frac{\partial p}{\partial T})_V = \frac{5\pi^2 k^2}{6\mu_0^2} (1 - \frac{5}{12} \theta^2) T$ .  
 $\kappa_T \equiv -\frac{1}{V} (\frac{\partial V}{\partial p})_T = \frac{3}{2n\mu_0} (1 - \frac{1}{12} \theta^2)$ .

# A9.36 Adiabatic process of strong degenerated Fermi gas.

## A9.36 Adiabatic process of strong degenerated Fermi gas.

- We have  $S = \frac{\pi k}{2} N \theta (1 - \frac{1}{10} \theta^2) \simeq \frac{\pi k}{2} N \theta = \frac{\pi^2 k^2}{2b} N T V^{\frac{2}{3}}$ ,  
where  $\theta = \frac{\pi}{\beta \mu_0} = \frac{\pi k T}{\mu_0}$ ,  $\mu_0 = (\frac{3N}{2a})^{\frac{2}{3}}$ ,  $a = 2\pi g V (\frac{2m}{h^2})^{\frac{3}{2}}$ ,  
 $b = \frac{h^2}{2m} (\frac{3N}{4\pi g})^{\frac{2}{3}}$ .

## A9.36 Adiabatic process of strong degenerated Fermi gas.

- We have  $S = \frac{\pi k}{2} N \theta (1 - \frac{1}{10} \theta^2) \simeq \frac{\pi k}{2} N \theta = \frac{\pi^2 k^2}{2b} N T V^{\frac{2}{3}}$ ,  
where  $\theta = \frac{\pi}{\beta \mu_0} = \frac{\pi k T}{\mu_0}$ ,  $\mu_0 = (\frac{3N}{2a})^{\frac{2}{3}}$ ,  $a = 2\pi g V (\frac{2m}{h^2})^{\frac{3}{2}}$ ,  
 $b = \frac{h^2}{2m} (\frac{3N}{4\pi g})^{\frac{2}{3}}$ .
- Reversible adiabatic process is in equal entropy.

## A9.36 Adiabatic process of strong degenerated Fermi gas.

- We have  $S = \frac{\pi k}{2} N \theta (1 - \frac{1}{10} \theta^2) \simeq \frac{\pi k}{2} N \theta = \frac{\pi^2 k^2}{2b} N T V^{\frac{2}{3}}$ ,  
where  $\theta = \frac{\pi}{\beta \mu_0} = \frac{\pi k T}{\mu_0}$ ,  $\mu_0 = (\frac{3N}{2a})^{\frac{2}{3}}$ ,  $a = 2\pi g V (\frac{2m}{h^2})^{\frac{3}{2}}$ ,  
 $b = \frac{h^2}{2m} (\frac{3N}{4\pi g})^{\frac{2}{3}}$ .
- Reversible adiabatic process is in equal entropy.  
 $\Rightarrow TV^{\frac{2}{3}} = \text{Const.}$ , or  $\theta = \text{Const.}$

## A9.36 Adiabatic process of strong degenerated Fermi gas.

- We have  $S = \frac{\pi k}{2} N \theta (1 - \frac{1}{10} \theta^2) \simeq \frac{\pi k}{2} N \theta = \frac{\pi^2 k^2}{2b} N T V^{\frac{2}{3}}$ ,  
where  $\theta = \frac{\pi}{\beta \mu_0} = \frac{\pi k T}{\mu_0}$ ,  $\mu_0 = (\frac{3N}{2a})^{\frac{2}{3}}$ ,  $a = 2\pi g V (\frac{2m}{h^2})^{\frac{3}{2}}$ ,  
 $b = \frac{h^2}{2m} (\frac{3N}{4\pi g})^{\frac{2}{3}}$ .
- Reversible adiabatic process is in equal entropy.  
 $\Rightarrow TV^{\frac{2}{3}} = \text{Const.}$ , or  $\theta = \text{Const.}$
- $p = \frac{2N}{5V} \mu_0 (1 + \frac{5}{12} \theta^2 - \frac{1}{16} \theta^4)$ ,

## A9.36 Adiabatic process of strong degenerated Fermi gas.

- We have  $S = \frac{\pi k}{2} N \theta (1 - \frac{1}{10} \theta^2) \simeq \frac{\pi k}{2} N \theta = \frac{\pi^2 k^2}{2b} N T V^{\frac{2}{3}}$ ,  
 where  $\theta = \frac{\pi}{\beta \mu_0} = \frac{\pi k T}{\mu_0}$ ,  $\mu_0 = (\frac{3N}{2a})^{\frac{2}{3}}$ ,  $a = 2\pi g V (\frac{2m}{h^2})^{\frac{3}{2}}$ ,  
 $b = \frac{h^2}{2m} (\frac{3N}{4\pi g})^{\frac{2}{3}}$ .
- Reversible adiabatic process is in equal entropy.  
 $\Rightarrow TV^{\frac{2}{3}} = \text{Const.}, \text{ or } \theta = \text{Const.}$
- $p = \frac{2N}{5V} \mu_0 (1 + \frac{5}{12} \theta^2 - \frac{1}{16} \theta^4)$ ,  
 $\Rightarrow p \propto \frac{2N}{5V} \mu_0 \propto V^{-5/3}$ .



## A9.36 Adiabatic process of strong degenerated Fermi gas.

- We have  $S = \frac{\pi k}{2} N \theta (1 - \frac{1}{10} \theta^2) \simeq \frac{\pi k}{2} N \theta = \frac{\pi^2 k^2}{2b} N T V^{\frac{2}{3}}$ ,  
where  $\theta = \frac{\pi}{\beta \mu_0} = \frac{\pi k T}{\mu_0}$ ,  $\mu_0 = (\frac{3N}{2a})^{\frac{2}{3}}$ ,  $a = 2\pi g V (\frac{2m}{h^2})^{\frac{3}{2}}$ ,  
 $b = \frac{h^2}{2m} (\frac{3N}{4\pi g})^{\frac{2}{3}}$ .

- Reversible adiabatic process is in equal entropy.

$$\Rightarrow T V^{\frac{2}{3}} = \text{Const.}, \text{ or } \theta = \text{Const.}$$

- $p = \frac{2N}{5V} \mu_0 (1 + \frac{5}{12} \theta^2 - \frac{1}{16} \theta^4)$ ,  
 $\Rightarrow p \propto \frac{2N}{5V} \mu_0 \propto V^{-5/3}$ .

Comparing with ideal gas:  $pV^\gamma = \text{Const.}$  ( $\gamma = 5/3$ ,  
monatomic molecular) (1.8.4)

## A9.36 Adiabatic process of strong degenerated Fermi gas.

- We have  $S = \frac{\pi k}{2} N \theta (1 - \frac{1}{10} \theta^2) \simeq \frac{\pi k}{2} N \theta = \frac{\pi^2 k^2}{2b} N T V^{\frac{2}{3}}$ ,  
where  $\theta = \frac{\pi}{\beta \mu_0} = \frac{\pi k T}{\mu_0}$ ,  $\mu_0 = (\frac{3N}{2a})^{\frac{2}{3}}$ ,  $a = 2\pi g V (\frac{2m}{h^2})^{\frac{3}{2}}$ ,  
 $b = \frac{h^2}{2m} (\frac{3N}{4\pi g})^{\frac{2}{3}}$ .

- Reversible adiabatic process is in equal entropy.

$$\Rightarrow T V^{\frac{2}{3}} = \text{Const.}, \text{ or } \theta = \text{Const.}$$

- $p = \frac{2N}{5V} \mu_0 (1 + \frac{5}{12} \theta^2 - \frac{1}{16} \theta^4)$ ,  
 $\Rightarrow p \propto \frac{2N}{5V} \mu_0 \propto V^{-5/3}$ .

Comparing with ideal gas:  $pV^\gamma = \text{Const.}$  ( $\gamma = 5/3$ ,  
monatomic molecular) (1.8.4)

- Using these two,  $p \propto T^{5/2}$ .

A9.43 Calculate the grand partition function  $\mathcal{Z}$  and the grand potential  $J$  for **relativistic** strong degenerated Fermion gas.

A9.43 Calculate the grand partition function  $\mathcal{Z}$  and the grand potential  $J$  for **relativistic** strong degenerated Fermion gas.

- $D(\varepsilon)d\varepsilon = g \cdot \frac{V4\pi p^2 dp}{h^3} = \frac{4\pi gV}{(hc)^3} \varepsilon^2 d\varepsilon = a\varepsilon^2 d\varepsilon,$   
where  $a = \frac{4\pi gV}{(hc)^3}.$

A9.43 Calculate the grand partition function  $\mathcal{Z}$  and the grand potential  $J$  for **relativistic** strong degenerated Fermion gas.

- $D(\varepsilon)d\varepsilon = g \cdot \frac{V4\pi p^2 dp}{h^3} = \frac{4\pi gV}{(hc)^3} \varepsilon^2 d\varepsilon = a\varepsilon^2 d\varepsilon,$   
where  $a = \frac{4\pi gV}{(hc)^3}.$   
 $\ln \mathcal{Z} = \sum \omega_l \ln(1 + e^{-\alpha - \beta \varepsilon_l})$

A9.43 Calculate the grand partition function  $\mathcal{Z}$  and the grand potential  $J$  for **relativistic** strong degenerated Fermion gas.

- $D(\varepsilon)d\varepsilon = g \cdot \frac{V4\pi p^2 dp}{h^3} = \frac{4\pi gV}{(hc)^3} \varepsilon^2 d\varepsilon = a\varepsilon^2 d\varepsilon,$

where  $a = \frac{4\pi gV}{(hc)^3}.$

$$\begin{aligned}\ln \mathcal{Z} &= \sum \omega_l \ln(1 + e^{-\alpha - \beta \varepsilon_l}) \\ &= \int_0^\infty D(\varepsilon) \ln(1 + e^{-\alpha - \beta \varepsilon}) d\varepsilon\end{aligned}$$

A9.43 Calculate the grand partition function  $\mathcal{Z}$  and the grand potential  $J$  for **relativistic** strong degenerated Fermion gas.

- $D(\varepsilon)d\varepsilon = g \cdot \frac{V4\pi p^2 dp}{h^3} = \frac{4\pi gV}{(hc)^3} \varepsilon^2 d\varepsilon = a\varepsilon^2 d\varepsilon,$

where  $a = \frac{4\pi gV}{(hc)^3}.$

$$\begin{aligned}\ln \mathcal{Z} &= \sum \omega_l \ln(1 + e^{-\alpha - \beta \varepsilon_l}) \\ &= \int_0^\infty D(\varepsilon) \ln(1 + e^{-\alpha - \beta \varepsilon}) d\varepsilon \\ &= a \int_0^\infty \varepsilon^2 \ln(1 + e^{-\alpha - \beta \varepsilon}) d\varepsilon\end{aligned}$$

A9.43 Calculate the grand partition function  $\mathcal{Z}$  and the grand potential  $J$  for **relativistic** strong degenerated Fermion gas.

- $D(\varepsilon)d\varepsilon = g \cdot \frac{V4\pi p^2 dp}{h^3} = \frac{4\pi gV}{(hc)^3} \varepsilon^2 d\varepsilon = a\varepsilon^2 d\varepsilon,$

where  $a = \frac{4\pi gV}{(hc)^3}.$

$$\ln \mathcal{Z} = \sum \omega_l \ln(1 + e^{-\alpha - \beta \varepsilon_l})$$

$$= \int_0^\infty D(\varepsilon) \ln(1 + e^{-\alpha - \beta \varepsilon}) d\varepsilon$$

$$= a \int_0^\infty \varepsilon^2 \ln(1 + e^{-\alpha - \beta \varepsilon}) d\varepsilon$$

$$= a \left\{ \frac{1}{3} \varepsilon^3 \ln(1 + e^{-\alpha - \beta \varepsilon}) \Big|_0^\infty + \frac{1}{3} \int_0^\infty \varepsilon^3 \frac{\beta e^{-\alpha - \beta \varepsilon}}{1 + e^{-\alpha - \beta \varepsilon}} d\varepsilon \right\}$$



A9.43 Calculate the grand partition function  $\mathcal{Z}$  and the grand potential  $J$  for **relativistic** strong degenerated Fermion gas.

$$\bullet D(\varepsilon)d\varepsilon = g \cdot \frac{V4\pi p^2 dp}{h^3} = \frac{4\pi gV}{(hc)^3} \varepsilon^2 d\varepsilon = a\varepsilon^2 d\varepsilon,$$

$$\text{where } a = \frac{4\pi gV}{(hc)^3}.$$

$$\ln \mathcal{Z} = \sum \omega_l \ln(1 + e^{-\alpha - \beta \varepsilon_l})$$

$$= \int_0^\infty D(\varepsilon) \ln(1 + e^{-\alpha - \beta \varepsilon}) d\varepsilon$$

$$= a \int_0^\infty \varepsilon^2 \ln(1 + e^{-\alpha - \beta \varepsilon}) d\varepsilon$$

$$= a \left\{ \frac{1}{3} \varepsilon^3 \ln(1 + e^{-\alpha - \beta \varepsilon}) \Big|_0^\infty + \frac{1}{3} \int_0^\infty \varepsilon^3 \frac{\beta e^{-\alpha - \beta \varepsilon}}{1 + e^{-\alpha - \beta \varepsilon}} d\varepsilon \right\}$$

..(A grand dic of phys problems & solutions 5, P<sub>332</sub>.)

A9.43 Calculate the grand partition function  $\mathcal{Z}$  and the grand potential  $J$  for **relativistic** strong degenerated Fermion gas.

$$\bullet D(\varepsilon)d\varepsilon = g \cdot \frac{V4\pi p^2 dp}{h^3} = \frac{4\pi gV}{(hc)^3} \varepsilon^2 d\varepsilon = a\varepsilon^2 d\varepsilon,$$

$$\text{where } a = \frac{4\pi gV}{(hc)^3}.$$

$$\ln \mathcal{Z} = \sum \omega_l \ln(1 + e^{-\alpha - \beta \varepsilon_l})$$

$$= \int_0^\infty D(\varepsilon) \ln(1 + e^{-\alpha - \beta \varepsilon}) d\varepsilon$$

$$= a \int_0^\infty \varepsilon^2 \ln(1 + e^{-\alpha - \beta \varepsilon}) d\varepsilon$$

$$= a \left\{ \frac{1}{3} \varepsilon^3 \ln(1 + e^{-\alpha - \beta \varepsilon}) \Big|_0^\infty + \frac{1}{3} \int_0^\infty \varepsilon^3 \frac{\beta e^{-\alpha - \beta \varepsilon}}{1 + e^{-\alpha - \beta \varepsilon}} d\varepsilon \right\}$$

..(A grand dic of phys problems & solutions 5, P<sub>332</sub>.)

$$= a \left[ \frac{1}{12} (-\alpha)^4 + \frac{\pi^2}{6} (-\alpha)^2 - \frac{7\pi^4}{180} \right] \beta^{-3}.$$

A9.43 Calculate the grand partition function  $\mathcal{Z}$  and the grand potential  $J$  for **relativistic** strong degenerated Fermion gas.

$$\bullet D(\varepsilon)d\varepsilon = g \cdot \frac{V4\pi p^2 dp}{h^3} = \frac{4\pi gV}{(hc)^3} \varepsilon^2 d\varepsilon = a\varepsilon^2 d\varepsilon,$$

$$\text{where } a = \frac{4\pi gV}{(hc)^3}.$$

$$\ln \mathcal{Z} = \sum \omega_l \ln(1 + e^{-\alpha - \beta \varepsilon_l})$$

$$= \int_0^\infty D(\varepsilon) \ln(1 + e^{-\alpha - \beta \varepsilon}) d\varepsilon$$

$$= a \int_0^\infty \varepsilon^2 \ln(1 + e^{-\alpha - \beta \varepsilon}) d\varepsilon$$

$$= a \left\{ \frac{1}{3} \varepsilon^3 \ln(1 + e^{-\alpha - \beta \varepsilon}) \Big|_0^\infty + \frac{1}{3} \int_0^\infty \varepsilon^3 \frac{\beta e^{-\alpha - \beta \varepsilon}}{1 + e^{-\alpha - \beta \varepsilon}} d\varepsilon \right\}$$

..(A grand dic of phys problems & solutions 5, P<sub>332</sub>.)

$$= a \left[ \frac{1}{12} (-\alpha)^4 + \frac{\pi^2}{6} (-\alpha)^2 - \frac{7\pi^4}{180} \right] \beta^{-3}.$$

$$\bullet J = -kT \ln \mathcal{Z} = -a \left( \frac{1}{12} \mu^4 + \frac{\pi^2}{6\beta^2} \mu^2 + \frac{7\pi^2}{180\beta^4} \right).$$

# Table of contents

- ① §6. Most probable distribution of nearly independent particles
- ② §7. Boltzmann statistics
- ③ §8. Bose statistics and Fermi statistics
  - Bose statistics
  - Fermi statistics
  - Fermion gas in astrophysics

A9.55 He white dwarf in  $T_{\text{WD}} = 10^7 \text{K}$  at center.

$M_{\text{WD}} \sim 1M_{\odot} \simeq 2 \times 10^{30} \text{kg}$ ,  $R_{\text{WD}} \sim 1R_{\oplus} \simeq 6 \times 10^6 \text{m}$ .

A9.55 He white dwarf in  $T_{\text{WD}} = 10^7 \text{K}$  at center.

$M_{\text{WD}} \sim 1M_{\odot} \simeq 2 \times 10^{30} \text{kg}$ ,  $R_{\text{WD}} \sim 1R_{\oplus} \simeq 6 \times 10^6 \text{m}$ .

- $kT \simeq 862.5 \text{eV}$ , ionized.

A9.55 He white dwarf in  $T_{\text{WD}} = 10^7 \text{K}$  at center.

$M_{\text{WD}} \sim 1M_{\odot} \simeq 2 \times 10^{30} \text{kg}$ ,  $R_{\text{WD}} \sim 1R_{\oplus} \simeq 6 \times 10^6 \text{m}$ .

- $kT \simeq 862.5 \text{eV}$ , ionized.
- Number density:  $n_e = 2 \cdot \frac{M_{\text{WD}}}{\frac{4}{3}\pi R_{\text{WD}}^3 \cdot m_{\text{He}}}$

A9.55 He white dwarf in  $T_{\text{WD}} = 10^7 \text{K}$  at center.

$M_{\text{WD}} \sim 1M_{\odot} \simeq 2 \times 10^{30} \text{kg}$ ,  $R_{\text{WD}} \sim 1R_{\oplus} \simeq 6 \times 10^6 \text{m}$ .

- $kT \simeq 862.5 \text{eV}$ , ionized.
- Number density:  $n_e = 2 \cdot \frac{M_{\text{WD}}}{\frac{4}{3}\pi R_{\text{WD}}^3 \cdot m_{\text{He}}}$   
 $\simeq \frac{2.2 \times 10^{30} \text{kg}}{\frac{4}{3}\pi (6 \times 10^6 \text{m})^3 \cdot 4 \times 1.67 \times 10^{-27} \text{kg}}$



A9.55 He white dwarf in  $T_{\text{WD}} = 10^7 \text{K}$  at center.

$M_{\text{WD}} \sim 1M_{\odot} \simeq 2 \times 10^{30} \text{kg}$ ,  $R_{\text{WD}} \sim 1R_{\oplus} \simeq 6 \times 10^6 \text{m}$ .

- $kT \simeq 862.5 \text{eV}$ , ionized.
- Number density:  $n_e = 2 \cdot \frac{M_{\text{WD}}}{\frac{4}{3}\pi R_{\text{WD}}^3 \cdot m_{\text{He}}}$   
 $\simeq \frac{2.2 \times 10^{30} \text{kg}}{\frac{4}{3}\pi (6 \times 10^6 \text{m})^3 \cdot 4 \times 1.67 \times 10^{-27} \text{kg}} \simeq 6.6 \times 10^{35} \text{m}^{-3}.$

A9.55 He white dwarf in  $T_{\text{WD}} = 10^7 \text{K}$  at center.

$M_{\text{WD}} \sim 1M_{\odot} \simeq 2 \times 10^{30} \text{kg}$ ,  $R_{\text{WD}} \sim 1R_{\oplus} \simeq 6 \times 10^6 \text{m}$ .

- $kT \simeq 862.5 \text{eV}$ , ionized.
- Number density:  $n_e = 2 \cdot \frac{M_{\text{WD}}}{\frac{4}{3}\pi R_{\text{WD}}^3 \cdot m_{\text{He}}}$   
 $\simeq \frac{2.2 \times 10^{30} \text{kg}}{\frac{4}{3}\pi (6 \times 10^6 \text{m})^3 \cdot 4 \times 1.67 \times 10^{-27} \text{kg}} \simeq 6.6 \times 10^{35} \text{m}^{-3}$ .
- i. Non-relativistic:  $n\lambda^3 = n\left(\frac{h^2}{2\pi m kT}\right)^{\frac{3}{2}}$

A9.55 He white dwarf in  $T_{\text{WD}} = 10^7 \text{K}$  at center.

$M_{\text{WD}} \sim 1M_{\odot} \simeq 2 \times 10^{30} \text{kg}$ ,  $R_{\text{WD}} \sim 1R_{\oplus} \simeq 6 \times 10^6 \text{m}$ .

- $kT \simeq 862.5 \text{eV}$ , ionized.
- Number density:  $n_e = 2 \cdot \frac{M_{\text{WD}}}{\frac{4}{3}\pi R_{\text{WD}}^3 \cdot m_{\text{He}}}$   
 $\simeq \frac{2.2 \times 10^{30} \text{kg}}{\frac{4}{3}\pi (6 \times 10^6 \text{m})^3 \cdot 4 \times 1.67 \times 10^{-27} \text{kg}} \simeq 6.6 \times 10^{35} \text{m}^{-3}$ .
- i. Non-relativistic:  $n\lambda^3 = n \left( \frac{h^2}{2\pi m k T} \right)^{\frac{3}{2}}$   
 $= 6.6 \times 10^{35} \text{m}^{-3} \left( \frac{(6.626 \times 10^{-34} \text{J}\cdot\text{s})^2}{2\pi \cdot 9.1 \times 10^{-31} \text{kg} \cdot 1.38 \times 10^{-23} \text{J/K} \cdot 10^7 \text{K}} \right)^{\frac{3}{2}}$

A9.55 He white dwarf in  $T_{\text{WD}} = 10^7 \text{K}$  at center.

$M_{\text{WD}} \sim 1M_{\odot} \simeq 2 \times 10^{30} \text{kg}$ ,  $R_{\text{WD}} \sim 1R_{\oplus} \simeq 6 \times 10^6 \text{m}$ .

- $kT \simeq 862.5 \text{eV}$ , ionized.
- Number density:  $n_e = 2 \cdot \frac{M_{\text{WD}}}{\frac{4}{3}\pi R_{\text{WD}}^3 \cdot m_{\text{He}}}$   
 $\simeq \frac{2.2 \times 10^{30} \text{kg}}{\frac{4}{3}\pi (6 \times 10^6 \text{m})^3 \cdot 4 \times 1.67 \times 10^{-27} \text{kg}} \simeq 6.6 \times 10^{35} \text{m}^{-3}$ .
- i. Non-relativistic:  $n\lambda^3 = n \left( \frac{h^2}{2\pi m k T} \right)^{\frac{3}{2}}$   
 $= 6.6 \times 10^{35} \text{m}^{-3} \left( \frac{(6.626 \times 10^{-34} \text{J}\cdot\text{s})^2}{2\pi \cdot 9.1 \times 10^{-31} \text{kg} \cdot 1.38 \times 10^{-23} \text{J/K} \cdot 10^7 \text{K}} \right)^{\frac{3}{2}}$   
 $\simeq 8660 \gg 1$ .

A9.55 He white dwarf in  $T_{\text{WD}} = 10^7 \text{K}$  at center.

$M_{\text{WD}} \sim 1M_{\odot} \simeq 2 \times 10^{30} \text{kg}$ ,  $R_{\text{WD}} \sim 1R_{\oplus} \simeq 6 \times 10^6 \text{m}$ .

- $kT \simeq 862.5 \text{eV}$ , ionized.
- Number density:  $n_e = 2 \cdot \frac{M_{\text{WD}}}{\frac{4}{3}\pi R_{\text{WD}}^3 \cdot m_{\text{He}}}$   
 $\simeq \frac{2.2 \times 10^{30} \text{kg}}{\frac{4}{3}\pi (6 \times 10^6 \text{m})^3 \cdot 4 \times 1.67 \times 10^{-27} \text{kg}} \simeq 6.6 \times 10^{35} \text{m}^{-3}$ .
- i. Non-relativistic:  $n\lambda^3 = n \left( \frac{h^2}{2\pi m k T} \right)^{\frac{3}{2}}$   
 $= 6.6 \times 10^{35} \text{m}^{-3} \left( \frac{(6.626 \times 10^{-34} \text{J}\cdot\text{s})^2}{2\pi \cdot 9.1 \times 10^{-31} \text{kg} \cdot 1.38 \times 10^{-23} \text{J/K} \cdot 10^7 \text{K}} \right)^{\frac{3}{2}}$   
 $\simeq 8660 \gg 1$ .
- ii. Relativistic:  $n\lambda^3 = n \left( \frac{h}{p} \right)^3$

A9.55 He white dwarf in  $T_{\text{WD}} = 10^7 \text{K}$  at center.

$M_{\text{WD}} \sim 1M_{\odot} \simeq 2 \times 10^{30} \text{kg}$ ,  $R_{\text{WD}} \sim 1R_{\oplus} \simeq 6 \times 10^6 \text{m}$ .

- $kT \simeq 862.5 \text{eV}$ , ionized.
- Number density:  $n_e = 2 \cdot \frac{M_{\text{WD}}}{\frac{4}{3}\pi R_{\text{WD}}^3 \cdot m_{\text{He}}}$   
 $\simeq \frac{2.2 \times 10^{30} \text{kg}}{\frac{4}{3}\pi (6 \times 10^6 \text{m})^3 \cdot 4 \times 1.67 \times 10^{-27} \text{kg}} \simeq 6.6 \times 10^{35} \text{m}^{-3}$ .
- i. Non-relativistic:  $n\lambda^3 = n\left(\frac{h^2}{2\pi m kT}\right)^{\frac{3}{2}}$   
 $= 6.6 \times 10^{35} \text{m}^{-3} \left( \frac{(6.626 \times 10^{-34} \text{J}\cdot\text{s})^2}{2\pi \cdot 9.1 \times 10^{-31} \text{kg} \cdot 1.38 \times 10^{-23} \text{J/K} \cdot 10^7 \text{K}} \right)^{\frac{3}{2}}$   
 $\simeq 8660 \gg 1$ .
- ii. Relativistic:  $n\lambda^3 = n\left(\frac{h}{p}\right)^3 = n\left(\frac{hc}{\varepsilon}\right)^3$

A9.55 He white dwarf in  $T_{\text{WD}} = 10^7 \text{K}$  at center.

$M_{\text{WD}} \sim 1M_{\odot} \simeq 2 \times 10^{30} \text{kg}$ ,  $R_{\text{WD}} \sim 1R_{\oplus} \simeq 6 \times 10^6 \text{m}$ .

- $kT \simeq 862.5 \text{eV}$ , ionized.
- Number density:  $n_e = 2 \cdot \frac{M_{\text{WD}}}{\frac{4}{3}\pi R_{\text{WD}}^3 \cdot m_{\text{He}}}$   
 $\simeq \frac{2.2 \times 10^{30} \text{kg}}{\frac{4}{3}\pi (6 \times 10^6 \text{m})^3 \cdot 4 \times 1.67 \times 10^{-27} \text{kg}} \simeq 6.6 \times 10^{35} \text{m}^{-3}$ .
- i. Non-relativistic:  $n\lambda^3 = n\left(\frac{h^2}{2\pi m kT}\right)^{\frac{3}{2}}$   
 $= 6.6 \times 10^{35} \text{m}^{-3} \left( \frac{(6.626 \times 10^{-34} \text{J}\cdot\text{s})^2}{2\pi \cdot 9.1 \times 10^{-31} \text{kg} \cdot 1.38 \times 10^{-23} \text{J/K} \cdot 10^7 \text{K}} \right)^{\frac{3}{2}}$   
 $\simeq 8660 \gg 1$ .
- ii. Relativistic:  $n\lambda^3 = n\left(\frac{h}{p}\right)^3 = n\left(\frac{hc}{\varepsilon}\right)^3 = n\left(\frac{hc}{kT_{\text{WD}}}\right)^3$

A9.55 He white dwarf in  $T_{\text{WD}} = 10^7 \text{K}$  at center.

$M_{\text{WD}} \sim 1M_{\odot} \simeq 2 \times 10^{30} \text{kg}$ ,  $R_{\text{WD}} \sim 1R_{\oplus} \simeq 6 \times 10^6 \text{m}$ .

- $kT \simeq 862.5 \text{eV}$ , ionized.
- Number density:  $n_e = 2 \cdot \frac{M_{\text{WD}}}{\frac{4}{3}\pi R_{\text{WD}}^3 \cdot m_{\text{He}}}$   
 $\simeq \frac{2.2 \times 10^{30} \text{kg}}{\frac{4}{3}\pi (6 \times 10^6 \text{m})^3 \cdot 4 \times 1.67 \times 10^{-27} \text{kg}} \simeq 6.6 \times 10^{35} \text{m}^{-3}$ .
- i. Non-relativistic:  $n\lambda^3 = n\left(\frac{h^2}{2\pi m kT}\right)^{\frac{3}{2}}$   
 $= 6.6 \times 10^{35} \text{m}^{-3} \left( \frac{(6.626 \times 10^{-34} \text{J}\cdot\text{s})^2}{2\pi \cdot 9.1 \times 10^{-31} \text{kg} \cdot 1.38 \times 10^{-23} \text{J/K} \cdot 10^7 \text{K}} \right)^{\frac{3}{2}}$   
 $\simeq 8660 \gg 1$ .
- ii. Relativistic:  $n\lambda^3 = n\left(\frac{h}{p}\right)^3 = n\left(\frac{hc}{\varepsilon}\right)^3 = n\left(\frac{hc}{kT_{\text{WD}}}\right)^3$   
 $\simeq 6.6 \times 10^{35} \text{m}^{-3} \left( \frac{6.626 \times 10^{-34} \text{J}\cdot\text{s} \cdot 3 \times 10^8 \text{m/s}}{1.38 \times 10^{-23} \text{J/K} \cdot 10^7 \text{K}} \right)^3$



A9.55 He white dwarf in  $T_{\text{WD}} = 10^7 \text{K}$  at center.

$M_{\text{WD}} \sim 1M_{\odot} \simeq 2 \times 10^{30} \text{kg}$ ,  $R_{\text{WD}} \sim 1R_{\oplus} \simeq 6 \times 10^6 \text{m}$ .

- $kT \simeq 862.5 \text{eV}$ , ionized.
- Number density:  $n_e = 2 \cdot \frac{M_{\text{WD}}}{\frac{4}{3}\pi R_{\text{WD}}^3 \cdot m_{\text{He}}}$   
 $\simeq \frac{2.2 \times 10^{30} \text{kg}}{\frac{4}{3}\pi (6 \times 10^6 \text{m})^3 \cdot 4 \times 1.67 \times 10^{-27} \text{kg}} \simeq 6.6 \times 10^{35} \text{m}^{-3}$ .
- i. Non-relativistic:  $n\lambda^3 = n\left(\frac{h^2}{2\pi m kT}\right)^{\frac{3}{2}}$   
 $= 6.6 \times 10^{35} \text{m}^{-3} \left( \frac{(6.626 \times 10^{-34} \text{J}\cdot\text{s})^2}{2\pi \cdot 9.1 \times 10^{-31} \text{kg} \cdot 1.38 \times 10^{-23} \text{J/K} \cdot 10^7 \text{K}} \right)^{\frac{3}{2}}$   
 $\simeq 8660 \gg 1$ .
- ii. Relativistic:  $n\lambda^3 = n\left(\frac{h}{p}\right)^3 = n\left(\frac{hc}{\varepsilon}\right)^3 = n\left(\frac{hc}{kT_{\text{WD}}}\right)^3$   
 $\simeq 6.6 \times 10^{35} \text{m}^{-3} \left( \frac{6.626 \times 10^{-34} \text{J}\cdot\text{s} \cdot 3 \times 10^8 \text{m/s}}{1.38 \times 10^{-23} \text{J/K} \cdot 10^7 \text{K}} \right)^3$   
 $\simeq 2 \times 10^9 \gg 1$ . (Only estimation if using  $kT_{\text{WD}}$ ).

A9.55 He white dwarf in  $T_{\text{WD}} = 10^7 \text{K}$  at center.

$M_{\text{WD}} \sim 1M_{\odot} \simeq 2 \times 10^{30} \text{kg}$ ,  $R_{\text{WD}} \sim 1R_{\oplus} \simeq 6 \times 10^6 \text{m}$ .

- $kT \simeq 862.5 \text{eV}$ , ionized.
- Number density:  $n_e = 2 \cdot \frac{M_{\text{WD}}}{\frac{4}{3}\pi R_{\text{WD}}^3 \cdot m_{\text{He}}}$   
 $\simeq \frac{2.2 \times 10^{30} \text{kg}}{\frac{4}{3}\pi (6 \times 10^6 \text{m})^3 \cdot 4 \times 1.67 \times 10^{-27} \text{kg}} \simeq 6.6 \times 10^{35} \text{m}^{-3}$ .
- i. Non-relativistic:  $n\lambda^3 = n\left(\frac{h^2}{2\pi m kT}\right)^{\frac{3}{2}}$   
 $= 6.6 \times 10^{35} \text{m}^{-3} \left( \frac{(6.626 \times 10^{-34} \text{J}\cdot\text{s})^2}{2\pi \cdot 9.1 \times 10^{-31} \text{kg} \cdot 1.38 \times 10^{-23} \text{J/K} \cdot 10^7 \text{K}} \right)^{\frac{3}{2}}$   
 $\simeq 8660 \gg 1$ .
- ii. Relativistic:  $n\lambda^3 = n\left(\frac{h}{p}\right)^3 = n\left(\frac{hc}{\varepsilon}\right)^3 = n\left(\frac{hc}{kT_{\text{WD}}}\right)^3$   
 $\simeq 6.6 \times 10^{35} \text{m}^{-3} \left( \frac{6.626 \times 10^{-34} \text{J}\cdot\text{s} \cdot 3 \times 10^8 \text{m/s}}{1.38 \times 10^{-23} \text{J/K} \cdot 10^7 \text{K}} \right)^3$   
 $\simeq 2 \times 10^9 \gg 1$ . (Only estimation if using  $kT_{\text{WD}}$ ).
- Strong degenerated.

... He white dwarf.

- Fermi energy (non-relativistic) (8.5.6):

$$\mu_0 = \frac{\hbar^2}{2m} (3\pi^2 n)^{2/3}$$

... He white dwarf.

- Fermi energy (non-relativistic) (8.5.6):

$$\begin{aligned}\mu_0 &= \frac{\hbar^2}{2m} (3\pi^2 n)^{2/3} \\ &\simeq \frac{(1.054 \times 10^{-34} \text{ J}\cdot\text{s})^2}{2 \times 9.1 \times 10^{-31} \text{ kg}} (3\pi^2 \cdot 6.6 \times 10^{35} \text{ m}^{-3})^{2/3}\end{aligned}$$

... He white dwarf.

- Fermi energy (non-relativistic) (8.5.6):

$$\begin{aligned}\mu_0 &= \frac{\hbar^2}{2m} (3\pi^2 n)^{2/3} \\ &\simeq \frac{(1.054 \times 10^{-34} \text{J}\cdot\text{s})^2}{2 \times 9.1 \times 10^{-31} \text{kg}} (3\pi^2 \cdot 6.6 \times 10^{35} \text{m}^{-3})^{2/3} \\ &\simeq 4.4 \times 10^{-14} \text{J} \simeq 2.7 \times 10^5 \text{eV} \\ (T_F = \frac{\mu_0}{k} &\simeq 3.2 \times 10^9 \text{K.})\end{aligned}$$

... He white dwarf.

- Fermi energy (non-relativistic) (8.5.6):

$$\mu_0 = \frac{\hbar^2}{2m} (3\pi^2 n)^{2/3}$$

$$\simeq \frac{(1.054 \times 10^{-34} \text{J}\cdot\text{s})^2}{2 \times 9.1 \times 10^{-31} \text{kg}} (3\pi^2 \cdot 6.6 \times 10^{35} \text{m}^{-3})^{2/3}$$

$$\simeq 4.4 \times 10^{-14} \text{J} \simeq 2.7 \times 10^5 \text{eV}$$

$$(T_F = \frac{\mu_0}{k} \simeq 3.2 \times 10^9 \text{K}.)$$

$$\mu_0 \sim m_e c^2 \simeq 0.511 \text{MeV}, \text{ roughly relativistic.}$$

... He white dwarf.

- Fermi energy (non-relativistic) (8.5.6):

$$\mu_0 = \frac{\hbar^2}{2m} (3\pi^2 n)^{2/3}$$

$$\simeq \frac{(1.054 \times 10^{-34} \text{ J}\cdot\text{s})^2}{2 \times 9.1 \times 10^{-31} \text{ kg}} (3\pi^2 \cdot 6.6 \times 10^{35} \text{ m}^{-3})^{2/3}$$

$$\simeq 4.4 \times 10^{-14} \text{ J} \simeq 2.7 \times 10^5 \text{ eV}$$

$$(T_F = \frac{\mu_0}{k} \simeq 3.2 \times 10^9 \text{ K.})$$

$$\mu_0 \sim m_e c^2 \simeq 0.511 \text{ MeV, roughly relativistic.}$$

- Fermi energy (relativistic).

## ... He white dwarf.

- Fermi energy (non-relativistic) (8.5.6):

$$\mu_0 = \frac{\hbar^2}{2m} (3\pi^2 n)^{2/3}$$

$$\simeq \frac{(1.054 \times 10^{-34} \text{ J}\cdot\text{s})^2}{2 \times 9.1 \times 10^{-31} \text{ kg}} (3\pi^2 \cdot 6.6 \times 10^{35} \text{ m}^{-3})^{2/3}$$

$$\simeq 4.4 \times 10^{-14} \text{ J} \simeq 2.7 \times 10^5 \text{ eV}$$

$$(T_F = \frac{\mu_0}{k} \simeq 3.2 \times 10^9 \text{ K.})$$

$$\mu_0 \sim m_e c^2 \simeq 0.511 \text{ MeV, roughly relativistic.}$$

- Fermi energy (relativistic).  $D(\varepsilon) d\varepsilon = 2 \cdot \frac{V 4\pi p^2 dp}{h^3}$



... He white dwarf.

- Fermi energy (non-relativistic) (8.5.6):

$$\mu_0 = \frac{\hbar^2}{2m} (3\pi^2 n)^{2/3}$$

$$\simeq \frac{(1.054 \times 10^{-34} \text{ J}\cdot\text{s})^2}{2 \times 9.1 \times 10^{-31} \text{ kg}} (3\pi^2 \cdot 6.6 \times 10^{35} \text{ m}^{-3})^{2/3}$$

$$\simeq 4.4 \times 10^{-14} \text{ J} \simeq 2.7 \times 10^5 \text{ eV}$$

$$(T_F = \frac{\mu_0}{k} \simeq 3.2 \times 10^9 \text{ K.})$$

$$\mu_0 \sim m_e c^2 \simeq 0.511 \text{ MeV, roughly relativistic.}$$

- Fermi energy (relativistic).  $D(\varepsilon) d\varepsilon = 2 \cdot \frac{V 4\pi p^2 dp}{h^3}$   
 $\simeq \frac{V 8\pi}{h^3 c^3} \varepsilon^2 d\varepsilon.$

... He white dwarf.

- Fermi energy (non-relativistic) (8.5.6):

$$\mu_0 = \frac{\hbar^2}{2m} (3\pi^2 n)^{2/3}$$

$$\simeq \frac{(1.054 \times 10^{-34} \text{ J}\cdot\text{s})^2}{2 \times 9.1 \times 10^{-31} \text{ kg}} (3\pi^2 \cdot 6.6 \times 10^{35} \text{ m}^{-3})^{2/3}$$

$$\simeq 4.4 \times 10^{-14} \text{ J} \simeq 2.7 \times 10^5 \text{ eV}$$

$$(T_F = \frac{\mu_0}{k} \simeq 3.2 \times 10^9 \text{ K.})$$

$$\mu_0 \sim m_e c^2 \simeq 0.511 \text{ MeV, roughly relativistic.}$$

- Fermi energy (relativistic).  $D(\varepsilon) d\varepsilon = 2 \cdot \frac{V 4\pi p^2 dp}{h^3}$

$$\simeq \frac{V 8\pi}{h^3 c^3} \varepsilon^2 d\varepsilon. \text{ As strong degeneracy indicates}$$

$$T_{\text{WD}} \ll T_F, \int_0^{\mu_0} D(\varepsilon) d\varepsilon = N,$$

## ... He white dwarf.

- Fermi energy (non-relativistic) (8.5.6):

$$\mu_0 = \frac{\hbar^2}{2m} (3\pi^2 n)^{2/3}$$

$$\simeq \frac{(1.054 \times 10^{-34} \text{ J}\cdot\text{s})^2}{2 \times 9.1 \times 10^{-31} \text{ kg}} (3\pi^2 \cdot 6.6 \times 10^{35} \text{ m}^{-3})^{2/3}$$

$$\simeq 4.4 \times 10^{-14} \text{ J} \simeq 2.7 \times 10^5 \text{ eV}$$

$$(T_F = \frac{\mu_0}{k} \simeq 3.2 \times 10^9 \text{ K.})$$

$$\mu_0 \sim m_e c^2 \simeq 0.511 \text{ MeV, roughly relativistic.}$$

- Fermi energy (relativistic).  $D(\varepsilon) d\varepsilon = 2 \cdot \frac{V 4\pi p^2 dp}{h^3}$

$$\simeq \frac{V 8\pi}{h^3 c^3} \varepsilon^2 d\varepsilon. \text{ As strong degeneracy indicates}$$

$$T_{\text{WD}} \ll T_F, \int_0^{\mu_0} D(\varepsilon) d\varepsilon = N, \Rightarrow \mu_0 = hc \left( \frac{3N}{8\pi V} \right)^{\frac{1}{3}}$$

## ... He white dwarf.

- Fermi energy (non-relativistic) (8.5.6):

$$\mu_0 = \frac{\hbar^2}{2m} (3\pi^2 n)^{2/3}$$

$$\simeq \frac{(1.054 \times 10^{-34} \text{ J}\cdot\text{s})^2}{2 \times 9.1 \times 10^{-31} \text{ kg}} (3\pi^2 \cdot 6.6 \times 10^{35} \text{ m}^{-3})^{2/3}$$

$$\simeq 4.4 \times 10^{-14} \text{ J} \simeq 2.7 \times 10^5 \text{ eV}$$

$$(T_F = \frac{\mu_0}{k} \simeq 3.2 \times 10^9 \text{ K.})$$

$$\mu_0 \sim m_e c^2 \simeq 0.511 \text{ MeV, roughly relativistic.}$$

- Fermi energy (relativistic).  $D(\varepsilon) d\varepsilon = 2 \cdot \frac{V 4\pi p^2 dp}{h^3}$

$$\simeq \frac{V 8\pi}{h^3 c^3} \varepsilon^2 d\varepsilon. \text{ As strong degeneracy indicates}$$

$$T_{\text{WD}} \ll T_F, \int_0^{\mu_0} D(\varepsilon) d\varepsilon = N, \Rightarrow \mu_0 = hc \left( \frac{3N}{8\pi V} \right)^{\frac{1}{3}}$$

$$= 6.626 \times 10^{-34} \text{ J} \cdot \text{s} \cdot 3 \times 10^8 \text{ m/s} \cdot \left( \frac{3}{8\pi} \cdot 6.6 \times 10^{35} \text{ m}^{-3} \right)^{\frac{1}{3}}$$

... He white dwarf.

- Fermi energy (non-relativistic) (8.5.6):

$$\mu_0 = \frac{\hbar^2}{2m} (3\pi^2 n)^{2/3}$$

$$\simeq \frac{(1.054 \times 10^{-34} \text{ J}\cdot\text{s})^2}{2 \times 9.1 \times 10^{-31} \text{ kg}} (3\pi^2 \cdot 6.6 \times 10^{35} \text{ m}^{-3})^{2/3}$$

$$\simeq 4.4 \times 10^{-14} \text{ J} \simeq 2.7 \times 10^5 \text{ eV}$$

$$(T_F = \frac{\mu_0}{k} \simeq 3.2 \times 10^9 \text{ K.})$$

$$\mu_0 \sim m_e c^2 \simeq 0.511 \text{ MeV, roughly relativistic.}$$

- Fermi energy (relativistic).  $D(\varepsilon)d\varepsilon = 2 \cdot \frac{V 4\pi p^2 dp}{h^3}$

$$\simeq \frac{V 8\pi}{h^3 c^3} \varepsilon^2 d\varepsilon. \text{ As strong degeneracy indicates}$$

$$T_{\text{WD}} \ll T_F, \int_0^{\mu_0} D(\varepsilon) d\varepsilon = N, \Rightarrow \mu_0 = hc \left( \frac{3N}{8\pi V} \right)^{\frac{1}{3}}$$

$$= 6.626 \times 10^{-34} \text{ J} \cdot \text{s} \cdot 3 \times 10^8 \text{ m/s} \cdot \left( \frac{3}{8\pi} \cdot 6.6 \times 10^{35} \text{ m}^{-3} \right)^{\frac{1}{3}}$$

$$\simeq 8.5 \times 10^{-14} \text{ J} \simeq 0.53 \text{ MeV. (This is kinetic energy, mildly relativistic.)}$$

... He white dwarf.

- Internal energy:  $U = \int_0^{\mu_0} \varepsilon D(\varepsilon) d\varepsilon$

... He white dwarf.

- Internal energy:  $U = \int_0^{\mu_0} \varepsilon D(\varepsilon) d\varepsilon = \int_0^{\mu_0} \varepsilon \frac{V 8\pi}{h^3 c^3} \varepsilon^2 d\varepsilon$

... He white dwarf.

- Internal energy:  $U = \int_0^{\mu_0} \varepsilon D(\varepsilon) d\varepsilon = \int_0^{\mu_0} \varepsilon \frac{V 8\pi}{h^3 c^3} \varepsilon^2 d\varepsilon$   
 $= \frac{3}{4} N \mu_0$



... He white dwarf.

- Internal energy:  $U = \int_0^{\mu_0} \varepsilon D(\varepsilon) d\varepsilon = \int_0^{\mu_0} \varepsilon \frac{V 8\pi}{h^3 c^3} \varepsilon^2 d\varepsilon$   
 $= \frac{3}{4} N \mu_0 \simeq \frac{3}{4} N h c \left( \frac{3N}{8\pi V} \right)^{\frac{1}{3}}$

... He white dwarf.

- Internal energy: 
$$U = \int_0^{\mu_0} \varepsilon D(\varepsilon) d\varepsilon = \int_0^{\mu_0} \varepsilon \frac{V 8\pi}{h^3 c^3} \varepsilon^2 d\varepsilon$$
$$= \frac{3}{4} N \mu_0 \simeq \frac{3}{4} N h c \left( \frac{3N}{8\pi V} \right)^{\frac{1}{3}}$$
$$\simeq \frac{3}{4} \cdot \frac{2.2 \times 10^{30} \text{kg}}{4 \times 1.67 \times 10^{-27} \text{kg}} \cdot 8.5 \times 10^{-14} \text{J} \simeq 3.8 \times 10^{43} \text{J}.$$

... He white dwarf.

- Internal energy:  $U = \int_0^{\mu_0} \varepsilon D(\varepsilon) d\varepsilon = \int_0^{\mu_0} \varepsilon \frac{V 8\pi}{h^3 c^3} \varepsilon^2 d\varepsilon$   
 $= \frac{3}{4} N \mu_0 \simeq \frac{3}{4} N h c \left( \frac{3N}{8\pi V} \right)^{\frac{1}{3}}$   
 $\simeq \frac{3}{4} \cdot \frac{2.2 \times 10^{30} \text{kg}}{4 \times 1.67 \times 10^{-27} \text{kg}} \cdot 8.5 \times 10^{-14} \text{J} \simeq 3.8 \times 10^{43} \text{J}.$
- Pressure:  $p = \frac{U}{3V}$

... He white dwarf.

- Internal energy:  $U = \int_0^{\mu_0} \varepsilon D(\varepsilon) d\varepsilon = \int_0^{\mu_0} \varepsilon \frac{V 8\pi}{h^3 c^3} \varepsilon^2 d\varepsilon$   
 $= \frac{3}{4} N \mu_0 \simeq \frac{3}{4} N h c \left( \frac{3N}{8\pi V} \right)^{\frac{1}{3}}$   
 $\simeq \frac{3}{4} \cdot \frac{2.2 \times 10^{30} \text{kg}}{4 \times 1.67 \times 10^{-27} \text{kg}} \cdot 8.5 \times 10^{-14} \text{J} \simeq 3.8 \times 10^{43} \text{J}.$
- Pressure:  $p = \frac{U}{3V} = \frac{N}{4V} \mu_0 = \frac{hc}{4} \left( \frac{3}{8\pi} \right)^{\frac{1}{3}} n^{\frac{4}{3}}$

... He white dwarf.

- Internal energy:  $U = \int_0^{\mu_0} \varepsilon D(\varepsilon) d\varepsilon = \int_0^{\mu_0} \varepsilon \frac{V 8\pi}{h^3 c^3} \varepsilon^2 d\varepsilon$   
 $= \frac{3}{4} N \mu_0 \simeq \frac{3}{4} N h c \left( \frac{3N}{8\pi V} \right)^{\frac{1}{3}}$   
 $\simeq \frac{3}{4} \cdot \frac{2.2 \times 10^{30} \text{kg}}{4 \times 1.67 \times 10^{-27} \text{kg}} \cdot 8.5 \times 10^{-14} \text{J} \simeq 3.8 \times 10^{43} \text{J}.$
- Pressure:  $p = \frac{U}{3V} = \frac{N}{4V} \mu_0 = \frac{hc}{4} \left( \frac{3}{8\pi} \right)^{\frac{1}{3}} n^{\frac{4}{3}}$   
 $\simeq \frac{3.8 \times 10^{43} \text{J}}{3 \cdot \frac{4}{3} \pi (6 \times 10^6 \text{m})^3} \simeq 1.4 \times 10^{22} \text{J} \cdot \text{m}^{-3}.$

## ... He white dwarf.

- Internal energy:  $U = \int_0^{\mu_0} \varepsilon D(\varepsilon) d\varepsilon = \int_0^{\mu_0} \varepsilon \frac{V 8\pi}{h^3 c^3} \varepsilon^2 d\varepsilon$   
 $= \frac{3}{4} N \mu_0 \simeq \frac{3}{4} N h c \left( \frac{3N}{8\pi V} \right)^{\frac{1}{3}}$   
 $\simeq \frac{3}{4} \cdot \frac{2.2 \times 10^{30} \text{kg}}{4 \times 1.67 \times 10^{-27} \text{kg}} \cdot 8.5 \times 10^{-14} \text{J} \simeq 3.8 \times 10^{43} \text{J}.$
- Pressure:  $p = \frac{U}{3V} = \frac{N}{4V} \mu_0 = \frac{hc}{4} \left( \frac{3}{8\pi} \right)^{\frac{1}{3}} n^{\frac{4}{3}}$   
 $\simeq \frac{3.8 \times 10^{43} \text{J}}{3 \cdot \frac{4}{3} \pi (6 \times 10^6 \text{m})^3} \simeq 1.4 \times 10^{22} \text{J} \cdot \text{m}^{-3}.$
- Stability: for a virtual adiabatic variance  $\delta R$ , 1.  
 $\delta E_{\text{tot}} = \delta(U + V_p) = 0$ , and 2.  $\delta^2 E_{\text{tot}} > 0$ .

## ... He white dwarf.

- Internal energy:  $U = \int_0^{\mu_0} \varepsilon D(\varepsilon) d\varepsilon = \int_0^{\mu_0} \varepsilon \frac{V 8\pi}{h^3 c^3} \varepsilon^2 d\varepsilon$   
 $= \frac{3}{4} N \mu_0 \simeq \frac{3}{4} N h c \left( \frac{3N}{8\pi V} \right)^{\frac{1}{3}}$   
 $\simeq \frac{3}{4} \cdot \frac{2.2 \times 10^{30} \text{kg}}{4 \times 1.67 \times 10^{-27} \text{kg}} \cdot 8.5 \times 10^{-14} \text{J} \simeq 3.8 \times 10^{43} \text{J}.$
- Pressure:  $p = \frac{U}{3V} = \frac{N}{4V} \mu_0 = \frac{hc}{4} \left( \frac{3}{8\pi} \right)^{\frac{1}{3}} n^{\frac{4}{3}}$   
 $\simeq \frac{3.8 \times 10^{43} \text{J}}{3 \cdot \frac{4}{3} \pi (6 \times 10^6 \text{m})^3} \simeq 1.4 \times 10^{22} \text{J} \cdot \text{m}^{-3}.$
- Stability: for a virtual adiabatic variance  $\delta R$ , 1.  
 $\delta E_{\text{tot}} = \delta(U + V_p) = 0$ , and 2.  $\delta^2 E_{\text{tot}} > 0$ .  
 Non-relativistic: (8.5.7)  $U = \frac{3N}{5} \mu_0 = \frac{3N}{5} \frac{\hbar^2}{2m} (3\pi^2 n)^{2/3}$

## ... He white dwarf.

- Internal energy:  $U = \int_0^{\mu_0} \varepsilon D(\varepsilon) d\varepsilon = \int_0^{\mu_0} \varepsilon \frac{V 8\pi}{h^3 c^3} \varepsilon^2 d\varepsilon$   
 $= \frac{3}{4} N \mu_0 \simeq \frac{3}{4} N h c \left( \frac{3N}{8\pi V} \right)^{\frac{1}{3}}$   
 $\simeq \frac{3}{4} \cdot \frac{2.2 \times 10^{30} \text{kg}}{4 \times 1.67 \times 10^{-27} \text{kg}} \cdot 8.5 \times 10^{-14} \text{J} \simeq 3.8 \times 10^{43} \text{J}.$
- Pressure:  $p = \frac{U}{3V} = \frac{N}{4V} \mu_0 = \frac{h c}{4} \left( \frac{3}{8\pi} \right)^{\frac{1}{3}} n^{\frac{4}{3}}$   
 $\simeq \frac{3.8 \times 10^{43} \text{J}}{3 \cdot \frac{4}{3} \pi (6 \times 10^6 \text{m})^3} \simeq 1.4 \times 10^{22} \text{J} \cdot \text{m}^{-3}.$
- Stability: for a virtual adiabatic variance  $\delta R$ , 1.  
 $\delta E_{\text{tot}} = \delta(U + V_p) = 0$ , and 2.  $\delta^2 E_{\text{tot}} > 0$ .  
 Non-relativistic: (8.5.7)  $U = \frac{3N}{5} \mu_0 = \frac{3N}{5} \frac{\hbar^2}{2m} (3\pi^2 n)^{2/3}$   
 $\propto \frac{M^{5/3}}{R^2};$



## ... He white dwarf.

- Internal energy:  $U = \int_0^{\mu_0} \varepsilon D(\varepsilon) d\varepsilon = \int_0^{\mu_0} \varepsilon \frac{V 8\pi}{h^3 c^3} \varepsilon^2 d\varepsilon$   
 $= \frac{3}{4} N \mu_0 \simeq \frac{3}{4} N h c \left( \frac{3N}{8\pi V} \right)^{\frac{1}{3}}$   
 $\simeq \frac{3}{4} \cdot \frac{2.2 \times 10^{30} \text{kg}}{4 \times 1.67 \times 10^{-27} \text{kg}} \cdot 8.5 \times 10^{-14} \text{J} \simeq 3.8 \times 10^{43} \text{J}.$
- Pressure:  $p = \frac{U}{3V} = \frac{N}{4V} \mu_0 = \frac{h c}{4} \left( \frac{3}{8\pi} \right)^{\frac{1}{3}} n^{\frac{4}{3}}$   
 $\simeq \frac{3.8 \times 10^{43} \text{J}}{3 \cdot \frac{4}{3} \pi (6 \times 10^6 \text{m})^3} \simeq 1.4 \times 10^{22} \text{J} \cdot \text{m}^{-3}.$
- Stability: for a virtual adiabatic variance  $\delta R$ , 1.  
 $\delta E_{\text{tot}} = \delta(U + V_p) = 0$ , and 2.  $\delta^2 E_{\text{tot}} > 0$ .  
 Non-relativistic: (8.5.7)  $U = \frac{3N}{5} \mu_0 = \frac{3N}{5} \frac{\hbar^2}{2m} (3\pi^2 n)^{2/3}$   
 $\propto \frac{M^{5/3}}{R^2}; V_p = -\frac{GM^2}{R} \propto -\frac{M^2}{R}.$

... He white dwarf.

- Internal energy:  $U = \int_0^{\mu_0} \varepsilon D(\varepsilon) d\varepsilon = \int_0^{\mu_0} \varepsilon \frac{V 8\pi}{h^3 c^3} \varepsilon^2 d\varepsilon$   
 $= \frac{3}{4} N \mu_0 \simeq \frac{3}{4} N h c \left( \frac{3N}{8\pi V} \right)^{\frac{1}{3}}$   
 $\simeq \frac{3}{4} \cdot \frac{2.2 \times 10^{30} \text{kg}}{4 \times 1.67 \times 10^{-27} \text{kg}} \cdot 8.5 \times 10^{-14} \text{J} \simeq 3.8 \times 10^{43} \text{J}.$
- Pressure:  $p = \frac{U}{3V} = \frac{N}{4V} \mu_0 = \frac{hc}{4} \left( \frac{3}{8\pi} \right)^{\frac{1}{3}} n^{\frac{4}{3}}$   
 $\simeq \frac{3.8 \times 10^{43} \text{J}}{3 \cdot \frac{4}{3} \pi (6 \times 10^6 \text{m})^3} \simeq 1.4 \times 10^{22} \text{J} \cdot \text{m}^{-3}.$
- Stability: for a virtual adiabatic variance  $\delta R$ , 1.  
 $\delta E_{\text{tot}} = \delta(U + V_p) = 0$ , and 2.  $\delta^2 E_{\text{tot}} > 0$ .  
 Non-relativistic: (8.5.7)  $U = \frac{3N}{5} \mu_0 = \frac{3N}{5} \frac{\hbar^2}{2m} (3\pi^2 n)^{2/3}$   
 $\propto \frac{M^{5/3}}{R^2}; V_p = -\frac{GM^2}{R} \propto -\frac{M^2}{R}.$   
 $E_{\text{tot}} = k_1 \frac{M^{5/3}}{R^2} - k_2 \frac{M^2}{R}.$

... He white dwarf.

- $E_{\text{tot}} = k_1 \frac{M^{5/3}}{R^2} - k_2 \frac{M^2}{R}.$

... He white dwarf.

- $E_{\text{tot}} = k_1 \frac{M^{5/3}}{R^2} - k_2 \frac{M^2}{R}.$   
 $\delta E_{\text{tot}} = 0 \Rightarrow \left(-2k_1 \frac{M^{5/3}}{R^3} + k_2 \frac{M^2}{R^2}\right) \delta R = 0$

... He white dwarf.

- $E_{\text{tot}} = k_1 \frac{M^{5/3}}{R^2} - k_2 \frac{M^2}{R}.$   
 $\delta E_{\text{tot}} = 0 \Rightarrow \left(-2k_1 \frac{M^{5/3}}{R^3} + k_2 \frac{M^2}{R^2}\right) \delta R = 0$   
 $\Rightarrow R = k_3 M^{-1/3}; \text{ Minimum.}$

... He white dwarf.

- $E_{\text{tot}} = k_1 \frac{M^{5/3}}{R^2} - k_2 \frac{M^2}{R}.$   
 $\delta E_{\text{tot}} = 0 \Rightarrow (-2k_1 \frac{M^{5/3}}{R^3} + k_2 \frac{M^2}{R^2}) \delta R = 0$   
 $\Rightarrow R = k_3 M^{-1/3}; \text{ Minimum.}$
- Relativistic:  $U = \frac{3}{4} N h c \left( \frac{3N}{8\pi V} \right)^{\frac{1}{3}} \propto \frac{M^{4/3}}{R}$

... He white dwarf.

- $E_{\text{tot}} = k_1 \frac{M^{5/3}}{R^2} - k_2 \frac{M^2}{R}.$   
 $\delta E_{\text{tot}} = 0 \Rightarrow (-2k_1 \frac{M^{5/3}}{R^3} + k_2 \frac{M^2}{R^2}) \delta R = 0$   
 $\Rightarrow R = k_3 M^{-1/3}; \text{ Minimum.}$
- Relativistic:  $U = \frac{3}{4} N h c \left( \frac{3N}{8\pi V} \right)^{\frac{1}{3}} \propto \frac{M^{4/3}}{R}$   
 $E_{\text{tot}} = k_4 \frac{M^{4/3}}{R} - k_2 \frac{M^2}{R} = \frac{k_5}{R}.$

... He white dwarf.

- $E_{\text{tot}} = k_1 \frac{M^{5/3}}{R^2} - k_2 \frac{M^2}{R}.$

$$\delta E_{\text{tot}} = 0 \Rightarrow \left(-2k_1 \frac{M^{5/3}}{R^3} + k_2 \frac{M^2}{R^2}\right) \delta R = 0$$

$$\Rightarrow R = k_3 M^{-1/3}; \text{ Minimum.}$$

- Relativistic:  $U = \frac{3}{4} N h c \left(\frac{3N}{8\pi V}\right)^{\frac{1}{3}} \propto \frac{M^{4/3}}{R}$

$$E_{\text{tot}} = k_4 \frac{M^{4/3}}{R} - k_2 \frac{M^2}{R} = \frac{k_5}{R}. \text{ No minimum! Not stable.}$$



... He white dwarf.



- $E_{\text{tot}} = k_1 \frac{M^{5/3}}{R^2} - k_2 \frac{M^2}{R}.$

$$\delta E_{\text{tot}} = 0 \Rightarrow \left(-2k_1 \frac{M^{5/3}}{R^3} + k_2 \frac{M^2}{R^2}\right) \delta R = 0$$

$$\Rightarrow R = k_3 M^{-1/3}; \text{ Minimum.}$$

- Relativistic:  $U = \frac{3}{4} N h c \left(\frac{3N}{8\pi V}\right)^{1/3} \propto \frac{M^{4/3}}{R}$

$$E_{\text{tot}} = k_4 \frac{M^{4/3}}{R} - k_2 \frac{M^2}{R} = \frac{k_5}{R}. \text{ No minimum! Not stable.}$$



... He white dwarf.

- $E_{\text{tot}} > 0, R \nearrow, \rightarrow$  non-relativistic;

... He white dwarf.

- $E_{\text{tot}} > 0, R \nearrow, \rightarrow$  non-relativistic;  
 $E_{\text{tot}} < 0, R \searrow, \rightarrow$  unstable.

... He white dwarf.

- $E_{\text{tot}} > 0, R \nearrow, \rightarrow$  non-relativistic;  
 $E_{\text{tot}} < 0, R \searrow$ , unstable.
- For critical relativistic case ( $E_{\text{tot}} = 0$ ):

... He white dwarf.

- $E_{\text{tot}} > 0, R \nearrow, \rightarrow$  non-relativistic;  
 $E_{\text{tot}} < 0, R \searrow$ , unstable.
- For critical relativistic case ( $E_{\text{tot}} = 0$ ):  
$$E_{\text{tot}} = \frac{3}{4}Nhc\left(\frac{3N}{8\pi V}\right)^{\frac{1}{3}} - \frac{GM^2}{R} = 0$$

... He white dwarf.

- $E_{\text{tot}} > 0, R \nearrow, \rightarrow$  non-relativistic;  
 $E_{\text{tot}} < 0, R \searrow$ , unstable.
- For critical relativistic case ( $E_{\text{tot}} = 0$ ):  

$$E_{\text{tot}} = \frac{3}{4}Nhc\left(\frac{3N}{8\pi V}\right)^{\frac{1}{3}} - \frac{GM^2}{R} = 0$$

$$\Rightarrow \frac{3}{4}Nhc\left(\frac{3N}{8\pi^{\frac{4}{3}}\pi}\right)^{\frac{1}{3}} = GM^2$$

... He white dwarf.

- $E_{\text{tot}} > 0, R \nearrow, \rightarrow$  non-relativistic;  
 $E_{\text{tot}} < 0, R \searrow$ , unstable.
- For critical relativistic case ( $E_{\text{tot}} = 0$ ):  

$$E_{\text{tot}} = \frac{3}{4}Nhc\left(\frac{3N}{8\pi V}\right)^{\frac{1}{3}} - \frac{GM^2}{R} = 0$$

$$\Rightarrow \frac{3}{4}Nhc\left(\frac{3N}{8\pi^{\frac{4}{3}}\pi}\right)^{\frac{1}{3}} = GM^2$$

$$\Rightarrow \frac{3}{4}N^{\frac{4}{3}}hc\left(\frac{3}{8\pi^{\frac{4}{3}}\pi}\right)^{\frac{1}{3}} = GM^2$$

... He white dwarf.

- $E_{\text{tot}} > 0, R \nearrow, \rightarrow$  non-relativistic;  
 $E_{\text{tot}} < 0, R \searrow$ , unstable.
- For critical relativistic case ( $E_{\text{tot}} = 0$ ):

$$E_{\text{tot}} = \frac{3}{4}Nhc\left(\frac{3N}{8\pi V}\right)^{\frac{1}{3}} - \frac{GM^2}{R} = 0$$

$$\Rightarrow \frac{3}{4}Nhc\left(\frac{3N}{8\pi^{\frac{4}{3}}\pi}\right)^{\frac{1}{3}} = GM^2$$

$$\Rightarrow \frac{3}{4}N^{\frac{4}{3}}hc\left(\frac{3}{8\pi^{\frac{4}{3}}\pi}\right)^{\frac{1}{3}} = GM^2$$

$$\Rightarrow \frac{3}{4}\left(\frac{M}{2m_p}\right)^{\frac{4}{3}}hc\left(\frac{3}{8\pi^{\frac{4}{3}}\pi}\right)^{\frac{1}{3}} = GM^2$$



... He white dwarf.

- $E_{\text{tot}} > 0$ ,  $R \nearrow$ ,  $\rightarrow$  non-relativistic;

$E_{\text{tot}} < 0$ ,  $R \searrow$ , unstable.

- For critical relativistic case ( $E_{\text{tot}} = 0$ ):

$$E_{\text{tot}} = \frac{3}{4}Nhc\left(\frac{3N}{8\pi V}\right)^{\frac{1}{3}} - \frac{GM^2}{R} = 0$$

$$\Rightarrow \frac{3}{4}Nhc\left(\frac{3N}{8\pi^{\frac{4}{3}}\pi}\right)^{\frac{1}{3}} = GM^2$$

$$\Rightarrow \frac{3}{4}N^{\frac{4}{3}}hc\left(\frac{3}{8\pi^{\frac{4}{3}}\pi}\right)^{\frac{1}{3}} = GM^2$$

$$\Rightarrow \frac{3}{4}\left(\frac{M}{2m_p}\right)^{\frac{4}{3}}hc\left(\frac{3}{8\pi^{\frac{4}{3}}\pi}\right)^{\frac{1}{3}} = GM^2$$

$$\Rightarrow M \simeq \frac{3}{64\pi m_p^2}\left(\frac{3hc}{2G}\right)^{\frac{3}{2}} \simeq 5 \times 10^{30}\text{kg} \simeq 2.5M_{\odot}.$$

... He white dwarf.

- $E_{\text{tot}} > 0$ ,  $R \nearrow$ ,  $\rightarrow$  non-relativistic;

$E_{\text{tot}} < 0$ ,  $R \searrow$ , unstable.

- For critical relativistic case ( $E_{\text{tot}} = 0$ ):

$$E_{\text{tot}} = \frac{3}{4}Nhc\left(\frac{3N}{8\pi V}\right)^{\frac{1}{3}} - \frac{GM^2}{R} = 0$$

$$\Rightarrow \frac{3}{4}Nhc\left(\frac{3N}{8\pi^{\frac{4}{3}}\pi}\right)^{\frac{1}{3}} = GM^2$$

$$\Rightarrow \frac{3}{4}N^{\frac{4}{3}}hc\left(\frac{3}{8\pi^{\frac{4}{3}}\pi}\right)^{\frac{1}{3}} = GM^2$$

$$\Rightarrow \frac{3}{4}\left(\frac{M}{2m_p}\right)^{\frac{4}{3}}hc\left(\frac{3}{8\pi^{\frac{4}{3}}\pi}\right)^{\frac{1}{3}} = GM^2$$

$$\Rightarrow M \simeq \frac{3}{64\pi m_p^2}\left(\frac{3hc}{2G}\right)^{\frac{3}{2}} \simeq 5 \times 10^{30}\text{kg} \simeq 2.5M_{\odot}.$$

- More precise results (H component; Density, EOS profile on radius; exact expression for mildly relativistic particle), Chandrasekhar mass limit:

... He white dwarf.

- $E_{\text{tot}} > 0$ ,  $R \nearrow$ ,  $\rightarrow$  non-relativistic;

$E_{\text{tot}} < 0$ ,  $R \searrow$ , unstable.

- For critical relativistic case ( $E_{\text{tot}} = 0$ ):

$$E_{\text{tot}} = \frac{3}{4}Nhc\left(\frac{3N}{8\pi V}\right)^{\frac{1}{3}} - \frac{GM^2}{R} = 0$$

$$\Rightarrow \frac{3}{4}Nhc\left(\frac{3N}{8\pi^{\frac{4}{3}}\pi}\right)^{\frac{1}{3}} = GM^2$$

$$\Rightarrow \frac{3}{4}N^{\frac{4}{3}}hc\left(\frac{3}{8\pi^{\frac{4}{3}}\pi}\right)^{\frac{1}{3}} = GM^2$$

$$\Rightarrow \frac{3}{4}\left(\frac{M}{2m_p}\right)^{\frac{4}{3}}hc\left(\frac{3}{8\pi^{\frac{4}{3}}\pi}\right)^{\frac{1}{3}} = GM^2$$

$$\Rightarrow M \simeq \frac{3}{64\pi m_p^2}\left(\frac{3hc}{2G}\right)^{\frac{3}{2}} \simeq 5 \times 10^{30}\text{kg} \simeq 2.5M_{\odot}.$$

- More precise results (H component; Density, EOS profile on radius; exact expression for mildly relativistic particle), Chandrasekhar mass limit:  $M \lesssim 1.4M_{\odot}$ .

# References

- A: "A grand dictionary of physics problems and solutions 5", Science press, 2005

# Table of contents

- ① §6. Most probable distribution of nearly independent particles
- ② §7. Boltzmann statistics
- ③ §8. Bose statistics and Fermi statistics
  - Bose statistics
  - Fermi statistics
  - Fermion gas in astrophysics