

Thermodynamics & Statistical Physics

Chapter 7. Boltzmann Statistics

Yuan-Chuan Zou
zouyc@hust.edu.cn

School of Physics, Huazhong University of Science and Technology

December 30, 2013

Table of contents

1 §7 Boltzmann Statistics

- 7.1 Thermal quantities in statistics
- 7.2 Equation of state of ideal gas
- 7.3 Maxwell speed distribution
- 7.4 Energy equipartition theorem
- 7.5 Internal energy and heat capacity of the ideal gas
- 7.6 Entropy of the ideal gas
- 7.7 Einstein's theory on heat capacity of solid
- 7.8 Paramagnetic solid
- 7.9 Negative temperature

7.1 Thermal quantities in statistics

7.1 Thermal quantities in statistics

- Internal energy: $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}.$

7.1 Thermal quantities in statistics

- Internal energy: $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}.$
- Introduce partition function: $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}.$

7.1 Thermal quantities in statistics

- Internal energy: $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}.$
- Introduce partition function: $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}.$
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)

7.1 Thermal quantities in statistics

- Internal energy: $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$.
- Introduce partition function: $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$.
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)
- $N = \sum a_l$

7.1 Thermal quantities in statistics

- Internal energy: $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$.
- Introduce partition function: $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$.
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)
- $N = \sum a_l = \sum \omega_l e^{-\alpha - \beta \varepsilon_l}$

7.1 Thermal quantities in statistics

- Internal energy: $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$.
- Introduce partition function: $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$.
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)
- $N = \sum a_l = \sum \omega_l e^{-\alpha - \beta \varepsilon_l} = e^{-\alpha} \sum \omega_l e^{-\beta \varepsilon_l}$

7.1 Thermal quantities in statistics

- Internal energy: $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$.
- Introduce partition function: $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$.
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)
- $N = \sum a_l = \sum \omega_l e^{-\alpha - \beta \varepsilon_l} = e^{-\alpha} \sum \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} Z_1$.

7.1 Thermal quantities in statistics

- Internal energy: $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$.
- Introduce partition function: $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$.
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)
- $N = \sum a_l = \sum \omega_l e^{-\alpha - \beta \varepsilon_l} = e^{-\alpha} \sum \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} Z_1$.
- $\therefore e^{\alpha} = \frac{Z_1}{N}$.

7.1 Thermal quantities in statistics

- Internal energy: $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$.
- Introduce partition function: $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$.
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)
- $N = \sum a_l = \sum \omega_l e^{-\alpha - \beta \varepsilon_l} = e^{-\alpha} \sum \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} Z_1$.
- $\therefore e^{\alpha} = \frac{Z_1}{N}$.
- $U = e^{-\alpha} \sum \varepsilon_l \omega_l e^{-\beta \varepsilon_l}$

7.1 Thermal quantities in statistics

- Internal energy: $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$.
- Introduce partition function: $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$.
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)
- $N = \sum a_l = \sum \omega_l e^{-\alpha - \beta \varepsilon_l} = e^{-\alpha} \sum \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} Z_1$.
- $\therefore e^{\alpha} = \frac{Z_1}{N}$.
- $U = e^{-\alpha} \sum \varepsilon_l \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} \sum -\frac{\partial}{\partial \beta} (\omega_l e^{-\beta \varepsilon_l})$

7.1 Thermal quantities in statistics

- Internal energy: $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$.
- Introduce partition function: $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$.
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)
- $N = \sum a_l = \sum \omega_l e^{-\alpha - \beta \varepsilon_l} = e^{-\alpha} \sum \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} Z_1$.
- $\therefore e^{\alpha} = \frac{Z_1}{N}$.
- $U = e^{-\alpha} \sum \varepsilon_l \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} \sum -\frac{\partial}{\partial \beta} (\omega_l e^{-\beta \varepsilon_l})$
 $= e^{-\alpha} \left(-\frac{\partial}{\partial \beta}\right) \sum (\omega_l e^{-\beta \varepsilon_l})$

7.1 Thermal quantities in statistics

- Internal energy: $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$.
- Introduce partition function: $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$.
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)
- $N = \sum a_l = \sum \omega_l e^{-\alpha - \beta \varepsilon_l} = e^{-\alpha} \sum \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} Z_1$.
- $\therefore e^{\alpha} = \frac{Z_1}{N}$.
- $U = e^{-\alpha} \sum \varepsilon_l \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} \sum -\frac{\partial}{\partial \beta} (\omega_l e^{-\beta \varepsilon_l})$
 $= e^{-\alpha} \left(-\frac{\partial}{\partial \beta}\right) \sum (\omega_l e^{-\beta \varepsilon_l}) = \frac{N}{Z_1} \left(-\frac{\partial}{\partial \beta}\right) Z_1$

7.1 Thermal quantities in statistics

- Internal energy: $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$.
- Introduce partition function: $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$.
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)
- $N = \sum a_l = \sum \omega_l e^{-\alpha - \beta \varepsilon_l} = e^{-\alpha} \sum \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} Z_1$.
- $\therefore e^{\alpha} = \frac{Z_1}{N}$.
- $U = e^{-\alpha} \sum \varepsilon_l \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} \sum -\frac{\partial}{\partial \beta} (\omega_l e^{-\beta \varepsilon_l})$
 $= e^{-\alpha} \left(-\frac{\partial}{\partial \beta}\right) \sum (\omega_l e^{-\beta \varepsilon_l}) = \frac{N}{Z_1} \left(-\frac{\partial}{\partial \beta}\right) Z_1 = -N \frac{\partial}{\partial \beta} \ln Z_1$.

7.1 Thermal quantities in statistics

- Internal energy: $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$.
- Introduce partition function: $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$.
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)
- $N = \sum a_l = \sum \omega_l e^{-\alpha - \beta \varepsilon_l} = e^{-\alpha} \sum \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} Z_1$.
- $\therefore e^{\alpha} = \frac{Z_1}{N}$.
- $U = e^{-\alpha} \sum \varepsilon_l \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} \sum -\frac{\partial}{\partial \beta} (\omega_l e^{-\beta \varepsilon_l})$
 $= e^{-\alpha} \left(-\frac{\partial}{\partial \beta}\right) \sum (\omega_l e^{-\beta \varepsilon_l}) = \frac{N}{Z_1} \left(-\frac{\partial}{\partial \beta}\right) Z_1 = -N \frac{\partial}{\partial \beta} \ln Z_1$.
- 1st law: $dU = \delta W + \delta Q$.

7.1 Thermal quantities in statistics

- Internal energy: $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$.
- Introduce partition function: $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$.
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)
- $N = \sum a_l = \sum \omega_l e^{-\alpha - \beta \varepsilon_l} = e^{-\alpha} \sum \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} Z_1$.
- $\therefore e^{\alpha} = \frac{Z_1}{N}$.
- $U = e^{-\alpha} \sum \varepsilon_l \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} \sum -\frac{\partial}{\partial \beta} (\omega_l e^{-\beta \varepsilon_l})$
 $= e^{-\alpha} \left(-\frac{\partial}{\partial \beta}\right) \sum (\omega_l e^{-\beta \varepsilon_l}) = \frac{N}{Z_1} \left(-\frac{\partial}{\partial \beta}\right) Z_1 = -N \frac{\partial}{\partial \beta} \ln Z_1$.
- 1st law: $dU = \delta W + \delta Q$. $\delta W = \sum_i Y_i dy_i$ (Y_i generalized force, dy_i generalized displacement)

7.1 Thermal quantities in statistics

- Internal energy: $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$.
- Introduce partition function: $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$.
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)
- $N = \sum a_l = \sum \omega_l e^{-\alpha - \beta \varepsilon_l} = e^{-\alpha} \sum \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} Z_1$.
- $\therefore e^{\alpha} = \frac{Z_1}{N}$.
- $U = e^{-\alpha} \sum \varepsilon_l \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} \sum -\frac{\partial}{\partial \beta} (\omega_l e^{-\beta \varepsilon_l})$
 $= e^{-\alpha} \left(-\frac{\partial}{\partial \beta}\right) \sum (\omega_l e^{-\beta \varepsilon_l}) = \frac{N}{Z_1} \left(-\frac{\partial}{\partial \beta}\right) Z_1 = -N \frac{\partial}{\partial \beta} \ln Z_1$.
- 1st law: $dU = \delta W + \delta Q$. $\delta W = \sum_i Y_i dy_i$ (Y_i generalized force, dy_i generalized displacement)
 A force exerted on the system, U increases. But $\{a_l\}$ does not change. What changes?

7.1 Thermal quantities in statistics

- Internal energy: $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$.
- Introduce partition function: $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$.
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)
- $N = \sum a_l = \sum \omega_l e^{-\alpha - \beta \varepsilon_l} = e^{-\alpha} \sum \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} Z_1$.
- $\therefore e^{\alpha} = \frac{Z_1}{N}$.
- $U = e^{-\alpha} \sum \varepsilon_l \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} \sum -\frac{\partial}{\partial \beta} (\omega_l e^{-\beta \varepsilon_l})$
 $= e^{-\alpha} \left(-\frac{\partial}{\partial \beta}\right) \sum (\omega_l e^{-\beta \varepsilon_l}) = \frac{N}{Z_1} \left(-\frac{\partial}{\partial \beta}\right) Z_1 = -N \frac{\partial}{\partial \beta} \ln Z_1$.
- 1st law: $dU = \delta W + \delta Q$. $\delta W = \sum_i Y_i dy_i$ (Y_i generalized force, dy_i generalized displacement)
 A force exerted on the system, U increases. But $\{a_l\}$ does not change. What changes? $\{\varepsilon_l\}$.

Thermal quantities in statistics

- For each particle, the generalized force is $\frac{\partial \varepsilon_l}{\partial y}$,

Thermal quantities in statistics

- For each particle, the generalized force is $\frac{\partial \varepsilon_l}{\partial y}$, the whole force (Y_i): $Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l$

Thermal quantities in statistics

- For each particle, the generalized force is $\frac{\partial \varepsilon_l}{\partial y}$, the whole force (Y_i): $Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$

Thermal quantities in statistics

- For each particle, the generalized force is $\frac{\partial \varepsilon_l}{\partial y}$, the whole force (Y_i):
$$Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$$
$$= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l}$$

Thermal quantities in statistics

- For each particle, the generalized force is $\frac{\partial \varepsilon_l}{\partial y}$, the whole force (Y_i):
$$Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$$
$$= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l} = e^{-\alpha} \sum \omega_l \left(-\frac{1}{\beta}\right) \frac{\partial e^{-\beta \varepsilon_l}}{\partial y}$$

Thermal quantities in statistics

- For each particle, the generalized force is $\frac{\partial \varepsilon_l}{\partial y}$, the whole force (Y_i):
$$Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$$
$$= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l} = e^{-\alpha} \sum \omega_l \left(-\frac{1}{\beta}\right) \frac{\partial e^{-\beta \varepsilon_l}}{\partial y}$$
$$= e^{-\alpha} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} \sum \omega_l e^{-\beta \varepsilon_l}$$

Thermal quantities in statistics

- For each particle, the generalized force is $\frac{\partial \varepsilon_l}{\partial y}$, the whole force (Y_i):
$$Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$$
$$= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l} = e^{-\alpha} \sum \omega_l \left(-\frac{1}{\beta}\right) \frac{\partial e^{-\beta \varepsilon_l}}{\partial y}$$
$$= e^{-\alpha} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} \sum \omega_l e^{-\beta \varepsilon_l} = \frac{N}{Z_1} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} Z_1$$

Thermal quantities in statistics

- For each particle, the generalized force is $\frac{\partial \varepsilon_l}{\partial y}$, the whole force (Y_i):
$$Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$$
$$= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l} = e^{-\alpha} \sum \omega_l \left(-\frac{1}{\beta}\right) \frac{\partial e^{-\beta \varepsilon_l}}{\partial y}$$
$$= e^{-\alpha} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} \sum \omega_l e^{-\beta \varepsilon_l} = \frac{N}{Z_1} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} Z_1 = -\frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y}.$$

Thermal quantities in statistics

- For each particle, the generalized force is $\frac{\partial \varepsilon_l}{\partial y}$, the whole force (Y_i):
$$Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$$
$$= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l} = e^{-\alpha} \sum \omega_l \left(-\frac{1}{\beta}\right) \frac{\partial e^{-\beta \varepsilon_l}}{\partial y}$$
$$= e^{-\alpha} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} \sum \omega_l e^{-\beta \varepsilon_l} = \frac{N}{Z_1} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} Z_1 = -\frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y}.$$
- For p and V , $\delta W = -p dV$, so $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V}$.

Thermal quantities in statistics

- For each particle, the generalized force is $\frac{\partial \varepsilon_l}{\partial y}$, the whole force (Y_i):
$$Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$$
$$= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l} = e^{-\alpha} \sum \omega_l \left(-\frac{1}{\beta}\right) \frac{\partial e^{-\beta \varepsilon_l}}{\partial y}$$
$$= e^{-\alpha} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} \sum \omega_l e^{-\beta \varepsilon_l} = \frac{N}{Z_1} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} Z_1 = -\frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y}.$$
- For p and V , $\delta W = -p dV$, so $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V}$.
- As $d\varepsilon_l = \sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i$,

Thermal quantities in statistics

- For each particle, the generalized force is $\frac{\partial \varepsilon_l}{\partial y}$, the whole force (Y_i):
$$Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$$
$$= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l} = e^{-\alpha} \sum \omega_l \left(-\frac{1}{\beta}\right) \frac{\partial e^{-\beta \varepsilon_l}}{\partial y}$$
$$= e^{-\alpha} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} \sum \omega_l e^{-\beta \varepsilon_l} = \frac{N}{Z_1} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} Z_1 = -\frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y}.$$
- For p and V , $\delta W = -p dV$, so $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V}$.
- As $d\varepsilon_l = \sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i$, then $\delta W = \sum_i Y_i dy_i$

Thermal quantities in statistics

- For each particle, the generalized force is $\frac{\partial \varepsilon_l}{\partial y}$, the whole force (Y_i): $Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$

$$= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l} = e^{-\alpha} \sum \omega_l \left(-\frac{1}{\beta}\right) \frac{\partial e^{-\beta \varepsilon_l}}{\partial y}$$

$$= e^{-\alpha} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} \sum \omega_l e^{-\beta \varepsilon_l} = \frac{N}{Z_1} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} Z_1 = -\frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y}.$$
- For p and V , $\delta W = -p dV$, so $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V}$.
- As $d\varepsilon_l = \sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i$, then $\delta W = \sum_i Y_i dy_i$

$$= \sum_i dy_i \sum \frac{\partial \varepsilon_l}{\partial y_i} a_l$$

Thermal quantities in statistics

- For each particle, the generalized force is $\frac{\partial \varepsilon_l}{\partial y}$, the whole force (Y_i):
$$Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$$
$$= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l} = e^{-\alpha} \sum \omega_l \left(-\frac{1}{\beta}\right) \frac{\partial e^{-\beta \varepsilon_l}}{\partial y}$$
$$= e^{-\alpha} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} \sum \omega_l e^{-\beta \varepsilon_l} = \frac{N}{Z_1} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} Z_1 = -\frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y}.$$
- For p and V , $\delta W = -p dV$, so $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V}$.
- As $d\varepsilon_l = \sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i$, then $\delta W = \sum_i Y_i dy_i$
$$= \sum_i dy_i \sum \frac{\partial \varepsilon_l}{\partial y_i} a_l = \sum \left(\sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i\right) a_l$$

Thermal quantities in statistics

- For each particle, the generalized force is $\frac{\partial \varepsilon_l}{\partial y}$, the whole force (Y_i):
$$Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$$
$$= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l} = e^{-\alpha} \sum \omega_l \left(-\frac{1}{\beta}\right) \frac{\partial e^{-\beta \varepsilon_l}}{\partial y}$$
$$= e^{-\alpha} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} \sum \omega_l e^{-\beta \varepsilon_l} = \frac{N}{Z_1} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} Z_1 = -\frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y}.$$
- For p and V , $\delta W = -p dV$, so $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V}$.
- As $d\varepsilon_l = \sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i$, then $\delta W = \sum_i Y_i dy_i$
$$= \sum_i dy_i \sum \frac{\partial \varepsilon_l}{\partial y_i} a_l = \sum \left(\sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i\right) a_l = \sum a_l d\varepsilon_l.$$

Thermal quantities in statistics

- For each particle, the generalized force is $\frac{\partial \varepsilon_l}{\partial y}$, the whole force (Y_i):
$$Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$$
$$= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l} = e^{-\alpha} \sum \omega_l \left(-\frac{1}{\beta}\right) \frac{\partial e^{-\beta \varepsilon_l}}{\partial y}$$
$$= e^{-\alpha} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} \sum \omega_l e^{-\beta \varepsilon_l} = \frac{N}{Z_1} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} Z_1 = -\frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y}.$$
- For p and V , $\delta W = -p dV$, so $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V}$.
- As $d\varepsilon_l = \sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i$, then $\delta W = \sum_i Y_i dy_i$
$$= \sum_i dy_i \sum \frac{\partial \varepsilon_l}{\partial y_i} a_l = \sum \left(\sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i\right) a_l = \sum a_l d\varepsilon_l.$$
- $\therefore U = \sum \varepsilon_l a_l,$

Thermal quantities in statistics

- For each particle, the generalized force is $\frac{\partial \varepsilon_l}{\partial y}$, the whole force (Y_i):
$$Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$$
$$= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l} = e^{-\alpha} \sum \omega_l \left(-\frac{1}{\beta}\right) \frac{\partial e^{-\beta \varepsilon_l}}{\partial y}$$
$$= e^{-\alpha} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} \sum \omega_l e^{-\beta \varepsilon_l} = \frac{N}{Z_1} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} Z_1 = -\frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y}.$$
- For p and V , $\delta W = -p dV$, so $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V}$.
- As $d\varepsilon_l = \sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i$, then $\delta W = \sum_i Y_i dy_i$
$$= \sum_i dy_i \sum \frac{\partial \varepsilon_l}{\partial y_i} a_l = \sum \left(\sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i\right) a_l = \sum a_l d\varepsilon_l.$$
- $\because U = \sum \varepsilon_l a_l, \therefore dU = \sum a_l d\varepsilon_l + \sum \varepsilon_l da_l$

Thermal quantities in statistics

- For each particle, the generalized force is $\frac{\partial \varepsilon_l}{\partial y}$, the whole force (Y_i):
$$Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$$
$$= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l} = e^{-\alpha} \sum \omega_l \left(-\frac{1}{\beta}\right) \frac{\partial e^{-\beta \varepsilon_l}}{\partial y}$$
$$= e^{-\alpha} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} \sum \omega_l e^{-\beta \varepsilon_l} = \frac{N}{Z_1} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} Z_1 = -\frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y}.$$
- For p and V , $\delta W = -p dV$, so $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V}$.
- As $d\varepsilon_l = \sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i$, then $\delta W = \sum_i Y_i dy_i$
$$= \sum_i dy_i \sum \frac{\partial \varepsilon_l}{\partial y_i} a_l = \sum \left(\sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i\right) a_l = \sum a_l d\varepsilon_l.$$
- $\because U = \sum \varepsilon_l a_l, \therefore dU = \sum a_l d\varepsilon_l + \sum \varepsilon_l da_l$
$$= \delta W + \delta Q.$$

Thermal quantities in statistics

- For each particle, the generalized force is $\frac{\partial \varepsilon_l}{\partial y}$, the whole force (Y_i):
$$Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$$
$$= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l} = e^{-\alpha} \sum \omega_l \left(-\frac{1}{\beta}\right) \frac{\partial e^{-\beta \varepsilon_l}}{\partial y}$$
$$= e^{-\alpha} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} \sum \omega_l e^{-\beta \varepsilon_l} = \frac{N}{Z_1} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} Z_1 = -\frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y}.$$
 - For p and V , $\delta W = -p dV$, so $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V}$.
 - As $d\varepsilon_l = \sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i$, then $\delta W = \sum_i Y_i dy_i$
$$= \sum_i dy_i \sum \frac{\partial \varepsilon_l}{\partial y_i} a_l = \sum \left(\sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i\right) a_l = \sum a_l d\varepsilon_l.$$
 - $\because U = \sum \varepsilon_l a_l, \therefore dU = \sum a_l d\varepsilon_l + \sum \varepsilon_l da_l$
$$= \delta W + \delta Q.$$
- δW changes the energy levels; δQ changes $\{a_l\}$.

Thermal quantities in statistics — $\beta = \frac{1}{kT}$

- Using the partition function,

Thermal quantities in statistics — $\beta = \frac{1}{kT}$

- Using the partition function,

$$\delta Q = dU - Y dy = -N d\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} dy$$

Thermal quantities in statistics — $\beta = \frac{1}{kT}$

- Using the partition function,

$$\mathfrak{d}Q = dU - Ydy = -Nd\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} dy$$

$$\beta \mathfrak{d}Q = \beta(dU - Ydy) = -N\beta d\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + N \frac{\partial \ln Z_1}{\partial y} dy$$

Thermal quantities in statistics — $\beta = \frac{1}{kT}$

- Using the partition function,

$$\mathfrak{d}Q = dU - Ydy = -Nd\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} dy$$

$$\begin{aligned}\beta \mathfrak{d}Q &= \beta(dU - Ydy) = -N\beta d\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + N \frac{\partial \ln Z_1}{\partial y} dy \\ &= \left[N \frac{\partial \ln Z_1}{\partial \beta} d\beta - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right)\right] + N \frac{\partial \ln Z_1}{\partial y} dy\end{aligned}$$

Thermal quantities in statistics — $\beta = \frac{1}{kT}$

- Using the partition function,

$$\mathfrak{d}Q = dU - Ydy = -Nd\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} dy$$

$$\beta \mathfrak{d}Q = \beta(dU - Ydy) = -N\beta d\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[N \frac{\partial \ln Z_1}{\partial \beta} d\beta - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right) \right] + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[N \frac{\partial \ln Z_1}{\partial \beta} d\beta + N \frac{\partial \ln Z_1}{\partial y} dy \right] - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right),$$

Thermal quantities in statistics — $\beta = \frac{1}{kT}$

- Using the partition function,

$$\mathfrak{d}Q = dU - Ydy = -Nd\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} dy$$

$$\beta \mathfrak{d}Q = \beta(dU - Ydy) = -N\beta d\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[N \frac{\partial \ln Z_1}{\partial \beta} d\beta - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right) \right] + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[N \frac{\partial \ln Z_1}{\partial \beta} d\beta + N \frac{\partial \ln Z_1}{\partial y} dy \right] - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right),$$

$$Z_1(\beta, \varepsilon_l), \text{ and } \varepsilon(y), \text{ so, } d \ln Z_1 = \frac{\partial \ln Z_1}{\partial \beta} d\beta + \frac{\partial \ln Z_1}{\partial y} dy.$$

Thermal quantities in statistics — $\beta = \frac{1}{kT}$

- Using the partition function,

$$\overline{dQ} = dU - Ydy = -Nd\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} dy$$

$$\beta \overline{dQ} = \beta(dU - Ydy) = -N\beta d\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[N \frac{\partial \ln Z_1}{\partial \beta} d\beta - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right) \right] + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[N \frac{\partial \ln Z_1}{\partial \beta} d\beta + N \frac{\partial \ln Z_1}{\partial y} dy \right] - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right),$$

$$Z_1(\beta, \varepsilon_l), \text{ and } \varepsilon(y), \text{ so, } d \ln Z_1 = \frac{\partial \ln Z_1}{\partial \beta} d\beta + \frac{\partial \ln Z_1}{\partial y} dy.$$

$$\beta \overline{dQ} = Nd \ln Z_1 - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right)$$

Thermal quantities in statistics — $\beta = \frac{1}{kT}$

- Using the partition function,

$$\mathfrak{d}Q = dU - Ydy = -Nd\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} dy$$

$$\beta \mathfrak{d}Q = \beta(dU - Ydy) = -N\beta d\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[N \frac{\partial \ln Z_1}{\partial \beta} d\beta - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right) \right] + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[N \frac{\partial \ln Z_1}{\partial \beta} d\beta + N \frac{\partial \ln Z_1}{\partial y} dy \right] - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right),$$

$$Z_1(\beta, \varepsilon_l), \text{ and } \varepsilon(y), \text{ so, } d \ln Z_1 = \frac{\partial \ln Z_1}{\partial \beta} d\beta + \frac{\partial \ln Z_1}{\partial y} dy.$$

$$\beta \mathfrak{d}Q = Nd \ln Z_1 - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right)$$

$$= Nd\left(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta}\right),$$

Thermal quantities in statistics — $\beta = \frac{1}{kT}$

- Using the partition function,

$$\mathfrak{d}Q = dU - Ydy = -Nd\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} dy$$

$$\beta \mathfrak{d}Q = \beta(dU - Ydy) = -N\beta d\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[N \frac{\partial \ln Z_1}{\partial \beta} d\beta - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right) \right] + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[N \frac{\partial \ln Z_1}{\partial \beta} d\beta + N \frac{\partial \ln Z_1}{\partial y} dy \right] - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right),$$

$$Z_1(\beta, \varepsilon_l), \text{ and } \varepsilon(y), \text{ so, } d \ln Z_1 = \frac{\partial \ln Z_1}{\partial \beta} d\beta + \frac{\partial \ln Z_1}{\partial y} dy.$$

$$\beta \mathfrak{d}Q = Nd \ln Z_1 - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right)$$

$$= Nd\left(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta}\right), \text{ exact differential.}$$

Thermal quantities in statistics — $\beta = \frac{1}{kT}$

- Using the partition function,

$$\mathfrak{d}Q = dU - Ydy = -Nd\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} dy$$

$$\beta \mathfrak{d}Q = \beta(dU - Ydy) = -N\beta d\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[N \frac{\partial \ln Z_1}{\partial \beta} d\beta - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right) \right] + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[N \frac{\partial \ln Z_1}{\partial \beta} d\beta + N \frac{\partial \ln Z_1}{\partial y} dy \right] - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right),$$

$$Z_1(\beta, \varepsilon_l), \text{ and } \varepsilon(y), \text{ so, } d \ln Z_1 = \frac{\partial \ln Z_1}{\partial \beta} d\beta + \frac{\partial \ln Z_1}{\partial y} dy.$$

$$\beta \mathfrak{d}Q = Nd \ln Z_1 - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right)$$

$$= Nd\left(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta}\right), \text{ exact differential.}$$

- $\frac{1}{T} \mathfrak{d}Q = dS$ is also an exact differential.

Thermal quantities in statistics — $\beta = \frac{1}{kT}$

- Using the partition function,

$$\mathfrak{d}Q = dU - Ydy = -Nd\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} dy$$

$$\beta \mathfrak{d}Q = \beta(dU - Ydy) = -N\beta d\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[N \frac{\partial \ln Z_1}{\partial \beta} d\beta - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right) \right] + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[N \frac{\partial \ln Z_1}{\partial \beta} d\beta + N \frac{\partial \ln Z_1}{\partial y} dy \right] - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right),$$

$$Z_1(\beta, \varepsilon_l), \text{ and } \varepsilon(y), \text{ so, } d \ln Z_1 = \frac{\partial \ln Z_1}{\partial \beta} d\beta + \frac{\partial \ln Z_1}{\partial y} dy.$$

$$\beta \mathfrak{d}Q = Nd \ln Z_1 - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right)$$

$$= Nd\left(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta}\right), \text{ exact differential.}$$

- $\frac{1}{T} \mathfrak{d}Q = dS$ is also an exact differential.
- $\therefore \beta = \frac{1}{kT}$, where k is a constant.

Thermal quantities in statistics — $\beta = \frac{1}{kT}$

- Using the partition function,

$$\mathfrak{d}Q = dU - Ydy = -Nd\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} dy$$

$$\beta \mathfrak{d}Q = \beta(dU - Ydy) = -N\beta d\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[N \frac{\partial \ln Z_1}{\partial \beta} d\beta - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right) \right] + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[N \frac{\partial \ln Z_1}{\partial \beta} d\beta + N \frac{\partial \ln Z_1}{\partial y} dy \right] - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right),$$

$$Z_1(\beta, \varepsilon_l), \text{ and } \varepsilon(y), \text{ so, } d \ln Z_1 = \frac{\partial \ln Z_1}{\partial \beta} d\beta + \frac{\partial \ln Z_1}{\partial y} dy.$$

$$\beta \mathfrak{d}Q = Nd \ln Z_1 - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right)$$

$$= Nd\left(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta}\right), \text{ exact differential.}$$

- $\frac{1}{T} \mathfrak{d}Q = dS$ is also an exact differential.

- $\therefore \beta = \frac{1}{kT}$, where k is a constant.

$k = 1.381 \times 10^{-23} \text{ J} \cdot \text{K}^{-1}$ can be measured.

Thermal quantities in statistics — entropy

- $dS = \frac{1}{T} dQ$

Thermal quantities in statistics — entropy

- $dS = \frac{1}{T}dQ = k\beta dQ$

Thermal quantities in statistics — entropy

- $dS = \frac{1}{T}dQ = k\beta dQ = kN d(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$.

Thermal quantities in statistics — entropy

- $dS = \frac{1}{T} \delta Q = k\beta \delta Q = kN d(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta}).$
 $\Rightarrow S = Nk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta}).$

Thermal quantities in statistics — entropy

- $dS = \frac{1}{T} \delta Q = k\beta \delta Q = kN d(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$.
 $\Rightarrow S = Nk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$.
- $N = e^{-\alpha} Z_1, \Rightarrow \ln Z_1 = \ln N + \alpha$.

Thermal quantities in statistics — entropy

- $dS = \frac{1}{T} \delta Q = k\beta \delta Q = kN d(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$.
 $\Rightarrow S = Nk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$.
- $N = e^{-\alpha} Z_1, \Rightarrow \ln Z_1 = \ln N + \alpha$.
- $S = Nk[\ln N + \alpha - \beta \frac{\partial \ln Z_1}{\partial \beta}]$

Thermal quantities in statistics — entropy

- $dS = \frac{1}{T} \delta Q = k\beta \delta Q = kN d(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$.
 $\Rightarrow S = Nk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$.
- $N = e^{-\alpha} Z_1, \Rightarrow \ln Z_1 = \ln N + \alpha$.
- $S = Nk[\ln N + \alpha - \beta \frac{\partial \ln Z_1}{\partial \beta}]$
 $= k[N \ln N + \alpha N + \beta U]$

Thermal quantities in statistics — entropy

- $dS = \frac{1}{T}dQ = k\beta dQ = kN d(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$.
 $\Rightarrow S = Nk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$.
- $N = e^{-\alpha} Z_1, \Rightarrow \ln Z_1 = \ln N + \alpha$.
- $S = Nk[\ln N + \alpha - \beta \frac{\partial \ln Z_1}{\partial \beta}]$
 $= k[N \ln N + \alpha N + \beta U] = k[N \ln N + \sum(\alpha + \beta \varepsilon_l) a_l]$

Thermal quantities in statistics — entropy

- $dS = \frac{1}{T}dQ = k\beta dQ = kN d(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$.
 $\Rightarrow S = Nk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$.
- $N = e^{-\alpha} Z_1, \Rightarrow \ln Z_1 = \ln N + \alpha$.
- $S = Nk[\ln N + \alpha - \beta \frac{\partial \ln Z_1}{\partial \beta}]$
 $= k[N \ln N + \alpha N + \beta U] = k[N \ln N + \sum(\alpha + \beta \varepsilon_l) a_l]$
- Boltzmann distribution: $\ln \frac{\omega_l}{a_l} = \alpha + \beta \varepsilon_l$

Thermal quantities in statistics — entropy

- $dS = \frac{1}{T}dQ = k\beta dQ = kN d(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$.
 $\Rightarrow S = Nk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$.
- $N = e^{-\alpha} Z_1, \Rightarrow \ln Z_1 = \ln N + \alpha$.
- $S = Nk[\ln N + \alpha - \beta \frac{\partial \ln Z_1}{\partial \beta}]$
 $= k[N \ln N + \alpha N + \beta U] = k[N \ln N + \sum(\alpha + \beta \varepsilon_l) a_l]$
- Boltzmann distribution: $\ln \frac{\omega_l}{a_l} = \alpha + \beta \varepsilon_l$
- $\therefore S = k[N \ln N + \sum \ln \frac{\omega_l}{a_l} a_l]$

Thermal quantities in statistics — entropy

- $dS = \frac{1}{T}dQ = k\beta dQ = kNd(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$.
 $\Rightarrow S = Nk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$.
- $N = e^{-\alpha} Z_1, \Rightarrow \ln Z_1 = \ln N + \alpha$.
- $S = Nk[\ln N + \alpha - \beta \frac{\partial \ln Z_1}{\partial \beta}]$
 $= k[N \ln N + \alpha N + \beta U] = k[N \ln N + \sum(\alpha + \beta \varepsilon_l) a_l]$
- Boltzmann distribution: $\ln \frac{\omega_l}{a_l} = \alpha + \beta \varepsilon_l$
- $\therefore S = k[N \ln N + \sum \ln \frac{\omega_l}{a_l} a_l] = k \ln \Omega_M$.

Thermal quantities in statistics — entropy

- $dS = \frac{1}{T} \delta Q = k\beta \delta Q = kN d(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$.
 $\Rightarrow S = Nk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$.
- $N = e^{-\alpha} Z_1, \Rightarrow \ln Z_1 = \ln N + \alpha$.
- $S = Nk[\ln N + \alpha - \beta \frac{\partial \ln Z_1}{\partial \beta}]$
 $= k[N \ln N + \alpha N + \beta U] = k[N \ln N + \sum (\alpha + \beta \varepsilon_l) a_l]$
- Boltzmann distribution: $\ln \frac{\omega_l}{a_l} = \alpha + \beta \varepsilon_l$
- $\therefore S = k[N \ln N + \sum \ln \frac{\omega_l}{a_l} a_l] = k \ln \Omega_M$.
- Entropy statistical meaning: number of micro-states.

Thermal quantities in statistics — entropy

- $dS = \frac{1}{T}dQ = k\beta dQ = kN d(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$.
 $\Rightarrow S = Nk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$.
- $N = e^{-\alpha} Z_1, \Rightarrow \ln Z_1 = \ln N + \alpha$.
- $S = Nk[\ln N + \alpha - \beta \frac{\partial \ln Z_1}{\partial \beta}]$
 $= k[N \ln N + \alpha N + \beta U] = k[N \ln N + \sum(\alpha + \beta \varepsilon_l) a_l]$
- Boltzmann distribution: $\ln \frac{\omega_l}{a_l} = \alpha + \beta \varepsilon_l$
- $\therefore S = k[N \ln N + \sum \ln \frac{\omega_l}{a_l} a_l] = k \ln \Omega_M$.
- Entropy statistical meaning: number of micro-states.
- $F = U - TS$

Thermal quantities in statistics — entropy

- $dS = \frac{1}{T}dQ = k\beta dQ = kN d(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$.
 $\Rightarrow S = Nk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$.
- $N = e^{-\alpha} Z_1, \Rightarrow \ln Z_1 = \ln N + \alpha$.
- $S = Nk[\ln N + \alpha - \beta \frac{\partial \ln Z_1}{\partial \beta}]$
 $= k[N \ln N + \alpha N + \beta U] = k[N \ln N + \sum(\alpha + \beta \varepsilon_l) a_l]$
- Boltzmann distribution: $\ln \frac{\omega_l}{a_l} = \alpha + \beta \varepsilon_l$
- $\therefore S = k[N \ln N + \sum \ln \frac{\omega_l}{a_l} a_l] = k \ln \Omega_M$.
- Entropy statistical meaning: number of micro-states.
- $F = U - TS = -N \frac{\partial \ln Z_1}{\partial \beta} - TNk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$

Thermal quantities in statistics — entropy

- $dS = \frac{1}{T}dQ = k\beta dQ = kN d(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$.
 $\Rightarrow S = Nk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$.
- $N = e^{-\alpha} Z_1, \Rightarrow \ln Z_1 = \ln N + \alpha$.
- $S = Nk[\ln N + \alpha - \beta \frac{\partial \ln Z_1}{\partial \beta}]$
 $= k[N \ln N + \alpha N + \beta U] = k[N \ln N + \sum(\alpha + \beta \varepsilon_l) a_l]$
- Boltzmann distribution: $\ln \frac{\omega_l}{a_l} = \alpha + \beta \varepsilon_l$
- $\therefore S = k[N \ln N + \sum \ln \frac{\omega_l}{a_l} a_l] = k \ln \Omega_M$.
- Entropy statistical meaning: number of micro-states.
- $F = U - TS = -N \frac{\partial \ln Z_1}{\partial \beta} - TNk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$
 $= -NkT \ln Z_1$.

Thermal quantities in statistics — entropy

- $dS = \frac{1}{T} \delta Q = k\beta \delta Q = kN d(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$.
 $\Rightarrow S = Nk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$.
- $N = e^{-\alpha} Z_1, \Rightarrow \ln Z_1 = \ln N + \alpha$.
- $S = Nk[\ln N + \alpha - \beta \frac{\partial \ln Z_1}{\partial \beta}]$
 $= k[N \ln N + \alpha N + \beta U] = k[N \ln N + \sum (\alpha + \beta \varepsilon_l) a_l]$
- Boltzmann distribution: $\ln \frac{\omega_l}{a_l} = \alpha + \beta \varepsilon_l$
- $\therefore S = k[N \ln N + \sum \ln \frac{\omega_l}{a_l} a_l] = k \ln \Omega_M$.
- Entropy statistical meaning: number of micro-states.
- $F = U - TS = -N \frac{\partial \ln Z_1}{\partial \beta} - TNk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$
 $= -NkT \ln Z_1$.
- Conclusion: Given the partition function
 $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$, thermal variables are determined.

Thermal quantities in statistics

- For classical particle system, $\omega_l \rightarrow \frac{\Delta\omega_l}{h_0^r}$,

Thermal quantities in statistics

- For classical particle system, $\omega_l \rightarrow \frac{\Delta\omega_l}{h_0^r}$,
the partition function: $Z_1 = \sum e^{-\beta\epsilon_l} \frac{\Delta\omega_l}{h_0^r}$.

Thermal quantities in statistics

- For classical particle system, $\omega_l \rightarrow \frac{\Delta\omega_l}{h_0^r}$,
the partition function: $Z_1 = \sum e^{-\beta\epsilon_l} \frac{\Delta\omega_l}{h_0^r}$.
- For continuous (q, p) , $\Delta\omega_l \rightarrow d\omega = dq_1 \dots dq_r dp_1 \dots dp_r$,

Thermal quantities in statistics

- For classical particle system, $\omega_l \rightarrow \frac{\Delta\omega_l}{h_0^r}$,
the partition function: $Z_1 = \sum e^{-\beta\epsilon_l} \frac{\Delta\omega_l}{h_0^r}$.
- For continuous (q, p) , $\Delta\omega_l \rightarrow d\omega = dq_1 \dots dq_r dp_1 \dots dp_r$,
the partition function: $Z_1 = \int e^{-\beta\epsilon} \frac{d\omega}{h_0^r}$

Thermal quantities in statistics

- For classical particle system, $\omega_l \rightarrow \frac{\Delta\omega_l}{h_0^r}$,
the partition function: $Z_1 = \sum e^{-\beta\epsilon_l} \frac{\Delta\omega_l}{h_0^r}$.
- For continuous (q, p) , $\Delta\omega_l \rightarrow d\omega = dq_1 \dots dq_r dp_1 \dots dp_r$,
the partition function: $Z_1 = \int e^{-\beta\epsilon} \frac{d\omega}{h_0^r}$
 $= \int \dots \int e^{-\beta\epsilon(p,q)} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r}$.

Table of contents

1 §7 Boltzmann Statistics

- 7.1 Thermal quantities in statistics
- 7.2 Equation of state of ideal gas
- 7.3 Maxwell speed distribution
- 7.4 Energy equipartition theorem
- 7.5 Internal energy and heat capacity of the ideal gas
- 7.6 Entropy of the ideal gas
- 7.7 Einstein's theory on heat capacity of solid
- 7.8 Paramagnetic solid
- 7.9 Negative temperature

7.2 Equation of state of ideal gas

- A simple application of Boltzmann statistics.

7.2 Equation of state of ideal gas

- A simple application of Boltzmann statistics.
- Object: ideal gas with monatomic molecule, nearly independent.

7.2 Equation of state of ideal gas

- A simple application of Boltzmann statistics.
- Object: ideal gas with monatomic molecule, nearly independent.
- Energy of each particle: $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$.

7.2 Equation of state of ideal gas

- A simple application of Boltzmann statistics.
- Object: ideal gas with monatomic molecule, nearly independent.
- Energy of each particle: $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$.
- Micro-states number: $\frac{dx dy dz dp_x dp_y dp_z}{h^3}$.

7.2 Equation of state of ideal gas

- A simple application of Boltzmann statistics.
- Object: ideal gas with monatomic molecule, nearly independent.
- Energy of each particle: $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$.
- Micro-states number: $\frac{dx dy dz dp_x dp_y dp_z}{h^3}$.
- Partition function:

$$Z_1 = \frac{1}{h^3} \int \dots \int e^{-\beta \varepsilon_l} dx dy dz dp_x dp_y dp_z$$

7.2 Equation of state of ideal gas

- A simple application of Boltzmann statistics.
- Object: ideal gas with monatomic molecule, nearly independent.
- Energy of each particle: $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$.
- Micro-states number: $\frac{dx dy dz dp_x dp_y dp_z}{h^3}$.
- Partition function:

$$\begin{aligned} Z_1 &= \frac{1}{h^3} \int \dots \int e^{-\beta \varepsilon_l} dx dy dz dp_x dp_y dp_z \\ &= \frac{1}{h^3} \iiint dx dy dz \iiint e^{-\frac{\beta}{2m}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z \end{aligned}$$

7.2 Equation of state of ideal gas

- A simple application of Boltzmann statistics.
- Object: ideal gas with monatomic molecule, nearly independent.
- Energy of each particle: $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$.
- Micro-states number: $\frac{dx dy dz dp_x dp_y dp_z}{h^3}$.
- Partition function:

$$\begin{aligned}
 Z_1 &= \frac{1}{h^3} \int \dots \int e^{-\beta \varepsilon_l} dx dy dz dp_x dp_y dp_z \\
 &= \frac{1}{h^3} \iiint dx dy dz \iiint e^{-\frac{\beta}{2m}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z \\
 &= \frac{V}{h^3} \int_{-\infty}^{+\infty} e^{-\frac{\beta}{2m}p_x^2} dp_x \int_{-\infty}^{+\infty} e^{-\frac{\beta}{2m}p_y^2} dp_y \int_{-\infty}^{+\infty} e^{-\frac{\beta}{2m}p_z^2} dp_z
 \end{aligned}$$

7.2 Equation of state of ideal gas

- A simple application of Boltzmann statistics.
- Object: ideal gas with monatomic molecule, nearly independent.
- Energy of each particle: $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$.
- Micro-states number: $\frac{dx dy dz dp_x dp_y dp_z}{h^3}$.
- Partition function:

$$\begin{aligned}
 Z_1 &= \frac{1}{h^3} \int \dots \int e^{-\beta \varepsilon_l} dx dy dz dp_x dp_y dp_z \\
 &= \frac{1}{h^3} \iiint dx dy dz \iiint e^{-\frac{\beta}{2m}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z \\
 &= \frac{V}{h^3} \int_{-\infty}^{+\infty} e^{-\frac{\beta}{2m}p_x^2} dp_x \int_{-\infty}^{+\infty} e^{-\frac{\beta}{2m}p_y^2} dp_y \int_{-\infty}^{+\infty} e^{-\frac{\beta}{2m}p_z^2} dp_z \\
 &= \left(\frac{2\pi m}{h^2 \beta} \right)^{\frac{3}{2}} V.
 \end{aligned}$$

Equation of state of ideal gas

- $$p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y}$$

Equation of state of ideal gas

- $$p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V}$$

Equation of state of ideal gas

$$\bullet p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V}$$

Equation of state of ideal gas

$$\bullet p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V}$$

Equation of state of ideal gas

$$\bullet p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V} = \frac{NkT}{V}.$$

Equation of state of ideal gas

- $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V} = \frac{NkT}{V}.$
- Comparing with $pV = nRT$, $k = \frac{n}{N}R$ is determined.

Equation of state of ideal gas

- $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V} = \frac{NkT}{V}.$
- Comparing with $pV = nRT$, $k = \frac{n}{N}R$ is determined.
- Classical limit condition: $\frac{\omega_l}{a_l} \gg 1$, or $e^\alpha \gg 1$.

Equation of state of ideal gas

- $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V} = \frac{NkT}{V}.$
- Comparing with $pV = nRT$, $k = \frac{n}{N}R$ is determined.
- Classical limit condition: $\frac{\omega_l}{a_l} \gg 1$, or $e^\alpha \gg 1$.
$$e^\alpha = \frac{Z_1}{N}$$

Equation of state of ideal gas

- $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V} = \frac{NkT}{V}.$
- Comparing with $pV = nRT$, $k = \frac{n}{N}R$ is determined.
- Classical limit condition: $\frac{\omega_l}{a_l} \gg 1$, or $e^\alpha \gg 1$.
$$e^\alpha = \frac{Z_1}{N} = \frac{V}{N} \left(\frac{2\pi m}{h^2 \beta} \right)^{\frac{3}{2}}$$

Equation of state of ideal gas

- $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V} = \frac{NkT}{V}.$
- Comparing with $pV = nRT$, $k = \frac{n}{N}R$ is determined.
- Classical limit condition: $\frac{\omega_l}{a_l} \gg 1$, or $e^\alpha \gg 1$.
$$e^\alpha = \frac{Z_1}{N} = \frac{V}{N} \left(\frac{2\pi m}{h^2 \beta} \right)^{\frac{3}{2}} = \frac{V}{N} \left(\frac{2\pi m kT}{h^2} \right)^{\frac{3}{2}} \gg 1.$$

Equation of state of ideal gas

- $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V} = \frac{NkT}{V}.$
- Comparing with $pV = nRT$, $k = \frac{n}{N}R$ is determined.
- Classical limit condition: $\frac{\omega_l}{a_l} \gg 1$, or $e^\alpha \gg 1$.

$$e^\alpha = \frac{Z_1}{N} = \frac{V}{N} \left(\frac{2\pi m}{h^2 \beta} \right)^{\frac{3}{2}} = \frac{V}{N} \left(\frac{2\pi m kT}{h^2} \right)^{\frac{3}{2}} \gg 1.$$
 I.e., 1. $N/V \rightarrow 0$, rare;

Equation of state of ideal gas

- $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V} = \frac{NkT}{V}.$
- Comparing with $pV = nRT$, $k = \frac{n}{N}R$ is determined.
- Classical limit condition: $\frac{\omega_l}{a_l} \gg 1$, or $e^\alpha \gg 1$.

$$e^\alpha = \frac{Z_1}{N} = \frac{V}{N} \left(\frac{2\pi m}{h^2 \beta} \right)^{\frac{3}{2}} = \frac{V}{N} \left(\frac{2\pi m kT}{h^2} \right)^{\frac{3}{2}} \gg 1.$$
 I.e., 1. $N/V \rightarrow 0$, rare; 2. high temperature;

Equation of state of ideal gas

- $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V} = \frac{NkT}{V}.$
- Comparing with $pV = nRT$, $k = \frac{n}{N}R$ is determined.
- Classical limit condition: $\frac{\omega_l}{a_l} \gg 1$, or $e^\alpha \gg 1$.

$$e^\alpha = \frac{Z_1}{N} = \frac{V}{N} \left(\frac{2\pi m}{h^2 \beta} \right)^{\frac{3}{2}} = \frac{V}{N} \left(\frac{2\pi m kT}{h^2} \right)^{\frac{3}{2}} \gg 1.$$

i.e., 1. $N/V \rightarrow 0$, rare; 2. high temperature; 3. high mass of the molecule.

Equation of state of ideal gas

- $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V} = \frac{NkT}{V}.$
- Comparing with $pV = nRT$, $k = \frac{n}{N}R$ is determined.
- Classical limit condition: $\frac{\omega_l}{a_l} \gg 1$, or $e^\alpha \gg 1$.

$$e^\alpha = \frac{Z_1}{N} = \frac{V}{N} \left(\frac{2\pi m}{h^2 \beta} \right)^{\frac{3}{2}} = \frac{V}{N} \left(\frac{2\pi m kT}{h^2} \right)^{\frac{3}{2}} \gg 1.$$

i.e., 1. $N/V \rightarrow 0$, rare; 2. high temperature; 3. high mass of the molecule.
- de Broglie wavelength: $\lambda = \frac{h}{p}$

Equation of state of ideal gas

- $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V} = \frac{NkT}{V}.$
- Comparing with $pV = nRT$, $k = \frac{n}{N}R$ is determined.
- Classical limit condition: $\frac{\omega_l}{a_l} \gg 1$, or $e^\alpha \gg 1$.

$$e^\alpha = \frac{Z_1}{N} = \frac{V}{N} \left(\frac{2\pi m}{h^2 \beta} \right)^{\frac{3}{2}} = \frac{V}{N} \left(\frac{2\pi m kT}{h^2} \right)^{\frac{3}{2}} \gg 1.$$

i.e., 1. $N/V \rightarrow 0$, rare; 2. high temperature; 3. high mass of the molecule.
- de Broglie wavelength: $\lambda = \frac{h}{p} = \frac{h}{\sqrt{2m\varepsilon}}$

Equation of state of ideal gas

- $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V} = \frac{NkT}{V}.$
- Comparing with $pV = nRT$, $k = \frac{n}{N}R$ is determined.
- Classical limit condition: $\frac{\omega_l}{a_l} \gg 1$, or $e^\alpha \gg 1$.

$$e^\alpha = \frac{Z_1}{N} = \frac{V}{N} \left(\frac{2\pi m}{h^2 \beta} \right)^{\frac{3}{2}} = \frac{V}{N} \left(\frac{2\pi mkT}{h^2} \right)^{\frac{3}{2}} \gg 1.$$

i.e., 1. $N/V \rightarrow 0$, rare; 2. high temperature; 3. high mass of the molecule.
- de Broglie wavelength: $\lambda = \frac{h}{p} = \frac{h}{\sqrt{2m\varepsilon}} \simeq \frac{h}{\sqrt{2\pi mkT}}.$

Equation of state of ideal gas

- $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V} = \frac{NkT}{V}.$
- Comparing with $pV = nRT$, $k = \frac{n}{N}R$ is determined.
- Classical limit condition: $\frac{\omega_l}{a_l} \gg 1$, or $e^\alpha \gg 1$.

$$e^\alpha = \frac{Z_1}{N} = \frac{V}{N} \left(\frac{2\pi m}{h^2 \beta} \right)^{\frac{3}{2}} = \frac{V}{N} \left(\frac{2\pi m kT}{h^2} \right)^{\frac{3}{2}} \gg 1.$$

I.e., 1. $N/V \rightarrow 0$, rare; 2. high temperature; 3. high mass of the molecule.
- de Broglie wavelength: $\lambda = \frac{h}{p} = \frac{h}{\sqrt{2m\varepsilon}} \simeq \frac{h}{\sqrt{2\pi m kT}}.$
 So, $e^\alpha \gg 1 \Rightarrow \frac{1}{n} \lambda^{-3} \gg 1$, or $n\lambda^3 \ll 1$, or $\lambda \ll \left(\frac{1}{n}\right)^{\frac{1}{3}}.$

Equation of state of ideal gas

$$\bullet p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V} = \frac{NkT}{V}.$$

$$\bullet \text{Comparing with } pV = nRT, k = \frac{n}{N}R \text{ is determined.}$$

$$\bullet \text{Classical limit condition: } \frac{\omega_l}{a_l} \gg 1, \text{ or } e^\alpha \gg 1.$$

$$e^\alpha = \frac{Z_1}{N} = \frac{V}{N} \left(\frac{2\pi m}{h^2 \beta} \right)^{\frac{3}{2}} = \frac{V}{N} \left(\frac{2\pi mkT}{h^2} \right)^{\frac{3}{2}} \gg 1.$$

I.e., 1. $N/V \rightarrow 0$, rare; 2. high temperature; 3. high mass of the molecule.

$$\bullet \text{de Broglie wavelength: } \lambda = \frac{h}{p} = \frac{h}{\sqrt{2m\varepsilon}} \simeq \frac{h}{\sqrt{2\pi mkT}}.$$

$$\text{So, } e^\alpha \gg 1 \Rightarrow \frac{1}{n} \lambda^{-3} \gg 1, \text{ or } n\lambda^3 \ll 1, \text{ or } \lambda \ll \left(\frac{1}{n}\right)^{\frac{1}{3}}.$$

Means, the wavelength should be much less than the distance of any two molecules. (Can be distinguished.)

Table of contents

1 §7 Boltzmann Statistics

- 7.1 Thermal quantities in statistics
- 7.2 Equation of state of ideal gas
- 7.3 Maxwell speed distribution
- 7.4 Energy equipartition theorem
- 7.5 Internal energy and heat capacity of the ideal gas
- 7.6 Entropy of the ideal gas
- 7.7 Einstein's theory on heat capacity of solid
- 7.8 Paramagnetic solid
- 7.9 Negative temperature

7.3 Maxwell speed distribution

7.3 Maxwell speed distribution

- Object: gas. Obeys classical limit,

7.3 Maxwell speed distribution

- Object: gas. Obeys classical limit,
so Boltzmann distribution: $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$.

7.3 Maxwell speed distribution

- Object: gas. Obeys classical limit,
so Boltzmann distribution: $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$.
- Simple case: monatomic molecules.

7.3 Maxwell speed distribution

- Object: gas. Obeys classical limit,
so Boltzmann distribution: $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$.
- Simple case: monatomic molecules.
- Energy: $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$;

7.3 Maxwell speed distribution

- Object: gas. Obeys classical limit,
so Boltzmann distribution: $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$.
- Simple case: monatomic molecules.
- Energy: $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$;
Degeneracy: $\omega = \frac{V}{h^3} dp_x dp_y dp_z$.

7.3 Maxwell speed distribution

- Object: gas. Obeys classical limit,
so Boltzmann distribution: $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$.
- Simple case: monatomic molecules.
- Energy: $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$;
Degeneracy: $\omega = \frac{V}{h^3} dp_x dp_y dp_z$.
- $a = \frac{V}{h^3} e^{-\alpha - \frac{1}{2mkT}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z$.

7.3 Maxwell speed distribution

- Object: gas. Obeys classical limit,
so Boltzmann distribution: $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$.
- Simple case: monatomic molecules.
- Energy: $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$;
Degeneracy: $\omega = \frac{V}{h^3} dp_x dp_y dp_z$.
- $a = \frac{V}{h^3} e^{-\alpha - \frac{1}{2mkT}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z$.
- $N = \sum a_l$

7.3 Maxwell speed distribution

- Object: gas. Obeys classical limit,
so Boltzmann distribution: $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$.
- Simple case: monatomic molecules.
- Energy: $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$;
Degeneracy: $\omega = \frac{V}{h^3} dp_x dp_y dp_z$.
- $a = \frac{V}{h^3} e^{-\alpha - \frac{1}{2mkT}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z$.
- $N = \sum a_l = \frac{V}{h^3} \iiint e^{-\alpha - \frac{1}{2mkT}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z$

7.3 Maxwell speed distribution

- Object: gas. Obeys classical limit,
so Boltzmann distribution: $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$.

- Simple case: monatomic molecules.

- Energy: $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$;

Degeneracy: $\omega = \frac{V}{h^3} dp_x dp_y dp_z$.

- $a = \frac{V}{h^3} e^{-\alpha - \frac{1}{2mkT}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z$.

- $N = \sum a_l = \frac{V}{h^3} \iiint e^{-\alpha - \frac{1}{2mkT}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z =$
 $e^{-\alpha} \frac{V}{h^3} (2mkT)^{\frac{3}{2}} \int e^{-\frac{p_x^2}{2mkT}} d\frac{p_x}{\sqrt{2mkT}} \int e^{-\frac{p_y^2}{2mkT}} d\frac{p_y}{\sqrt{2mkT}} \int e^{-\frac{p_z^2}{2mkT}} d\frac{p_z}{\sqrt{2mkT}}$

7.3 Maxwell speed distribution

- Object: gas. Obeys classical limit,
so Boltzmann distribution: $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$.

- Simple case: monatomic molecules.

- Energy: $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$;

Degeneracy: $\omega = \frac{V}{h^3} dp_x dp_y dp_z$.

- $a = \frac{V}{h^3} e^{-\alpha - \frac{1}{2mkT}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z$.

- $$N = \sum a_l = \frac{V}{h^3} \iiint e^{-\alpha - \frac{1}{2mkT}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z =$$

$$e^{-\alpha} \frac{V}{h^3} (2mkT)^{\frac{3}{2}} \int e^{-\frac{p_x^2}{2mkT}} d\frac{p_x}{\sqrt{2mkT}} \int e^{-\frac{p_y^2}{2mkT}} d\frac{p_y}{\sqrt{2mkT}} \int e^{-\frac{p_z^2}{2mkT}} d\frac{p_z}{\sqrt{2mkT}}$$

$$= e^{-\alpha} \frac{V}{h^3} (2mkT)^{\frac{3}{2}} (\sqrt{\pi})^3$$

7.3 Maxwell speed distribution

- Object: gas. Obeys classical limit,
so Boltzmann distribution: $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$.

- Simple case: monatomic molecules.

- Energy: $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$;

Degeneracy: $\omega = \frac{V}{h^3} dp_x dp_y dp_z$.

- $a = \frac{V}{h^3} e^{-\alpha - \frac{1}{2mkT}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z$.

- $$N = \sum a_l = \frac{V}{h^3} \iiint e^{-\alpha - \frac{1}{2mkT}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z =$$

$$e^{-\alpha} \frac{V}{h^3} (2mkT)^{\frac{3}{2}} \int e^{-\frac{p_x^2}{2mkT}} d\frac{p_x}{\sqrt{2mkT}} \int e^{-\frac{p_y^2}{2mkT}} d\frac{p_y}{\sqrt{2mkT}} \int e^{-\frac{p_z^2}{2mkT}} d\frac{p_z}{\sqrt{2mkT}}$$

$$= e^{-\alpha} \frac{V}{h^3} (2mkT)^{\frac{3}{2}} (\sqrt{\pi})^3 \Rightarrow e^{-\alpha} = \frac{N}{V} \left(\frac{h^2}{2\pi mkT} \right)^{\frac{3}{2}}.$$

7.3 Maxwell speed distribution

- Object: gas. Obeys classical limit,
so Boltzmann distribution: $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$.
- Simple case: monatomic molecules.
- Energy: $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$;
Degeneracy: $\omega = \frac{V}{h^3} dp_x dp_y dp_z$.
- $a = \frac{V}{h^3} e^{-\alpha - \frac{1}{2mkT}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z$.
- $N = \sum a_l = \frac{V}{h^3} \iiint e^{-\alpha - \frac{1}{2mkT}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z =$

$$e^{-\alpha} \frac{V}{h^3} (2mkT)^{\frac{3}{2}} \int e^{-\frac{p_x^2}{2mkT}} d\frac{p_x}{\sqrt{2mkT}} \int e^{-\frac{p_y^2}{2mkT}} d\frac{p_y}{\sqrt{2mkT}} \int e^{-\frac{p_z^2}{2mkT}} d\frac{p_z}{\sqrt{2mkT}}$$

$$= e^{-\alpha} \frac{V}{h^3} (2mkT)^{\frac{3}{2}} (\sqrt{\pi})^3 \Rightarrow e^{-\alpha} = \frac{N}{V} \left(\frac{h^2}{2\pi mkT} \right)^{\frac{3}{2}}.$$
- $\therefore a = N \left(\frac{1}{2\pi mkT} \right)^{\frac{3}{2}} e^{-\frac{1}{2mkT}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z$.

Maxwell speed distribution

- Turn to velocity space: $v_x = \frac{p_x}{m}$,

Maxwell speed distribution

- Turn to velocity space: $v_x = \frac{p_x}{m}$,
 $a \equiv f(v_x, v_y, v_z) dv_x dv_y dv_z$
 $= n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z.$

Maxwell speed distribution

- Turn to velocity space: $v_x = \frac{p_x}{m}$,
 $a \equiv f(v_x, v_y, v_z) dv_x dv_y dv_z$
 $= n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z.$
 $f(v_x, v_y, v_z)$ is the Maxwell velocity distribution.

Maxwell speed distribution

- Turn to velocity space: $v_x = \frac{p_x}{m}$,

$$a \equiv \int f(v_x, v_y, v_z) dv_x dv_y dv_z$$
$$= n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z.$$

$f(v_x, v_y, v_z)$ is the Maxwell velocity distribution.

Obeys $\iiint f(v_x, v_y, v_z) dv_x dv_y dv_z = n$.

Maxwell speed distribution

- Turn to velocity space: $v_x = \frac{p_x}{m}$,
 $a \equiv f(v_x, v_y, v_z)dv_x dv_y dv_z$
 $= n\left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z.$
 $f(v_x, v_y, v_z)$ is the Maxwell velocity distribution.
Obeys $\iiint f(v_x, v_y, v_z)dv_x dv_y dv_z = n.$
- Convert to spheric polar coordinates $(v, \theta, \varphi),$
 $dv_x dv_y dv_z = v^2 \sin \theta dv d\theta d\varphi.$

Maxwell speed distribution

- Turn to velocity space: $v_x = \frac{p_x}{m}$,
 $a \equiv f(v_x, v_y, v_z) dv_x dv_y dv_z$
 $= n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z.$
 $f(v_x, v_y, v_z)$ is the Maxwell velocity distribution.
 Obeys $\iiint f(v_x, v_y, v_z) dv_x dv_y dv_z = n.$
- Convert to spheric polar coordinates (v, θ, φ) ,
 $dv_x dv_y dv_z = v^2 \sin \theta dv d\theta d\varphi.$
 For isotropic velocity, $\int_0^\pi \sin \theta d\theta \int_0^{2\pi} d\varphi = 4\pi,$

Maxwell speed distribution

- Turn to velocity space: $v_x = \frac{p_x}{m}$,

$$a \equiv \int f(v_x, v_y, v_z) dv_x dv_y dv_z$$

$$= n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z.$$

$f(v_x, v_y, v_z)$ is the Maxwell velocity distribution.

Obeys $\iiint f(v_x, v_y, v_z) dv_x dv_y dv_z = n$.

- Convert to spheric polar coordinates (v, θ, φ) ,

$$dv_x dv_y dv_z = v^2 \sin \theta dv d\theta d\varphi.$$

For isotropic velocity, $\int_0^\pi \sin \theta d\theta \int_0^{2\pi} d\varphi = 4\pi$, the number of molecules:

$$f(v) dv = 4\pi n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2 dv.$$

Maxwell speed distribution

- Turn to velocity space: $v_x = \frac{p_x}{m}$,

$$a \equiv \int f(v_x, v_y, v_z) dv_x dv_y dv_z$$

$$= n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z.$$

$f(v_x, v_y, v_z)$ is the Maxwell velocity distribution.

Obeys $\iiint f(v_x, v_y, v_z) dv_x dv_y dv_z = n$.

- Convert to spheric polar coordinates (v, θ, φ) ,

$$dv_x dv_y dv_z = v^2 \sin \theta dv d\theta d\varphi.$$

For isotropic velocity, $\int_0^\pi \sin \theta d\theta \int_0^{2\pi} d\varphi = 4\pi$, the number of molecules:

$$f(v) dv = 4\pi n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2 dv.$$

Maxwell speed distribution. Obeys:

$$\int 4\pi n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2 dv = n.$$

Maxwell speed distribution

$$f(v) = 4\pi n \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2$$

- The most probable speed v_m : $\frac{df(v)}{dv} = 0$.

Maxwell speed distribution

$$f(v) = 4\pi n \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2$$

- The most probable speed v_m : $\frac{df(v)}{dv} = 0$.

$$\frac{d(e^{-\frac{m}{2kT}v^2} v^2)}{dv} = e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{kT}\right) \cdot 2v \cdot v^2 + e^{-\frac{m}{2kT}v^2} \cdot 2v$$

Maxwell speed distribution

$$f(v) = 4\pi n \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2$$

- The most probable speed v_m : $\frac{df(v)}{dv} = 0$.

$$\begin{aligned}\frac{d(e^{-\frac{m}{2kT}v^2} v^2)}{dv} &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{kT}\right) \cdot 2v \cdot v^2 + e^{-\frac{m}{2kT}v^2} \cdot 2v \\ &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2kT}v^2 + 1\right) \cdot 2v\end{aligned}$$

Maxwell speed distribution

$$f(v) = 4\pi n \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2$$

- The most probable speed v_m : $\frac{df(v)}{dv} = 0$.

$$\begin{aligned}\frac{d(e^{-\frac{m}{2kT}v^2} v^2)}{dv} &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2kT}\right) \cdot 2v \cdot v^2 + e^{-\frac{m}{2kT}v^2} \cdot 2v \\ &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2kT}v^2 + 1\right) \cdot 2v = 0,\end{aligned}$$

Maxwell speed distribution

$$f(v) = 4\pi n \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2$$

- The most probable speed v_m : $\frac{df(v)}{dv} = 0$.

$$\begin{aligned}\frac{d(e^{-\frac{m}{2kT}v^2} v^2)}{dv} &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2kT}\right) \cdot 2v \cdot v^2 + e^{-\frac{m}{2kT}v^2} \cdot 2v \\ &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2kT}v^2 + 1\right) \cdot 2v = 0, \Rightarrow v_m^2 = \frac{2kT}{m}.\end{aligned}$$

Maxwell speed distribution

$$f(v) = 4\pi n \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2$$

- The most probable speed v_m : $\frac{df(v)}{dv} = 0$.

$$\begin{aligned}\frac{d(e^{-\frac{m}{2kT}v^2} v^2)}{dv} &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2kT}\right) \cdot 2v \cdot v^2 + e^{-\frac{m}{2kT}v^2} \cdot 2v \\ &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2kT}v^2 + 1\right) \cdot 2v = 0, \Rightarrow v_m^2 = \frac{2kT}{m}.\end{aligned}$$

- Average speed $\bar{v} = \int v f(v) / n dv$

Maxwell speed distribution

$$f(v) = 4\pi n \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2$$

- The most probable speed v_m : $\frac{df(v)}{dv} = 0$.

$$\begin{aligned}\frac{d(e^{-\frac{m}{2kT}v^2} v^2)}{dv} &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2kT}\right) \cdot 2v \cdot v^2 + e^{-\frac{m}{2kT}v^2} \cdot 2v \\ &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2\pi kT}v^2 + 1\right) \cdot 2v = 0, \Rightarrow v_m^2 = \frac{2kT}{m}.\end{aligned}$$

- Average speed $\bar{v} = \int v f(v) / n dv$
$$= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \int_0^\infty v e^{-\frac{m}{2kT}v^2} v^2 dv$$

Maxwell speed distribution

$$f(v) = 4\pi n \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2$$

- The most probable speed v_m : $\frac{df(v)}{dv} = 0$.

$$\begin{aligned}\frac{d(e^{-\frac{m}{2kT}v^2} v^2)}{dv} &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2kT}\right) \cdot 2v \cdot v^2 + e^{-\frac{m}{2kT}v^2} \cdot 2v \\ &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2\pi kT}v^2 + 1\right) \cdot 2v = 0, \Rightarrow v_m^2 = \frac{2kT}{m}.\end{aligned}$$

- Average speed $\bar{v} = \int v f(v) / n dv$

$$\begin{aligned}&= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \int_0^\infty v e^{-\frac{m}{2kT}v^2} v^2 dv \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \int_0^\infty e^{-\frac{m}{2kT}v^2} v^2 dv^2\end{aligned}$$

Maxwell speed distribution

$$f(v) = 4\pi n \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2$$

- The most probable speed v_m : $\frac{df(v)}{dv} = 0$.

$$\begin{aligned} \frac{d(e^{-\frac{m}{2kT}v^2} v^2)}{dv} &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2kT}\right) \cdot 2v \cdot v^2 + e^{-\frac{m}{2kT}v^2} \cdot 2v \\ &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2\pi kT}v^2 + 1\right) \cdot 2v = 0, \Rightarrow v_m^2 = \frac{2kT}{m}. \end{aligned}$$

- Average speed $\bar{v} = \int v f(v) / n dv$

$$\begin{aligned} &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \int_0^\infty v e^{-\frac{m}{2kT}v^2} v^2 dv \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \int_0^\infty e^{-\frac{m}{2kT}v^2} v^2 dv^2 \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(-\frac{2kT}{m}\right) \int_0^\infty v^2 de^{-\frac{m}{2kT}v^2} \end{aligned}$$

Maxwell speed distribution

$$f(v) = 4\pi n \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2$$

- The most probable speed v_m : $\frac{df(v)}{dv} = 0$.

$$\begin{aligned} \frac{d(e^{-\frac{m}{2kT}v^2} v^2)}{dv} &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2kT}\right) \cdot 2v \cdot v^2 + e^{-\frac{m}{2kT}v^2} \cdot 2v \\ &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2kT}v^2 + 1\right) \cdot 2v = 0, \Rightarrow v_m^2 = \frac{2kT}{m}. \end{aligned}$$

- Average speed $\bar{v} = \int v f(v) / n dv$

$$\begin{aligned} &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \int_0^\infty v e^{-\frac{m}{2kT}v^2} v^2 dv \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \int_0^\infty e^{-\frac{m}{2kT}v^2} v^2 dv^2 \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(-\frac{2kT}{m}\right) \int_0^\infty v^2 de^{-\frac{m}{2kT}v^2} \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(-\frac{2kT}{m}\right) [(v^2 e^{-\frac{m}{2kT}v^2})|_0^\infty - \int_0^\infty e^{-\frac{m}{2kT}v^2} dv^2] \end{aligned}$$

Maxwell speed distribution

$$f(v) = 4\pi n \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2$$

- The most probable speed v_m : $\frac{df(v)}{dv} = 0$.

$$\begin{aligned} \frac{d(e^{-\frac{m}{2kT}v^2} v^2)}{dv} &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2kT}\right) \cdot 2v \cdot v^2 + e^{-\frac{m}{2kT}v^2} \cdot 2v \\ &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2\pi kT}v^2 + 1\right) \cdot 2v = 0, \Rightarrow v_m^2 = \frac{2kT}{m}. \end{aligned}$$

- Average speed $\bar{v} = \int v f(v) / n dv$

$$\begin{aligned} &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \int_0^\infty v e^{-\frac{m}{2kT}v^2} v^2 dv \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \int_0^\infty e^{-\frac{m}{2kT}v^2} v^2 dv^2 \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(-\frac{2kT}{m}\right) \int_0^\infty v^2 de^{-\frac{m}{2kT}v^2} \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(-\frac{2kT}{m}\right) [(v^2 e^{-\frac{m}{2kT}v^2})|_0^\infty - \int_0^\infty e^{-\frac{m}{2kT}v^2} dv^2] \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(\frac{2kT}{m}\right) \int_0^\infty e^{-\frac{m}{2kT}v^2} dv^2 \end{aligned}$$

Maxwell speed distribution

$$f(v) = 4\pi n \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2$$

- The most probable speed v_m : $\frac{df(v)}{dv} = 0$.

$$\begin{aligned} \frac{d(e^{-\frac{m}{2kT}v^2} v^2)}{dv} &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2kT}\right) \cdot 2v \cdot v^2 + e^{-\frac{m}{2kT}v^2} \cdot 2v \\ &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2\pi kT}v^2 + 1\right) \cdot 2v = 0, \Rightarrow v_m^2 = \frac{2kT}{m}. \end{aligned}$$

- Average speed $\bar{v} = \int v f(v) / n dv$

$$\begin{aligned} &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \int_0^\infty v e^{-\frac{m}{2kT}v^2} v^2 dv \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \int_0^\infty e^{-\frac{m}{2kT}v^2} v^2 dv^2 \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(-\frac{2kT}{m}\right) \int_0^\infty v^2 de^{-\frac{m}{2kT}v^2} \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(-\frac{2kT}{m}\right) [(v^2 e^{-\frac{m}{2kT}v^2})|_0^\infty - \int_0^\infty e^{-\frac{m}{2kT}v^2} dv^2] \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(\frac{2kT}{m}\right) \int_0^\infty e^{-\frac{m}{2kT}v^2} dv^2 \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(\frac{2kT}{m}\right)^2 \int_0^\infty e^{-\frac{m}{2kT}v^2} d\left(\frac{m}{2kT}v^2\right) \end{aligned}$$

Maxwell speed distribution

$$f(v) = 4\pi n \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2$$

- The most probable speed v_m : $\frac{df(v)}{dv} = 0$.

$$\begin{aligned} \frac{d(e^{-\frac{m}{2kT}v^2} v^2)}{dv} &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2kT}\right) \cdot 2v \cdot v^2 + e^{-\frac{m}{2kT}v^2} \cdot 2v \\ &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2\pi kT}v^2 + 1\right) \cdot 2v = 0, \Rightarrow v_m^2 = \frac{2kT}{m}. \end{aligned}$$

- Average speed $\bar{v} = \int v f(v) / n dv$

$$\begin{aligned} &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \int_0^\infty v e^{-\frac{m}{2kT}v^2} v^2 dv \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \int_0^\infty e^{-\frac{m}{2kT}v^2} v^2 dv^2 \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(-\frac{2kT}{m}\right) \int_0^\infty v^2 de^{-\frac{m}{2kT}v^2} \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(-\frac{2kT}{m}\right) [(v^2 e^{-\frac{m}{2kT}v^2})|_0^\infty - \int_0^\infty e^{-\frac{m}{2kT}v^2} dv^2] \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(\frac{2kT}{m}\right) \int_0^\infty e^{-\frac{m}{2kT}v^2} dv^2 \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(\frac{2kT}{m}\right)^2 \int_0^\infty e^{-\frac{m}{2kT}v^2} d\left(\frac{m}{2kT}v^2\right) \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(\frac{2kT}{m}\right)^2 \int_0^\infty e^{-u} du \end{aligned}$$

Maxwell speed distribution

$$f(v) = 4\pi n \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2$$

- The most probable speed v_m : $\frac{df(v)}{dv} = 0$.

$$\begin{aligned} \frac{d(e^{-\frac{m}{2kT}v^2} v^2)}{dv} &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2kT}\right) \cdot 2v \cdot v^2 + e^{-\frac{m}{2kT}v^2} \cdot 2v \\ &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2kT}v^2 + 1\right) \cdot 2v = 0, \Rightarrow v_m^2 = \frac{2kT}{m}. \end{aligned}$$

- Average speed $\bar{v} = \int v f(v) / n dv$

$$\begin{aligned} &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \int_0^\infty v e^{-\frac{m}{2kT}v^2} v^2 dv \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \int_0^\infty e^{-\frac{m}{2kT}v^2} v^2 dv^2 \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(-\frac{2kT}{m}\right) \int_0^\infty v^2 de^{-\frac{m}{2kT}v^2} \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(-\frac{2kT}{m}\right) [(v^2 e^{-\frac{m}{2kT}v^2})|_0^\infty - \int_0^\infty e^{-\frac{m}{2kT}v^2} dv^2] \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(\frac{2kT}{m}\right) \int_0^\infty e^{-\frac{m}{2kT}v^2} dv^2 \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(\frac{2kT}{m}\right)^2 \int_0^\infty e^{-\frac{m}{2kT}v^2} d\left(\frac{m}{2kT}v^2\right) \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(\frac{2kT}{m}\right)^2 = \sqrt{\frac{8kT}{\pi m}}. \end{aligned}$$

Maxwell speed distribution

- Root-mean-square speed:

Maxwell speed distribution

- Root-mean-square speed:

$$v_s^2 \equiv \overline{v^2}$$

Maxwell speed distribution

- Root-mean-square speed:

$$v_s^2 \equiv \bar{v^2} = \int v^2 f(v) / n dv$$

Maxwell speed distribution

- Root-mean-square speed:

$$v_s^2 \equiv \bar{v^2} = \int v^2 f(v) / n dv = \frac{3kT}{m}.$$

Maxwell speed distribution

- Root-mean-square speed:

$$v_s^2 \equiv \bar{v^2} = \int v^2 f(v) / n dv = \frac{3kT}{m}.$$

Useful in calculating average energy $\bar{\varepsilon} = \frac{1}{2} m \bar{v^2}$.

Maxwell speed distribution

- Root-mean-square speed:

$$v_s^2 \equiv \bar{v^2} = \int v^2 f(v) / n dv = \frac{3kT}{m}.$$

Useful in calculating average energy $\bar{\varepsilon} = \frac{1}{2} m \bar{v^2}$.

Example: N_2 at 0°C .

Maxwell speed distribution

- Root-mean-square speed:

$$v_s^2 \equiv \bar{v^2} = \int v^2 f(v) / n dv = \frac{3kT}{m}.$$

Useful in calculating average energy $\bar{\varepsilon} = \frac{1}{2} m \bar{v^2}$.

Example: N_2 at 0°C . $m = 28 \times 1.67 \times 10^{-27} \text{kg}$,

Maxwell speed distribution

- Root-mean-square speed:

$$v_s^2 \equiv \bar{v^2} = \int v^2 f(v) n dv = \frac{3kT}{m}.$$

Useful in calculating average energy $\bar{\varepsilon} = \frac{1}{2} m \bar{v^2}$.

Example: N_2 at 0°C . $m = 28 \times 1.67 \times 10^{-27} \text{kg}$,

$$v_s = \left(\frac{3kT}{m} \right)^{\frac{1}{2}}$$

Maxwell speed distribution

- Root-mean-square speed:

$$v_s^2 \equiv \bar{v^2} = \int v^2 f(v) / n dv = \frac{3kT}{m}.$$

Useful in calculating average energy $\bar{\epsilon} = \frac{1}{2} m \bar{v^2}$.

Example: N_2 at 0°C . $m = 28 \times 1.67 \times 10^{-27} \text{kg}$,

$$v_s = \left(\frac{3kT}{m} \right)^{\frac{1}{2}} = \left(\frac{3 \cdot 1.38 \times 10^{-23} \text{JK}^{-1} \cdot 273 \text{K}}{28 \times 1.67 \times 10^{-27} \text{kg}} \right)^{\frac{1}{2}}$$

Maxwell speed distribution

- Root-mean-square speed:

$$v_s^2 \equiv \bar{v^2} = \int v^2 f(v) / n dv = \frac{3kT}{m}.$$

Useful in calculating average energy $\bar{\epsilon} = \frac{1}{2} m \bar{v^2}$.

Example: N_2 at 0°C . $m = 28 \times 1.67 \times 10^{-27} \text{kg}$,

$$v_s = \left(\frac{3kT}{m} \right)^{\frac{1}{2}} = \left(\frac{3 \cdot 1.38 \times 10^{-23} \text{JK}^{-1} \cdot 273 \text{K}}{28 \times 1.67 \times 10^{-27} \text{kg}} \right)^{\frac{1}{2}} \simeq 493 \text{m/s}.$$

Maxwell speed distribution

- Root-mean-square speed:

$$v_s^2 \equiv \bar{v^2} = \int v^2 f(v) / n dv = \frac{3kT}{m}.$$


Useful in calculating average energy $\bar{\varepsilon} = \frac{1}{2} m \bar{v^2}$.

Example: N_2 at 0°C . $m = 28 \times 1.67 \times 10^{-27} \text{kg}$,

$$v_s = \left(\frac{3kT}{m} \right)^{\frac{1}{2}} = \left(\frac{3 \cdot 1.38 \times 10^{-23} \text{JK}^{-1} \cdot 273 \text{K}}{28 \times 1.67 \times 10^{-27} \text{kg}} \right)^{\frac{1}{2}} \simeq 493 \text{m/s}.$$

Application 1: frequency of colliding.

Γ : Number of colliding molecules on the surface in unit time unit area.



Maxwell speed distribution

- Root-mean-square speed:

$$v_s^2 \equiv \bar{v^2} = \int v^2 f(v) / n dv = \frac{3kT}{m}.$$


Useful in calculating average energy $\bar{\varepsilon} = \frac{1}{2} m \bar{v^2}$.

Example: N_2 at 0°C . $m = 28 \times 1.67 \times 10^{-27} \text{kg}$,

$$v_s = \left(\frac{3kT}{m} \right)^{\frac{1}{2}} = \left(\frac{3 \cdot 1.38 \times 10^{-23} \text{JK}^{-1} \cdot 273 \text{K}}{28 \times 1.67 \times 10^{-27} \text{kg}} \right)^{\frac{1}{2}} \simeq 493 \text{m/s}.$$

Application 1: frequency of colliding.

Γ : Number of colliding molecules on the surface in unit time unit area.



For a given velocity, (v_x, v_y, v_z) , all particles inside the oblique cylinder can collide.

Maxwell speed distribution

- Root-mean-square speed:

$$v_s^2 \equiv \bar{v^2} = \int v^2 f(v) / n dv = \frac{3kT}{m}.$$


Useful in calculating average energy $\bar{\varepsilon} = \frac{1}{2} m \bar{v^2}$.

Example: N_2 at 0°C . $m = 28 \times 1.67 \times 10^{-27} \text{kg}$,

$$v_s = \left(\frac{3kT}{m} \right)^{\frac{1}{2}} = \left(\frac{3 \cdot 1.38 \times 10^{-23} \text{JK}^{-1} \cdot 273 \text{K}}{28 \times 1.67 \times 10^{-27} \text{kg}} \right)^{\frac{1}{2}} \simeq 493 \text{m/s}.$$

Application 1: frequency of colliding.

Γ : Number of colliding molecules on the surface in unit time unit area.



For a given velocity, (v_x, v_y, v_z) , all particles inside the oblique cylinder can collide. The number is

$$f(v_x, v_y, v_z) dv_x dv_y dv_z \cdot v_x dt dA.$$

Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- In unit time unit area, all velocity:

$$\Gamma = \iiint f(v_x, v_y, v_z) v_x dv_x dv_y dv_z$$

Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- In unit time unit area, all velocity:

$$\begin{aligned}\Gamma &= \iiint f(v_x, v_y, v_z) v_x dv_x dv_y dv_z \\ &= n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} \iiint v_x e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z\end{aligned}$$

Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- In unit time unit area, all velocity:

$$\begin{aligned} \Gamma &= \iiint f(v_x, v_y, v_z) v_x dv_x dv_y dv_z \\ &= n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} \iiint v_x e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z \\ &= n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} \int_0^\infty v_x e^{-\frac{m}{2kT}v_x^2} dv_x \iint_{-\infty}^\infty e^{-\frac{m}{2kT}(v_y^2 + v_z^2)} dv_y dv_z \end{aligned}$$

Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- In unit time unit area, all velocity:

$$\begin{aligned} \Gamma &= \iiint f(v_x, v_y, v_z) v_x dv_x dv_y dv_z \\ &= n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} \iiint v_x e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z \\ &= n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} \int_0^\infty v_x e^{-\frac{m}{2kT}v_x^2} dv_x \iint_{-\infty}^\infty e^{-\frac{m}{2kT}(v_y^2 + v_z^2)} dv_y dv_z \\ &= n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} \cdot \left(\frac{1}{2} \frac{2kT}{m} \right) \cdot \left(\frac{2kT}{m} \pi \right) \end{aligned}$$

Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- In unit time unit area, all velocity:

$$\begin{aligned} \Gamma &= \iiint f(v_x, v_y, v_z) v_x dv_x dv_y dv_z \\ &= n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} \iiint v_x e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z \\ &= n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} \int_0^\infty v_x e^{-\frac{m}{2kT}v_x^2} dv_x \iint_{-\infty}^\infty e^{-\frac{m}{2kT}(v_y^2 + v_z^2)} dv_y dv_z \\ &= n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} \cdot \left(\frac{1}{2} \frac{2kT}{m} \right) \cdot \left(\frac{2kT}{m} \pi \right) \\ &= n \sqrt{\frac{kT}{2\pi m}} \end{aligned}$$

Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- In unit time unit area, all velocity:

$$\begin{aligned} \Gamma &= \iiint f(v_x, v_y, v_z) v_x dv_x dv_y dv_z \\ &= n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} \iiint v_x e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z \\ &= n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} \int_0^\infty v_x e^{-\frac{m}{2kT}v_x^2} dv_x \iint_{-\infty}^\infty e^{-\frac{m}{2kT}(v_y^2 + v_z^2)} dv_y dv_z \\ &= n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} \cdot \left(\frac{1}{2} \frac{2kT}{m} \right) \cdot \left(\frac{2kT}{m} \pi \right) \\ &= n \sqrt{\frac{kT}{2\pi m}} \\ &= \frac{1}{4} n \bar{v}. \end{aligned}$$

Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- Application 2: Pressure.

Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- Application 2: Pressure.

$$p = \frac{F}{A}$$

Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- Application 2: Pressure.

$$p = \frac{F}{A} = \frac{\Delta p_x}{\Delta t A}$$

Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- Application 2: Pressure.

$$p = \frac{F}{A} = \frac{\Delta p_x}{\Delta t A} = \frac{\sum m v_x \cdot 2}{dA dt}$$

Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- Application 2: Pressure.

$$p = \frac{F}{A} = \frac{\Delta p_x}{\Delta t A} = \frac{\sum m v_x \cdot 2}{dA dt}$$

Number of particles $f(v_x, v_y, v_z) v_x dv_x dv_y dv_z dA dt$.

Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- Application 2: Pressure.

$$p = \frac{F}{A} = \frac{\Delta p_x}{\Delta t A} = \frac{\sum m v_x \cdot 2}{dA dt}$$

Number of particles $f(v_x, v_y, v_z) v_x dv_x dv_y dv_z dA dt$.

- $\therefore p = \iiint f(v_x, v_y, v_z) v_x \cdot 2m v_x dv_x dv_y dv_z$

Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- Application 2: Pressure.

$$p = \frac{F}{A} = \frac{\Delta p_x}{\Delta t A} = \frac{\sum m v_x \cdot 2}{dA dt}$$

Number of particles $f(v_x, v_y, v_z) v_x dv_x dv_y dv_z dA dt$.

- $\therefore p = \iiint f(v_x, v_y, v_z) v_x \cdot 2m v_x dv_x dv_y dv_z$
 $= n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} \iiint 2m v_x^2 e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z$

Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- Application 2: Pressure.

$$p = \frac{F}{A} = \frac{\Delta p_x}{\Delta t A} = \frac{\sum m v_x \cdot 2}{dA dt}$$

Number of particles $f(v_x, v_y, v_z) v_x dv_x dv_y dv_z dA dt$.

$$\begin{aligned} \bullet \therefore p &= \iiint f(v_x, v_y, v_z) v_x \cdot 2m v_x dv_x dv_y dv_z \\ &= n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} \iiint 2m v_x^2 e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z \\ &= 2mn \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} \left(\frac{2kT}{m} \right)^{\frac{3}{2}} \int_0^\infty \left(\frac{m}{2kT} v_x^2 \right) e^{-\frac{m}{2kT} v_x^2} d\left(\sqrt{\frac{m}{2kT}} v_x \right) \cdot \\ &\quad \frac{2kT}{m} \iint_{-\infty}^\infty e^{-\frac{m}{2kT}(v_y^2 + v_z^2)} d\left(\sqrt{\frac{m}{2kT}} v_y \right) d\left(\sqrt{\frac{m}{2kT}} v_z \right) \end{aligned}$$

Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- Application 2: Pressure.

$$p = \frac{F}{A} = \frac{\Delta p_x}{\Delta t A} = \frac{\sum m v_x \cdot 2}{dA dt}$$

Number of particles $f(v_x, v_y, v_z) v_x dv_x dv_y dv_z dA dt$.

- $\therefore p = \iiint f(v_x, v_y, v_z) v_x \cdot 2m v_x dv_x dv_y dv_z$
 $= n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} \iiint 2m v_x^2 e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z$
 $= 2mn \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} \left(\frac{2kT}{m} \right)^{\frac{3}{2}} \int_0^\infty \left(\frac{m}{2kT} v_x^2 \right) e^{-\frac{m}{2kT} v_x^2} d\left(\sqrt{\frac{m}{2kT}} v_x \right) \cdot$
 $\frac{2kT}{m} \int_{-\infty}^\infty e^{-\frac{m}{2kT}(v_y^2 + v_z^2)} d\left(\sqrt{\frac{m}{2kT}} v_y \right) d\left(\sqrt{\frac{m}{2kT}} v_z \right)$
 $= 2mn \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} \left(\frac{2kT}{m} \right)^{\frac{3}{2}} \cdot \frac{\sqrt{\pi}}{4} \cdot \frac{2kT}{m} \pi$

Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- Application 2: Pressure.

$$p = \frac{F}{A} = \frac{\Delta p_x}{\Delta t A} = \frac{\sum m v_x \cdot 2}{dA dt}$$

Number of particles $f(v_x, v_y, v_z) v_x dv_x dv_y dv_z dA dt$.

- $\therefore p = \iiint f(v_x, v_y, v_z) v_x \cdot 2m v_x dv_x dv_y dv_z$
 $= n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} \iiint 2m v_x^2 e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z$
 $= 2mn \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} \left(\frac{2kT}{m} \right)^{\frac{3}{2}} \int_0^\infty \left(\frac{m}{2kT} v_x^2 \right) e^{-\frac{m}{2kT} v_x^2} d\left(\sqrt{\frac{m}{2kT}} v_x \right) \cdot$
 $\frac{2kT}{m} \iint_{-\infty}^\infty e^{-\frac{m}{2kT}(v_y^2 + v_z^2)} d\left(\sqrt{\frac{m}{2kT}} v_y \right) d\left(\sqrt{\frac{m}{2kT}} v_z \right)$
 $= 2mn \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} \left(\frac{2kT}{m} \right)^{\frac{3}{2}} \cdot \frac{\sqrt{\pi}}{4} \cdot \frac{2kT}{m} \pi$
 $= nkT.$

Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- Application 2: Pressure.

$$p = \frac{F}{A} = \frac{\Delta p_x}{\Delta t A} = \frac{\sum m v_x \cdot 2}{dA dt}$$

Number of particles $f(v_x, v_y, v_z) v_x dv_x dv_y dv_z dA dt$.

- $\therefore p = \iiint f(v_x, v_y, v_z) v_x \cdot 2m v_x dv_x dv_y dv_z$
 $= n \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} \iiint 2m v_x^2 e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z$
 $= 2mn \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} \left(\frac{2kT}{m} \right)^{\frac{3}{2}} \int_0^\infty \left(\frac{m}{2kT} v_x^2 \right) e^{-\frac{m}{2kT} v_x^2} d\left(\sqrt{\frac{m}{2kT}} v_x \right) \cdot$
 $\frac{2kT}{m} \iint_{-\infty}^\infty e^{-\frac{m}{2kT}(v_y^2 + v_z^2)} d\left(\sqrt{\frac{m}{2kT}} v_y \right) d\left(\sqrt{\frac{m}{2kT}} v_z \right)$
 $= 2mn \left(\frac{m}{2\pi kT} \right)^{\frac{3}{2}} \left(\frac{2kT}{m} \right)^{\frac{3}{2}} \cdot \frac{\sqrt{\pi}}{4} \cdot \frac{2kT}{m} \pi$
 $= nkT.$

Same as eq (7.2.5).

Table of contents

1 §7 Boltzmann Statistics

- 7.1 Thermal quantities in statistics
- 7.2 Equation of state of ideal gas
- 7.3 Maxwell speed distribution
- 7.4 Energy equipartition theorem
- 7.5 Internal energy and heat capacity of the ideal gas
- 7.6 Entropy of the ideal gas
- 7.7 Einstein's theory on heat capacity of solid
- 7.8 Paramagnetic solid
- 7.9 Negative temperature

7.4 Energy equipartition theorem

7.4 Energy equipartition theorem

- Theorem: any degree of freedom x which appears in the total energy ε only as a simple quadratic term, has an average energy of $\frac{1}{2}kT$ in thermal equilibrium.

7.4 Energy equipartition theorem

- Theorem: any degree of freedom x which appears in the total energy ε only as a simple quadratic term, has an average energy of $\frac{1}{2}kT$ in thermal equilibrium.
- Example: monatomic ideal gas.

7.4 Energy equipartition theorem

- Theorem: any degree of freedom x which appears in the total energy ε only as a simple quadratic term, has an average energy of $\frac{1}{2}kT$ in thermal equilibrium.
- Example: monatomic ideal gas.
$$\bar{\varepsilon} = \overline{\frac{1}{2}mv^2} = \overline{\frac{1}{2}m(v_x^2 + v_y^2 + v_z^2)}$$

7.4 Energy equipartition theorem

- Theorem: any degree of freedom x which appears in the total energy ε only as a simple quadratic term, has an average energy of $\frac{1}{2}kT$ in thermal equilibrium.
- Example: monatomic ideal gas.

$$\begin{aligned}\bar{\varepsilon} &= \overline{\frac{1}{2}mv^2} = \overline{\frac{1}{2}m(v_x^2 + v_y^2 + v_z^2)} \\ &= \frac{1}{2}m\overline{v^2}\end{aligned}$$

7.4 Energy equipartition theorem

- Theorem: any degree of freedom x which appears in the total energy ε only as a simple quadratic term, has an average energy of $\frac{1}{2}kT$ in thermal equilibrium.
- Example: monatomic ideal gas.

$$\begin{aligned}\bar{\varepsilon} &= \overline{\frac{1}{2}mv^2} = \overline{\frac{1}{2}m(v_x^2 + v_y^2 + v_z^2)} \\ &= \frac{1}{2}m\overline{v^2} = \frac{1}{2}m \cdot \frac{3kT}{m}\end{aligned}$$

7.4 Energy equipartition theorem

- Theorem: any degree of freedom x which appears in the total energy ε only as a simple quadratic term, has an average energy of $\frac{1}{2}kT$ in thermal equilibrium.
- Example: monatomic ideal gas.

$$\begin{aligned}\bar{\varepsilon} &= \overline{\frac{1}{2}mv^2} = \overline{\frac{1}{2}m(v_x^2 + v_y^2 + v_z^2)} \\ &= \frac{1}{2}m\overline{v^2} = \frac{1}{2}m \cdot \frac{3kT}{m} = \frac{3}{2}kT.\end{aligned}$$

7.4 Energy equipartition theorem

- Theorem: any degree of freedom x which appears in the total energy ε only as a simple quadratic term, has an average energy of $\frac{1}{2}kT$ in thermal equilibrium.
- Example: monatomic ideal gas.

$$\begin{aligned}\bar{\varepsilon} &= \overline{\frac{1}{2}mv^2} = \overline{\frac{1}{2}m(v_x^2 + v_y^2 + v_z^2)} \\ &= \frac{1}{2}m\overline{v^2} = \frac{1}{2}m \cdot \frac{3kT}{m} = \frac{3}{2}kT.\end{aligned}$$

- Proof: $\varepsilon = \varepsilon_k + \varepsilon_p$,

7.4 Energy equipartition theorem

- Theorem: any degree of freedom x which appears in the total energy ε only as a simple quadratic term, has an average energy of $\frac{1}{2}kT$ in thermal equilibrium.
- Example: monatomic ideal gas.

$$\begin{aligned}\bar{\varepsilon} &= \overline{\frac{1}{2}mv^2} = \overline{\frac{1}{2}m(v_x^2 + v_y^2 + v_z^2)} \\ &= \frac{1}{2}m\overline{v^2} = \frac{1}{2}m \cdot \frac{3kT}{m} = \frac{3}{2}kT.\end{aligned}$$

- Proof: $\varepsilon = \varepsilon_k + \varepsilon_p$, where $\varepsilon_k = \frac{1}{2} \sum a_i p_i^2$.

7.4 Energy equipartition theorem

- Theorem: any degree of freedom x which appears in the total energy ε only as a simple quadratic term, has an average energy of $\frac{1}{2}kT$ in thermal equilibrium.
- Example: monatomic ideal gas.

$$\begin{aligned}\bar{\varepsilon} &= \overline{\frac{1}{2}mv^2} = \overline{\frac{1}{2}m(v_x^2 + v_y^2 + v_z^2)} \\ &= \frac{1}{2}m\overline{v^2} = \frac{1}{2}m \cdot \frac{3kT}{m} = \frac{3}{2}kT.\end{aligned}$$

- Proof: $\varepsilon = \varepsilon_k + \varepsilon_p$, where $\varepsilon_k = \frac{1}{2} \sum a_i p_i^2$.

Considering the 1st term,

$$\overline{\frac{1}{2}a_1 p_1^2}$$

7.4 Energy equipartition theorem

- Theorem: any degree of freedom x which appears in the total energy ε only as a simple quadratic term, has an average energy of $\frac{1}{2}kT$ in thermal equilibrium.
- Example: monatomic ideal gas.

$$\begin{aligned}\bar{\varepsilon} &= \overline{\frac{1}{2}mv^2} = \overline{\frac{1}{2}m(v_x^2 + v_y^2 + v_z^2)} \\ &= \frac{1}{2}m\overline{v^2} = \frac{1}{2}m \cdot \frac{3kT}{m} = \frac{3}{2}kT.\end{aligned}$$

- Proof: $\varepsilon = \varepsilon_k + \varepsilon_p$, where $\varepsilon_k = \frac{1}{2} \sum a_i p_i^2$.

Considering the 1st term,

$$\overline{\frac{1}{2}a_1 p_1^2} = \frac{1}{N} \sum \frac{1}{2}a_1 p_1^2 \cdot a_l$$

7.4 Energy equipartition theorem

- Theorem: any degree of freedom x which appears in the total energy ε only as a simple quadratic term, has an average energy of $\frac{1}{2}kT$ in thermal equilibrium.
- Example: monatomic ideal gas.

$$\begin{aligned}\bar{\varepsilon} &= \overline{\frac{1}{2}mv^2} = \overline{\frac{1}{2}m(v_x^2 + v_y^2 + v_z^2)} \\ &= \frac{1}{2}m\overline{v^2} = \frac{1}{2}m \cdot \frac{3kT}{m} = \frac{3}{2}kT.\end{aligned}$$

- Proof: $\varepsilon = \varepsilon_k + \varepsilon_p$, where $\varepsilon_k = \frac{1}{2} \sum a_i p_i^2$.

Considering the 1st term,

$$\begin{aligned}\overline{\frac{1}{2}a_1 p_1^2} &= \frac{1}{N} \sum \frac{1}{2}a_1 p_1^2 \cdot a_l \\ &= \frac{1}{N} \sum \frac{1}{2}a_1 p_1^2 \cdot \omega_l e^{-\alpha - \beta \varepsilon_l}\end{aligned}$$

7.4 Energy equipartition theorem

- Theorem: any degree of freedom x which appears in the total energy ε only as a simple quadratic term, has an average energy of $\frac{1}{2}kT$ in thermal equilibrium.
- Example: monatomic ideal gas.

$$\begin{aligned}\bar{\varepsilon} &= \overline{\frac{1}{2}mv^2} = \overline{\frac{1}{2}m(v_x^2 + v_y^2 + v_z^2)} \\ &= \frac{1}{2}m\overline{v^2} = \frac{1}{2}m \cdot \frac{3kT}{m} = \frac{3}{2}kT.\end{aligned}$$

- Proof: $\varepsilon = \varepsilon_k + \varepsilon_p$, where $\varepsilon_k = \frac{1}{2} \sum a_i p_i^2$.

Considering the 1st term,

$$\begin{aligned}\overline{\frac{1}{2}a_1 p_1^2} &= \frac{1}{N} \sum \frac{1}{2}a_1 p_1^2 \cdot a_l \\ &= \frac{1}{N} \sum \frac{1}{2}a_1 p_1^2 \cdot \omega_l e^{-\alpha - \beta \varepsilon_l} \\ &= \frac{1}{N} \int \frac{1}{2}a_1 p_1^2 \cdot e^{-\alpha - \beta \varepsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r}\end{aligned}$$

7.4 Energy equipartition theorem

- Theorem: any degree of freedom x which appears in the total energy ε only as a simple quadratic term, has an average energy of $\frac{1}{2}kT$ in thermal equilibrium.
- Example: monatomic ideal gas.

$$\begin{aligned}\bar{\varepsilon} &= \overline{\frac{1}{2}mv^2} = \overline{\frac{1}{2}m(v_x^2 + v_y^2 + v_z^2)} \\ &= \frac{1}{2}m\overline{v^2} = \frac{1}{2}m \cdot \frac{3kT}{m} = \frac{3}{2}kT.\end{aligned}$$

- Proof: $\varepsilon = \varepsilon_k + \varepsilon_p$, where $\varepsilon_k = \frac{1}{2} \sum a_i p_i^2$.

Considering the 1st term,

$$\begin{aligned}\overline{\frac{1}{2}a_1 p_1^2} &= \frac{1}{N} \sum \frac{1}{2}a_1 p_1^2 \cdot a_l \\ &= \frac{1}{N} \sum \frac{1}{2}a_1 p_1^2 \cdot \omega_l e^{-\alpha - \beta \varepsilon_l} \\ &= \frac{1}{N} \int \frac{1}{2}a_1 p_1^2 \cdot e^{-\alpha - \beta \varepsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r} \\ &= \frac{1}{Z} \int \frac{1}{2}a_1 p_1^2 \cdot e^{-\beta \varepsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h^r} \dots\end{aligned}$$

Energy equipartition theorem

- $\overline{\frac{1}{2}a_1p_1^2} = \frac{1}{Z_1} \int \frac{1}{2}a_1p_1^2 \cdot e^{-\beta\epsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r}$

Energy equipartition theorem

- $$\overline{\frac{1}{2}a_1p_1^2} = \frac{1}{Z_1} \int \frac{1}{2}a_1p_1^2 \cdot e^{-\beta\epsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r}$$

$$= \int_{-\infty}^{\infty} \frac{1}{2}a_1p_1^2 e^{-\beta\frac{1}{2}a_1p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\epsilon - \frac{1}{2}a_1p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r}$$

Energy equipartition theorem

$$\begin{aligned}
 \bullet \quad \overline{\frac{1}{2}a_1 p_1^2} &= \frac{1}{Z_1} \int \frac{1}{2}a_1 p_1^2 \cdot e^{-\beta \varepsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{2}a_1 p_1^2 e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{4}a_1 p_1 e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1^2 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r}
 \end{aligned}$$

Energy equipartition theorem

$$\begin{aligned}
 \bullet \quad \overline{\frac{1}{2}a_1 p_1^2} &= \frac{1}{Z_1} \int \frac{1}{2}a_1 p_1^2 \cdot e^{-\beta \varepsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{2}a_1 p_1^2 e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{4}a_1 p_1 e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1^2 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{4}a_1 p_1 \cdot \left(\frac{-2}{\beta a_1}\right) de^{-\beta \frac{1}{2}a_1 p_1^2} \cdot \\
 &\quad \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r}
 \end{aligned}$$

Energy equipartition theorem

$$\begin{aligned}
 \bullet \quad \overline{\frac{1}{2}a_1 p_1^2} &= \frac{1}{Z_1} \int \frac{1}{2}a_1 p_1^2 \cdot e^{-\beta \varepsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{2}a_1 p_1^2 e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{4}a_1 p_1 e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1^2 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{4}a_1 p_1 \cdot \left(\frac{-2}{\beta a_1}\right) de^{-\beta \frac{1}{2}a_1 p_1^2} \cdot \\
 &\quad \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \frac{1}{2\beta} \int_{-\infty}^{\infty} e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r}
 \end{aligned}$$

Energy equipartition theorem

$$\begin{aligned}
 \bullet \quad \overline{\frac{1}{2}a_1 p_1^2} &= \frac{1}{Z_1} \int \frac{1}{2}a_1 p_1^2 \cdot e^{-\beta \varepsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{2}a_1 p_1^2 e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{4}a_1 p_1 e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1^2 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{4}a_1 p_1 \cdot \left(\frac{-2}{\beta a_1}\right) de^{-\beta \frac{1}{2}a_1 p_1^2} \cdot \\
 &\quad \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \frac{1}{2\beta} \int_{-\infty}^{\infty} e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \frac{1}{2\beta} \frac{1}{Z_1} \int e^{-\beta \varepsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r}
 \end{aligned}$$

Energy equipartition theorem

$$\begin{aligned}
 \bullet \quad \overline{\frac{1}{2}a_1 p_1^2} &= \frac{1}{Z_1} \int \frac{1}{2}a_1 p_1^2 \cdot e^{-\beta \varepsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{2}a_1 p_1^2 e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{4}a_1 p_1 e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1^2 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{4}a_1 p_1 \cdot \left(\frac{-2}{\beta a_1}\right) de^{-\beta \frac{1}{2}a_1 p_1^2} \cdot \\
 &\quad \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \frac{1}{2\beta} \int_{-\infty}^{\infty} e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \frac{1}{2\beta} \frac{1}{Z_1} \int e^{-\beta \varepsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r} \\
 &= \frac{1}{2\beta}
 \end{aligned}$$

Energy equipartition theorem

$$\begin{aligned}
 \bullet \quad \overline{\frac{1}{2}a_1 p_1^2} &= \frac{1}{Z_1} \int \frac{1}{2}a_1 p_1^2 \cdot e^{-\beta \varepsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{2}a_1 p_1^2 e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{4}a_1 p_1 e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1^2 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{4}a_1 p_1 \cdot \left(\frac{-2}{\beta a_1}\right) de^{-\beta \frac{1}{2}a_1 p_1^2} \cdot \\
 &\quad \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \frac{1}{2\beta} \int_{-\infty}^{\infty} e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \frac{1}{2\beta} \frac{1}{Z_1} \int e^{-\beta \varepsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r} \\
 &= \frac{1}{2\beta} = \frac{1}{2}kT.
 \end{aligned}$$

Energy equipartition theorem

- $$\begin{aligned}
 \overline{\frac{1}{2}a_1p_1^2} &= \frac{1}{Z_1} \int \frac{1}{2}a_1p_1^2 \cdot e^{-\beta\varepsilon} \frac{dq_1\dots dq_r dp_1\dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{2}a_1p_1^2 e^{-\beta\frac{1}{2}a_1p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1p_1^2)} \frac{dq_1\dots dq_r dp_2\dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{4}a_1p_1 e^{-\beta\frac{1}{2}a_1p_1^2} dp_1^2 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1p_1^2)} \frac{dq_1\dots dq_r dp_2\dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{4}a_1p_1 \cdot \left(\frac{-2}{\beta a_1}\right) de^{-\beta\frac{1}{2}a_1p_1^2} \cdot \\
 &\quad \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1p_1^2)} \frac{dq_1\dots dq_r dp_2\dots dp_r}{h_0^r} \\
 &= \frac{1}{2\beta} \int_{-\infty}^{\infty} e^{-\beta\frac{1}{2}a_1p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1p_1^2)} \frac{dq_1\dots dq_r dp_2\dots dp_r}{h_0^r} \\
 &= \frac{1}{2\beta} \frac{1}{Z_1} \int e^{-\beta\varepsilon} \frac{dq_1\dots dq_r dp_1\dots dp_r}{h_0^r} \\
 &= \frac{1}{2\beta} = \frac{1}{2}kT.
 \end{aligned}$$
- Similar for the other quadratic term.

Energy equipartition theorem

- $$\begin{aligned} \overline{\frac{1}{2}a_1 p_1^2} &= \frac{1}{Z_1} \int \frac{1}{2}a_1 p_1^2 \cdot e^{-\beta \varepsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r} \\ &= \int_{-\infty}^{\infty} \frac{1}{2}a_1 p_1^2 e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\ &= \int_{-\infty}^{\infty} \frac{1}{4}a_1 p_1 e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1^2 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\ &= \int_{-\infty}^{\infty} \frac{1}{4}a_1 p_1 \cdot \left(\frac{-2}{\beta a_1}\right) de^{-\beta \frac{1}{2}a_1 p_1^2} \cdot \\ &\quad \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\ &= \frac{1}{2\beta} \int_{-\infty}^{\infty} e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\ &= \frac{1}{2\beta} \frac{1}{Z_1} \int e^{-\beta \varepsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r} \\ &= \frac{1}{2\beta} = \frac{1}{2}kT. \end{aligned}$$
- Similar for the other quadratic term.
- Application: internal energy U , thermal capacity C_V .

Application 1. Monatomic molecules

Application 1. Monatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$

Application 1. Monatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$
 $\Rightarrow \bar{\varepsilon} = \frac{3}{2}kT$ (3 quadratic terms).

Application 1. Monatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$
 $\Rightarrow \bar{\varepsilon} = \frac{3}{2}kT$ (3 quadratic terms).
- $U = N\bar{\varepsilon} = \frac{3}{2}NkT$.

Application 1. Monatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$
 $\Rightarrow \bar{\varepsilon} = \frac{3}{2}kT$ (3 quadratic terms).
- $U = N\bar{\varepsilon} = \frac{3}{2}NkT.$
- $C_V = \frac{dU}{dT} = \frac{3}{2}Nk.$

Application 1. Monatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$
 $\Rightarrow \bar{\varepsilon} = \frac{3}{2}kT$ (3 quadratic terms).
- $U = N\bar{\varepsilon} = \frac{3}{2}NkT$.
- $C_V = \frac{dU}{dT} = \frac{3}{2}Nk$.
- As $pV = nRT = NkT$, $C_p - C_V = nR = Nk$,

Application 1. Monatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$
 $\Rightarrow \bar{\varepsilon} = \frac{3}{2}kT$ (3 quadratic terms).
- $U = N\bar{\varepsilon} = \frac{3}{2}NkT$.
- $C_V = \frac{dU}{dT} = \frac{3}{2}Nk$.
- As $pV = nRT = NkT$, $C_p - C_V = nR = Nk$,
 $\therefore C_p = \frac{5}{2}Nk$, and $\gamma = \frac{C_p}{C_V} = \frac{5}{3}$.

Application 1. Monatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$
 $\Rightarrow \bar{\varepsilon} = \frac{3}{2}kT$ (3 quadratic terms).
- $U = N\bar{\varepsilon} = \frac{3}{2}NkT$.
- $C_V = \frac{dU}{dT} = \frac{3}{2}Nk$.
- As $pV = nRT = NkT$, $C_p - C_V = nR = Nk$,
 $\therefore C_p = \frac{5}{2}Nk$, and $\gamma = \frac{C_p}{C_V} = \frac{5}{3}$.
- Table 7.2.

Application 1. Monatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$
 $\Rightarrow \bar{\varepsilon} = \frac{3}{2}kT$ (3 quadratic terms).
- $U = N\bar{\varepsilon} = \frac{3}{2}NkT$.
- $C_V = \frac{dU}{dT} = \frac{3}{2}Nk$.
- As $pV = nRT = NkT$, $C_p - C_V = nR = Nk$,
 $\therefore C_p = \frac{5}{2}Nk$, and $\gamma = \frac{C_p}{C_V} = \frac{5}{3}$.
- Table 7.2.

Did not consider the contribution of internal structure (electrons and protons), but consistent with the experiments. Why?

Application 1. Monatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$
 $\Rightarrow \bar{\varepsilon} = \frac{3}{2}kT$ (3 quadratic terms).
- $U = N\bar{\varepsilon} = \frac{3}{2}NkT$.
- $C_V = \frac{dU}{dT} = \frac{3}{2}Nk$.
- As $pV = nRT = NkT$, $C_p - C_V = nR = Nk$,
 $\therefore C_p = \frac{5}{2}Nk$, and $\gamma = \frac{C_p}{C_V} = \frac{5}{3}$.
- Table 7.2.

Did not consider the contribution of internal structure (electrons and protons), but consistent with the experiments. Why?

- Quantum effect. Energy level are discrete.

Application 1. Monatomic molecules

- Such as potassium (K), the ionization energy $418.8 \text{ kJ mol}^{-1} = \frac{418.8 \times 10^3 \text{ J}}{6.02 \times 10^{23}} \simeq 7 \times 10^{-19} \text{ J}$.

Application 1. Monatomic molecules

- Such as potassium (K), the ionization energy $418.8 \text{ kJ mol}^{-1} = \frac{418.8 \times 10^3 \text{ J}}{6.02 \times 10^{23}} \simeq 7 \times 10^{-19} \text{ J}$.

The transition energy (ground state to 1st excited state) is $\frac{3}{4} \times 7 \times 10^{-19} \text{ J} \simeq 5 \times 10^{-19} \text{ J}$.

Application 1. Monatomic molecules

- Such as potassium (K), the ionization energy $418.8 \text{ kJ mol}^{-1} = \frac{418.8 \times 10^3 \text{ J}}{6.02 \times 10^{23}} \simeq 7 \times 10^{-19} \text{ J}$.
The transition energy (ground state to 1st excited state) is $\frac{3}{4} \times 7 \times 10^{-19} \text{ J} \simeq 5 \times 10^{-19} \text{ J}$.
- Typical energy for $T = 1000 \text{ K}$,
 $kT = 1.38 \times 10^{-23} \text{ J K}^{-1} \times 10^3 \text{ K} \simeq 1.38 \times 10^{-20} \text{ J}$.

Application 1. Monatomic molecules

- Such as potassium (K), the ionization energy $418.8 \text{ kJ mol}^{-1} = \frac{418.8 \times 10^3 \text{ J}}{6.02 \times 10^{23}} \simeq 7 \times 10^{-19} \text{ J}$.
The transition energy (ground state to 1st excited state) is $\frac{3}{4} \times 7 \times 10^{-19} \text{ J} \simeq 5 \times 10^{-19} \text{ J}$.
- Typical energy for $T = 1000 \text{ K}$,
 $kT = 1.38 \times 10^{-23} \text{ J K}^{-1} \times 10^3 \text{ K} \simeq 1.38 \times 10^{-20} \text{ J}$.
Smaller than the energy required by the transition energy.

Application 1. Monatomic molecules

- Such as potassium (K), the ionization energy $418.8 \text{ kJ mol}^{-1} = \frac{418.8 \times 10^3 \text{ J}}{6.02 \times 10^{23}} \simeq 7 \times 10^{-19} \text{ J}$.
The transition energy (ground state to 1st excited state) is $\frac{3}{4} \times 7 \times 10^{-19} \text{ J} \simeq 5 \times 10^{-19} \text{ J}$.
- Typical energy for $T = 1000 \text{ K}$,
 $kT = 1.38 \times 10^{-23} \text{ J K}^{-1} \times 10^3 \text{ K} \simeq 1.38 \times 10^{-20} \text{ J}$.
Smaller than the energy required by the transition energy.
Therefore, the contribution of electrons should not be considered.

Application 2. diatomic molecules

Application 2. diatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2\theta}p_\varphi^2) + \frac{1}{2m_\mu}p_r^2 + u(r)$
where $m_\mu = \frac{m_1 m_2}{m_1 + m_2}$ is the reduced mass, I is the moment of inertia. $\frac{1}{2m_\mu}p_r^2$ is the energy because of relative motion.

Application 2. diatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2\theta}p_\varphi^2) + \frac{1}{2m_\mu}p_r^2 + u(r)$
where $m_\mu = \frac{m_1 m_2}{m_1 + m_2}$ is the reduced mass, I is the moment of inertia. $\frac{1}{2m_\mu}p_r^2$ is the energy because of relative motion.
- Not considering p_r^2 , there are 5 quadratic terms.

Application 2. diatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2\theta}p_\varphi^2) + \frac{1}{2m_\mu}p_r^2 + u(r)$
where $m_\mu = \frac{m_1 m_2}{m_1 + m_2}$ is the reduced mass, I is the moment of inertia. $\frac{1}{2m_\mu}p_r^2$ is the energy because of relative motion.
- Not considering p_r^2 , there are 5 quadratic terms.
 $\bar{\varepsilon} = \frac{5}{2}kT$.

Application 2. diatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2\theta}p_\varphi^2) + \frac{1}{2m_\mu}p_r^2 + u(r)$
where $m_\mu = \frac{m_1 m_2}{m_1 + m_2}$ is the reduced mass, I is the moment of inertia. $\frac{1}{2m_\mu}p_r^2$ is the energy because of relative motion.
- Not considering p_r^2 , there are 5 quadratic terms.
 $\bar{\varepsilon} = \frac{5}{2}kT.$
- $U = \frac{5}{2}NkT,$

Application 2. diatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2\theta}p_\varphi^2) + \frac{1}{2m_\mu}p_r^2 + u(r)$
where $m_\mu = \frac{m_1 m_2}{m_1 + m_2}$ is the reduced mass, I is the moment of inertia. $\frac{1}{2m_\mu}p_r^2$ is the energy because of relative motion.
- Not considering p_r^2 , there are 5 quadratic terms.
 $\bar{\varepsilon} = \frac{5}{2}kT.$
- $U = \frac{5}{2}NkT, C_V = \frac{5}{2}Nk,$

Application 2. diatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2\theta}p_\varphi^2) + \frac{1}{2m_\mu}p_r^2 + u(r)$
where $m_\mu = \frac{m_1 m_2}{m_1 + m_2}$ is the reduced mass, I is the moment of inertia. $\frac{1}{2m_\mu}p_r^2$ is the energy because of relative motion.
- Not considering p_r^2 , there are 5 quadratic terms.
 $\bar{\varepsilon} = \frac{5}{2}kT$.
- $U = \frac{5}{2}NkT$, $C_V = \frac{5}{2}Nk$, $C_p = \frac{7}{2}Nk$,

Application 2. diatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2\theta}p_\varphi^2) + \frac{1}{2m_\mu}p_r^2 + u(r)$
 where $m_\mu = \frac{m_1 m_2}{m_1 + m_2}$ is the reduced mass, I is the moment of inertia. $\frac{1}{2m_\mu}p_r^2$ is the energy because of relative motion.
- Not considering p_r^2 , there are 5 quadratic terms.
 $\bar{\varepsilon} = \frac{5}{2}kT$.
- $U = \frac{5}{2}NkT$, $C_V = \frac{5}{2}Nk$, $C_p = \frac{7}{2}Nk$,
 $\gamma = 1.4$, see table 7.3.

Application 2. diatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2\theta}p_\varphi^2) + \frac{1}{2m_\mu}p_r^2 + u(r)$
 where $m_\mu = \frac{m_1 m_2}{m_1 + m_2}$ is the reduced mass, I is the moment of inertia. $\frac{1}{2m_\mu}p_r^2$ is the energy because of relative motion.
- Not considering p_r^2 , there are 5 quadratic terms.
 $\bar{\varepsilon} = \frac{5}{2}kT$.
- $U = \frac{5}{2}NkT$, $C_V = \frac{5}{2}Nk$, $C_p = \frac{7}{2}Nk$,
 $\gamma = 1.4$, see table 7.3.
- i. Why no p_r^2 ?

Application 2. diatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2\theta}p_\varphi^2) + \frac{1}{2m_\mu}p_r^2 + u(r)$
 where $m_\mu = \frac{m_1 m_2}{m_1 + m_2}$ is the reduced mass, I is the moment of inertia. $\frac{1}{2m_\mu}p_r^2$ is the energy because of relative motion.
- Not considering p_r^2 , there are 5 quadratic terms.
 $\bar{\varepsilon} = \frac{5}{2}kT$.
- $U = \frac{5}{2}NkT$, $C_V = \frac{5}{2}Nk$, $C_p = \frac{7}{2}Nk$,
 $\gamma = 1.4$, see table 7.3.
- i. Why no p_r^2 ? Discrete energy level.

Application 2. diatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2\theta}p_\varphi^2) + \frac{1}{2m_\mu}p_r^2 + u(r)$
 where $m_\mu = \frac{m_1 m_2}{m_1 + m_2}$ is the reduced mass, I is the moment of inertia. $\frac{1}{2m_\mu}p_r^2$ is the energy because of relative motion.
- Not considering p_r^2 , there are 5 quadratic terms.
 $\bar{\varepsilon} = \frac{5}{2}kT$.
- $U = \frac{5}{2}NkT$, $C_V = \frac{5}{2}Nk$, $C_p = \frac{7}{2}Nk$,
 $\gamma = 1.4$, see table 7.3.
- i. Why no p_r^2 ? Discrete energy level.
 ii. Why H_2 not good in low temperature?

Application 2. diatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2\theta}p_\varphi^2) + \frac{1}{2m_\mu}p_r^2 + u(r)$
 where $m_\mu = \frac{m_1 m_2}{m_1 + m_2}$ is the reduced mass, I is the moment of inertia. $\frac{1}{2m_\mu}p_r^2$ is the energy because of relative motion.
- Not considering p_r^2 , there are 5 quadratic terms.
 $\bar{\varepsilon} = \frac{5}{2}kT$.
- $U = \frac{5}{2}NkT$, $C_V = \frac{5}{2}Nk$, $C_p = \frac{7}{2}Nk$,
 $\gamma = 1.4$, see table 7.3.
- i. Why no p_r^2 ? Discrete energy level.
 ii. Why H_2 not good in low temperature?
 Eq. (7.2.6), $\frac{V}{N}(\frac{2\pi mkT}{h^2})^{3/2}$, classical limit.

Application 3. Atoms in solid

Application 3. Atoms in solid

- 1. Atoms are trapped.

Application 3. Atoms in solid

- 1. Atoms are trapped.
- 2. Atom can vibrate near its equilibrium position.

Application 3. Atoms in solid

- 1. Atoms are trapped.
- 2. Atom can vibrate near its equilibrium position.
- 3. Vibrations of each atom are independent.

Application 3. Atoms in solid

- 1. Atoms are trapped.
- 2. Atom can vibrate near its equilibrium position.
- 3. Vibrations of each atom are independent.
- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2}m\omega^2(x^2 + y^2 + z^2).$

Application 3. Atoms in solid

- 1. Atoms are trapped.
- 2. Atom can vibrate near its equilibrium position.
- 3. Vibrations of each atom are independent.
- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2}m\omega^2(x^2 + y^2 + z^2)$.
- 6 quadratic terms,

Application 3. Atoms in solid

- 1. Atoms are trapped.
- 2. Atom can vibrate near its equilibrium position.
- 3. Vibrations of each atom are independent.
- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2}m\omega^2(x^2 + y^2 + z^2)$.
- 6 quadratic terms,
 $\bar{\varepsilon} = 3kT, U = 3NkT, C_V = 3Nk$.
Consistent with the experiment at high temperature.

Application 4. Equilibrium radiation

Application 4. Equilibrium radiation

- What is the microscopic unit of thermal radiation?

Application 4. Equilibrium radiation

- What is the microscopic unit of thermal radiation?
- View 1: consist of particles – bosons – photons (§8.4).

Application 4. Equilibrium radiation

- What is the microscopic unit of thermal radiation?
- View 1: consist of particles – bosons – photons (§8.4).
- View 2: consist of electron-magnetic waves with different frequencies (discrete).

Application 4. Equilibrium radiation

- What is the microscopic unit of thermal radiation?
- View 1: consist of particles – bosons – photons (§8.4).
- View 2: consist of electron-magnetic waves with different frequencies (discrete).

Each monochromatic wave with **a certain frequency** acts like a classical particle.

Application 4. Equilibrium radiation

- What is the microscopic unit of thermal radiation?
- View 1: consist of particles – bosons – photons (§8.4).
- View 2: consist of electron-magnetic waves with different frequencies (discrete).

Each monochromatic wave with **a certain frequency** acts like a classical particle.

Each wave is equivalent to a harmonic oscillation.

Application 4. Equilibrium radiation

- What is the microscopic unit of thermal radiation?
- View 1: consist of particles – bosons – photons (§8.4).
- View 2: consist of electron-magnetic waves with different frequencies (discrete).

Each monochromatic wave with **a certain frequency** acts like a classical particle.

Each wave is equivalent to a harmonic oscillation.

$$\varepsilon = \frac{p^2}{2m} + \frac{1}{2}m\omega^2 q^2.$$

Application 4. Equilibrium radiation

- What is the microscopic unit of thermal radiation?
- View 1: consist of particles – bosons – photons (§8.4).
- View 2: consist of electron-magnetic waves with different frequencies (discrete).

Each monochromatic wave with **a certain frequency** acts like a classical particle.

Each wave is equivalent to a harmonic oscillation.

$$\varepsilon = \frac{p^2}{2m} + \frac{1}{2}m\omega^2 q^2.$$

- Two quadratic terms, $\bar{\varepsilon} = kT$.

Application 4. Equilibrium radiation

- What is the microscopic unit of thermal radiation?
- View 1: consist of particles – bosons – photons (§8.4).
- View 2: consist of electron-magnetic waves with different frequencies (discrete).

Each monochromatic wave with **a certain frequency** acts like a classical particle.

Each wave is equivalent to a harmonic oscillation.

$$\varepsilon = \frac{p^2}{2m} + \frac{1}{2}m\omega^2 q^2.$$

- Two quadratic terms, $\bar{\varepsilon} = kT$.
- Using the periodic boundary condition:

Application 4. Equilibrium radiation

- $k_x = \frac{2\pi}{L}n_x, n_x = 0, \pm 1, \pm 2, \dots$
 $k_y = \frac{2\pi}{L}n_y, n_y = 0, \pm 1, \pm 2, \dots$
 $k_z = \frac{2\pi}{L}n_z, n_z = 0, \pm 1, \pm 2, \dots$

Application 4. Equilibrium radiation

- $k_x = \frac{2\pi}{L}n_x, n_x = 0, \pm 1, \pm 2, \dots$
 $k_y = \frac{2\pi}{L}n_y, n_y = 0, \pm 1, \pm 2, \dots$
 $k_z = \frac{2\pi}{L}n_z, n_z = 0, \pm 1, \pm 2, \dots$
- $dk_x dk_y dk_z = \left(\frac{2\pi}{L}\right)^3 \Delta n_x \Delta n_y \Delta n_z,$

Application 4. Equilibrium radiation

- $k_x = \frac{2\pi}{L}n_x, n_x = 0, \pm 1, \pm 2, \dots$
 $k_y = \frac{2\pi}{L}n_y, n_y = 0, \pm 1, \pm 2, \dots$
 $k_z = \frac{2\pi}{L}n_z, n_z = 0, \pm 1, \pm 2, \dots$
- $dk_x dk_y dk_z = \left(\frac{2\pi}{L}\right)^3 \Delta n_x \Delta n_y \Delta n_z,$
i.e., number of oscillations:
$$\Delta n_x \Delta n_y \Delta n_z = \frac{V}{(2\pi)^3} dk_x dk_y dk_z.$$

Application 4. Equilibrium radiation

- $k_x = \frac{2\pi}{L}n_x$, $n_x = 0, \pm 1, \pm 2, \dots$
 $k_y = \frac{2\pi}{L}n_y$, $n_y = 0, \pm 1, \pm 2, \dots$
 $k_z = \frac{2\pi}{L}n_z$, $n_z = 0, \pm 1, \pm 2, \dots$
- $dk_x dk_y dk_z = \left(\frac{2\pi}{L}\right)^3 \Delta n_x \Delta n_y \Delta n_z$,
i.e., number of oscillations:
 $\Delta n_x \Delta n_y \Delta n_z = \frac{V}{(2\pi)^3} dk_x dk_y dk_z$.
- In spheric polar coordinate, isotropic radiation,
together with the polarization:

Application 4. Equilibrium radiation

- $k_x = \frac{2\pi}{L}n_x$, $n_x = 0, \pm 1, \pm 2, \dots$
 $k_y = \frac{2\pi}{L}n_y$, $n_y = 0, \pm 1, \pm 2, \dots$
 $k_z = \frac{2\pi}{L}n_z$, $n_z = 0, \pm 1, \pm 2, \dots$
- $dk_x dk_y dk_z = \left(\frac{2\pi}{L}\right)^3 \Delta n_x \Delta n_y \Delta n_z$,
i.e., number of oscillations:
$$\Delta n_x \Delta n_y \Delta n_z = \frac{V}{(2\pi)^3} dk_x dk_y dk_z.$$
- In spheric polar coordinate, isotropic radiation,
together with the polarization:
$$\Delta n_x \Delta n_y \Delta n_z = 2 \frac{V}{(2\pi)^3} k^2 dk \cdot 4\pi$$

Application 4. Equilibrium radiation

- $k_x = \frac{2\pi}{L}n_x$, $n_x = 0, \pm 1, \pm 2, \dots$
 $k_y = \frac{2\pi}{L}n_y$, $n_y = 0, \pm 1, \pm 2, \dots$
 $k_z = \frac{2\pi}{L}n_z$, $n_z = 0, \pm 1, \pm 2, \dots$
- $dk_x dk_y dk_z = \left(\frac{2\pi}{L}\right)^3 \Delta n_x \Delta n_y \Delta n_z$,
 i.e., number of oscillations:
 $\Delta n_x \Delta n_y \Delta n_z = \frac{V}{(2\pi)^3} dk_x dk_y dk_z$.
- In spheric polar coordinate, isotropic radiation,
 together with the polarization:
 $\Delta n_x \Delta n_y \Delta n_z = 2 \frac{V}{(2\pi)^3} k^2 dk \cdot 4\pi = 2 \frac{V}{(2\pi)^3} \left(\frac{\omega}{c}\right)^2 d\frac{\omega}{c} \cdot 4\pi$

Application 4. Equilibrium radiation

- $k_x = \frac{2\pi}{L}n_x$, $n_x = 0, \pm 1, \pm 2, \dots$
 $k_y = \frac{2\pi}{L}n_y$, $n_y = 0, \pm 1, \pm 2, \dots$
 $k_z = \frac{2\pi}{L}n_z$, $n_z = 0, \pm 1, \pm 2, \dots$
- $dk_x dk_y dk_z = \left(\frac{2\pi}{L}\right)^3 \Delta n_x \Delta n_y \Delta n_z$,
 i.e., number of oscillations:
 $\Delta n_x \Delta n_y \Delta n_z = \frac{V}{(2\pi)^3} dk_x dk_y dk_z$.
- In spheric polar coordinate, isotropic radiation,
 together with the polarization:
 $\Delta n_x \Delta n_y \Delta n_z = 2 \frac{V}{(2\pi)^3} k^2 dk \cdot 4\pi = 2 \frac{V}{(2\pi)^3} \left(\frac{\omega}{c}\right)^2 d\frac{\omega}{c} \cdot 4\pi$
 $= \frac{V}{\pi^2 c^3} \omega^2 d\omega \equiv D(\omega) d\omega$.

Application 4. Equilibrium radiation

- $k_x = \frac{2\pi}{L}n_x$, $n_x = 0, \pm 1, \pm 2, \dots$
 $k_y = \frac{2\pi}{L}n_y$, $n_y = 0, \pm 1, \pm 2, \dots$
 $k_z = \frac{2\pi}{L}n_z$, $n_z = 0, \pm 1, \pm 2, \dots$
- $dk_x dk_y dk_z = \left(\frac{2\pi}{L}\right)^3 \Delta n_x \Delta n_y \Delta n_z$,
 i.e., number of oscillations:

$$\Delta n_x \Delta n_y \Delta n_z = \frac{V}{(2\pi)^3} dk_x dk_y dk_z.$$
- In spheric polar coordinate, isotropic radiation,
 together with the polarization:

$$\Delta n_x \Delta n_y \Delta n_z = 2 \frac{V}{(2\pi)^3} k^2 dk \cdot 4\pi = 2 \frac{V}{(2\pi)^3} \left(\frac{\omega}{c}\right)^2 d\frac{\omega}{c} \cdot 4\pi$$

$$= \frac{V}{\pi^2 c^3} \omega^2 d\omega \equiv D(\omega) d\omega.$$
- As for each vibration $\bar{\epsilon} = kT$,

Application 4. Equilibrium radiation

- $k_x = \frac{2\pi}{L}n_x$, $n_x = 0, \pm 1, \pm 2, \dots$
 $k_y = \frac{2\pi}{L}n_y$, $n_y = 0, \pm 1, \pm 2, \dots$
 $k_z = \frac{2\pi}{L}n_z$, $n_z = 0, \pm 1, \pm 2, \dots$
- $dk_x dk_y dk_z = \left(\frac{2\pi}{L}\right)^3 \Delta n_x \Delta n_y \Delta n_z$,
 i.e., number of oscillations:

$$\Delta n_x \Delta n_y \Delta n_z = \frac{V}{(2\pi)^3} dk_x dk_y dk_z.$$
- In spheric polar coordinate, isotropic radiation,
 together with the polarization:

$$\Delta n_x \Delta n_y \Delta n_z = 2 \frac{V}{(2\pi)^3} k^2 dk \cdot 4\pi = 2 \frac{V}{(2\pi)^3} \left(\frac{\omega}{c}\right)^2 d\frac{\omega}{c} \cdot 4\pi$$

$$= \frac{V}{\pi^2 c^3} \omega^2 d\omega \equiv D(\omega) d\omega.$$
- As for each vibration $\bar{\varepsilon} = kT$, the internal energy

$$U_\omega d\omega = \frac{V}{\pi^2 c^3} kT \omega^2 d\omega.$$

Application 4. Equilibrium radiation

- $k_x = \frac{2\pi}{L}n_x$, $n_x = 0, \pm 1, \pm 2, \dots$
 $k_y = \frac{2\pi}{L}n_y$, $n_y = 0, \pm 1, \pm 2, \dots$
 $k_z = \frac{2\pi}{L}n_z$, $n_z = 0, \pm 1, \pm 2, \dots$
- $dk_x dk_y dk_z = \left(\frac{2\pi}{L}\right)^3 \Delta n_x \Delta n_y \Delta n_z$,
 i.e., number of oscillations:

$$\Delta n_x \Delta n_y \Delta n_z = \frac{V}{(2\pi)^3} dk_x dk_y dk_z.$$
- In spheric polar coordinate, isotropic radiation,
 together with the polarization:

$$\Delta n_x \Delta n_y \Delta n_z = 2 \frac{V}{(2\pi)^3} k^2 dk \cdot 4\pi = 2 \frac{V}{(2\pi)^3} \left(\frac{\omega}{c}\right)^2 d\frac{\omega}{c} \cdot 4\pi$$

$$= \frac{V}{\pi^2 c^3} \omega^2 d\omega \equiv D(\omega) d\omega.$$
- As for each vibration $\bar{\varepsilon} = kT$, the internal energy

$$U_\omega d\omega = \frac{V}{\pi^2 c^3} kT \omega^2 d\omega.$$
 Only suitable for low frequency.

Application 4. Equilibrium radiation

- Theory: $U_\omega d\omega = \frac{V}{\pi^2 c^3} kT \omega^2 d\omega.$

Application 4. Equilibrium radiation

- Theory: $U_{\omega}d\omega = \frac{V}{\pi^2 c^3} kT \omega^2 d\omega.$
Experiment: $U_{\omega}d\omega = \frac{V}{2\pi^3 c^3} \frac{h\omega^3 d\omega}{e^{\frac{h\omega}{2\pi kT}} - 1}.$

Application 4. Equilibrium radiation

- Theory: $U_\omega d\omega = \frac{V}{\pi^2 c^3} kT \omega^2 d\omega$.
Experiment: $U_\omega d\omega = \frac{V}{2\pi^3 c^3} \frac{h\omega^3 d\omega}{e^{\frac{h\omega}{2\pi kT}} - 1}$.
- Planck's idea: $D(\omega) d\omega = \frac{V}{\pi^2 c^3} \omega^2 d\omega$ still obeys.

Application 4. Equilibrium radiation

- Theory: $U_\omega d\omega = \frac{V}{\pi^2 c^3} kT \omega^2 d\omega$.
Experiment: $U_\omega d\omega = \frac{V}{2\pi^3 c^3} \frac{h\omega^3 d\omega}{e^{\frac{h\omega}{2\pi kT}} - 1}$.
- Planck's idea: $D(\omega) d\omega = \frac{V}{\pi^2 c^3} \omega^2 d\omega$ still obeys. But the energy of the oscillator is NOT continuous, supposed to be multiples of $\hbar\omega$, i.e., $\varepsilon_n(\omega) = n\hbar\omega$.

Application 4. Equilibrium radiation

- Theory: $U_\omega d\omega = \frac{V}{\pi^2 c^3} kT \omega^2 d\omega$.
Experiment: $U_\omega d\omega = \frac{V}{2\pi^3 c^3} \frac{h\omega^3 d\omega}{e^{\frac{h\omega}{2\pi kT}} - 1}$.
- Planck's idea: $D(\omega)d\omega = \frac{V}{\pi^2 c^3} \omega^2 d\omega$ still obeys. But the energy of the oscillator is NOT continuous, supposed to be multiples of $\hbar\omega$, i.e., $\varepsilon_n(\omega) = n\hbar\omega$.
- The most probable Boltzmann distribution:
 $a_n(\omega) = e^{-\alpha - \beta\varepsilon_n},$

Application 4. Equilibrium radiation

- Theory: $U_\omega d\omega = \frac{V}{\pi^2 c^3} kT \omega^2 d\omega$.
Experiment: $U_\omega d\omega = \frac{V}{2\pi^3 c^3} \frac{h\omega^3 d\omega}{e^{\frac{h\omega}{2\pi kT}} - 1}$.
- Planck's idea: $D(\omega)d\omega = \frac{V}{\pi^2 c^3} \omega^2 d\omega$ still obeys. But the energy of the oscillator is NOT continuous, supposed to be multiples of $\hbar\omega$, i.e., $\varepsilon_n(\omega) = n\hbar\omega$.
- The most probable Boltzmann distribution:
 $a_n(\omega) = e^{-\alpha - \beta\varepsilon_n},$
degeneracy $\omega_n(\omega) = 2$ (polarization),

Application 4. Equilibrium radiation

- Theory: $U_\omega d\omega = \frac{V}{\pi^2 c^3} kT \omega^2 d\omega$.
Experiment: $U_\omega d\omega = \frac{V}{2\pi^3 c^3} \frac{h\omega^3 d\omega}{e^{\frac{h\omega}{2\pi kT}} - 1}$.
- Planck's idea: $D(\omega)d\omega = \frac{V}{\pi^2 c^3} \omega^2 d\omega$ still obeys. But the energy of the oscillator is NOT continuous, supposed to be multiples of $\hbar\omega$, i.e., $\varepsilon_n(\omega) = n\hbar\omega$.
- The most probable Boltzmann distribution:
 $a_n(\omega) = e^{-\alpha - \beta\varepsilon_n},$
 degeneracy $\omega_n(\omega) = 2$ (polarization),
 partition function: $Z_1 = \sum_{n=0}^{\infty} \omega_n e^{-\beta\varepsilon_n}$

Application 4. Equilibrium radiation

- Theory: $U_\omega d\omega = \frac{V}{\pi^2 c^3} kT \omega^2 d\omega$.
Experiment: $U_\omega d\omega = \frac{V}{2\pi^3 c^3} \frac{h\omega^3 d\omega}{e^{\frac{h\omega}{2\pi kT}} - 1}$.
- Planck's idea: $D(\omega)d\omega = \frac{V}{\pi^2 c^3} \omega^2 d\omega$ still obeys. But the energy of the oscillator is NOT continuous, supposed to be multiples of $\hbar\omega$, i.e., $\varepsilon_n(\omega) = n\hbar\omega$.
- The most probable Boltzmann distribution:
 $a_n(\omega) = e^{-\alpha - \beta\varepsilon_n},$
 degeneracy $\omega_n(\omega) = 2$ (polarization),
 partition function: $Z_1 = \sum_{n=0}^{\infty} \omega_n e^{-\beta\varepsilon_n} = \sum_{n=0}^{\infty} 2e^{-\beta n\hbar\omega}$

Application 4. Equilibrium radiation

- Theory: $U_\omega d\omega = \frac{V}{\pi^2 c^3} kT \omega^2 d\omega$.
Experiment: $U_\omega d\omega = \frac{V}{2\pi^3 c^3} \frac{h\omega^3 d\omega}{e^{\frac{h\omega}{2\pi kT}} - 1}$.
- Planck's idea: $D(\omega)d\omega = \frac{V}{\pi^2 c^3} \omega^2 d\omega$ still obeys. But the energy of the oscillator is NOT continuous, supposed to be multiples of $\hbar\omega$, i.e., $\varepsilon_n(\omega) = n\hbar\omega$.
- The most probable Boltzmann distribution:
 $a_n(\omega) = e^{-\alpha - \beta\varepsilon_n},$
 degeneracy $\omega_n(\omega) = 2$ (polarization),
 partition function: $Z_1 = \sum_{n=0}^{\infty} \omega_n e^{-\beta\varepsilon_n} = \sum_{n=0}^{\infty} 2e^{-\beta n\hbar\omega}$
 $= \frac{2}{1 - e^{-\beta\hbar\omega}}.$

Application 4. Equilibrium radiation

$$Z_1 = \sum_{n=0}^{\infty} 2e^{-\beta\varepsilon_n}, Z_1 = \frac{2}{1-e^{-\beta\hbar\omega}}$$

- The average energy of an oscillator with frequency ω :

Application 4. Equilibrium radiation

$$Z_1 = \sum_{n=0}^{\infty} 2e^{-\beta\varepsilon_n}, Z_1 = \frac{2}{1-e^{-\beta\hbar\omega}}$$

- The average energy of an oscillator with frequency ω :

$$\bar{\varepsilon} = \frac{\sum a_n \varepsilon_n}{\sum a_n}$$

Application 4. Equilibrium radiation

$$Z_1 = \sum_{n=0}^{\infty} 2e^{-\beta\varepsilon_n}, Z_1 = \frac{2}{1-e^{-\beta\hbar\omega}}$$

- The average energy of an oscillator with frequency ω :

$$\bar{\varepsilon} = \frac{\sum a_n \varepsilon_n}{\sum a_n} = \frac{\sum e^{-\alpha-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\alpha-\beta\varepsilon_n}}$$

Application 4. Equilibrium radiation

$$Z_1 = \sum_{n=0}^{\infty} 2e^{-\beta\varepsilon_n}, Z_1 = \frac{2}{1-e^{-\beta\hbar\omega}}$$

- The average energy of an oscillator with frequency ω :

$$\bar{\varepsilon} = \frac{\sum a_n \varepsilon_n}{\sum a_n} = \frac{\sum e^{-\alpha-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\alpha-\beta\varepsilon_n}} = \frac{\sum e^{-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\beta\varepsilon_n}}$$

Application 4. Equilibrium radiation

$$Z_1 = \sum_{n=0}^{\infty} 2e^{-\beta\varepsilon_n}, Z_1 = \frac{2}{1-e^{-\beta\hbar\omega}}$$

- The average energy of an oscillator with frequency ω :

$$\bar{\varepsilon} = \frac{\sum a_n \varepsilon_n}{\sum a_n} = \frac{\sum e^{-\alpha-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\alpha-\beta\varepsilon_n}} = \frac{\sum e^{-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\beta\varepsilon_n}} = -\frac{\partial}{\partial\beta} \ln Z_1$$

Application 4. Equilibrium radiation

$$Z_1 = \sum_{n=0}^{\infty} 2e^{-\beta\varepsilon_n}, Z_1 = \frac{2}{1-e^{-\beta\hbar\omega}}$$

- The average energy of an oscillator with frequency ω :

$$\begin{aligned}\bar{\varepsilon} &= \frac{\sum a_n \varepsilon_n}{\sum a_n} = \frac{\sum e^{-\alpha-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\alpha-\beta\varepsilon_n}} = \frac{\sum e^{-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\beta\varepsilon_n}} = -\frac{\partial}{\partial\beta} \ln Z_1 \\ &= -\frac{\partial}{\partial\beta} \ln \frac{2}{1-e^{-\beta\hbar\omega}}\end{aligned}$$

Application 4. Equilibrium radiation

$$Z_1 = \sum_{n=0}^{\infty} 2e^{-\beta\varepsilon_n}, Z_1 = \frac{2}{1-e^{-\beta\hbar\omega}}$$

- The average energy of an oscillator with frequency ω :

$$\begin{aligned}\bar{\varepsilon} &= \frac{\sum a_n \varepsilon_n}{\sum a_n} = \frac{\sum e^{-\alpha-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\alpha-\beta\varepsilon_n}} = \frac{\sum e^{-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\beta\varepsilon_n}} = -\frac{\partial}{\partial\beta} \ln Z_1 \\ &= -\frac{\partial}{\partial\beta} \ln \frac{2}{1-e^{-\beta\hbar\omega}} = \frac{\partial}{\partial\beta} \ln(1 - e^{-\beta\hbar\omega})\end{aligned}$$

Application 4. Equilibrium radiation

$$Z_1 = \sum_{n=0}^{\infty} 2e^{-\beta\varepsilon_n}, Z_1 = \frac{2}{1-e^{-\beta\hbar\omega}}$$

- The average energy of an oscillator with frequency ω :

$$\begin{aligned}\bar{\varepsilon} &= \frac{\sum a_n \varepsilon_n}{\sum a_n} = \frac{\sum e^{-\alpha-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\alpha-\beta\varepsilon_n}} = \frac{\sum e^{-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\beta\varepsilon_n}} = -\frac{\partial}{\partial\beta} \ln Z_1 \\ &= -\frac{\partial}{\partial\beta} \ln \frac{2}{1-e^{-\beta\hbar\omega}} = \frac{\partial}{\partial\beta} \ln(1 - e^{-\beta\hbar\omega}) = \frac{\hbar\omega e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}\end{aligned}$$

Application 4. Equilibrium radiation

$$Z_1 = \sum_{n=0}^{\infty} 2e^{-\beta\varepsilon_n}, Z_1 = \frac{2}{1-e^{-\beta\hbar\omega}}$$

- The average energy of an oscillator with frequency ω :

$$\begin{aligned}\bar{\varepsilon} &= \frac{\sum a_n \varepsilon_n}{\sum a_n} = \frac{\sum e^{-\alpha-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\alpha-\beta\varepsilon_n}} = \frac{\sum e^{-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\beta\varepsilon_n}} = -\frac{\partial}{\partial\beta} \ln Z_1 \\ &= -\frac{\partial}{\partial\beta} \ln \frac{2}{1-e^{-\beta\hbar\omega}} = \frac{\partial}{\partial\beta} \ln(1 - e^{-\beta\hbar\omega}) = \frac{\hbar\omega e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \\ &= \frac{\hbar\omega}{e^{\beta\hbar\omega}-1}.\end{aligned}$$

Application 4. Equilibrium radiation

$$Z_1 = \sum_{n=0}^{\infty} 2e^{-\beta\varepsilon_n}, Z_1 = \frac{2}{1-e^{-\beta\hbar\omega}}$$

- The average energy of an oscillator with frequency ω :

$$\begin{aligned}\bar{\varepsilon} &= \frac{\sum a_n \varepsilon_n}{\sum a_n} = \frac{\sum e^{-\alpha-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\alpha-\beta\varepsilon_n}} = \frac{\sum e^{-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\beta\varepsilon_n}} = -\frac{\partial}{\partial\beta} \ln Z_1 \\ &= -\frac{\partial}{\partial\beta} \ln \frac{2}{1-e^{-\beta\hbar\omega}} = \frac{\partial}{\partial\beta} \ln(1 - e^{-\beta\hbar\omega}) = \frac{\hbar\omega e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \\ &= \frac{\hbar\omega}{e^{\beta\hbar\omega}-1}.\end{aligned}$$

- The internal energy: $U_\omega d\omega = \bar{\varepsilon} D(\omega) d\omega$

Application 4. Equilibrium radiation

$$Z_1 = \sum_{n=0}^{\infty} 2e^{-\beta\varepsilon_n}, Z_1 = \frac{2}{1-e^{-\beta\hbar\omega}}$$

- The average energy of an oscillator with frequency ω :

$$\begin{aligned}\bar{\varepsilon} &= \frac{\sum a_n \varepsilon_n}{\sum a_n} = \frac{\sum e^{-\alpha-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\alpha-\beta\varepsilon_n}} = \frac{\sum e^{-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\beta\varepsilon_n}} = -\frac{\partial}{\partial\beta} \ln Z_1 \\ &= -\frac{\partial}{\partial\beta} \ln \frac{2}{1-e^{-\beta\hbar\omega}} = \frac{\partial}{\partial\beta} \ln(1 - e^{-\beta\hbar\omega}) = \frac{\hbar\omega e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \\ &= \frac{\hbar\omega}{e^{\beta\hbar\omega}-1}.\end{aligned}$$

- The internal energy: $U_\omega d\omega = \bar{\varepsilon} D(\omega) d\omega$

$$= \frac{\hbar\omega}{e^{\beta\hbar\omega}-1} \frac{V}{\pi^2 c^3} \omega^2 d\omega$$

Application 4. Equilibrium radiation

$$Z_1 = \sum_{n=0}^{\infty} 2e^{-\beta\varepsilon_n}, Z_1 = \frac{2}{1-e^{-\beta\hbar\omega}}$$

- The average energy of an oscillator with frequency ω :

$$\begin{aligned}\bar{\varepsilon} &= \frac{\sum a_n \varepsilon_n}{\sum a_n} = \frac{\sum e^{-\alpha-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\alpha-\beta\varepsilon_n}} = \frac{\sum e^{-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\beta\varepsilon_n}} = -\frac{\partial}{\partial\beta} \ln Z_1 \\ &= -\frac{\partial}{\partial\beta} \ln \frac{2}{1-e^{-\beta\hbar\omega}} = \frac{\partial}{\partial\beta} \ln(1 - e^{-\beta\hbar\omega}) = \frac{\hbar\omega e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \\ &= \frac{\hbar\omega}{e^{\beta\hbar\omega}-1}.\end{aligned}$$

- The internal energy: $U_\omega d\omega = \bar{\varepsilon} D(\omega) d\omega$

$$= \frac{\hbar\omega}{e^{\beta\hbar\omega}-1} \frac{V}{\pi^2 c^3} \omega^2 d\omega = \frac{V}{2\pi^3 c^3} \frac{\hbar\omega^3}{e^{\frac{\hbar\omega}{2\pi kT}}-1} d\omega.$$

Application 4. Equilibrium radiation

$$Z_1 = \sum_{n=0}^{\infty} 2e^{-\beta\epsilon_n}, Z_1 = \frac{2}{1-e^{-\beta\hbar\omega}}$$

- The average energy of an oscillator with frequency ω :

$$\begin{aligned}\bar{\epsilon} &= \frac{\sum a_n \epsilon_n}{\sum a_n} = \frac{\sum e^{-\alpha-\beta\epsilon_n} \epsilon_n}{\sum e^{-\alpha-\beta\epsilon_n}} = \frac{\sum e^{-\beta\epsilon_n} \epsilon_n}{\sum e^{-\beta\epsilon_n}} = -\frac{\partial}{\partial\beta} \ln Z_1 \\ &= -\frac{\partial}{\partial\beta} \ln \frac{2}{1-e^{-\beta\hbar\omega}} = \frac{\partial}{\partial\beta} \ln(1 - e^{-\beta\hbar\omega}) = \frac{\hbar\omega e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \\ &= \frac{\hbar\omega}{e^{\beta\hbar\omega}-1}.\end{aligned}$$

- The internal energy: $U_\omega d\omega = \bar{\epsilon} D(\omega) d\omega$

$$= \frac{\hbar\omega}{e^{\beta\hbar\omega}-1} \frac{V}{\pi^2 c^3} \omega^2 d\omega = \frac{V}{2\pi^3 c^3} \frac{\hbar\omega^3}{e^{\frac{\hbar\omega}{2\pi kT}}-1} d\omega.$$

Exactly the black body radiation formula!

Table of contents

1 §7 Boltzmann Statistics

- 7.1 Thermal quantities in statistics
- 7.2 Equation of state of ideal gas
- 7.3 Maxwell speed distribution
- 7.4 Energy equipartition theorem
- 7.5 Internal energy and heat capacity of the ideal gas
- 7.6 Entropy of the ideal gas
- 7.7 Einstein's theory on heat capacity of solid
- 7.8 Paramagnetic solid
- 7.9 Negative temperature

7.5 Internal energy and heat capacity of the ideal gas

7.5 Internal energy and heat capacity of the ideal gas

- Object: diatomic molecules.

7.5 Internal energy and heat capacity of the ideal gas

- Object: diatomic molecules.
- Method: directly use the partition function.

7.5 Internal energy and heat capacity of the ideal gas

- Object: diatomic molecules.
- Method: directly use the partition function.
- Reason: equipartition theorem only suitable for classical limits. However, in normal pressure and temperature, relative motion is not in classical limits.

7.5 Internal energy and heat capacity of the ideal gas

- Object: diatomic molecules.
- Method: directly use the partition function.
- Reason: equipartition theorem only suitable for classical limits. However, in normal pressure and temperature, relative motion is not in classical limits.
- Energy: $\varepsilon = \varepsilon^t + \varepsilon^v + \varepsilon^r$, where t for translational motion, v for vibration, and r for rotation.

7.5 Internal energy and heat capacity of the ideal gas

- Object: diatomic molecules.
- Method: directly use the partition function.
- Reason: equipartition theorem only suitable for classical limits. However, in normal pressure and temperature, relative motion is not in classical limits.
- Energy: $\varepsilon = \varepsilon^t + \varepsilon^v + \varepsilon^r$, where t for translational motion, v for vibration, and r for rotation.
- Partition function: $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$

7.5 Internal energy and heat capacity of the ideal gas

- Object: diatomic molecules.
- Method: directly use the partition function.
- Reason: equipartition theorem only suitable for classical limits. However, in normal pressure and temperature, relative motion is not in classical limits.
- Energy: $\varepsilon = \varepsilon^t + \varepsilon^v + \varepsilon^r$, where t for translational motion, v for vibration, and r for rotation.
- Partition function:
$$Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$$
$$= \sum \omega^t \omega^v \omega^r e^{-\beta(\varepsilon^t + \varepsilon^v + \varepsilon^r)}$$

7.5 Internal energy and heat capacity of the ideal gas

- Object: diatomic molecules.
- Method: directly use the partition function.
- Reason: equipartition theorem only suitable for classical limits. However, in normal pressure and temperature, relative motion is not in classical limits.
- Energy: $\varepsilon = \varepsilon^t + \varepsilon^v + \varepsilon^r$, where t for translational motion, v for vibration, and r for rotation.
- Partition function:
$$\begin{aligned} Z_1 &= \sum \omega_l e^{-\beta \varepsilon_l} \\ &= \sum \omega^t \omega^v \omega^r e^{-\beta(\varepsilon^t + \varepsilon^v + \varepsilon^r)} \\ &= \sum \omega^t e^{-\beta \varepsilon^t} \sum \omega^v e^{-\beta \varepsilon^v} \sum \omega^r e^{-\beta \varepsilon^r} \end{aligned}$$

7.5 Internal energy and heat capacity of the ideal gas

- Object: diatomic molecules.
- Method: directly use the partition function.
- Reason: equipartition theorem only suitable for classical limits. However, in normal pressure and temperature, relative motion is not in classical limits.
- Energy: $\varepsilon = \varepsilon^t + \varepsilon^v + \varepsilon^r$, where t for translational motion, v for vibration, and r for rotation.
- Partition function:
$$\begin{aligned} Z_1 &= \sum \omega_l e^{-\beta \varepsilon_l} \\ &= \sum \omega^t \omega^v \omega^r e^{-\beta(\varepsilon^t + \varepsilon^v + \varepsilon^r)} \\ &= \sum \omega^t e^{-\beta \varepsilon^t} \sum \omega^v e^{-\beta \varepsilon^v} \sum \omega^r e^{-\beta \varepsilon^r} = Z_1^t \cdot Z_1^v \cdot Z_1^r. \end{aligned}$$

Internal energy and heat capacity of the ideal gas

- Internal energy: $U = -N \frac{\partial}{\partial \beta} \ln Z_1$

Internal energy and heat capacity of the ideal gas

- Internal energy:
$$U = -N \frac{\partial}{\partial \beta} \ln Z_1$$
$$= -N \frac{\partial}{\partial \beta} (\ln Z_1^t + \ln Z_1^v + \ln Z_1^r)$$

Internal energy and heat capacity of the ideal gas

- Internal energy:
$$U = -N \frac{\partial}{\partial \beta} \ln Z_1$$
$$= -N \frac{\partial}{\partial \beta} (\ln Z_1^t + \ln Z_1^v + \ln Z_1^r) = U^t + U^v + U^r.$$

Internal energy and heat capacity of the ideal gas

- Internal energy: $U = -N \frac{\partial}{\partial \beta} \ln Z_1$
 $= -N \frac{\partial}{\partial \beta} (\ln Z_1^t + \ln Z_1^v + \ln Z_1^r) = U^t + U^v + U^r.$
- Heat capacity at constant volume:
 $C_V = C_V^t + C_V^v + C_V^r.$

Internal energy and heat capacity of the ideal gas

- Internal energy: $U = -N \frac{\partial}{\partial \beta} \ln Z_1$
 $= -N \frac{\partial}{\partial \beta} (\ln Z_1^t + \ln Z_1^v + \ln Z_1^r) = U^t + U^v + U^r.$
- Heat capacity at constant volume:
 $C_V = C_V^t + C_V^v + C_V^r.$
- i. Translational motion part:

Internal energy and heat capacity of the ideal gas

- Internal energy: $U = -N \frac{\partial}{\partial \beta} \ln Z_1$
 $= -N \frac{\partial}{\partial \beta} (\ln Z_1^t + \ln Z_1^v + \ln Z_1^r) = U^t + U^v + U^r.$
- Heat capacity at constant volume:
 $C_V = C_V^t + C_V^v + C_V^r.$
- i. Translational motion part: (7.2.4) $Z_1^t = V \left(\frac{2\pi m}{h^2 \beta} \right)^{3/2}.$

Internal energy and heat capacity of the ideal gas

- Internal energy: $U = -N \frac{\partial}{\partial \beta} \ln Z_1$
 $= -N \frac{\partial}{\partial \beta} (\ln Z_1^t + \ln Z_1^v + \ln Z_1^r) = U^t + U^v + U^r.$
- Heat capacity at constant volume:
 $C_V = C_V^t + C_V^v + C_V^r.$
- i. Translational motion part: (7.2.4) $Z_1^t = V \left(\frac{2\pi m}{h^2 \beta} \right)^{3/2}.$
 $U^t = \frac{3}{2} N k T,$

Internal energy and heat capacity of the ideal gas

- Internal energy: $U = -N \frac{\partial}{\partial \beta} \ln Z_1$
 $= -N \frac{\partial}{\partial \beta} (\ln Z_1^t + \ln Z_1^v + \ln Z_1^r) = U^t + U^v + U^r.$
- Heat capacity at constant volume:
 $C_V = C_V^t + C_V^v + C_V^r.$
- i. Translational motion part: (7.2.4) $Z_1^t = V \left(\frac{2\pi m}{h^2 \beta} \right)^{3/2}.$
 $U^t = \frac{3}{2} N k T, C_V^t = \frac{3}{2} N k.$

Internal energy and heat capacity of the ideal gas

- Internal energy: $U = -N \frac{\partial}{\partial \beta} \ln Z_1$
 $= -N \frac{\partial}{\partial \beta} (\ln Z_1^t + \ln Z_1^v + \ln Z_1^r) = U^t + U^v + U^r.$
- Heat capacity at constant volume:
 $C_V = C_V^t + C_V^v + C_V^r.$
- i. Translational motion part: (7.2.4) $Z_1^t = V \left(\frac{2\pi m}{h^2 \beta} \right)^{3/2}.$
 $U^t = \frac{3}{2} N k T, C_V^t = \frac{3}{2} N k.$
- ii. Vibration part:

Internal energy and heat capacity of the ideal gas

- Internal energy: $U = -N \frac{\partial}{\partial \beta} \ln Z_1$
 $= -N \frac{\partial}{\partial \beta} (\ln Z_1^t + \ln Z_1^v + \ln Z_1^r) = U^t + U^v + U^r.$
- Heat capacity at constant volume:
 $C_V = C_V^t + C_V^v + C_V^r.$
- i. Translational motion part: (7.2.4) $Z_1^t = V \left(\frac{2\pi m}{h^2 \beta} \right)^{3/2}.$
 $U^t = \frac{3}{2} N k T, C_V^t = \frac{3}{2} N k.$
- ii. Vibration part: taken as linear harmonic vibration,
 $\varepsilon_n = (n + \frac{1}{2}) \hbar \omega.$

Internal energy and heat capacity of the ideal gas

- Internal energy: $U = -N \frac{\partial}{\partial \beta} \ln Z_1$
 $= -N \frac{\partial}{\partial \beta} (\ln Z_1^t + \ln Z_1^v + \ln Z_1^r) = U^t + U^v + U^r.$
- Heat capacity at constant volume:
 $C_V = C_V^t + C_V^v + C_V^r.$
- i. Translational motion part: (7.2.4) $Z_1^t = V \left(\frac{2\pi m}{h^2 \beta} \right)^{3/2}.$
 $U^t = \frac{3}{2} N k T, C_V^t = \frac{3}{2} N k.$
- ii. Vibration part: taken as linear harmonic vibration,
 $\varepsilon_n = (n + \frac{1}{2}) \hbar \omega.$

Partition function: $Z_1^v = \sum_{n=0}^{\infty} e^{-\beta \hbar \omega (n + \frac{1}{2})}$

Internal energy and heat capacity of the ideal gas

- Internal energy: $U = -N \frac{\partial}{\partial \beta} \ln Z_1$
 $= -N \frac{\partial}{\partial \beta} (\ln Z_1^t + \ln Z_1^v + \ln Z_1^r) = U^t + U^v + U^r.$
- Heat capacity at constant volume:
 $C_V = C_V^t + C_V^v + C_V^r.$
- i. Translational motion part: (7.2.4) $Z_1^t = V \left(\frac{2\pi m}{h^2 \beta} \right)^{3/2}.$
 $U^t = \frac{3}{2} N k T, C_V^t = \frac{3}{2} N k.$
- ii. Vibration part: taken as linear harmonic vibration,
 $\varepsilon_n = (n + \frac{1}{2}) \hbar \omega.$

Partition function: $Z_1^v = \sum_{n=0}^{\infty} e^{-\beta \hbar \omega (n + \frac{1}{2})}$

$$= e^{-\frac{1}{2} \beta \hbar \omega} \sum_{n=0}^{\infty} (e^{-\beta \hbar \omega})^n$$

Internal energy and heat capacity of the ideal gas

- Internal energy: $U = -N \frac{\partial}{\partial \beta} \ln Z_1$
 $= -N \frac{\partial}{\partial \beta} (\ln Z_1^t + \ln Z_1^v + \ln Z_1^r) = U^t + U^v + U^r.$
- Heat capacity at constant volume:
 $C_V = C_V^t + C_V^v + C_V^r.$
- i. Translational motion part: (7.2.4) $Z_1^t = V \left(\frac{2\pi m}{h^2 \beta} \right)^{3/2}.$
 $U^t = \frac{3}{2} N k T, C_V^t = \frac{3}{2} N k.$
- ii. Vibration part: taken as linear harmonic vibration,
 $\varepsilon_n = (n + \frac{1}{2}) \hbar \omega.$

Partition function: $Z_1^v = \sum_{n=0}^{\infty} e^{-\beta \hbar \omega (n + \frac{1}{2})}$

$$= e^{-\frac{1}{2} \beta \hbar \omega} \sum_{n=0}^{\infty} (e^{-\beta \hbar \omega})^n = \frac{e^{-\frac{1}{2} \beta \hbar \omega}}{1 - e^{-\beta \hbar \omega}}.$$

Internal energy and heat capacity of the ideal gas

- $Z_1^v = \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}.$

Internal energy and heat capacity of the ideal gas

- $Z_1^v = \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}.$
- $U^v = -N \frac{\partial}{\partial \beta} \ln Z_1^v$

Internal energy and heat capacity of the ideal gas

- $Z_1^v = \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}.$
- $U^v = -N \frac{\partial}{\partial \beta} \ln Z_1^v = -N \frac{\partial}{\partial \beta} \ln\left(\frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}\right)$

Internal energy and heat capacity of the ideal gas

- $Z_1^v = \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}.$
- $$U^v = -N \frac{\partial}{\partial \beta} \ln Z_1^v = -N \frac{\partial}{\partial \beta} \ln\left(\frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}\right)$$
$$= -N \frac{\partial}{\partial \beta} \left[-\frac{1}{2}\beta\hbar\omega - \ln(1 - e^{-\beta\hbar\omega}) \right]$$

Internal energy and heat capacity of the ideal gas

- $Z_1^v = \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}.$
- $$\begin{aligned} U^v &= -N \frac{\partial}{\partial \beta} \ln Z_1^v = -N \frac{\partial}{\partial \beta} \ln \left(\frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \right) \\ &= -N \frac{\partial}{\partial \beta} \left[-\frac{1}{2}\beta\hbar\omega - \ln(1 - e^{-\beta\hbar\omega}) \right] \\ &= \frac{1}{2}N\hbar\omega + N \frac{\hbar\omega e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \end{aligned}$$

Internal energy and heat capacity of the ideal gas

- $Z_1^v = \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}.$
- $$\begin{aligned} U^v &= -N \frac{\partial}{\partial \beta} \ln Z_1^v = -N \frac{\partial}{\partial \beta} \ln \left(\frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \right) \\ &= -N \frac{\partial}{\partial \beta} \left[-\frac{1}{2}\beta\hbar\omega - \ln(1 - e^{-\beta\hbar\omega}) \right] \\ &= \frac{1}{2}N\hbar\omega + N \frac{\hbar\omega e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \\ &= \frac{1}{2}N\hbar\omega + \frac{N\hbar\omega}{e^{\beta\hbar\omega}-1}. \end{aligned}$$

Internal energy and heat capacity of the ideal gas

- $Z_1^v = \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}.$
- $$\begin{aligned} U^v &= -N \frac{\partial}{\partial \beta} \ln Z_1^v = -N \frac{\partial}{\partial \beta} \ln \left(\frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \right) \\ &= -N \frac{\partial}{\partial \beta} \left[-\frac{1}{2}\beta\hbar\omega - \ln(1 - e^{-\beta\hbar\omega}) \right] \\ &= \frac{1}{2}N\hbar\omega + N \frac{\hbar\omega e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \\ &= \frac{1}{2}N\hbar\omega + \frac{N\hbar\omega}{e^{\beta\hbar\omega}-1}. \end{aligned}$$
- $\frac{1}{2}N\hbar\omega$ zero-point energy;

Internal energy and heat capacity of the ideal gas

- $Z_1^v = \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}.$
- $$\begin{aligned} U^v &= -N \frac{\partial}{\partial \beta} \ln Z_1^v = -N \frac{\partial}{\partial \beta} \ln \left(\frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \right) \\ &= -N \frac{\partial}{\partial \beta} \left[-\frac{1}{2}\beta\hbar\omega - \ln(1 - e^{-\beta\hbar\omega}) \right] \\ &= \frac{1}{2}N\hbar\omega + N \frac{\hbar\omega e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \\ &= \frac{1}{2}N\hbar\omega + \frac{N\hbar\omega}{e^{\beta\hbar\omega}-1}. \end{aligned}$$
- $\frac{1}{2}N\hbar\omega$ zero-point energy;
 $\frac{N\hbar\omega}{e^{\beta\hbar\omega}-1}$ energy excited by heat.

Internal energy and heat capacity of the ideal gas

- Define a typical temperature: θ_v : $k\theta_v = \hbar\omega$.

Internal energy and heat capacity of the ideal gas

- Define a typical temperature: θ_v : $k\theta_v = \hbar\omega$. Then
$$U^v = \frac{1}{2}Nk\theta_v + \frac{Nk\theta_v}{e^{\theta_v/T} - 1},$$

Internal energy and heat capacity of the ideal gas

- Define a typical temperature: θ_v : $k\theta_v = \hbar\omega$. Then

$$U^v = \frac{1}{2}Nk\theta_v + \frac{Nk\theta_v}{e^{\theta_v/T} - 1},$$

$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 \frac{e^{\theta_v/T}}{(e^{\theta_v/T} - 1)^2}.$$

Internal energy and heat capacity of the ideal gas

- Define a typical temperature: θ_v : $k\theta_v = \hbar\omega$. Then
$$U^v = \frac{1}{2}Nk\theta_v + \frac{Nk\theta_v}{e^{\theta_v/T}-1},$$
$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 \frac{e^{\theta_v/T}}{(e^{\theta_v/T}-1)^2}.$$
- θ_v is determined by the property of the material.

Internal energy and heat capacity of the ideal gas

- Define a typical temperature: θ_v : $k\theta_v = \hbar\omega$. Then
$$U^v = \frac{1}{2}Nk\theta_v + \frac{Nk\theta_v}{e^{\theta_v/T}-1},$$
$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 \frac{e^{\theta_v/T}}{(e^{\theta_v/T}-1)^2}.$$
- θ_v is determined by the property of the material.
- If $T \ll \theta_v$,

Internal energy and heat capacity of the ideal gas

- Define a typical temperature: θ_v : $k\theta_v = \hbar\omega$. Then
$$U^v = \frac{1}{2}Nk\theta_v + \frac{Nk\theta_v}{e^{\theta_v/T}-1},$$
$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 \frac{e^{\theta_v/T}}{(e^{\theta_v/T}-1)^2}.$$
- θ_v is determined by the property of the material.
- If $T \ll \theta_v$, $U^v = \frac{1}{2}Nk\theta_v + Nk\theta_v e^{-\theta_v/T}$

Internal energy and heat capacity of the ideal gas

- Define a typical temperature: θ_v : $k\theta_v = \hbar\omega$. Then
$$U^v = \frac{1}{2}Nk\theta_v + \frac{Nk\theta_v}{e^{\theta_v/T}-1},$$
$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 \frac{e^{\theta_v/T}}{(e^{\theta_v/T}-1)^2}.$$
- θ_v is determined by the property of the material.
- If $T \ll \theta_v$, $U^v = \frac{1}{2}Nk\theta_v + Nk\theta_v e^{-\theta_v/T}$
$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 e^{-\theta_v/T}$$

Internal energy and heat capacity of the ideal gas

- Define a typical temperature: θ_v : $k\theta_v = \hbar\omega$. Then
$$U^v = \frac{1}{2}Nk\theta_v + \frac{Nk\theta_v}{e^{\theta_v/T}-1},$$
$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 \frac{e^{\theta_v/T}}{(e^{\theta_v/T}-1)^2}.$$
- θ_v is determined by the property of the material.
- If $T \ll \theta_v$, $U^v = \frac{1}{2}Nk\theta_v + Nk\theta_v e^{-\theta_v/T}$
$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 e^{-\theta_v/T} \rightarrow 0.$$

Internal energy and heat capacity of the ideal gas

- Define a typical temperature: θ_v : $k\theta_v = \hbar\omega$. Then

$$U^v = \frac{1}{2}Nk\theta_v + \frac{Nk\theta_v}{e^{\theta_v/T} - 1},$$

$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 \frac{e^{\theta_v/T}}{(e^{\theta_v/T} - 1)^2}.$$

- θ_v is determined by the property of the material.

- If $T \ll \theta_v$, $U^v = \frac{1}{2}Nk\theta_v + Nk\theta_v e^{-\theta_v/T}$

$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 e^{-\theta_v/T} \rightarrow 0.$$

Reason: $\hbar\omega \gg kT$.

Internal energy and heat capacity of the ideal gas

- Define a typical temperature: θ_v : $k\theta_v = \hbar\omega$. Then
$$U^v = \frac{1}{2}Nk\theta_v + \frac{Nk\theta_v}{e^{\theta_v/T} - 1},$$
$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 \frac{e^{\theta_v/T}}{(e^{\theta_v/T} - 1)^2}.$$
- θ_v is determined by the property of the material.
- If $T \ll \theta_v$, $U^v = \frac{1}{2}Nk\theta_v + Nk\theta_v e^{-\theta_v/T}$
$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 e^{-\theta_v/T} \rightarrow 0.$$
Reason: $\hbar\omega \gg kT$.
- iii. Rotational part.

Internal energy and heat capacity of the ideal gas

- Define a typical temperature: θ_v : $k\theta_v = \hbar\omega$. Then

$$U^v = \frac{1}{2}Nk\theta_v + \frac{Nk\theta_v}{e^{\theta_v/T} - 1},$$

$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 \frac{e^{\theta_v/T}}{(e^{\theta_v/T} - 1)^2}.$$
- θ_v is determined by the property of the material.
- If $T \ll \theta_v$, $U^v = \frac{1}{2}Nk\theta_v + Nk\theta_v e^{-\theta_v/T}$

$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 e^{-\theta_v/T} \rightarrow 0.$$
 Reason: $\hbar\omega \gg kT$.
- iii. Rotational part.
- iii.1. Diatomic molecule with different nuclei.

Internal energy and heat capacity of the ideal gas

- Define a typical temperature: θ_v : $k\theta_v = \hbar\omega$. Then

$$U^v = \frac{1}{2}Nk\theta_v + \frac{Nk\theta_v}{e^{\theta_v/T} - 1},$$

$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 \frac{e^{\theta_v/T}}{(e^{\theta_v/T} - 1)^2}.$$

- θ_v is determined by the property of the material.

- If $T \ll \theta_v$, $U^v = \frac{1}{2}Nk\theta_v + Nk\theta_v e^{-\theta_v/T}$

$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 e^{-\theta_v/T} \rightarrow 0.$$

Reason: $\hbar\omega \gg kT$.

- iii. Rotational part.
- iii.1. Diatomic molecule with different nuclei.

$$\varepsilon^r = \frac{l(l+1)}{2I} \hbar^2, \quad l = 0, 1, 2, \dots$$

Internal energy and heat capacity of the ideal gas

- Define a typical temperature: θ_v : $k\theta_v = \hbar\omega$. Then

$$U^v = \frac{1}{2}Nk\theta_v + \frac{Nk\theta_v}{e^{\theta_v/T} - 1},$$

$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 \frac{e^{\theta_v/T}}{(e^{\theta_v/T} - 1)^2}.$$

- θ_v is determined by the property of the material.

- If $T \ll \theta_v$, $U^v = \frac{1}{2}Nk\theta_v + Nk\theta_v e^{-\theta_v/T}$

$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 e^{-\theta_v/T} \rightarrow 0.$$

Reason: $\hbar\omega \gg kT$.

- iii. Rotational part.
- iii.1. Diatomic molecule with different nuclei.

$$\varepsilon^r = \frac{l(l+1)}{2I} \hbar^2, \quad l = 0, 1, 2, \dots$$

Degeneracy: $\omega_l = 2l + 1$.

Internal energy and heat capacity of the ideal gas

- Partition function: $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$.

Internal energy and heat capacity of the ideal gas

- Partition function: $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$.
- Define typical temperature θ_r : $k\theta_r = \frac{\hbar^2}{2I}$.

Internal energy and heat capacity of the ideal gas

- Partition function: $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$.
- Define typical temperature θ_r : $k\theta_r = \frac{\hbar^2}{2I}$.
- $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$.

Internal energy and heat capacity of the ideal gas

- Partition function: $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$.
- Define typical temperature θ_r : $k\theta_r = \frac{\hbar^2}{2I}$.
- $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$.
- If $\theta_r/T \ll 1$, ε^r is approximately continuous.

Internal energy and heat capacity of the ideal gas

- Partition function: $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$.
- Define typical temperature θ_r : $k\theta_r = \frac{\hbar^2}{2I}$.
- $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$.
- If $\theta_r/T \ll 1$, ε^r is approximately continuous. Set $x = l(l+1)\frac{\theta_r}{T}$, then $dx = (2l+1)\frac{\theta_r}{T}\Delta l$.

Internal energy and heat capacity of the ideal gas

- Partition function: $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$.
- Define typical temperature θ_r : $k\theta_r = \frac{\hbar^2}{2I}$.
- $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$.
- If $\theta_r/T \ll 1$, ε^r is approximately continuous. Set $x = l(l+1)\frac{\theta_r}{T}$, then $dx = (2l+1)\frac{\theta_r}{T}\Delta l$.

$$Z_1^r = \int_0^{\infty} (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)}$$

Internal energy and heat capacity of the ideal gas

- Partition function: $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$.
- Define typical temperature θ_r : $k\theta_r = \frac{\hbar^2}{2I}$.
- $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$.
- If $\theta_r/T \ll 1$, ε^r is approximately continuous. Set $x = l(l+1)\frac{\theta_r}{T}$, then $dx = (2l+1)\frac{\theta_r}{T}\Delta l$.

$$Z_1^r = \int_0^{\infty} (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{T}{\theta_r} \int_0^{\infty} e^{-x} dx$$

Internal energy and heat capacity of the ideal gas

- Partition function: $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$.
- Define typical temperature θ_r : $k\theta_r = \frac{\hbar^2}{2I}$.
- $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$.
- If $\theta_r/T \ll 1$, ε^r is approximately continuous. Set $x = l(l+1)\frac{\theta_r}{T}$, then $dx = (2l+1)\frac{\theta_r}{T}\Delta l$.

$$Z_1^r = \int_0^{\infty} (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{T}{\theta_r} \int_0^{\infty} e^{-x} dx = \frac{T}{\theta_r}$$

Internal energy and heat capacity of the ideal gas

- Partition function: $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$.
- Define typical temperature θ_r : $k\theta_r = \frac{\hbar^2}{2I}$.
- $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$.
- If $\theta_r/T \ll 1$, ε^r is approximately continuous. Set $x = l(l+1)\frac{\theta_r}{T}$, then $dx = (2l+1)\frac{\theta_r}{T}\Delta l$.

$$Z_1^r = \int_0^{\infty} (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{T}{\theta_r} \int_0^{\infty} e^{-x} dx = \frac{T}{\theta_r}$$

$$= T \frac{2Ik}{\hbar^2}$$

Internal energy and heat capacity of the ideal gas

- Partition function: $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$.
- Define typical temperature θ_r : $k\theta_r = \frac{\hbar^2}{2I}$.
- $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$.
- If $\theta_r/T \ll 1$, ε^r is approximately continuous. Set $x = l(l+1)\frac{\theta_r}{T}$, then $dx = (2l+1)\frac{\theta_r}{T}\Delta l$.

$$Z_1^r = \int_0^{\infty} (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{T}{\theta_r} \int_0^{\infty} e^{-x} dx = \frac{T}{\theta_r}$$

$$= T \frac{2Ik}{\hbar^2} = \frac{2I}{\beta \hbar^2}.$$

Internal energy and heat capacity of the ideal gas

- Partition function: $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$.
- Define typical temperature θ_r : $k\theta_r = \frac{\hbar^2}{2I}$.
- $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$.
- If $\theta_r/T \ll 1$, ε^r is approximately continuous. Set $x = l(l+1)\frac{\theta_r}{T}$, then $dx = (2l+1)\frac{\theta_r}{T}\Delta l$.

$$Z_1^r = \int_0^{\infty} (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{T}{\theta_r} \int_0^{\infty} e^{-x} dx = \frac{T}{\theta_r}$$

$$= T \frac{2Ik}{\hbar^2} = \frac{2I}{\beta \hbar^2}.$$
- $U^r = -N \frac{\partial}{\partial \beta} \ln Z_1^r$

Internal energy and heat capacity of the ideal gas

- Partition function: $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$.
- Define typical temperature θ_r : $k\theta_r = \frac{\hbar^2}{2I}$.
- $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$.
- If $\theta_r/T \ll 1$, ε^r is approximately continuous. Set $x = l(l+1)\frac{\theta_r}{T}$, then $dx = (2l+1)\frac{\theta_r}{T}\Delta l$.

$$Z_1^r = \int_0^{\infty} (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{T}{\theta_r} \int_0^{\infty} e^{-x} dx = \frac{T}{\theta_r}$$

$$= T \frac{2Ik}{\hbar^2} = \frac{2I}{\beta\hbar^2}.$$
- $U^r = -N \frac{\partial}{\partial \beta} \ln Z_1^r = N \left(\frac{2I}{\beta\hbar^2} \right)^{-1} \cdot \frac{2I}{\beta^2\hbar^2}$

Internal energy and heat capacity of the ideal gas

- Partition function: $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$.
- Define typical temperature θ_r : $k\theta_r = \frac{\hbar^2}{2I}$.
- $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$.
- If $\theta_r/T \ll 1$, ε^r is approximately continuous. Set $x = l(l+1)\frac{\theta_r}{T}$, then $dx = (2l+1)\frac{\theta_r}{T}\Delta l$.

$$Z_1^r = \int_0^{\infty} (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{T}{\theta_r} \int_0^{\infty} e^{-x} dx = \frac{T}{\theta_r}$$

$$= T \frac{2Ik}{\hbar^2} = \frac{2I}{\beta \hbar^2}.$$
- $U^r = -N \frac{\partial}{\partial \beta} \ln Z_1^r = N \left(\frac{2I}{\beta \hbar^2} \right)^{-1} \cdot \frac{2I}{\beta^2 \hbar^2} = \frac{N}{\beta}$

Internal energy and heat capacity of the ideal gas

- Partition function: $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$.
- Define typical temperature θ_r : $k\theta_r = \frac{\hbar^2}{2I}$.
- $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$.
- If $\theta_r/T \ll 1$, ε^r is approximately continuous. Set $x = l(l+1)\frac{\theta_r}{T}$, then $dx = (2l+1)\frac{\theta_r}{T}\Delta l$.

$$Z_1^r = \int_0^{\infty} (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{T}{\theta_r} \int_0^{\infty} e^{-x} dx = \frac{T}{\theta_r}$$

$$= T \frac{2Ik}{\hbar^2} = \frac{2I}{\beta \hbar^2}.$$
- $U^r = -N \frac{\partial}{\partial \beta} \ln Z_1^r = N \left(\frac{2I}{\beta \hbar^2} \right)^{-1} \cdot \frac{2I}{\beta^2 \hbar^2} = \frac{N}{\beta} = NkT.$

Internal energy and heat capacity of the ideal gas

- Partition function: $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$.
- Define typical temperature θ_r : $k\theta_r = \frac{\hbar^2}{2I}$.
- $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$.
- If $\theta_r/T \ll 1$, ε^r is approximately continuous. Set $x = l(l+1)\frac{\theta_r}{T}$, then $dx = (2l+1)\frac{\theta_r}{T}\Delta l$.

$$Z_1^r = \int_0^{\infty} (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{T}{\theta_r} \int_0^{\infty} e^{-x} dx = \frac{T}{\theta_r}$$

$$= T \frac{2Ik}{\hbar^2} = \frac{2I}{\beta \hbar^2}.$$
- $U^r = -N \frac{\partial}{\partial \beta} \ln Z_1^r = N \left(\frac{2I}{\beta \hbar^2} \right)^{-1} \cdot \frac{2I}{\beta^2 \hbar^2} = \frac{N}{\beta} = NkT.$
- $C_V^r = Nk.$

Internal energy and heat capacity of the ideal gas

- Partition function: $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$.
- Define typical temperature θ_r : $k\theta_r = \frac{\hbar^2}{2I}$.
- $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$.
- If $\theta_r/T \ll 1$, ε^r is approximately continuous. Set $x = l(l+1)\frac{\theta_r}{T}$, then $dx = (2l+1)\frac{\theta_r}{T}\Delta l$.

$$Z_1^r = \int_0^{\infty} (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{T}{\theta_r} \int_0^{\infty} e^{-x} dx = \frac{T}{\theta_r}$$

$$= T \frac{2Ik}{\hbar^2} = \frac{2I}{\beta \hbar^2}.$$
- $U^r = -N \frac{\partial}{\partial \beta} \ln Z_1^r = N \left(\frac{2I}{\beta \hbar^2} \right)^{-1} \cdot \frac{2I}{\beta^2 \hbar^2} = \frac{N}{\beta} = NkT.$
- $C_V^r = Nk.$
- At normal temperature (Table 7.5), $\frac{\theta_r}{T} \ll 1$ holds.

Internal energy and heat capacity of the ideal gas

- iii.2. Diatomic molecule with the same nuclei. H_2 as an example.

Internal energy and heat capacity of the ideal gas

- iii.2. Diatomic molecule with the same nuclei. H_2 as an example.
- At normal temperature, there are $3/4$ orthohydrogen (l odd) and $1/4$ parahydrogen (l even).

Internal energy and heat capacity of the ideal gas

- iii.2. Diatomic molecule with the same nuclei. H_2 as an example.
- At normal temperature, there are 3/4 orthohydrogen (l odd) and 1/4 parahydrogen (l even).
- Orthohydrogen: $Z_{1o}^r = \sum_{l=1,3,\dots} (2l+1)e^{-\frac{l(l+1)\theta_r}{T}};$

Internal energy and heat capacity of the ideal gas

- iii.2. Diatomic molecule with the same nuclei. H_2 as an example.
- At normal temperature, there are 3/4 orthohydrogen (l odd) and 1/4 parahydrogen (l even).
- Orthohydrogen: $Z_{1o}^r = \sum_{l=1,3,\dots} (2l+1)e^{-\frac{l(l+1)\theta_r}{T}}$;

$$\text{Parahydrogen: } Z_{1p}^r = \sum_{l=0,2,\dots} (2l+1)e^{-\frac{l(l+1)\theta_r}{T}}.$$

Internal energy and heat capacity of the ideal gas

- iii.2. Diatomic molecule with the same nuclei. H_2 as an example.
- At normal temperature, there are 3/4 orthohydrogen (l odd) and 1/4 parahydrogen (l even).
- Orthohydrogen: $Z_{1o}^r = \sum_{l=1,3,\dots} (2l+1)e^{-\frac{l(l+1)\theta_r}{T}}$;
Parahydrogen: $Z_{1p}^r = \sum_{l=0,2,\dots} (2l+1)e^{-\frac{l(l+1)\theta_r}{T}}$.
- $U^r = U_o^r + U_p^r$

Internal energy and heat capacity of the ideal gas

- iii.2. Diatomic molecule with the same nuclei. H_2 as an example.
- At normal temperature, there are 3/4 orthohydrogen (l odd) and 1/4 parahydrogen (l even).

- Orthohydrogen:
$$Z_{1o}^r = \sum_{l=1,3,\dots} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}};$$

Parahydrogen:
$$Z_{1p}^r = \sum_{l=0,2,\dots} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}.$$

- $$U^r = U_o^r + U_p^r = -\frac{3}{4}N \frac{\partial}{\partial \beta} \ln Z_{1o}^r - \frac{1}{4}N \frac{\partial}{\partial \beta} \ln Z_{1p}^r.$$

Internal energy and heat capacity of the ideal gas

- iii.2. Diatomic molecule with the same nuclei. H_2 as an example.
- At normal temperature, there are 3/4 orthohydrogen (l odd) and 1/4 parahydrogen (l even).

- Orthohydrogen: $Z_{1o}^r = \sum_{l=1,3,\dots} (2l+1)e^{-\frac{l(l+1)\theta_r}{T}};$

Parahydrogen: $Z_{1p}^r = \sum_{l=0,2,\dots} (2l+1)e^{-\frac{l(l+1)\theta_r}{T}}.$

- $U^r = U_o^r + U_p^r = -\frac{3}{4}N \frac{\partial}{\partial \beta} \ln Z_{1o}^r - \frac{1}{4}N \frac{\partial}{\partial \beta} \ln Z_{1p}^r.$
- If $\theta_r \ll T$, the hydrogen molecule is in high l state,
 $l \sim l+1,$

Internal energy and heat capacity of the ideal gas

- iii.2. Diatomic molecule with the same nuclei. H_2 as an example.
- At normal temperature, there are 3/4 orthohydrogen (l odd) and 1/4 parahydrogen (l even).
- Orthohydrogen: $Z_{1o}^r = \sum_{l=1,3,\dots} (2l+1)e^{-\frac{l(l+1)\theta_r}{T}}$;

$$\text{Parahydrogen: } Z_{1p}^r = \sum_{l=0,2,\dots} (2l+1)e^{-\frac{l(l+1)\theta_r}{T}}.$$

- $U^r = U_o^r + U_p^r = -\frac{3}{4}N \frac{\partial}{\partial \beta} \ln Z_{1o}^r - \frac{1}{4}N \frac{\partial}{\partial \beta} \ln Z_{1p}^r.$
- If $\theta_r \ll T$, the hydrogen molecule is in high l state, $l \sim l+1$, then $\sum_{l=0,2,\dots}^{\infty} \dots \simeq \sum_{l=1,3,\dots}^{\infty} \dots \simeq \frac{1}{2} \sum_{l=0,1,\dots}^{\infty} \dots$

Internal energy and heat capacity of the ideal gas

- $Z_{1o}^r \simeq Z_{1p}^r \simeq \frac{1}{2} \sum_{l=0,1,\dots} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$

Internal energy and heat capacity of the ideal gas

- $$\begin{aligned} Z_{1o}^r &\simeq Z_{1p}^r \simeq \frac{1}{2} \sum_{l=0,1,\dots} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}} \\ &\simeq \frac{1}{2} \int_0^\infty (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} \end{aligned}$$

Internal energy and heat capacity of the ideal gas

- $$\begin{aligned} Z_{1o}^r &\simeq Z_{1p}^r \simeq \frac{1}{2} \sum_{l=0,1,\dots} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}} \\ &\simeq \frac{1}{2} \int_0^\infty (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{1}{2} \frac{T}{\theta_r} \end{aligned}$$

Internal energy and heat capacity of the ideal gas

- $$\begin{aligned} Z_{1o}^r &\simeq Z_{1p}^r \simeq \frac{1}{2} \sum_{l=0,1,\dots} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}} \\ &\simeq \frac{1}{2} \int_0^\infty (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{1}{2} \frac{T}{\theta_r} = \frac{I}{\beta \hbar^2}. \end{aligned}$$

Internal energy and heat capacity of the ideal gas

- $Z_{1o}^r \simeq Z_{1p}^r \simeq \frac{1}{2} \sum_{l=0,1,\dots} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$
 $\simeq \frac{1}{2} \int_0^\infty (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{1}{2} \frac{T}{\theta_r} = \frac{I}{\beta \hbar^2}.$
- $U^r = -\frac{3}{4} N \frac{\partial}{\partial \beta} \ln Z_{1o}^r - \frac{1}{4} N \frac{\partial}{\partial \beta} \ln Z_{1p}^r$

Internal energy and heat capacity of the ideal gas

- $Z_{1o}^r \simeq Z_{1p}^r \simeq \frac{1}{2} \sum_{l=0,1,\dots} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$
 $\simeq \frac{1}{2} \int_0^\infty (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{1}{2} \frac{T}{\theta_r} = \frac{I}{\beta \hbar^2}.$
- $U^r = -\frac{3}{4} N \frac{\partial}{\partial \beta} \ln Z_{1o}^r - \frac{1}{4} N \frac{\partial}{\partial \beta} \ln Z_{1p}^r \simeq -N \frac{\partial}{\partial \beta} \ln Z_{1o}^r$

Internal energy and heat capacity of the ideal gas

- $Z_{1o}^r \simeq Z_{1p}^r \simeq \frac{1}{2} \sum_{l=0,1,\dots} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$
 $\simeq \frac{1}{2} \int_0^\infty (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{1}{2} \frac{T}{\theta_r} = \frac{I}{\beta \hbar^2}.$
- $U^r = -\frac{3}{4} N \frac{\partial}{\partial \beta} \ln Z_{1o}^r - \frac{1}{4} N \frac{\partial}{\partial \beta} \ln Z_{1p}^r \simeq -N \frac{\partial}{\partial \beta} \ln Z_{1o}^r$
 $= -N \frac{\partial}{\partial \beta} \ln \frac{I}{\beta \hbar^2}$

Internal energy and heat capacity of the ideal gas

- $$Z_{1o}^r \simeq Z_{1p}^r \simeq \frac{1}{2} \sum_{l=0,1,\dots} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$$

$$\simeq \frac{1}{2} \int_0^\infty (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{1}{2} \frac{T}{\theta_r} = \frac{I}{\beta \hbar^2}.$$
- $$U^r = -\frac{3}{4} N \frac{\partial}{\partial \beta} \ln Z_{1o}^r - \frac{1}{4} N \frac{\partial}{\partial \beta} \ln Z_{1p}^r \simeq -N \frac{\partial}{\partial \beta} \ln Z_{1o}^r$$

$$= -N \frac{\partial}{\partial \beta} \ln \frac{I}{\beta \hbar^2} = \frac{N}{\beta}$$

Internal energy and heat capacity of the ideal gas

- $Z_{1o}^r \simeq Z_{1p}^r \simeq \frac{1}{2} \sum_{l=0,1,\dots} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$
 $\simeq \frac{1}{2} \int_0^\infty (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{1}{2} \frac{T}{\theta_r} = \frac{I}{\beta \hbar^2}.$
- $U^r = -\frac{3}{4} N \frac{\partial}{\partial \beta} \ln Z_{1o}^r - \frac{1}{4} N \frac{\partial}{\partial \beta} \ln Z_{1p}^r \simeq -N \frac{\partial}{\partial \beta} \ln Z_{1o}^r$
 $= -N \frac{\partial}{\partial \beta} \ln \frac{I}{\beta \hbar^2} = \frac{N}{\beta} = NkT.$

Internal energy and heat capacity of the ideal gas

- $Z_{1o}^r \simeq Z_{1p}^r \simeq \frac{1}{2} \sum_{l=0,1,\dots} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$
 $\simeq \frac{1}{2} \int_0^\infty (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{1}{2} \frac{T}{\theta_r} = \frac{I}{\beta \hbar^2}.$
- $U^r = -\frac{3}{4} N \frac{\partial}{\partial \beta} \ln Z_{1o}^r - \frac{1}{4} N \frac{\partial}{\partial \beta} \ln Z_{1p}^r \simeq -N \frac{\partial}{\partial \beta} \ln Z_{1o}^r$
 $= -N \frac{\partial}{\partial \beta} \ln \frac{I}{\beta \hbar^2} = \frac{N}{\beta} = NkT.$
- $C_V^r = Nk.$

Internal energy and heat capacity of the ideal gas

- $$Z_{1o}^r \simeq Z_{1p}^r \simeq \frac{1}{2} \sum_{l=0,1,\dots} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$$

$$\simeq \frac{1}{2} \int_0^\infty (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{1}{2} \frac{T}{\theta_r} = \frac{I}{\beta \hbar^2}.$$
- $$U^r = -\frac{3}{4} N \frac{\partial}{\partial \beta} \ln Z_{1o}^r - \frac{1}{4} N \frac{\partial}{\partial \beta} \ln Z_{1p}^r \simeq -N \frac{\partial}{\partial \beta} \ln Z_{1o}^r$$

$$= -N \frac{\partial}{\partial \beta} \ln \frac{I}{\beta \hbar^2} = \frac{N}{\beta} = NkT.$$
- $$C_V^r = Nk.$$
- If $\theta_r \ll T$ does not hold, the summation is necessary.

General processes for **classical** statistics

General processes for **classical** statistics

- Object as an example: diatomic molecules.

General processes for **classical** statistics

- Object as an example: diatomic molecules.

- $\varepsilon =$

$$\frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2) + \frac{1}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2).$$

General processes for **classical** statistics

- Object as an example: diatomic molecules.

- $\varepsilon =$

$$\frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2) + \frac{1}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2).$$

- $Z_1 = \int \dots \int e^{-\beta \varepsilon(q,p)} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r},$

General processes for **classical** statistics

- Object as an example: diatomic molecules.

- $\varepsilon =$

$$\frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2) + \frac{1}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2).$$

- $Z_1 = \int \dots \int e^{-\beta \varepsilon(q,p)} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r},$
 $Z_1^t = \int e^{-\frac{\beta}{2m}(p_x^2 + p_y^2 + p_z^2)} \frac{dx dy dz dp_x dp_y dp_z}{h_0^3},$

General processes for **classical** statistics

- Object as an example: diatomic molecules.

- $\varepsilon =$

$$\frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2) + \frac{1}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2).$$

- $Z_1 = \int \dots \int e^{-\beta \varepsilon(q,p)} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r},$

$$Z_1^t = \int e^{-\frac{\beta}{2m}(p_x^2 + p_y^2 + p_z^2)} \frac{dx dy dz dp_x dp_y dp_z}{h_0^3},$$

$$Z_1^r = \int e^{-\frac{\beta}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2)} \frac{dp_\theta dp_\varphi d\theta d\varphi}{h_0^2},$$

General processes for **classical** statistics

- Object as an example: diatomic molecules.

- $\varepsilon =$

$$\frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2) + \frac{1}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2).$$

- $Z_1 = \int \dots \int e^{-\beta \varepsilon(q,p)} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r},$

$$Z_1^t = \int e^{-\frac{\beta}{2m}(p_x^2 + p_y^2 + p_z^2)} \frac{dx dy dz dp_x dp_y dp_z}{h_0^3},$$

$$Z_1^r = \int e^{-\frac{\beta}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2)} \frac{dp_\theta dp_\varphi d\theta d\varphi}{h_0^2},$$

$$Z_1^v = \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0}.$$

General processes for **classical** statistics

- Object as an example: diatomic molecules.

- $\varepsilon =$

$$\frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2) + \frac{1}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2).$$

- $Z_1 = \int \dots \int e^{-\beta \varepsilon(q,p)} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r},$

$$Z_1^t = \int e^{-\frac{\beta}{2m}(p_x^2 + p_y^2 + p_z^2)} \frac{dx dy dz dp_x dp_y dp_z}{h_0^3},$$

$$Z_1^r = \int e^{-\frac{\beta}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2)} \frac{dp_\theta dp_\varphi d\theta d\varphi}{h_0^2},$$

$$Z_1^v = \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0}.$$

- Integration:

General processes for **classical** statistics

- Object as an example: diatomic molecules.

- $\varepsilon =$

$$\frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2) + \frac{1}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2).$$

- $Z_1 = \int \dots \int e^{-\beta \varepsilon(q,p)} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r},$

$$Z_1^t = \int e^{-\frac{\beta}{2m}(p_x^2 + p_y^2 + p_z^2)} \frac{dx dy dz dp_x dp_y dp_z}{h_0^3},$$

$$Z_1^r = \int e^{-\frac{\beta}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2)} \frac{dp_\theta dp_\varphi d\theta d\varphi}{h_0^2},$$

$$Z_1^v = \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0}.$$

- Integration:

$$Z_1^t = V \left(\frac{2\pi m}{h_0^2 \beta} \right)^{3/2}.$$

General processes for classical statistics

- $$Z_1^r = \int e^{-\frac{\beta}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2)} \frac{dp_\theta dp_\varphi d\theta d\varphi}{h_0^2}$$

General processes for classical statistics

- $$Z_1^r = \int e^{-\frac{\beta}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2)} \frac{dp_\theta dp_\varphi d\theta d\varphi}{h_0^2}$$
$$= \frac{1}{h_0^2} \int_0^{2\pi} d\varphi \int_0^\pi d\theta \int_{-\infty}^\infty e^{-\frac{\beta}{2I} p_\theta^2} dp_\theta \int_{-\infty}^\infty e^{-\frac{\beta}{2I \sin^2 \theta} p_\varphi^2} dp_\varphi$$

General processes for classical statistics

$$\begin{aligned}
 \bullet \quad Z_1^r &= \int e^{-\frac{\beta}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2)} \frac{dp_\theta dp_\varphi d\theta d\varphi}{h_0^2} \\
 &= \frac{1}{h_0^2} \int_0^{2\pi} d\varphi \int_0^\pi d\theta \int_{-\infty}^{\infty} e^{-\frac{\beta}{2I} p_\theta^2} dp_\theta \int_{-\infty}^{\infty} e^{-\frac{\beta}{2I \sin^2 \theta} p_\varphi^2} dp_\varphi \\
 &= \\
 &\frac{2\pi}{h_0^2} \int_0^\pi d\theta \sqrt{\frac{2I}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2I \sin^2 \theta}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2I \sin^2 \theta} p_\varphi^2} d\sqrt{\frac{\beta}{2I \sin^2 \theta} p_\varphi}
 \end{aligned}$$

General processes for classical statistics

$$\begin{aligned}
 \bullet \quad Z_1^r &= \int e^{-\frac{\beta}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2)} \frac{dp_\theta dp_\varphi d\theta d\varphi}{h_0^2} \\
 &= \frac{1}{h_0^2} \int_0^{2\pi} d\varphi \int_0^\pi d\theta \int_{-\infty}^{\infty} e^{-\frac{\beta}{2I} p_\theta^2} dp_\theta \int_{-\infty}^{\infty} e^{-\frac{\beta}{2I \sin^2 \theta} p_\varphi^2} dp_\varphi \\
 &= \\
 &\frac{2\pi}{h_0^2} \int_0^\pi d\theta \sqrt{\frac{2I}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2I \sin^2 \theta}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2I \sin^2 \theta} p_\varphi^2} d\sqrt{\frac{\beta}{2I \sin^2 \theta}} p_\varphi \\
 &= \frac{2\pi}{h_0^2} \int_0^\pi d\theta \sqrt{\frac{2I}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2I \sin^2 \theta}{\beta}} \sqrt{\pi}
 \end{aligned}$$

General processes for classical statistics

$$\begin{aligned}
 \bullet \quad Z_1^r &= \int e^{-\frac{\beta}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2)} \frac{dp_\theta dp_\varphi d\theta d\varphi}{h_0^2} \\
 &= \frac{1}{h_0^2} \int_0^{2\pi} d\varphi \int_0^\pi d\theta \int_{-\infty}^{\infty} e^{-\frac{\beta}{2I} p_\theta^2} dp_\theta \int_{-\infty}^{\infty} e^{-\frac{\beta}{2I \sin^2 \theta} p_\varphi^2} dp_\varphi \\
 &= \\
 &\frac{2\pi}{h_0^2} \int_0^\pi d\theta \sqrt{\frac{2I}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2I \sin^2 \theta}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2I \sin^2 \theta} p_\varphi^2} d\sqrt{\frac{\beta}{2I \sin^2 \theta} p_\varphi} \\
 &= \frac{2\pi}{h_0^2} \int_0^\pi d\theta \sqrt{\frac{2I}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2I \sin^2 \theta}{\beta}} \sqrt{\pi} \\
 &= \frac{2\pi^2}{h_0^2} \frac{2I}{\beta} \int_0^\pi \sin \theta d\theta
 \end{aligned}$$

General processes for classical statistics

$$\begin{aligned}
 \bullet \quad Z_1^r &= \int e^{-\frac{\beta}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2)} \frac{dp_\theta dp_\varphi d\theta d\varphi}{h_0^2} \\
 &= \frac{1}{h_0^2} \int_0^{2\pi} d\varphi \int_0^\pi d\theta \int_{-\infty}^{\infty} e^{-\frac{\beta}{2I} p_\theta^2} dp_\theta \int_{-\infty}^{\infty} e^{-\frac{\beta}{2I \sin^2 \theta} p_\varphi^2} dp_\varphi \\
 &= \\
 &\frac{2\pi}{h_0^2} \int_0^\pi d\theta \sqrt{\frac{2I}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2I \sin^2 \theta}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2I \sin^2 \theta} p_\varphi^2} d\sqrt{\frac{\beta}{2I \sin^2 \theta}} p_\varphi \\
 &= \frac{2\pi}{h_0^2} \int_0^\pi d\theta \sqrt{\frac{2I}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2I \sin^2 \theta}{\beta}} \sqrt{\pi} \\
 &= \frac{2\pi^2}{h_0^2} \frac{2I}{\beta} \int_0^\pi \sin \theta d\theta \\
 &= \frac{8\pi^2 I}{h_0^2 \beta}.
 \end{aligned}$$

General processes for classical statistics

- $$Z_1^v = \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0}$$

General processes for classical statistics

- $$Z_1^v = \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0}$$
$$= \frac{1}{h_0} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} dp_r \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} dr$$

General processes for classical statistics

$$\begin{aligned}
 \bullet \quad Z_1^v &= \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0} \\
 &= \frac{1}{h_0} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} dp_r \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} dr \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} d\sqrt{\frac{\beta}{2m_\mu}} p_r \cdot \\
 &\quad \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} d\sqrt{\frac{m_\mu \omega^2 \beta}{2}} r
 \end{aligned}$$

General processes for classical statistics

$$\begin{aligned}
 \bullet \quad Z_1^v &= \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0} \\
 &= \frac{1}{h_0} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} dp_r \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} dr \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} d\sqrt{\frac{\beta}{2m_\mu}} p_r \cdot \\
 &\quad \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} d\sqrt{\frac{m_\mu \omega^2 \beta}{2}} r \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \sqrt{\pi}
 \end{aligned}$$

General processes for classical statistics

$$\begin{aligned}
 \bullet \quad Z_1^v &= \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0} \\
 &= \frac{1}{h_0} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} dp_r \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} dr \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} d\sqrt{\frac{\beta}{2m_\mu}} p_r \cdot \\
 &\quad \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} d\sqrt{\frac{m_\mu \omega^2 \beta}{2}} r \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \sqrt{\pi} \\
 &= \frac{2\pi}{h_0 \beta \omega}.
 \end{aligned}$$

General processes for classical statistics

- $$\begin{aligned}
 Z_1^v &= \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0} \\
 &= \frac{1}{h_0} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} dp_r \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} dr \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} d\sqrt{\frac{\beta}{2m_\mu}} p_r \cdot \\
 &\quad \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} d\sqrt{\frac{m_\mu \omega^2 \beta}{2}} r \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \sqrt{\pi} \\
 &= \frac{2\pi}{h_0 \beta \omega}.
 \end{aligned}$$
- $$U^t = -N \frac{\partial}{\partial \beta} \ln Z_1^t$$

General processes for classical statistics

- $$\begin{aligned}
 Z_1^v &= \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0} \\
 &= \frac{1}{h_0} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} dp_r \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} dr \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} d\sqrt{\frac{\beta}{2m_\mu}} p_r \cdot \\
 &\quad \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} d\sqrt{\frac{m_\mu \omega^2 \beta}{2}} r \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \sqrt{\pi} \\
 &= \frac{2\pi}{h_0 \beta \omega}.
 \end{aligned}$$
- $$U^t = -N \frac{\partial}{\partial \beta} \ln Z_1^t = -N \frac{\partial}{\partial \beta} \ln [V (\frac{2\pi m}{h_0^2 \beta})^{3/2}]$$

General processes for classical statistics

- $$\begin{aligned}
 Z_1^v &= \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0} \\
 &= \frac{1}{h_0} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} dp_r \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} dr \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} d\sqrt{\frac{\beta}{2m_\mu}} p_r \cdot \\
 &\quad \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} d\sqrt{\frac{m_\mu \omega^2 \beta}{2}} r \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \sqrt{\pi} \\
 &= \frac{2\pi}{h_0 \beta \omega}.
 \end{aligned}$$
- $$U^t = -N \frac{\partial}{\partial \beta} \ln Z_1^t = -N \frac{\partial}{\partial \beta} \ln [V (\frac{2\pi m}{h_0^2 \beta})^{3/2}] = \frac{3}{2} N k T.$$

General processes for classical statistics

- $$\begin{aligned}
 Z_1^v &= \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0} \\
 &= \frac{1}{h_0} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} dp_r \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} dr \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} d\sqrt{\frac{\beta}{2m_\mu}} p_r \cdot \\
 &\quad \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} d\sqrt{\frac{m_\mu \omega^2 \beta}{2}} r \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \sqrt{\pi} \\
 &= \frac{2\pi}{h_0 \beta \omega}.
 \end{aligned}$$
- $$\begin{aligned}
 U^t &= -N \frac{\partial}{\partial \beta} \ln Z_1^t = -N \frac{\partial}{\partial \beta} \ln [V (\frac{2\pi m}{h_0^2 \beta})^{3/2}] = \frac{3}{2} N k T. \\
 U^r &= -N \frac{\partial}{\partial \beta} \ln Z_1^r
 \end{aligned}$$

General processes for classical statistics

- $$\begin{aligned}
 Z_1^v &= \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0} \\
 &= \frac{1}{h_0} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} dp_r \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} dr \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} d\sqrt{\frac{\beta}{2m_\mu}} p_r \cdot \\
 &\quad \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} d\sqrt{\frac{m_\mu \omega^2 \beta}{2}} r \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \sqrt{\pi} \\
 &= \frac{2\pi}{h_0 \beta \omega}.
 \end{aligned}$$
- $$\begin{aligned}
 U^t &= -N \frac{\partial}{\partial \beta} \ln Z_1^t = -N \frac{\partial}{\partial \beta} \ln [V (\frac{2\pi m}{h_0^2 \beta})^{3/2}] = \frac{3}{2} N k T. \\
 U^r &= -N \frac{\partial}{\partial \beta} \ln Z_1^r = -N \frac{\partial}{\partial \beta} \ln \frac{8\pi^2 I}{h_0^2 \beta}
 \end{aligned}$$

General processes for classical statistics

- $$\begin{aligned}
 Z_1^v &= \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0} \\
 &= \frac{1}{h_0} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} dp_r \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} dr \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} d\sqrt{\frac{\beta}{2m_\mu}} p_r \cdot \\
 &\quad \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} d\sqrt{\frac{m_\mu \omega^2 \beta}{2}} r \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \sqrt{\pi} \\
 &= \frac{2\pi}{h_0 \beta \omega}.
 \end{aligned}$$
- $$\begin{aligned}
 U^t &= -N \frac{\partial}{\partial \beta} \ln Z_1^t = -N \frac{\partial}{\partial \beta} \ln [V (\frac{2\pi m}{h_0^2 \beta})^{3/2}] = \frac{3}{2} N k T. \\
 U^r &= -N \frac{\partial}{\partial \beta} \ln Z_1^r = -N \frac{\partial}{\partial \beta} \ln \frac{8\pi^2 I}{h_0^2 \beta} = N k T.
 \end{aligned}$$

General processes for classical statistics

- $$\begin{aligned}
 Z_1^v &= \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0} \\
 &= \frac{1}{h_0} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} dp_r \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} dr \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} d\sqrt{\frac{\beta}{2m_\mu}} p_r \cdot \\
 &\quad \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} d\sqrt{\frac{m_\mu \omega^2 \beta}{2}} r \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \sqrt{\pi} \\
 &= \frac{2\pi}{h_0 \beta \omega}.
 \end{aligned}$$
- $$\begin{aligned}
 U^t &= -N \frac{\partial}{\partial \beta} \ln Z_1^t = -N \frac{\partial}{\partial \beta} \ln [V (\frac{2\pi m}{h_0^2 \beta})^{3/2}] = \frac{3}{2} N k T. \\
 U^r &= -N \frac{\partial}{\partial \beta} \ln Z_1^r = -N \frac{\partial}{\partial \beta} \ln \frac{8\pi^2 I}{h_0^2 \beta} = N k T. \\
 U^v &= -N \frac{\partial}{\partial \beta} \ln Z_1^v
 \end{aligned}$$

General processes for classical statistics

- $$\begin{aligned}
 Z_1^v &= \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0} \\
 &= \frac{1}{h_0} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} dp_r \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} dr \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} d\sqrt{\frac{\beta}{2m_\mu}} p_r \cdot \\
 &\quad \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} d\sqrt{\frac{m_\mu \omega^2 \beta}{2}} r \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \sqrt{\pi} \\
 &= \frac{2\pi}{h_0 \beta \omega}.
 \end{aligned}$$
- $$\begin{aligned}
 U^t &= -N \frac{\partial}{\partial \beta} \ln Z_1^t = -N \frac{\partial}{\partial \beta} \ln [V (\frac{2\pi m}{h_0^2 \beta})^{3/2}] = \frac{3}{2} N k T. \\
 U^r &= -N \frac{\partial}{\partial \beta} \ln Z_1^r = -N \frac{\partial}{\partial \beta} \ln \frac{8\pi^2 I}{h_0^2 \beta} = N k T. \\
 U^v &= -N \frac{\partial}{\partial \beta} \ln Z_1^v = -N \frac{\partial}{\partial \beta} \ln \frac{2\pi}{h_0 \beta \omega}
 \end{aligned}$$

General processes for classical statistics

- $$\begin{aligned}
 Z_1^v &= \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0} \\
 &= \frac{1}{h_0} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} dp_r \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} dr \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} d\sqrt{\frac{\beta}{2m_\mu}} p_r \cdot \\
 &\quad \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} d\sqrt{\frac{m_\mu \omega^2 \beta}{2}} r \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \sqrt{\pi} \\
 &= \frac{2\pi}{h_0 \beta \omega}.
 \end{aligned}$$
- $$\begin{aligned}
 U^t &= -N \frac{\partial}{\partial \beta} \ln Z_1^t = -N \frac{\partial}{\partial \beta} \ln [V (\frac{2\pi m}{h_0^2 \beta})^{3/2}] = \frac{3}{2} N k T. \\
 U^r &= -N \frac{\partial}{\partial \beta} \ln Z_1^r = -N \frac{\partial}{\partial \beta} \ln \frac{8\pi^2 I}{h_0^2 \beta} = N k T. \\
 U^v &= -N \frac{\partial}{\partial \beta} \ln Z_1^v = -N \frac{\partial}{\partial \beta} \ln \frac{2\pi}{h_0 \beta \omega} = N k T.
 \end{aligned}$$

Table of contents

1 §7 Boltzmann Statistics

- 7.1 Thermal quantities in statistics
- 7.2 Equation of state of ideal gas
- 7.3 Maxwell speed distribution
- 7.4 Energy equipartition theorem
- 7.5 Internal energy and heat capacity of the ideal gas
- **7.6 Entropy of the ideal gas**
- 7.7 Einstein's theory on heat capacity of solid
- 7.8 Paramagnetic solid
- 7.9 Negative temperature

7.6 Entropy of the ideal gas

7.6 Entropy of the ideal gas

- Object: Monatomic molecules (translational motion).

7.6 Entropy of the ideal gas

- Object: Monatomic molecules (translational motion).
- (7.1.13): $S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1)$.

7.6 Entropy of the ideal gas

- Object: Monatomic molecules (translational motion).
- (7.1.13): $S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1)$.
- (7.5.21): $Z_1^t = V(\frac{2\pi m}{h_0^2 \beta})^{3/2}$.

7.6 Entropy of the ideal gas

- Object: Monatomic molecules (translational motion).
- (7.1.13): $S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1)$.
- (7.5.21): $Z_1^t = V(\frac{2\pi m}{h_0^2 \beta})^{3/2}$.
- $S = Nk\{\ln[V(\frac{2\pi m}{h_0^2 \beta})^{3/2}] - \beta \frac{\partial}{\partial \beta} \ln[V(\frac{2\pi m}{h_0^2 \beta})^{3/2}]\}$

7.6 Entropy of the ideal gas

- Object: Monatomic molecules (translational motion).
- (7.1.13): $S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1)$.
- (7.5.21): $Z_1^t = V(\frac{2\pi m}{h_0^2 \beta})^{3/2}$.
- $$S = Nk\{\ln[V(\frac{2\pi m}{h_0^2 \beta})^{3/2}] - \beta \frac{\partial}{\partial \beta} \ln[V(\frac{2\pi m}{h_0^2 \beta})^{3/2}]\}$$
$$= Nk\{\ln V + \frac{3}{2} \ln(\frac{2\pi m}{h_0^2 \beta}) - \beta \frac{\partial}{\partial \beta} [\ln V + \frac{3}{2} \ln(\frac{2\pi m}{h_0^2 \beta})]\}$$

7.6 Entropy of the ideal gas

- Object: Monatomic molecules (translational motion).
- (7.1.13): $S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1)$.
- (7.5.21): $Z_1^t = V(\frac{2\pi m}{h_0^2 \beta})^{3/2}$.
- $$\begin{aligned} S &= Nk \left\{ \ln \left[V \left(\frac{2\pi m}{h_0^2 \beta} \right)^{3/2} \right] - \beta \frac{\partial}{\partial \beta} \ln \left[V \left(\frac{2\pi m}{h_0^2 \beta} \right)^{3/2} \right] \right\} \\ &= Nk \left\{ \ln V + \frac{3}{2} \ln \left(\frac{2\pi m}{h_0^2 \beta} \right) - \beta \frac{\partial}{\partial \beta} \left[\ln V + \frac{3}{2} \ln \left(\frac{2\pi m}{h_0^2 \beta} \right) \right] \right\} \\ &= Nk \left[\ln V + \frac{3}{2} \ln \left(\frac{2\pi m}{h_0^2 \beta} \right) - 0 - \frac{3}{2} \beta \cdot \left(-\frac{1}{\beta} \right) \right] \end{aligned}$$

7.6 Entropy of the ideal gas

- Object: Monatomic molecules (translational motion).
- (7.1.13): $S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1)$.
- (7.5.21): $Z_1^t = V(\frac{2\pi m}{h_0^2 \beta})^{3/2}$.
- $$\begin{aligned} S &= Nk \left\{ \ln \left[V \left(\frac{2\pi m}{h_0^2 \beta} \right)^{3/2} \right] - \beta \frac{\partial}{\partial \beta} \ln \left[V \left(\frac{2\pi m}{h_0^2 \beta} \right)^{3/2} \right] \right\} \\ &= Nk \left\{ \ln V + \frac{3}{2} \ln \left(\frac{2\pi m}{h_0^2 \beta} \right) - \beta \frac{\partial}{\partial \beta} \left[\ln V + \frac{3}{2} \ln \left(\frac{2\pi m}{h_0^2 \beta} \right) \right] \right\} \\ &= Nk \left[\ln V + \frac{3}{2} \ln \left(\frac{2\pi m}{h_0^2 \beta} \right) - 0 - \frac{3}{2} \beta \cdot \left(-\frac{1}{\beta} \right) \right] \\ &= Nk \left[\ln V + \frac{3}{2} \ln \left(\frac{2\pi m}{h_0^2 \beta} \right) + \frac{3}{2} \right] \end{aligned}$$

7.6 Entropy of the ideal gas

- Object: Monatomic molecules (translational motion).
- (7.1.13): $S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1)$.
- (7.5.21): $Z_1^t = V(\frac{2\pi m}{h_0^2 \beta})^{3/2}$.
- $$\begin{aligned} S &= Nk \left\{ \ln \left[V \left(\frac{2\pi m}{h_0^2 \beta} \right)^{3/2} \right] - \beta \frac{\partial}{\partial \beta} \ln \left[V \left(\frac{2\pi m}{h_0^2 \beta} \right)^{3/2} \right] \right\} \\ &= Nk \left\{ \ln V + \frac{3}{2} \ln \left(\frac{2\pi m}{h_0^2 \beta} \right) - \beta \frac{\partial}{\partial \beta} \left[\ln V + \frac{3}{2} \ln \left(\frac{2\pi m}{h_0^2 \beta} \right) \right] \right\} \\ &= Nk \left[\ln V + \frac{3}{2} \ln \left(\frac{2\pi m}{h_0^2 \beta} \right) - 0 - \frac{3}{2} \beta \cdot \left(-\frac{1}{\beta} \right) \right] \\ &= Nk \left[\ln V + \frac{3}{2} \ln \left(\frac{2\pi m}{h_0^2 \beta} \right) + \frac{3}{2} \right] \\ &= \frac{3}{2} Nk \ln T + Nk \ln V + \frac{3}{2} Nk \left[1 + \ln \frac{2\pi m k}{h_0^2} \right]. \end{aligned}$$

7.6 Entropy of the ideal gas

- Object: Monatomic molecules (translational motion).
- (7.1.13): $S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1)$.
- (7.5.21): $Z_1^t = V(\frac{2\pi m}{h_0^2 \beta})^{3/2}$.
- $$\begin{aligned} S &= Nk \left\{ \ln \left[V \left(\frac{2\pi m}{h_0^2 \beta} \right)^{3/2} \right] - \beta \frac{\partial}{\partial \beta} \ln \left[V \left(\frac{2\pi m}{h_0^2 \beta} \right)^{3/2} \right] \right\} \\ &= Nk \left\{ \ln V + \frac{3}{2} \ln \left(\frac{2\pi m}{h_0^2 \beta} \right) - \beta \frac{\partial}{\partial \beta} \left[\ln V + \frac{3}{2} \ln \left(\frac{2\pi m}{h_0^2 \beta} \right) \right] \right\} \\ &= Nk \left[\ln V + \frac{3}{2} \ln \left(\frac{2\pi m}{h_0^2 \beta} \right) - 0 - \frac{3}{2} \beta \cdot \left(-\frac{1}{\beta} \right) \right] \\ &= Nk \left[\ln V + \frac{3}{2} \ln \left(\frac{2\pi m}{h_0^2 \beta} \right) + \frac{3}{2} \right] \\ &= \frac{3}{2} Nk \ln T + Nk \ln V + \frac{3}{2} Nk \left[1 + \ln \frac{2\pi m k}{h_0^2} \right]. \end{aligned}$$
- Problem 1. S depends on h_0 ;

7.6 Entropy of the ideal gas

- Object: Monatomic molecules (translational motion).
- (7.1.13): $S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1)$.
- (7.5.21): $Z_1^t = V(\frac{2\pi m}{h_0^2 \beta})^{3/2}$.
- $$\begin{aligned} S &= Nk \left\{ \ln \left[V \left(\frac{2\pi m}{h_0^2 \beta} \right)^{3/2} \right] - \beta \frac{\partial}{\partial \beta} \ln \left[V \left(\frac{2\pi m}{h_0^2 \beta} \right)^{3/2} \right] \right\} \\ &= Nk \left\{ \ln V + \frac{3}{2} \ln \left(\frac{2\pi m}{h_0^2 \beta} \right) - \beta \frac{\partial}{\partial \beta} \left[\ln V + \frac{3}{2} \ln \left(\frac{2\pi m}{h_0^2 \beta} \right) \right] \right\} \\ &= Nk \left[\ln V + \frac{3}{2} \ln \left(\frac{2\pi m}{h_0^2 \beta} \right) - 0 - \frac{3}{2} \beta \cdot \left(-\frac{1}{\beta} \right) \right] \\ &= Nk \left[\ln V + \frac{3}{2} \ln \left(\frac{2\pi m}{h_0^2 \beta} \right) + \frac{3}{2} \right] \\ &= \frac{3}{2} Nk \ln T + Nk \ln V + \frac{3}{2} Nk \left[1 + \ln \frac{2\pi m k}{h_0^2} \right]. \end{aligned}$$
- Problem 1. S depends on h_0 ;
- Problem 2. S is not extensive.

Entropy of the ideal gas

- Using (7.1.13') instead:

$$S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) - k \ln N!$$

Entropy of the ideal gas

- Using (7.1.13') instead:

$$S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) - k \ln N!$$

Reason: Monatomic molecules are identical particles.

Entropy of the ideal gas

- Using (7.1.13') instead:

$$S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) - k \ln N!$$

Reason: Monatomic molecules are identical particles.

$$S = k \ln \Omega_{\text{M.B.}} \rightarrow S = k \ln \frac{\Omega_{\text{M.B.}}}{N!} = k \ln \Omega_{\text{M.B.}} - k \ln N!.$$

Entropy of the ideal gas

- Using (7.1.13') instead:

$$S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) - k \ln N!$$

Reason: Monatomic molecules are identical particles.

$$S = k \ln \Omega_{\text{M.B.}} \rightarrow S = k \ln \frac{\Omega_{\text{M.B.}}}{N!} = k \ln \Omega_{\text{M.B.}} - k \ln N!.$$

- $$S = \frac{3}{2}Nk \ln T + Nk \ln V + \frac{3}{2}Nk \left[1 + \ln \frac{2\pi mk}{h^2} \right] - k \ln N!$$

Entropy of the ideal gas

- Using (7.1.13') instead:

$$S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) - k \ln N!$$

Reason: Monatomic molecules are identical particles.

$$S = k \ln \Omega_{\text{M.B.}} \rightarrow S = k \ln \frac{\Omega_{\text{M.B.}}}{N!} = k \ln \Omega_{\text{M.B.}} - k \ln N!.$$

$$\begin{aligned} \bullet S &= \frac{3}{2}Nk \ln T + Nk \ln V + \frac{3}{2}Nk \left[1 + \ln \frac{2\pi mk}{h^2} \right] - k \ln N! \\ &= \frac{3}{2}Nk \ln T + Nk \ln V + \frac{3}{2}Nk \left[1 + \ln \frac{2\pi mk}{h^2} \right] - kN(\ln N - 1) \end{aligned}$$

Entropy of the ideal gas

- Using (7.1.13') instead:

$$S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) - k \ln N!$$

Reason: Monatomic molecules are identical particles.

$$S = k \ln \Omega_{\text{M.B.}} \rightarrow S = k \ln \frac{\Omega_{\text{M.B.}}}{N!} = k \ln \Omega_{\text{M.B.}} - k \ln N!.$$

$$\begin{aligned} \bullet S &= \frac{3}{2}Nk \ln T + Nk \ln V + \frac{3}{2}Nk \left[1 + \ln \frac{2\pi mk}{h^2} \right] - k \ln N! \\ &= \frac{3}{2}Nk \ln T + Nk \ln V + \frac{3}{2}Nk \left[1 + \ln \frac{2\pi mk}{h^2} \right] - kN(\ln N - 1) \\ &= \frac{3}{2}Nk \ln T + Nk \ln \frac{V}{N} + \frac{3}{2}Nk \left[\frac{5}{2} + \ln \frac{2\pi mk}{h^2} \right]. \end{aligned}$$

Entropy of the ideal gas

- Using (7.1.13') instead:

$$S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) - k \ln N!$$

Reason: Monatomic molecules are identical particles.

$$S = k \ln \Omega_{\text{M.B.}} \rightarrow S = k \ln \frac{\Omega_{\text{M.B.}}}{N!} = k \ln \Omega_{\text{M.B.}} - k \ln N!.$$

$$\begin{aligned} \bullet S &= \frac{3}{2}Nk \ln T + Nk \ln V + \frac{3}{2}Nk \left[1 + \ln \frac{2\pi mk}{h^2} \right] - k \ln N! \\ &= \frac{3}{2}Nk \ln T + Nk \ln V + \frac{3}{2}Nk \left[1 + \ln \frac{2\pi mk}{h^2} \right] - kN(\ln N - 1) \\ &= \frac{3}{2}Nk \ln T + Nk \ln \frac{V}{N} + \frac{3}{2}Nk \left[\frac{5}{2} + \ln \frac{2\pi mk}{h^2} \right]. \end{aligned}$$

- Not depend on h_0 , extensive.

Entropy of the ideal gas

- Compare with the experiment.

Entropy of the ideal gas

- Compare with the experiment.
- Object: gas and condensed phase in equilibrium.

Entropy of the ideal gas

- Compare with the experiment.
- Object: gas and condensed phase in equilibrium.
- For gas, $pV = NkT$,

Entropy of the ideal gas

- Compare with the experiment.
- Object: gas and condensed phase in equilibrium.
- For gas, $pV = NkT$,
$$\frac{S_{\text{gas}}}{Nk} = \frac{3}{2} \ln T + \ln \frac{V}{N} + \frac{3}{2} \left[\frac{5}{2} + \ln \frac{2\pi mk}{h^2} \right]$$

Entropy of the ideal gas

- Compare with the experiment.
- Object: gas and condensed phase in equilibrium.

- For gas, $pV = NkT$,

$$\begin{aligned}\frac{S_{\text{gas}}}{Nk} &= \frac{3}{2} \ln T + \ln \frac{V}{N} + \frac{3}{2} \left[\frac{5}{2} + \ln \frac{2\pi mk}{h^2} \right] \\ &= \frac{5}{2} \ln T + \ln \frac{V}{NkT} + \ln k + \frac{3}{2} \left[\frac{5}{2} + \ln \frac{2\pi mk}{h^2} \right]\end{aligned}$$

Entropy of the ideal gas

- Compare with the experiment.
- Object: gas and condensed phase in equilibrium.
- For gas, $pV = NkT$,

$$\begin{aligned}\frac{S_{\text{gas}}}{Nk} &= \frac{3}{2} \ln T + \ln \frac{V}{N} + \frac{3}{2} \left[\frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] \\ &= \frac{5}{2} \ln T + \ln \frac{V}{NkT} + \ln k + \frac{3}{2} \left[\frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] \\ &= \frac{5}{2} \ln T + \ln \frac{1}{p} + \ln k + \frac{3}{2} \left[\frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right]\end{aligned}$$

Entropy of the ideal gas

- Compare with the experiment.
- Object: gas and condensed phase in equilibrium.

- For gas, $pV = NkT$,

$$\begin{aligned}\frac{S_{\text{gas}}}{Nk} &= \frac{3}{2} \ln T + \ln \frac{V}{N} + \frac{3}{2} \left[\frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] \\ &= \frac{5}{2} \ln T + \ln \frac{V}{NkT} + \ln k + \frac{3}{2} \left[\frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] \\ &= \frac{5}{2} \ln T + \ln \frac{1}{p} + \ln k + \frac{3}{2} \left[\frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right]\end{aligned}$$

- i.e., $\ln p = \frac{5}{2} \ln T + \ln k + \frac{3}{2} \left[\frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] - \frac{S_{\text{gas}}}{Nk}$

Entropy of the ideal gas

- Compare with the experiment.
- Object: gas and condensed phase in equilibrium.

- For gas, $pV = NkT$,

$$\begin{aligned}\frac{S_{\text{gas}}}{Nk} &= \frac{3}{2} \ln T + \ln \frac{V}{N} + \frac{3}{2} \left[\frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] \\ &= \frac{5}{2} \ln T + \ln \frac{V}{NkT} + \ln k + \frac{3}{2} \left[\frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] \\ &= \frac{5}{2} \ln T + \ln \frac{1}{p} + \ln k + \frac{3}{2} \left[\frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right]\end{aligned}$$

- i.e., $\ln p = \frac{5}{2} \ln T + \ln k + \frac{3}{2} \left[\frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] - \frac{S_{\text{gas}}}{Nk}$

- $dS = \frac{\delta Q}{T}$, \rightarrow Latent $L = T(S_{\text{gas}} - S_{\text{con}})$.

Entropy of the ideal gas

- Compare with the experiment.
- Object: gas and condensed phase in equilibrium.

- For gas, $pV = NkT$,

$$\begin{aligned}\frac{S_{\text{gas}}}{Nk} &= \frac{3}{2} \ln T + \ln \frac{V}{N} + \frac{3}{2} \left[\frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] \\ &= \frac{5}{2} \ln T + \ln \frac{V}{NkT} + \ln k + \frac{3}{2} \left[\frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] \\ &= \frac{5}{2} \ln T + \ln \frac{1}{p} + \ln k + \frac{3}{2} \left[\frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right]\end{aligned}$$

- i.e., $\ln p = \frac{5}{2} \ln T + \ln k + \frac{3}{2} \left[\frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] - \frac{S_{\text{gas}}}{Nk}$

- $dS = \frac{\delta Q}{T}$, \rightarrow Latent $L = T(S_{\text{gas}} - S_{\text{con}})$.

- If T is low enough, $S_{\text{gas}} \gg S_{\text{con}}$,

Entropy of the ideal gas

- Compare with the experiment.
- Object: gas and condensed phase in equilibrium.
- For gas, $pV = NkT$,

$$\begin{aligned}\frac{S_{\text{gas}}}{Nk} &= \frac{3}{2} \ln T + \ln \frac{V}{N} + \frac{3}{2} \left[\frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] \\ &= \frac{5}{2} \ln T + \ln \frac{V}{NkT} + \ln k + \frac{3}{2} \left[\frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] \\ &= \frac{5}{2} \ln T + \ln \frac{1}{p} + \ln k + \frac{3}{2} \left[\frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right]\end{aligned}$$
- I.e., $\ln p = \frac{5}{2} \ln T + \ln k + \frac{3}{2} \left[\frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] - \frac{S_{\text{gas}}}{Nk}$
- $dS = \frac{\delta Q}{T}$, \rightarrow Latent $L = T(S_{\text{gas}} - S_{\text{con}})$.
- If T is low enough, $S_{\text{gas}} \gg S_{\text{con}}$, $L = TS_{\text{gas}}$.

Entropy of the ideal gas

- Compare with the experiment.
- Object: gas and condensed phase in equilibrium.
- For gas, $pV = NkT$,

$$\begin{aligned}\frac{S_{\text{gas}}}{Nk} &= \frac{3}{2} \ln T + \ln \frac{V}{N} + \frac{3}{2} \left[\frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] \\ &= \frac{5}{2} \ln T + \ln \frac{V}{NkT} + \ln k + \frac{3}{2} \left[\frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] \\ &= \frac{5}{2} \ln T + \ln \frac{1}{p} + \ln k + \frac{3}{2} \left[\frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right]\end{aligned}$$
- I.e., $\ln p = \frac{5}{2} \ln T + \ln k + \frac{3}{2} \left[\frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] - \frac{S_{\text{gas}}}{Nk}$
- $dS = \frac{\delta Q}{T}$, \rightarrow Latent $L = T(S_{\text{gas}} - S_{\text{con}})$.
- If T is low enough, $S_{\text{gas}} \gg S_{\text{con}}$, $L = TS_{\text{gas}}$. Then
 $\ln p = \frac{5}{2} \ln T + \ln k + \frac{3}{2} \left[\frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] - \frac{L}{RT}$, comparable.

Chemical potential of the monatomic ideal gas

Chemical potential of the monatomic ideal gas

- $dF = -SdT - pdV + \mu dN, \quad (3.2.9)$

Chemical potential of the monatomic ideal gas

- $$dF = -SdT - pdV + \mu dN, \quad (3.2.9)$$
$$\mu = \left(\frac{\partial F}{\partial N}\right)_{T,V}.$$

Chemical potential of the monatomic ideal gas

- $dF = -SdT - pdV + \mu dN$, (3.2.9)
 $\mu = \left(\frac{\partial F}{\partial N}\right)_{T,V}.$
- (7.1.16'),
 $F = -NkT \ln Z_1 + kT \ln N!$

Chemical potential of the monatomic ideal gas

- $dF = -SdT - pdV + \mu dN, \quad (3.2.9)$

$$\mu = \left(\frac{\partial F}{\partial N}\right)_{T,V}.$$

- (7.1.16'),

$$F = -NkT \ln Z_1 + kT \ln N!$$

$$= -NkT \ln Z_1 + kTN(\ln N - 1)$$

Chemical potential of the monatomic ideal gas

- $dF = -SdT - pdV + \mu dN, \quad (3.2.9)$

$$\mu = \left(\frac{\partial F}{\partial N}\right)_{T,V}.$$

- (7.1.16'),

$$F = -NkT \ln Z_1 + kT \ln N!$$

$$= -NkT \ln Z_1 + kTN(\ln N - 1)$$

$$= NkT \ln \frac{N}{Z_1} - NkT.$$

Chemical potential of the monatomic ideal gas

- $dF = -SdT - pdV + \mu dN, \quad (3.2.9)$

$$\mu = \left(\frac{\partial F}{\partial N}\right)_{T,V}.$$

- (7.1.16'),

$$F = -NkT \ln Z_1 + kT \ln N!$$

$$= -NkT \ln Z_1 + kTN(\ln N - 1)$$

$$= NkT \ln \frac{N}{Z_1} - NkT.$$

$$(Z_1^t = V(\frac{2\pi m}{h^2\beta})^{3/2} \quad (7.2.4))$$

Chemical potential of the monatomic ideal gas

- $dF = -SdT - pdV + \mu dN, \quad (3.2.9)$

$$\mu = \left(\frac{\partial F}{\partial N}\right)_{T,V}.$$

- (7.1.16'),

$$F = -NkT \ln Z_1 + kT \ln N!$$

$$= -NkT \ln Z_1 + kTN(\ln N - 1)$$

$$= NkT \ln \frac{N}{Z_1} - NkT.$$

$$(Z_1^t = V(\frac{2\pi m}{h^2\beta})^{3/2} \quad (7.2.4))$$

- $\mu = \left(\frac{\partial F}{\partial N}\right)_{T,V}$

Chemical potential of the monatomic ideal gas

- $dF = -SdT - pdV + \mu dN, \quad (3.2.9)$

$$\mu = \left(\frac{\partial F}{\partial N}\right)_{T,V}.$$

- (7.1.16'),

$$F = -NkT \ln Z_1 + kT \ln N!$$

$$= -NkT \ln Z_1 + kTN(\ln N - 1)$$

$$= NkT \ln \frac{N}{Z_1} - NkT.$$

$$(Z_1^t = V(\frac{2\pi m}{h^2\beta})^{3/2} \quad (7.2.4))$$

- $\mu = \left(\frac{\partial F}{\partial N}\right)_{T,V} = kT \ln \frac{N}{Z_1} + NkT \frac{Z_1}{N} \frac{1}{Z_1} - kT$

Chemical potential of the monatomic ideal gas

- $dF = -SdT - pdV + \mu dN, \quad (3.2.9)$

$$\mu = \left(\frac{\partial F}{\partial N}\right)_{T,V}.$$

- (7.1.16'),

$$F = -NkT \ln Z_1 + kT \ln N!$$

$$= -NkT \ln Z_1 + kTN(\ln N - 1)$$

$$= NkT \ln \frac{N}{Z_1} - NkT.$$

$$(Z_1^t = V(\frac{2\pi m}{h^2\beta})^{3/2} \quad (7.2.4))$$

- $\mu = \left(\frac{\partial F}{\partial N}\right)_{T,V} = kT \ln \frac{N}{Z_1} + NkT \frac{Z_1}{N} \frac{1}{Z_1} - kT$
 $= kT \ln \frac{N}{Z_1}$

Chemical potential of the monatomic ideal gas

- $dF = -SdT - pdV + \mu dN, \quad (3.2.9)$

$$\mu = \left(\frac{\partial F}{\partial N}\right)_{T,V}.$$

- (7.1.16'),

$$F = -NkT \ln Z_1 + kT \ln N!$$

$$= -NkT \ln Z_1 + kTN(\ln N - 1)$$

$$= NkT \ln \frac{N}{Z_1} - NkT.$$

$$(Z_1^t = V(\frac{2\pi m}{h^2\beta})^{3/2} \quad (7.2.4))$$

- $\mu = \left(\frac{\partial F}{\partial N}\right)_{T,V} = kT \ln \frac{N}{Z_1} + NkT \frac{Z_1}{N} \frac{1}{Z_1} - kT$

$$= kT \ln \frac{N}{Z_1}$$

$$= kT \ln \left[\frac{N}{V} \left(\frac{h^2}{2\pi m kT} \right)^{3/2} \right].$$

Table of contents

1 §7 Boltzmann Statistics

- 7.1 Thermal quantities in statistics
- 7.2 Equation of state of ideal gas
- 7.3 Maxwell speed distribution
- 7.4 Energy equipartition theorem
- 7.5 Internal energy and heat capacity of the ideal gas
- 7.6 Entropy of the ideal gas
- 7.7 Einstein's theory on heat capacity of solid
- 7.8 Paramagnetic solid
- 7.9 Negative temperature

7.7 Einstein's theory on heat capacity of solid

7.7 Einstein's theory on heat capacity of solid

- Energy equipartition theorem: $U = 3NkT$,
 $C_V = 3Nk$.

7.7 Einstein's theory on heat capacity of solid

- Energy equipartition theorem: $U = 3NkT$,
 $C_V = 3Nk$.
Only suitable for high temperature.

7.7 Einstein's theory on heat capacity of solid

- Energy equipartition theorem: $U = 3NkT$,
 $C_V = 3Nk$.
Only suitable for high temperature.
- General: taken the thermal motion as $3N$ harmonic oscillators.

7.7 Einstein's theory on heat capacity of solid

- Energy equipartition theorem: $U = 3NkT$,
 $C_V = 3Nk$.
Only suitable for high temperature.
- General: taken the thermal motion as $3N$ harmonic oscillators.
- Energy level: $\varepsilon_n = (n + \frac{1}{2})\hbar\omega$, $n = 0, 1, 2, \dots$

7.7 Einstein's theory on heat capacity of solid

- Energy equipartition theorem: $U = 3NkT$,
 $C_V = 3Nk$.
Only suitable for high temperature.
- General: taken the thermal motion as $3N$ harmonic oscillators.
- Energy level: $\varepsilon_n = (n + \frac{1}{2})\hbar\omega$, $n = 0, 1, 2, \dots$
- Local, distinguishable, semi-independent \rightarrow Boltzmann distribution:

7.7 Einstein's theory on heat capacity of solid

- Energy equipartition theorem: $U = 3NkT$,
 $C_V = 3Nk$.
Only suitable for high temperature.
- General: taken the thermal motion as $3N$ harmonic oscillators.
- Energy level: $\varepsilon_n = (n + \frac{1}{2})\hbar\omega$, $n = 0, 1, 2, \dots$
- Local, distinguishable, semi-independent \rightarrow Boltzmann distribution:

$$Z_1 = \sum_{n=0}^{\infty} \omega_l e^{-\beta\hbar\omega(n+\frac{1}{2})}$$

7.7 Einstein's theory on heat capacity of solid

- Energy equipartition theorem: $U = 3NkT$,
 $C_V = 3Nk$.
Only suitable for high temperature.
- General: taken the thermal motion as $3N$ harmonic oscillators.
- Energy level: $\varepsilon_n = (n + \frac{1}{2})\hbar\omega$, $n = 0, 1, 2, \dots$
- Local, distinguishable, semi-independent \rightarrow Boltzmann distribution:

$$Z_1 = \sum_{n=0}^{\infty} \omega_l e^{-\beta\hbar\omega(n+\frac{1}{2})} = \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}.$$

Einstein's theory on heat capacity of solid

- Internal energy: $U = -3N \frac{\partial}{\partial \beta} \ln Z_1$

Einstein's theory on heat capacity of solid

- Internal energy: $U = -3N \frac{\partial}{\partial \beta} \ln Z_1$
 $= -3N \frac{\partial}{\partial \beta} \ln \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}$

Einstein's theory on heat capacity of solid

- Internal energy:
$$\begin{aligned} U &= -3N \frac{\partial}{\partial \beta} \ln Z_1 \\ &= -3N \frac{\partial}{\partial \beta} \ln \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1 - e^{-\beta\hbar\omega}} \\ &= -3N \frac{\partial}{\partial \beta} [\ln e^{-\frac{1}{2}\beta\hbar\omega} - \ln(1 - e^{-\beta\hbar\omega})] \end{aligned}$$

Einstein's theory on heat capacity of solid

- Internal energy:
$$\begin{aligned} U &= -3N \frac{\partial}{\partial \beta} \ln Z_1 \\ &= -3N \frac{\partial}{\partial \beta} \ln \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \\ &= -3N \frac{\partial}{\partial \beta} [\ln e^{-\frac{1}{2}\beta\hbar\omega} - \ln(1 - e^{-\beta\hbar\omega})] \\ &= 3N \left(\frac{1}{2}\hbar\omega + \frac{e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \hbar\omega \right) \end{aligned}$$

Einstein's theory on heat capacity of solid

- Internal energy:
$$\begin{aligned} U &= -3N \frac{\partial}{\partial \beta} \ln Z_1 \\ &= -3N \frac{\partial}{\partial \beta} \ln \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \\ &= -3N \frac{\partial}{\partial \beta} [\ln e^{-\frac{1}{2}\beta\hbar\omega} - \ln(1 - e^{-\beta\hbar\omega})] \\ &= 3N \left(\frac{1}{2}\hbar\omega + \frac{e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \hbar\omega \right) = \frac{3}{2}N\hbar\omega + \frac{3N\hbar\omega}{e^{\beta\hbar\omega} - 1}. \end{aligned}$$

Einstein's theory on heat capacity of solid

- Internal energy: $U = -3N \frac{\partial}{\partial \beta} \ln Z_1$
$$= -3N \frac{\partial}{\partial \beta} \ln \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1 - e^{-\beta\hbar\omega}}$$
$$= -3N \frac{\partial}{\partial \beta} [\ln e^{-\frac{1}{2}\beta\hbar\omega} - \ln(1 - e^{-\beta\hbar\omega})]$$
$$= 3N \left(\frac{1}{2}\hbar\omega + \frac{e^{-\beta\hbar\omega}}{1 - e^{-\beta\hbar\omega}} \hbar\omega \right) = \frac{3}{2}N\hbar\omega + \frac{3N\hbar\omega}{e^{\beta\hbar\omega} - 1}.$$
- Introduce typical temperature θ_E : $k\theta_E = \hbar\omega$.

Einstein's theory on heat capacity of solid

- Internal energy: $U = -3N \frac{\partial}{\partial \beta} \ln Z_1$
$$= -3N \frac{\partial}{\partial \beta} \ln \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1 - e^{-\beta\hbar\omega}}$$
$$= -3N \frac{\partial}{\partial \beta} [\ln e^{-\frac{1}{2}\beta\hbar\omega} - \ln(1 - e^{-\beta\hbar\omega})]$$
$$= 3N \left(\frac{1}{2}\hbar\omega + \frac{e^{-\beta\hbar\omega}}{1 - e^{-\beta\hbar\omega}} \hbar\omega \right) = \frac{3}{2}N\hbar\omega + \frac{3N\hbar\omega}{e^{\beta\hbar\omega} - 1}.$$
- Introduce typical temperature θ_E : $k\theta_E = \hbar\omega$.
$$U = \frac{3}{2}Nk\theta_E + \frac{3Nk\theta_E}{e^{\frac{\theta_E}{T}} - 1}.$$

Einstein's theory on heat capacity of solid

- Internal energy: $U = -3N \frac{\partial}{\partial \beta} \ln Z_1$
$$= -3N \frac{\partial}{\partial \beta} \ln \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1 - e^{-\beta\hbar\omega}}$$
$$= -3N \frac{\partial}{\partial \beta} [\ln e^{-\frac{1}{2}\beta\hbar\omega} - \ln(1 - e^{-\beta\hbar\omega})]$$
$$= 3N \left(\frac{1}{2}\hbar\omega + \frac{e^{-\beta\hbar\omega}}{1 - e^{-\beta\hbar\omega}} \hbar\omega \right) = \frac{3}{2}N\hbar\omega + \frac{3N\hbar\omega}{e^{\beta\hbar\omega} - 1}.$$
- Introduce typical temperature θ_E : $k\theta_E = \hbar\omega$.
$$U = \frac{3}{2}Nk\theta_E + \frac{3Nk\theta_E}{e^{\frac{\theta_E}{T}} - 1}.$$
$$C_V = \left(\frac{\partial U}{\partial T} \right)_V$$

Einstein's theory on heat capacity of solid

- Internal energy: $U = -3N \frac{\partial}{\partial \beta} \ln Z_1$

$$= -3N \frac{\partial}{\partial \beta} \ln \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}$$

$$= -3N \frac{\partial}{\partial \beta} [\ln e^{-\frac{1}{2}\beta\hbar\omega} - \ln(1 - e^{-\beta\hbar\omega})]$$

$$= 3N \left(\frac{1}{2}\hbar\omega + \frac{e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \hbar\omega \right) = \frac{3}{2}N\hbar\omega + \frac{3N\hbar\omega}{e^{\beta\hbar\omega}-1}.$$
- Introduce typical temperature θ_E : $k\theta_E = \hbar\omega$.

$$U = \frac{3}{2}Nk\theta_E + \frac{3Nk\theta_E}{e^{\frac{\theta_E}{T}}-1}.$$

$$C_V = \left(\frac{\partial U}{\partial T}\right)_V = 0 - \frac{3Nk\theta_E}{(e^{\frac{\theta_E}{T}}-1)^2} \cdot e^{\frac{\theta_E}{T}} \cdot \left(-\frac{\theta_E}{T^2}\right)$$

Einstein's theory on heat capacity of solid

- Internal energy: $U = -3N \frac{\partial}{\partial \beta} \ln Z_1$

$$= -3N \frac{\partial}{\partial \beta} \ln \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}$$

$$= -3N \frac{\partial}{\partial \beta} [\ln e^{-\frac{1}{2}\beta\hbar\omega} - \ln(1 - e^{-\beta\hbar\omega})]$$

$$= 3N \left(\frac{1}{2}\hbar\omega + \frac{e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \hbar\omega \right) = \frac{3}{2}N\hbar\omega + \frac{3N\hbar\omega}{e^{\beta\hbar\omega}-1}.$$
- Introduce typical temperature θ_E : $k\theta_E = \hbar\omega$.

$$U = \frac{3}{2}Nk\theta_E + \frac{3Nk\theta_E}{e^{\frac{\theta_E}{T}}-1}.$$

$$C_V = \left(\frac{\partial U}{\partial T}\right)_V = 0 - \frac{3Nk\theta_E}{(e^{\frac{\theta_E}{T}}-1)^2} \cdot e^{\frac{\theta_E}{T}} \cdot \left(-\frac{\theta_E}{T^2}\right)$$

$$= 3Nk\left(\frac{\theta_E}{T}\right)^2 \frac{e^{\frac{\theta_E}{T}}}{(e^{\frac{\theta_E}{T}}-1)^2}.$$

Einstein's theory on heat capacity of solid

- $$C_V = 3Nk\left(\frac{\theta_E}{T}\right)^2 \frac{e^{\frac{\theta_E}{T}}}{(e^{\frac{\theta_E}{T}} - 1)^2},$$

Einstein's theory on heat capacity of solid

- $$C_V = 3Nk \left(\frac{\theta_E}{T} \right)^2 \frac{e^{-\frac{\theta_E}{T}}}{(e^{-\frac{\theta_E}{T}} - 1)^2},$$
$$\theta_E \sim 1320\text{K for diamond.}$$

Einstein's theory on heat capacity of solid

- $C_V = 3Nk\left(\frac{\theta_E}{T}\right)^2 \frac{e^{\frac{\theta_E}{T}}}{(e^{\frac{\theta_E}{T}} - 1)^2},$
 $\theta_E \sim 1320\text{K}$ for diamond.
- High temperature: $T \gg \theta_E$, $e^{\frac{\theta_E}{T}} \simeq 1 + \frac{\theta_E}{T}.$

Einstein's theory on heat capacity of solid

- $C_V = 3Nk \left(\frac{\theta_E}{T}\right)^2 \frac{e^{\frac{\theta_E}{T}}}{(e^{\frac{\theta_E}{T}} - 1)^2},$
 $\theta_E \sim 1320\text{K}$ for diamond.
- High temperature: $T \gg \theta_E$, $e^{\frac{\theta_E}{T}} \simeq 1 + \frac{\theta_E}{T}.$
 $C_V \simeq 3Nk$, equipartition theorem.

Einstein's theory on heat capacity of solid

- $C_V = 3Nk \left(\frac{\theta_E}{T} \right)^2 \frac{e^{\frac{\theta_E}{T}}}{(e^{\frac{\theta_E}{T}} - 1)^2},$
 $\theta_E \sim 1320\text{K}$ for diamond.
- High temperature: $T \gg \theta_E$, $e^{\frac{\theta_E}{T}} \simeq 1 + \frac{\theta_E}{T}$.
 $C_V \simeq 3Nk$, equipartition theorem.
- Low temperature: $T \ll \theta_E$, $e^{\frac{\theta_E}{T}} - 1 \simeq e^{\frac{\theta_E}{T}}$.

Einstein's theory on heat capacity of solid

- $C_V = 3Nk \left(\frac{\theta_E}{T}\right)^2 \frac{e^{\frac{\theta_E}{T}}}{(e^{\frac{\theta_E}{T}} - 1)^2}$,
 $\theta_E \sim 1320\text{K}$ for diamond.
- High temperature: $T \gg \theta_E$, $e^{\frac{\theta_E}{T}} \simeq 1 + \frac{\theta_E}{T}$.
 $C_V \simeq 3Nk$, equipartition theorem.
- Low temperature: $T \ll \theta_E$, $e^{\frac{\theta_E}{T}} - 1 \simeq e^{\frac{\theta_E}{T}}$.
 $C_V \simeq 3Nk \left(\frac{\theta_E}{T}\right)^2 e^{-\frac{\theta_E}{T}} \rightarrow 0$.

Table of contents

1 §7 Boltzmann Statistics

- 7.1 Thermal quantities in statistics
- 7.2 Equation of state of ideal gas
- 7.3 Maxwell speed distribution
- 7.4 Energy equipartition theorem
- 7.5 Internal energy and heat capacity of the ideal gas
- 7.6 Entropy of the ideal gas
- 7.7 Einstein's theory on heat capacity of solid
- **7.8 Paramagnetic solid**
- 7.9 Negative temperature

7.8 Paramagnetic solid

7.8 Paramagnetic solid

- Object: paramagnetic particles fixed on the crystal lattice.

7.8 Paramagnetic solid

- Object: paramagnetic particles fixed on the crystal lattice.
- Simplification: Magnetic quantum number $\frac{1}{2}$.

7.8 Paramagnetic solid

- Object: paramagnetic particles fixed on the crystal lattice.
- Simplification: Magnetic quantum number $\frac{1}{2}$.
- In the external magnetic field B , only two choices of quantum number $\pm\frac{1}{2}$ (parallel or anti-parallel to B).

7.8 Paramagnetic solid

- Object: paramagnetic particles fixed on the crystal lattice.
- Simplification: Magnetic quantum number $\frac{1}{2}$.
- In the external magnetic field B , only two choices of quantum number $\pm\frac{1}{2}$ (parallel or anti-parallel to B).
- Energy level: $\pm\mu B$. (μ is the magnetic momentum.)

7.8 Paramagnetic solid

- Object: paramagnetic particles fixed on the crystal lattice.
- Simplification: Magnetic quantum number $\frac{1}{2}$.
- In the external magnetic field B , only two choices of quantum number $\pm\frac{1}{2}$ (parallel or anti-parallel to B).
- Energy level: $\pm\mu B$. (μ is the magnetic momentum.)
- Partition function: $Z_1 = \sum \omega_l e^{-\beta \epsilon_l}$

7.8 Paramagnetic solid

- Object: paramagnetic particles fixed on the crystal lattice.
- Simplification: Magnetic quantum number $\frac{1}{2}$.
- In the external magnetic field B , only two choices of quantum number $\pm\frac{1}{2}$ (parallel or anti-parallel to B).
- Energy level: $\pm\mu B$. (μ is the magnetic momentum.)
- Partition function: $Z_1 = \sum \omega_l e^{-\beta\epsilon_l} = e^{\beta\mu B} + e^{-\beta\mu B}$.

7.8 Paramagnetic solid

- Object: paramagnetic particles fixed on the crystal lattice.
- Simplification: Magnetic quantum number $\frac{1}{2}$.
- In the external magnetic field B , only two choices of quantum number $\pm\frac{1}{2}$ (parallel or anti-parallel to B).
- Energy level: $\pm\mu B$. (μ is the magnetic momentum.)
- Partition function: $Z_1 = \sum \omega_l e^{-\beta \epsilon_l} = e^{\beta \mu B} + e^{-\beta \mu B}$.
- Remind eq.(2.7.19), $\text{d}W = -\mu_0 m \text{d}H$.

7.8 Paramagnetic solid

- Object: paramagnetic particles fixed on the crystal lattice.
- Simplification: Magnetic quantum number $\frac{1}{2}$.
- In the external magnetic field B , only two choices of quantum number $\pm\frac{1}{2}$ (parallel or anti-parallel to B).
- Energy level: $\pm\mu B$. (μ is the magnetic momentum.)
- Partition function: $Z_1 = \sum \omega_l e^{-\beta \epsilon_l} = e^{\beta \mu B} + e^{-\beta \mu B}$.
- Remind eq.(2.7.19), $\text{d}W = -\mu_0 m \text{d}H$. ($-\mu_0 m$ generalized force, H generalized displacement.)

7.8 Paramagnetic solid

- Object: paramagnetic particles fixed on the crystal lattice.
- Simplification: Magnetic quantum number $\frac{1}{2}$.
- In the external magnetic field B , only two choices of quantum number $\pm\frac{1}{2}$ (parallel or anti-parallel to B).
- Energy level: $\pm\mu B$. (μ is the magnetic momentum.)
- Partition function: $Z_1 = \sum \omega_l e^{-\beta \epsilon_l} = e^{\beta \mu B} + e^{-\beta \mu B}$.
- Remind eq.(2.7.19), $\text{d}W = -\mu_0 m \text{d}H$. ($-\mu_0 m$ generalized force, H generalized displacement.)
eq(7.1.6), $Y = -\frac{N}{\beta} \frac{\partial}{\partial y} \ln Z_1$.

7.8 Paramagnetic solid

- Object: paramagnetic particles fixed on the crystal lattice.
- Simplification: Magnetic quantum number $\frac{1}{2}$.
- In the external magnetic field B , only two choices of quantum number $\pm\frac{1}{2}$ (parallel or anti-parallel to B).
- Energy level: $\pm\mu B$. (μ is the magnetic momentum.)
- Partition function: $Z_1 = \sum \omega_l e^{-\beta \epsilon_l} = e^{\beta \mu B} + e^{-\beta \mu B}$.
- Remind eq.(2.7.19), $\text{d}W = -\mu_0 m \text{d}H$. ($-\mu_0 m$ generalized force, H generalized displacement.)
eq(7.1.6), $Y = -\frac{N}{\beta} \frac{\partial}{\partial y} \ln Z_1$.
 $\therefore -\mu_0 m = -\frac{N}{\beta} \frac{\partial}{\partial H} \ln Z_1$.

Paramagnetic solid

- $-\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1,$

Paramagnetic solid

- $$-\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1,$$
$$\Rightarrow M = \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1$$

Paramagnetic solid

- $-\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1,$
 $\Rightarrow M = \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B})$

Paramagnetic solid

- $$\begin{aligned} -\mu_0 \frac{m}{V} &= -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1, \\ \Rightarrow M &= \frac{n}{\beta} \frac{\partial}{\partial(\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B}) \\ &= \frac{n}{\beta} \frac{\beta\mu e^{\beta\mu B} - \beta\mu e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} \end{aligned}$$

Paramagnetic solid

- $$\begin{aligned} -\mu_0 \frac{m}{V} &= -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1, \\ \Rightarrow M &= \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B}) \\ &= \frac{n}{\beta} \frac{\beta\mu e^{\beta\mu B} - \beta\mu e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = n\mu \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} \end{aligned}$$

Paramagnetic solid

$$\begin{aligned}
 \bullet \quad & -\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1, \\
 \Rightarrow \quad & M = \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B}) \\
 & = \frac{n}{\beta} \frac{\beta\mu e^{\beta\mu B} - \beta\mu e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = n\mu \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} = n\mu \frac{e^{\frac{2\mu B}{kT}} - 1}{e^{\frac{2\mu B}{kT}} + 1}.
 \end{aligned}$$

Paramagnetic solid

- $$-\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1,$$

$$\Rightarrow M = \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B})$$

$$= \frac{n}{\beta} \frac{\beta\mu e^{\beta\mu B} - \beta\mu e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = n\mu \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} = n\mu \frac{e^{\frac{2\mu B}{kT}} - 1}{e^{\frac{2\mu B}{kT}} + 1}.$$
- $$\frac{\mu B}{kT} \ll 1,$$

Paramagnetic solid

- $$-\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1,$$

$$\Rightarrow M = \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B})$$

$$= \frac{n}{\beta} \frac{\beta\mu e^{\beta\mu B} - \beta\mu e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = n\mu \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} = n\mu \frac{e^{\frac{2\mu B}{kT}} - 1}{e^{\frac{2\mu B}{kT}} + 1}.$$
- $$\frac{\mu B}{kT} \ll 1, \quad M \simeq n\mu \frac{\frac{2\mu B}{kT}}{2}$$

Paramagnetic solid

- $$-\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1,$$

$$\Rightarrow M = \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B})$$

$$= \frac{n}{\beta} \frac{\beta\mu e^{\beta\mu B} - \beta\mu e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = n\mu \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} = n\mu \frac{e^{\frac{2\mu B}{kT}} - 1}{e^{\frac{2\mu B}{kT}} + 1}.$$
- $$\frac{\mu B}{kT} \ll 1, \quad M \simeq n\mu \frac{\frac{2\mu B}{kT}}{2} = \frac{n\mu^2}{kT} B, \text{ Curie's law.}$$

Paramagnetic solid

- $$-\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1,$$

$$\Rightarrow M = \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B})$$

$$= \frac{n}{\beta} \frac{\beta\mu e^{\beta\mu B} - \beta\mu e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = n\mu \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} = n\mu \frac{e^{\frac{2\mu B}{kT}} - 1}{e^{\frac{2\mu B}{kT}} + 1}.$$
- $$\frac{\mu B}{kT} \ll 1, \quad M \simeq n\mu \frac{\frac{2\mu B}{kT}}{2} = \frac{n\mu^2}{kT} B, \text{ Curie's law.}$$
- $$\frac{\mu B}{kT} \gg 1,$$

Paramagnetic solid

- $$-\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1,$$

$$\Rightarrow M = \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B})$$

$$= \frac{n}{\beta} \frac{\beta\mu e^{\beta\mu B} - \beta\mu e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = n\mu \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} = n\mu \frac{e^{\frac{2\mu B}{kT}} - 1}{e^{\frac{2\mu B}{kT}} + 1}.$$
- $$\frac{\mu B}{kT} \ll 1, M \simeq n\mu \frac{\frac{2\mu B}{kT}}{2} = \frac{n\mu^2}{kT} B, \text{ Curie's law.}$$
- $$\frac{\mu B}{kT} \gg 1, M \simeq n\mu, \text{ saturation.}$$

Paramagnetic solid

- $$-\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1,$$

$$\Rightarrow M = \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B})$$

$$= \frac{n}{\beta} \frac{\beta\mu e^{\beta\mu B} - \beta\mu e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = n\mu \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} = n\mu \frac{e^{\frac{2\mu B}{kT}} - 1}{e^{\frac{2\mu B}{kT}} + 1}.$$
- $$\frac{\mu B}{kT} \ll 1, M \simeq n\mu \frac{\frac{2\mu B}{kT}}{2} = \frac{n\mu^2}{kT} B, \text{ Curie's law.}$$
- $$\frac{\mu B}{kT} \gg 1, M \simeq n\mu, \text{ saturation.}$$
- Internal energy density: $u = -n \frac{\partial}{\partial \beta} \ln Z_1$

Paramagnetic solid

- $$-\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1,$$

$$\Rightarrow M = \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B})$$

$$= \frac{n}{\beta} \frac{\beta\mu e^{\beta\mu B} - \beta\mu e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = n\mu \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} = n\mu \frac{e^{\frac{2\mu B}{kT}} - 1}{e^{\frac{2\mu B}{kT}} + 1}.$$
- $$\frac{\mu B}{kT} \ll 1, M \simeq n\mu \frac{\frac{2\mu B}{kT}}{2} = \frac{n\mu^2}{kT} B, \text{ Curie's law.}$$
- $$\frac{\mu B}{kT} \gg 1, M \simeq n\mu, \text{ saturation.}$$
- Internal energy density: $u = -n \frac{\partial}{\partial \beta} \ln Z_1$

$$= -n \frac{\partial}{\partial \beta} \ln(e^{\beta\mu B} + e^{-\beta\mu B})$$

Paramagnetic solid

- $$-\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1,$$

$$\Rightarrow M = \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B})$$

$$= \frac{n}{\beta} \frac{\beta\mu e^{\beta\mu B} - \beta\mu e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = n\mu \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} = n\mu \frac{e^{\frac{2\mu B}{kT}} - 1}{e^{\frac{2\mu B}{kT}} + 1}.$$
- $$\frac{\mu B}{kT} \ll 1, M \simeq n\mu \frac{\frac{2\mu B}{kT}}{2} = \frac{n\mu^2}{kT} B, \text{ Curie's law.}$$
- $$\frac{\mu B}{kT} \gg 1, M \simeq n\mu, \text{ saturation.}$$
- Internal energy density: $u = -n \frac{\partial}{\partial \beta} \ln Z_1$

$$= -n \frac{\partial}{\partial \beta} \ln(e^{\beta\mu B} + e^{-\beta\mu B}) = -n \frac{\mu B e^{\beta\mu B} - \mu B e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}}$$

Paramagnetic solid

- $$-\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1,$$

$$\Rightarrow M = \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B})$$

$$= \frac{n}{\beta} \frac{\beta\mu e^{\beta\mu B} - \beta\mu e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = n\mu \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} = n\mu \frac{e^{\frac{2\mu B}{kT}} - 1}{e^{\frac{2\mu B}{kT}} + 1}.$$
- $$\frac{\mu B}{kT} \ll 1, M \simeq n\mu \frac{\frac{2\mu B}{kT}}{2} = \frac{n\mu^2}{kT} B, \text{ Curie's law.}$$
- $$\frac{\mu B}{kT} \gg 1, M \simeq n\mu, \text{ saturation.}$$
- Internal energy density: $u = -n \frac{\partial}{\partial \beta} \ln Z_1$

$$= -n \frac{\partial}{\partial \beta} \ln(e^{\beta\mu B} + e^{-\beta\mu B}) = -n \frac{\mu B e^{\beta\mu B} - \mu B e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}}$$

$$= -n\mu B \frac{e^{\beta\mu B} - e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}}$$

Paramagnetic solid

- $$-\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1,$$

$$\Rightarrow M = \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B})$$

$$= \frac{n}{\beta} \frac{\beta\mu e^{\beta\mu B} - \beta\mu e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = n\mu \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} = n\mu \frac{e^{\frac{2\mu B}{kT}} - 1}{e^{\frac{2\mu B}{kT}} + 1}.$$
- $$\frac{\mu B}{kT} \ll 1, M \simeq n\mu \frac{\frac{2\mu B}{kT}}{2} = \frac{n\mu^2}{kT} B, \text{ Curie's law.}$$
- $$\frac{\mu B}{kT} \gg 1, M \simeq n\mu, \text{ saturation.}$$
- Internal energy density: $u = -n \frac{\partial}{\partial \beta} \ln Z_1$

$$= -n \frac{\partial}{\partial \beta} \ln(e^{\beta\mu B} + e^{-\beta\mu B}) = -n \frac{\mu B e^{\beta\mu B} - \mu B e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}}$$

$$= -n\mu B \frac{e^{\beta\mu B} - e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = -MB,$$

Paramagnetic solid

- $$-\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1,$$

$$\Rightarrow M = \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B})$$

$$= \frac{n}{\beta} \frac{\beta\mu e^{\beta\mu B} - \beta\mu e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = n\mu \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} = n\mu \frac{e^{\frac{2\mu B}{kT}} - 1}{e^{\frac{2\mu B}{kT}} + 1}.$$
- $$\frac{\mu B}{kT} \ll 1, M \simeq n\mu \frac{\frac{2\mu B}{kT}}{2} = \frac{n\mu^2}{kT} B, \text{ Curie's law.}$$
- $$\frac{\mu B}{kT} \gg 1, M \simeq n\mu, \text{ saturation.}$$
- Internal energy density: $u = -n \frac{\partial}{\partial \beta} \ln Z_1$

$$= -n \frac{\partial}{\partial \beta} \ln(e^{\beta\mu B} + e^{-\beta\mu B}) = -n \frac{\mu B e^{\beta\mu B} - \mu B e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}}$$

$$= -n\mu B \frac{e^{\beta\mu B} - e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = -MB,$$

potential energy in the external field.

Paramagnetic solid

- Entropy density eq.(7.1.13):

$$s = nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1)$$

Paramagnetic solid

- Entropy density eq.(7.1.13):

$$\begin{aligned}s &= nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) \\&= nk[\ln(e^{\beta\mu B} + e^{-\beta\mu B}) - \beta\mu B \frac{e^{\beta\mu B} - e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}}]\end{aligned}$$

Paramagnetic solid

- Entropy density eq.(7.1.13):

$$\begin{aligned} s &= nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) \\ &= nk[\ln(e^{\beta\mu B} + e^{-\beta\mu B}) - \beta\mu B \frac{e^{\beta\mu B} - e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}}] \\ &= nk \left[\ln(e^{2\beta\mu B} + 1) - \beta\mu B - \beta\mu B \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} \right] \end{aligned}$$

Paramagnetic solid

- Entropy density eq.(7.1.13):

$$\begin{aligned} s &= nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) \\ &= nk[\ln(e^{\beta\mu B} + e^{-\beta\mu B}) - \beta\mu B \frac{e^{\beta\mu B} - e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}}] \\ &= nk \left[\ln(e^{2\beta\mu B} + 1) - \beta\mu B - \beta\mu B \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} \right] \\ &= nk \left[\ln(e^{2\beta\mu B} + 1) - \beta\mu B \frac{2e^{2\beta\mu B}}{e^{2\beta\mu B} + 1} \right] \end{aligned}$$

Paramagnetic solid

- Entropy density eq.(7.1.13):

$$\begin{aligned} s &= nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) \\ &= nk[\ln(e^{\beta\mu B} + e^{-\beta\mu B}) - \beta\mu B \frac{e^{\beta\mu B} - e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}}] \\ &= nk \left[\ln(e^{2\beta\mu B} + 1) - \beta\mu B - \beta\mu B \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} \right] \\ &= nk \left[\ln(e^{2\beta\mu B} + 1) - \beta\mu B \frac{2e^{2\beta\mu B}}{e^{2\beta\mu B} + 1} \right] \\ &= nk \left[\ln(e^{\frac{2\mu B}{kT}} + 1) - \frac{\mu B}{kT} \frac{2e^{\frac{2\mu B}{kT}}}{e^{\frac{2\mu B}{kT}} + 1} \right]. \end{aligned}$$

Paramagnetic solid

- Entropy density eq.(7.1.13):

$$\begin{aligned}
 s &= nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) \\
 &= nk[\ln(e^{\beta\mu B} + e^{-\beta\mu B}) - \beta\mu B \frac{e^{\beta\mu B} - e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}}] \\
 &= nk \left[\ln(e^{2\beta\mu B} + 1) - \beta\mu B - \beta\mu B \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} \right] \\
 &= nk \left[\ln(e^{2\beta\mu B} + 1) - \beta\mu B \frac{2e^{2\beta\mu B}}{e^{2\beta\mu B} + 1} \right] \\
 &= nk \left[\ln(e^{\frac{2\mu B}{kT}} + 1) - \frac{\mu B}{kT} \frac{2e^{\frac{2\mu B}{kT}}}{e^{\frac{2\mu B}{kT}} + 1} \right].
 \end{aligned}$$

- $\frac{\mu B}{kT} \ll 1$, $s = nk \ln 2 = k \ln 2^n$,

Paramagnetic solid

- Entropy density eq.(7.1.13):

$$s = nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1)$$

$$= nk[\ln(e^{\beta\mu B} + e^{-\beta\mu B}) - \beta\mu B \frac{e^{\beta\mu B} - e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}}]$$

$$= nk \left[\ln(e^{2\beta\mu B} + 1) - \beta\mu B - \beta\mu B \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} \right]$$

$$= nk \left[\ln(e^{2\beta\mu B} + 1) - \beta\mu B \frac{2e^{2\beta\mu B}}{e^{2\beta\mu B} + 1} \right]$$

$$= nk \left[\ln\left(e^{\frac{2\mu B}{kT}} + 1\right) - \frac{\mu B}{kT} \frac{2e^{\frac{2\mu B}{kT}}}{e^{\frac{2\mu B}{kT}} + 1} \right].$$

- $\frac{\mu B}{kT} \ll 1$, $s = nk \ln 2 = k \ln 2^n$, high temperature, two states are both possible, total number of states: 2^n .

Paramagnetic solid

- Entropy density eq.(7.1.13):

$$s = nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1)$$

$$= nk[\ln(e^{\beta\mu B} + e^{-\beta\mu B}) - \beta\mu B \frac{e^{\beta\mu B} - e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}}]$$

$$= nk \left[\ln(e^{2\beta\mu B} + 1) - \beta\mu B - \beta\mu B \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} \right]$$

$$= nk \left[\ln(e^{2\beta\mu B} + 1) - \beta\mu B \frac{2e^{2\beta\mu B}}{e^{2\beta\mu B} + 1} \right]$$

$$= nk \left[\ln\left(e^{\frac{2\mu B}{kT}} + 1\right) - \frac{\mu B}{kT} \frac{2e^{\frac{2\mu B}{kT}}}{e^{\frac{2\mu B}{kT}} + 1} \right].$$

- $\frac{\mu B}{kT} \ll 1$, $s = nk \ln 2 = k \ln 2^n$, high temperature, two states are both possible, total number of states: 2^n .
- $\frac{\mu B}{kT} \gg 1$, $s \simeq 0$,

Paramagnetic solid

- Entropy density eq.(7.1.13):

$$s = nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1)$$

$$= nk[\ln(e^{\beta\mu B} + e^{-\beta\mu B}) - \beta\mu B \frac{e^{\beta\mu B} - e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}}]$$

$$= nk \left[\ln(e^{2\beta\mu B} + 1) - \beta\mu B - \beta\mu B \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} \right]$$

$$= nk \left[\ln(e^{2\beta\mu B} + 1) - \beta\mu B \frac{2e^{2\beta\mu B}}{e^{2\beta\mu B} + 1} \right]$$

$$= nk \left[\ln\left(e^{\frac{2\mu B}{kT}} + 1\right) - \frac{\mu B}{kT} \frac{2e^{\frac{2\mu B}{kT}}}{e^{\frac{2\mu B}{kT}} + 1} \right].$$

- $\frac{\mu B}{kT} \ll 1$, $s = nk \ln 2 = k \ln 2^n$, high temperature, two states are both possible, total number of states: 2^n .
- $\frac{\mu B}{kT} \gg 1$, $s \simeq 0$, only one micro-state: all magnetic moments are aligned along the external field.

Table of contents

1 §7 Boltzmann Statistics

- 7.1 Thermal quantities in statistics
- 7.2 Equation of state of ideal gas
- 7.3 Maxwell speed distribution
- 7.4 Energy equipartition theorem
- 7.5 Internal energy and heat capacity of the ideal gas
- 7.6 Entropy of the ideal gas
- 7.7 Einstein's theory on heat capacity of solid
- 7.8 Paramagnetic solid
- 7.9 Negative temperature

7.9 Negative temperature

7.9 Negative temperature

- Basic function of thermodynamics: $dU = TdS - pdV$.

7.9 Negative temperature

- Basic function of thermodynamics: $dU = TdS - pdV$.
More general: $dU = TdS + Ydy$,

7.9 Negative temperature

- Basic function of thermodynamics: $dU = TdS - pdV$.
More general: $dU = TdS + Ydy$,
 $\Rightarrow \left(\frac{\partial S}{\partial U}\right)_y = \frac{1}{T}$.

7.9 Negative temperature

- Basic function of thermodynamics: $dU = TdS - pdV$.

More general: $dU = TdS + Ydy$,

$$\Rightarrow \left(\frac{\partial S}{\partial U}\right)_y = \frac{1}{T}.$$

If U increases but S decreases (micro-states), T is negative.

7.9 Negative temperature

- Basic function of thermodynamics: $dU = TdS - pdV$.
More general: $dU = TdS + Ydy$,
 $\Rightarrow \left(\frac{\partial S}{\partial U}\right)_y = \frac{1}{T}$.
If U increases but S decreases (micro-states), T is negative.
- Example: isolated nuclear spin system.

7.9 Negative temperature

- Basic function of thermodynamics: $dU = TdS - pdV$.
More general: $dU = TdS + Ydy$,
 $\Rightarrow (\frac{\partial S}{\partial U})_y = \frac{1}{T}$.
If U increases but S decreases (micro-states), T is negative.
- Example: isolated nuclear spin system.
- In external magnetic field B , the energy
 $\pm\varepsilon = \pm\mu B = \pm\frac{e\hbar}{2m}B$.

7.9 Negative temperature

- Basic function of thermodynamics: $dU = TdS - pdV$.
More general: $dU = TdS + Ydy$,
 $\Rightarrow \left(\frac{\partial S}{\partial U}\right)_y = \frac{1}{T}$.
If U increases but S decreases (micro-states), T is negative.
- Example: isolated nuclear spin system.
- In external magnetic field B , the energy
 $\pm\varepsilon = \pm\mu B = \pm\frac{e\hbar}{2m}B$.
- Number of spins: N , where N_+ with ε , N_- with $-\varepsilon$.

7.9 Negative temperature

- Basic function of thermodynamics: $dU = TdS - pdV$.
More general: $dU = TdS + Ydy$,
 $\Rightarrow \left(\frac{\partial S}{\partial U}\right)_y = \frac{1}{T}$.
If U increases but S decreases (micro-states), T is negative.
- Example: isolated nuclear spin system.
- In external magnetic field B , the energy
 $\pm\varepsilon = \pm\mu B = \pm\frac{e\hbar}{2m}B$.
- Number of spins: N , where N_+ with ε , N_- with $-\varepsilon$.
- $N = N_+ + N_-$, $E = N_+\varepsilon + N_-(-\varepsilon)$

7.9 Negative temperature

- Basic function of thermodynamics: $dU = TdS - pdV$.
More general: $dU = TdS + Ydy$,
 $\Rightarrow (\frac{\partial S}{\partial U})_y = \frac{1}{T}$.
If U increases but S decreases (micro-states), T is negative.
- Example: isolated nuclear spin system.
- In external magnetic field B , the energy
 $\pm\varepsilon = \pm\mu B = \pm\frac{e\hbar}{2m}B$.
- Number of spins: N , where N_+ with ε , N_- with $-\varepsilon$.
- $N = N_+ + N_-$, $E = N_+\varepsilon + N_-(-\varepsilon)$
 $\Rightarrow N_+ = \frac{N}{2}(1 + \frac{E}{N\varepsilon})$, $N_- = \frac{N}{2}(1 - \frac{E}{N\varepsilon})$.

Negative temperature $N_{\pm} = \frac{N}{2} \left(1 \pm \frac{E}{N\varepsilon} \right)$

- Entropy $S = k \ln \Omega$

Negative temperature $N_{\pm} = \frac{N}{2} \left(1 \pm \frac{E}{N\varepsilon} \right)$

- Entropy $S = k \ln \Omega = k \ln \frac{N!}{N_+! N_-!}$

Negative temperature $N_{\pm} = \frac{N}{2} \left(1 \pm \frac{E}{N\varepsilon} \right)$

- Entropy $S = k \ln \Omega = k \ln \frac{N!}{N_+! N_-!}$
 $\simeq k[N(\ln N - 1) - N_+(\ln N_+ - 1) - N_-(\ln N_- - 1)]$

Negative temperature $N_{\pm} = \frac{N}{2} (1 \pm \frac{E}{N\varepsilon})$

- Entropy $S = k \ln \Omega = k \ln \frac{N!}{N_+! N_-!}$
 $\simeq k[N(\ln N - 1) - N_+(\ln N_+ - 1) - N_-(\ln N_- - 1)]$
 $= k(N \ln N - N_+ \ln N_+ - N_- \ln N_-)$

Negative temperature

$$N_{\pm} = \frac{N}{2} \left(1 \pm \frac{E}{N\varepsilon} \right)$$

- Entropy $S = k \ln \Omega = k \ln \frac{N!}{N_+! N_-!}$

$$\begin{aligned} &\simeq k[N(\ln N - 1) - N_+(\ln N_+ - 1) - N_-(\ln N_- - 1)] \\ &= k(N \ln N - N_+ \ln N_+ - N_- \ln N_-) \\ &= k\left\{ N \ln N - \frac{N}{2} \left(1 + \frac{E}{N\varepsilon} \right) \ln \left[\frac{N}{2} \left(1 + \frac{E}{N\varepsilon} \right) \right] \right. \\ &\quad \left. - \frac{N}{2} \left(1 - \frac{E}{N\varepsilon} \right) \ln \left[\frac{N}{2} \left(1 - \frac{E}{N\varepsilon} \right) \right] \right\} \end{aligned}$$

Negative temperature

$$N_{\pm} = \frac{N}{2} \left(1 \pm \frac{E}{N\varepsilon} \right)$$

- Entropy $S = k \ln \Omega = k \ln \frac{N!}{N_+! N_-!}$

$$\begin{aligned} &\simeq k [N(\ln N - 1) - N_+(\ln N_+ - 1) - N_-(\ln N_- - 1)] \\ &= k (N \ln N - N_+ \ln N_+ - N_- \ln N_-) \\ &= k \left\{ N \ln N - \frac{N}{2} \left(1 + \frac{E}{N\varepsilon} \right) \ln \left[\frac{N}{2} \left(1 + \frac{E}{N\varepsilon} \right) \right] \right. \\ &\quad \left. - \frac{N}{2} \left(1 - \frac{E}{N\varepsilon} \right) \ln \left[\frac{N}{2} \left(1 - \frac{E}{N\varepsilon} \right) \right] \right\} \\ &= k \left\{ N \ln N - \frac{N}{2} \left(1 + \frac{E}{N\varepsilon} \right) \left[\ln \frac{N}{2} + \ln \left(1 + \frac{E}{N\varepsilon} \right) \right] \right. \\ &\quad \left. - \frac{N}{2} \left(1 - \frac{E}{N\varepsilon} \right) \left[\ln \frac{N}{2} + \ln \left(1 - \frac{E}{N\varepsilon} \right) \right] \right\} \end{aligned}$$

Negative temperature $N_{\pm} = \frac{N}{2}(1 \pm \frac{E}{N\varepsilon})$

- Entropy $S = k \ln \Omega = k \ln \frac{N!}{N_+! N_-!}$

$$\begin{aligned} &\simeq k[N(\ln N - 1) - N_+(\ln N_+ - 1) - N_-(\ln N_- - 1)] \\ &= k(N \ln N - N_+ \ln N_+ - N_- \ln N_-) \\ &= k\left\{N \ln N - \frac{N}{2}\left(1 + \frac{E}{N\varepsilon}\right) \ln\left[\frac{N}{2}\left(1 + \frac{E}{N\varepsilon}\right)\right] \right. \\ &\quad \left. - \frac{N}{2}\left(1 - \frac{E}{N\varepsilon}\right) \ln\left[\frac{N}{2}\left(1 - \frac{E}{N\varepsilon}\right)\right]\right\} \\ &= k\left\{N \ln N - \frac{N}{2}\left(1 + \frac{E}{N\varepsilon}\right) \left[\ln \frac{N}{2} + \ln\left(1 + \frac{E}{N\varepsilon}\right)\right] \right. \\ &\quad \left. - \frac{N}{2}\left(1 - \frac{E}{N\varepsilon}\right) \left[\ln \frac{N}{2} + \ln\left(1 - \frac{E}{N\varepsilon}\right)\right]\right\} \\ &= k\left\{N \ln N - \frac{N}{2} \ln \frac{N}{2} \left[\left(1 + \frac{E}{N\varepsilon}\right) + \left(1 - \frac{E}{N\varepsilon}\right)\right] \right. \\ &\quad \left. - \frac{N}{2}\left(1 + \frac{E}{N\varepsilon}\right) \ln\left(1 + \frac{E}{N\varepsilon}\right) - \frac{N}{2}\left(1 - \frac{E}{N\varepsilon}\right) \ln\left(1 - \frac{E}{N\varepsilon}\right)\right\} \end{aligned}$$

Negative temperature

$$N_{\pm} = \frac{N}{2} \left(1 \pm \frac{E}{N\varepsilon} \right)$$

- Entropy $S = k \ln \Omega = k \ln \frac{N!}{N_+! N_-!}$

$$\begin{aligned} &\simeq k[N(\ln N - 1) - N_+(\ln N_+ - 1) - N_-(\ln N_- - 1)] \\ &= k(N \ln N - N_+ \ln N_+ - N_- \ln N_-) \\ &= k\left\{ N \ln N - \frac{N}{2} \left(1 + \frac{E}{N\varepsilon} \right) \ln \left[\frac{N}{2} \left(1 + \frac{E}{N\varepsilon} \right) \right] \right. \\ &\quad \left. - \frac{N}{2} \left(1 - \frac{E}{N\varepsilon} \right) \ln \left[\frac{N}{2} \left(1 - \frac{E}{N\varepsilon} \right) \right] \right\} \\ &= k\left\{ N \ln N - \frac{N}{2} \left(1 + \frac{E}{N\varepsilon} \right) \left[\ln \frac{N}{2} + \ln \left(1 + \frac{E}{N\varepsilon} \right) \right] \right. \\ &\quad \left. - \frac{N}{2} \left(1 - \frac{E}{N\varepsilon} \right) \left[\ln \frac{N}{2} + \ln \left(1 - \frac{E}{N\varepsilon} \right) \right] \right\} \\ &= k\left\{ N \ln N - \frac{N}{2} \ln \frac{N}{2} \left[\left(1 + \frac{E}{N\varepsilon} \right) + \left(1 - \frac{E}{N\varepsilon} \right) \right] \right. \\ &\quad \left. - \frac{N}{2} \left(1 + \frac{E}{N\varepsilon} \right) \ln \left(1 + \frac{E}{N\varepsilon} \right) - \frac{N}{2} \left(1 - \frac{E}{N\varepsilon} \right) \ln \left(1 - \frac{E}{N\varepsilon} \right) \right\} \\ &= k\left\{ N \ln N - N \ln \frac{N}{2} \right. \\ &\quad \left. - \frac{N}{2} \left(1 + \frac{E}{N\varepsilon} \right) \ln \left(1 + \frac{E}{N\varepsilon} \right) - \frac{N}{2} \left(1 - \frac{E}{N\varepsilon} \right) \ln \left(1 - \frac{E}{N\varepsilon} \right) \right\} \end{aligned}$$

Negative temperature

$$N_{\pm} = \frac{N}{2} \left(1 \pm \frac{E}{N\varepsilon}\right)$$

- Entropy $S = k \ln \Omega = k \ln \frac{N!}{N_+! N_-!}$

$$\begin{aligned} &\simeq k[N(\ln N - 1) - N_+(\ln N_+ - 1) - N_-(\ln N_- - 1)] \\ &= k(N \ln N - N_+ \ln N_+ - N_- \ln N_-) \\ &= k\left\{N \ln N - \frac{N}{2}\left(1 + \frac{E}{N\varepsilon}\right) \ln\left[\frac{N}{2}\left(1 + \frac{E}{N\varepsilon}\right)\right] \right. \\ &\quad \left. - \frac{N}{2}\left(1 - \frac{E}{N\varepsilon}\right) \ln\left[\frac{N}{2}\left(1 - \frac{E}{N\varepsilon}\right)\right]\right\} \\ &= k\left\{N \ln N - \frac{N}{2}\left(1 + \frac{E}{N\varepsilon}\right) \left[\ln \frac{N}{2} + \ln\left(1 + \frac{E}{N\varepsilon}\right)\right] \right. \\ &\quad \left. - \frac{N}{2}\left(1 - \frac{E}{N\varepsilon}\right) \left[\ln \frac{N}{2} + \ln\left(1 - \frac{E}{N\varepsilon}\right)\right]\right\} \\ &= k\left\{N \ln N - \frac{N}{2} \ln \frac{N}{2} \left[\left(1 + \frac{E}{N\varepsilon}\right) + \left(1 - \frac{E}{N\varepsilon}\right)\right] \right. \\ &\quad \left. - \frac{N}{2}\left(1 + \frac{E}{N\varepsilon}\right) \ln\left(1 + \frac{E}{N\varepsilon}\right) - \frac{N}{2}\left(1 - \frac{E}{N\varepsilon}\right) \ln\left(1 - \frac{E}{N\varepsilon}\right)\right\} \\ &= k\left\{N \ln N - N \ln \frac{N}{2} \right. \\ &\quad \left. - \frac{N}{2}\left(1 + \frac{E}{N\varepsilon}\right) \ln\left(1 + \frac{E}{N\varepsilon}\right) - \frac{N}{2}\left(1 - \frac{E}{N\varepsilon}\right) \ln\left(1 - \frac{E}{N\varepsilon}\right)\right\} \\ &= Nk\left[\ln 2 - \frac{1}{2}\left(1 + \frac{E}{N\varepsilon}\right) \ln\left(1 + \frac{E}{N\varepsilon}\right) - \frac{1}{2}\left(1 - \frac{E}{N\varepsilon}\right) \ln\left(1 - \frac{E}{N\varepsilon}\right)\right]. \end{aligned}$$

Negative temperature

Eq.(1.4.7) $\delta W = V H dB$, B generalized displacement.

Negative temperature

Eq.(1.4.7) $\delta W = V H dB$, B generalized displacement.

$$\frac{1}{T} = \left(\frac{\partial S}{\partial E} \right)_B$$

Negative temperature

Eq.(1.4.7) $\delta W = V H dB$, B generalized displacement.

$$\begin{aligned} \frac{1}{T} &= \left(\frac{\partial S}{\partial E} \right)_B \\ &= \frac{\partial}{\partial E} \left\{ Nk \left[\ln 2 - \frac{1}{2} \left(1 + \frac{E}{N\epsilon} \right) \ln \left(1 + \frac{E}{N\epsilon} \right) - \frac{1}{2} \left(1 - \frac{E}{N\epsilon} \right) \ln \left(1 - \frac{E}{N\epsilon} \right) \right] \right\} \end{aligned}$$

Negative temperature

Eq.(1.4.7) $\delta W = V H dB$, B generalized displacement.

$$\begin{aligned}
 \frac{1}{T} &= \left(\frac{\partial S}{\partial E} \right)_B \\
 &= \frac{\partial}{\partial E} \left\{ Nk \left[\ln 2 - \frac{1}{2} \left(1 + \frac{E}{N\varepsilon} \right) \ln \left(1 + \frac{E}{N\varepsilon} \right) - \frac{1}{2} \left(1 - \frac{E}{N\varepsilon} \right) \ln \left(1 - \frac{E}{N\varepsilon} \right) \right] \right\} \\
 &= -\frac{Nk}{2} \left[\left(1 + \frac{E}{N\varepsilon} \right) \cdot \frac{\frac{1}{N\varepsilon}}{1 + \frac{E}{N\varepsilon}} + \frac{1}{N\varepsilon} \ln \left(1 + \frac{E}{N\varepsilon} \right) + \left(1 - \frac{E}{N\varepsilon} \right) \cdot \right. \\
 &\quad \left. \frac{-\frac{1}{N\varepsilon}}{1 - \frac{E}{N\varepsilon}} - \frac{1}{N\varepsilon} \ln \left(1 - \frac{E}{N\varepsilon} \right) \right]
 \end{aligned}$$

Negative temperature

Eq.(1.4.7) $\delta W = V H dB$, B generalized displacement.

$$\begin{aligned}
 \frac{1}{T} &= \left(\frac{\partial S}{\partial E} \right)_B \\
 &= \frac{\partial}{\partial E} \left\{ Nk \left[\ln 2 - \frac{1}{2} \left(1 + \frac{E}{N\varepsilon} \right) \ln \left(1 + \frac{E}{N\varepsilon} \right) - \frac{1}{2} \left(1 - \frac{E}{N\varepsilon} \right) \ln \left(1 - \frac{E}{N\varepsilon} \right) \right] \right\} \\
 &= -\frac{Nk}{2} \left[\left(1 + \frac{E}{N\varepsilon} \right) \cdot \frac{\frac{1}{N\varepsilon}}{1 + \frac{E}{N\varepsilon}} + \frac{1}{N\varepsilon} \ln \left(1 + \frac{E}{N\varepsilon} \right) + \left(1 - \frac{E}{N\varepsilon} \right) \cdot \right. \\
 &\quad \left. \frac{-\frac{1}{N\varepsilon}}{1 - \frac{E}{N\varepsilon}} - \frac{1}{N\varepsilon} \ln \left(1 - \frac{E}{N\varepsilon} \right) \right] \\
 &= -\frac{Nk}{2} \left[\frac{1}{N\varepsilon} + \frac{1}{N\varepsilon} \ln \left(1 + \frac{E}{N\varepsilon} \right) - \frac{1}{N\varepsilon} - \frac{1}{N\varepsilon} \ln \left(1 - \frac{E}{N\varepsilon} \right) \right]
 \end{aligned}$$

Negative temperature

Eq.(1.4.7) $\delta W = V H dB$, B generalized displacement.

$$\begin{aligned}
 \frac{1}{T} &= \left(\frac{\partial S}{\partial E} \right)_B \\
 &= \frac{\partial}{\partial E} \left\{ Nk \left[\ln 2 - \frac{1}{2} \left(1 + \frac{E}{N\varepsilon} \right) \ln \left(1 + \frac{E}{N\varepsilon} \right) - \frac{1}{2} \left(1 - \frac{E}{N\varepsilon} \right) \ln \left(1 - \frac{E}{N\varepsilon} \right) \right] \right\} \\
 &= -\frac{Nk}{2} \left[\left(1 + \frac{E}{N\varepsilon} \right) \cdot \frac{\frac{1}{N\varepsilon}}{1 + \frac{E}{N\varepsilon}} + \frac{1}{N\varepsilon} \ln \left(1 + \frac{E}{N\varepsilon} \right) + \left(1 - \frac{E}{N\varepsilon} \right) \cdot \right. \\
 &\quad \left. -\frac{\frac{1}{N\varepsilon}}{1 - \frac{E}{N\varepsilon}} - \frac{1}{N\varepsilon} \ln \left(1 - \frac{E}{N\varepsilon} \right) \right] \\
 &= -\frac{Nk}{2} \left[\frac{1}{N\varepsilon} + \frac{1}{N\varepsilon} \ln \left(1 + \frac{E}{N\varepsilon} \right) - \frac{1}{N\varepsilon} - \frac{1}{N\varepsilon} \ln \left(1 - \frac{E}{N\varepsilon} \right) \right] \\
 &= -\frac{k}{2\varepsilon} \left[\ln \left(1 + \frac{E}{N\varepsilon} \right) - \ln \left(1 - \frac{E}{N\varepsilon} \right) \right]
 \end{aligned}$$

Negative temperature

Eq.(1.4.7) $\delta W = V H dB$, B generalized displacement.

$$\begin{aligned}
 \frac{1}{T} &= \left(\frac{\partial S}{\partial E} \right)_B \\
 &= \frac{\partial}{\partial E} \left\{ Nk \left[\ln 2 - \frac{1}{2} \left(1 + \frac{E}{N\varepsilon} \right) \ln \left(1 + \frac{E}{N\varepsilon} \right) - \frac{1}{2} \left(1 - \frac{E}{N\varepsilon} \right) \ln \left(1 - \frac{E}{N\varepsilon} \right) \right] \right\} \\
 &= -\frac{Nk}{2} \left[\left(1 + \frac{E}{N\varepsilon} \right) \cdot \frac{\frac{1}{N\varepsilon}}{1 + \frac{E}{N\varepsilon}} + \frac{1}{N\varepsilon} \ln \left(1 + \frac{E}{N\varepsilon} \right) + \left(1 - \frac{E}{N\varepsilon} \right) \cdot \right. \\
 &\quad \left. -\frac{\frac{1}{N\varepsilon}}{1 - \frac{E}{N\varepsilon}} - \frac{1}{N\varepsilon} \ln \left(1 - \frac{E}{N\varepsilon} \right) \right] \\
 &= -\frac{Nk}{2} \left[\frac{1}{N\varepsilon} + \frac{1}{N\varepsilon} \ln \left(1 + \frac{E}{N\varepsilon} \right) - \frac{1}{N\varepsilon} - \frac{1}{N\varepsilon} \ln \left(1 - \frac{E}{N\varepsilon} \right) \right] \\
 &= -\frac{k}{2\varepsilon} \left[\ln \left(1 + \frac{E}{N\varepsilon} \right) - \ln \left(1 - \frac{E}{N\varepsilon} \right) \right] = \frac{k}{2\varepsilon} \ln \frac{N\varepsilon - E}{N\varepsilon + E}.
 \end{aligned}$$

Negative temperature

Eq.(1.4.7) $\text{d}W = V H \text{d}B$, B generalized displacement.

$$\begin{aligned}
 \frac{1}{T} &= \left(\frac{\partial S}{\partial E} \right)_B \\
 &= \frac{\partial}{\partial E} \left\{ Nk \left[\ln 2 - \frac{1}{2} \left(1 + \frac{E}{N\epsilon} \right) \ln \left(1 + \frac{E}{N\epsilon} \right) - \frac{1}{2} \left(1 - \frac{E}{N\epsilon} \right) \ln \left(1 - \frac{E}{N\epsilon} \right) \right] \right\} \\
 &= -\frac{Nk}{2} \left[\left(1 + \frac{E}{N\epsilon} \right) \cdot \frac{\frac{1}{N\epsilon}}{1 + \frac{E}{N\epsilon}} + \frac{1}{N\epsilon} \ln \left(1 + \frac{E}{N\epsilon} \right) + \left(1 - \frac{E}{N\epsilon} \right) \cdot \right. \\
 &\quad \left. -\frac{\frac{1}{N\epsilon}}{1 - \frac{E}{N\epsilon}} - \frac{1}{N\epsilon} \ln \left(1 - \frac{E}{N\epsilon} \right) \right] \\
 &= -\frac{Nk}{2} \left[\frac{1}{N\epsilon} + \frac{1}{N\epsilon} \ln \left(1 + \frac{E}{N\epsilon} \right) - \frac{1}{N\epsilon} - \frac{1}{N\epsilon} \ln \left(1 - \frac{E}{N\epsilon} \right) \right] \\
 &= -\frac{k}{2\epsilon} \left[\ln \left(1 + \frac{E}{N\epsilon} \right) - \ln \left(1 - \frac{E}{N\epsilon} \right) \right] = \frac{k}{2\epsilon} \ln \frac{N\epsilon - E}{N\epsilon + E}.
 \end{aligned}$$





Table of contents

1 §7 Boltzmann Statistics

- 7.1 Thermal quantities in statistics
- 7.2 Equation of state of ideal gas
- 7.3 Maxwell speed distribution
- 7.4 Energy equipartition theorem
- 7.5 Internal energy and heat capacity of the ideal gas
- 7.6 Entropy of the ideal gas
- 7.7 Einstein's theory on heat capacity of solid
- 7.8 Paramagnetic solid
- 7.9 Negative temperature