

Thermodynamics & Statistical Physics

Chapter 9. Ensemble theory

Yuan-Chuan Zou
zouyc@hust.edu.cn

School of Physics, Huazhong University of Science and Technology

December 30, 2013

Table of contents

1 §9. Ensemble theory

- 9.1 Phase space and Liouville's theorem
- 9.2 The microcanonical ensemble
- 9.3 Thermodynamics in microcanonical ensemble theory
- 9.4 Canonical ensemble
- 9.5 Thermodynamics in canonical ensemble
- 9.6 EOS of real gas
- 9.7 Thermal capacity of solid
- 9.9 Ising model
- 9.10 Grand canonical ensemble
- 9.11 Thermodynamics in grand canonical ensemble
- 9.12 Applications of the grand canonical ensemble

9.1 Phase space and Liouville's theorem

9.1 Phase space and Liouville's theorem

- Chapters 6,7,8 are not suitable to interacting particles.

9.1 Phase space and Liouville's theorem

- Chapters 6,7,8 are not suitable to interacting particles. Because of potential energy, any movement of any particle changes states of other particles.

9.1 Phase space and Liouville's theorem

- Chapters 6,7,8 are not suitable to interacting particles. Because of potential energy, any movement of any particle changes states of other particles. No fixed energy or momentum for a particle, varying with time.

9.1 Phase space and Liouville's theorem

- Chapters 6,7,8 are not suitable to interacting particles. Because of potential energy, any movement of any particle changes states of other particles. No fixed energy or momentum for a particle, varying with time.
- Particle can not be separated from the system.

9.1 Phase space and Liouville's theorem

- Chapters 6,7,8 are not suitable to interacting particles. Because of potential energy, any movement of any particle changes states of other particles. No fixed energy or momentum for a particle, varying with time.
- Particle can not be separated from the system.
- Use state of system instead of state of particles.

9.1 Phase space and Liouville's theorem

- Chapters 6,7,8 are not suitable to interacting particles. Because of potential energy, any movement of any particle changes states of other particles. No fixed energy or momentum for a particle, varying with time.
- Particle can not be separated from the system.
- Use state of system instead of state of particles.
- Ensemble theory,

9.1 Phase space and Liouville's theorem

- Chapters 6,7,8 are not suitable to interacting particles. Because of potential energy, any movement of any particle changes states of other particles. No fixed energy or momentum for a particle, varying with time.
- Particle can not be separated from the system.
- Use state of system instead of state of particles.
- Ensemble theory, by using Hamiltonian.

9.1 Phase space and Liouville's theorem

- Chapters 6,7,8 are not suitable to interacting particles. Because of potential energy, any movement of any particle changes states of other particles. No fixed energy or momentum for a particle, varying with time.
- Particle can not be separated from the system.
- Use state of system instead of state of particles.
- Ensemble theory, by using Hamiltonian.
- Consider N particles system, each one with r freedom.

9.1 Phase space and Liouville's theorem

- Chapters 6,7,8 are not suitable to interacting particles. Because of potential energy, any movement of any particle changes states of other particles. No fixed energy or momentum for a particle, varying with time.
- Particle can not be separated from the system.
- Use state of system instead of state of particles.
- Ensemble theory, by using Hamiltonian.
- Consider N particles system, each one with r freedom.
- For a particle j , it has: $(q_{j1}, \dots, q_{jr}; p_{j1}, \dots, p_{jr})$.

9.1 Phase space and Liouville's theorem

- Chapters 6,7,8 are not suitable to interacting particles. Because of potential energy, any movement of any particle changes states of other particles. No fixed energy or momentum for a particle, varying with time.
- Particle can not be separated from the system.
- Use state of system instead of state of particles.
- Ensemble theory, by using Hamiltonian.
- Consider N particles system, each one with r freedom.
- For a particle j , it has: $(q_{j1}, \dots, q_{jr}; p_{j1}, \dots, p_{jr})$.
- For the system, to describe the micro-state, using the **phase-space point**:

$$(q_{11}, \dots, q_{1r}, \dots, q_{N1}, \dots, q_{Nr}; p_{11}, \dots, p_{1r}, \dots, p_{N1}, \dots, p_{Nr})$$

Phase space and Liouville's theorem

- There are $2Nr$ quantities in $(q_{11}, \dots, q_{1r}, \dots, q_{N1}, \dots, q_{Nr}; p_{11}, \dots, p_{1r}, \dots, p_{N1}, \dots, p_{Nr})$.

Phase space and Liouville's theorem

- There are $2Nr$ quantities in $(q_{11}, \dots, q_{1r}, \dots, q_{N1}, \dots, q_{Nr}; p_{11}, \dots, p_{1r}, \dots, p_{N1}, \dots, p_{Nr})$.
The change of any quantity means the change of the state of the system, and means the movement of the phase-space point.

Phase space and Liouville's theorem

- There are $2Nr$ quantities in $(q_{11}, \dots, q_{1r}, \dots, q_{N1}, \dots, q_{Nr}; p_{11}, \dots, p_{1r}, \dots, p_{N1}, \dots, p_{Nr})$.
The change of any quantity means the change of the state of the system, and means the movement of the phase-space point.
- Define $f = Nr$, simplifies the phase-space point: $(q_1, \dots, q_f; p_1, \dots, p_f)$.

Phase space and Liouville's theorem

- There are $2Nr$ quantities in $(q_{11}, \dots, q_{1r}, \dots, q_{N1}, \dots, q_{Nr}; p_{11}, \dots, p_{1r}, \dots, p_{N1}, \dots, p_{Nr})$.
The change of any quantity means the change of the state of the system, and means the movement of the phase-space point.
- Define $f = Nr$, simplifies the phase-space point: $(q_1, \dots, q_f; p_1, \dots, p_f)$.
- Any movement of the particles obeys Hamiltonian canonical equation: $\dot{q}_i = \frac{\partial H}{\partial p_i}, \dot{p}_i = -\frac{\partial H}{\partial q_i}, i = 1, 2, \dots, f$.

Phase space and Liouville's theorem

- There are $2Nr$ quantities in $(q_{11}, \dots, q_{1r}, \dots, q_{N1}, \dots, q_{Nr}; p_{11}, \dots, p_{1r}, \dots, p_{N1}, \dots, p_{Nr})$.
The change of any quantity means the change of the state of the system, and means the movement of the phase-space point.
- Define $f = Nr$, simplifies the phase-space point: $(q_1, \dots, q_f; p_1, \dots, p_f)$.
- Any movement of the particles obeys Hamiltonian canonical equation: $\dot{q}_i = \frac{\partial H}{\partial p_i}, \dot{p}_i = -\frac{\partial H}{\partial q_i}, i = 1, 2, \dots, f$.
- (\dot{q}_i, \dot{p}_i) determines the trajectory of the phase-space point.

Phase space and Liouville's theorem

- Consider a lot of (\mathcal{N}) identical systems, with different initial state. Let them develop.

Phase space and Liouville's theorem

- Consider a lot of (\mathcal{N}) identical systems, with different initial state. Let them develop.
- At a certain time t , one can count the number of points at any given volume $d\Omega = dq_1 \dots dq_f dp_1 \dots dp_f$,

Phase space and Liouville's theorem

- Consider a lot of (\mathcal{N}) identical systems, with different initial state. Let them develop.
- At a certain time t , one can count the number of points at any given volume $d\Omega = dq_1 \dots dq_f dp_1 \dots dp_f$, the number is $\rho(q_1, \dots q_f, p_1, \dots p_f, t) d\Omega$,

Phase space and Liouville's theorem

- Consider a lot of (\mathcal{N}) identical systems, with different initial state. Let them develop.
- At a certain time t , one can count the number of points at any given volume $d\Omega = dq_1 \dots dq_f dp_1 \dots dp_f$, the number is $\rho(q_1, \dots q_f, p_1, \dots p_f, t)d\Omega$, where ρ is the number density of points, or number of density of micro-states.

Phase space and Liouville's theorem

- Consider a lot of (\mathcal{N}) identical systems, with different initial state. Let them develop.
- At a certain time t , one can count the number of points at any given volume $d\Omega = dq_1 \dots dq_f dp_1 \dots dp_f$, the number is $\rho(q_1, \dots, q_f, p_1, \dots, p_f, t) d\Omega$, where ρ is the number density of points, or number of density of micro-states. Obviously, $\mathcal{N} = \int \rho d\Omega$.

Phase space and Liouville's theorem

- Consider a lot of (\mathcal{N}) identical systems, with different initial state. Let them develop.
- At a certain time t , one can count the number of points at any given volume $d\Omega = dq_1 \dots dq_f dp_1 \dots dp_f$, the number is $\rho(q_1, \dots, q_f, p_1, \dots, p_f, t) d\Omega$, where ρ is the number density of points, or number of density of micro-states. Obviously, $\mathcal{N} = \int \rho d\Omega$.
- The behavior of ρ depends on t :
 $\rho(q + dq, p + dp, t + dt)$

Phase space and Liouville's theorem

- Consider a lot of (\mathcal{N}) identical systems, with different initial state. Let them develop.
- At a certain time t , one can count the number of points at any given volume $d\Omega = dq_1 \dots dq_f dp_1 \dots dp_f$, the number is $\rho(q_1, \dots q_f, p_1, \dots p_f, t)d\Omega$, where ρ is the number density of points, or number of density of micro-states. Obviously, $\mathcal{N} = \int \rho d\Omega$.

- The behavior of ρ depends on t :

$$\begin{aligned} & \rho(q + dq, p + dp, t + dt) \\ &= \rho(q, p, t) + \frac{\partial \rho}{\partial q_1} dq_1 + \dots + \frac{\partial \rho}{\partial p_f} dp_f + \frac{\partial \rho}{\partial t} dt \end{aligned}$$

Phase space and Liouville's theorem

- Consider a lot of (\mathcal{N}) identical systems, with different initial state. Let them develop.
- At a certain time t , one can count the number of points at any given volume $d\Omega = dq_1 \dots dq_f dp_1 \dots dp_f$, the number is $\rho(q_1, \dots q_f, p_1, \dots p_f, t)d\Omega$, where ρ is the number density of points, or number of density of micro-states. Obviously, $\mathcal{N} = \int \rho d\Omega$.

- The behavior of ρ depends on t :

$$\begin{aligned}
 & \rho(q + dq, p + dp, t + dt) \\
 &= \rho(q, p, t) + \frac{\partial \rho}{\partial q_1} dq_1 + \dots + \frac{\partial \rho}{\partial p_f} dp_f + \frac{\partial \rho}{\partial t} dt \\
 &= \rho(t) + \left(\frac{\partial \rho}{\partial t} + \frac{\partial \rho}{\partial q_1} \dot{q}_1 + \dots + \frac{\partial \rho}{\partial p_f} \dot{p}_f \right) dt.
 \end{aligned}$$

Phase space and Liouville's theorem

- $\rho(q + dq, p + dp, t + dt)$
 $= \rho(t) + \left(\frac{\partial \rho}{\partial t} + \frac{\partial \rho}{\partial q_1} \dot{q}_1 + \dots + \frac{\partial \rho}{\partial p_f} \dot{p}_f \right) dt$

Phase space and Liouville's theorem

- $\rho(q + dq, p + dp, t + dt)$
 $= \rho(t) + (\frac{\partial \rho}{\partial t} + \frac{\partial \rho}{\partial q_1} \dot{q}_1 + \dots + \frac{\partial \rho}{\partial p_f} \dot{p}_f) dt$
 $= \rho(t) + [\frac{\partial \rho}{\partial t} + \sum (\frac{\partial \rho}{\partial q_i} \dot{q}_i + \frac{\partial \rho}{\partial p_i} \dot{p}_i)] dt$

Phase space and Liouville's theorem

- $\rho(q + dq, p + dp, t + dt)$
 $= \rho(t) + (\frac{\partial \rho}{\partial t} + \frac{\partial \rho}{\partial q_1} \dot{q}_1 + \dots + \frac{\partial \rho}{\partial p_f} \dot{p}_f) dt$
 $= \rho(t) + [\frac{\partial \rho}{\partial t} + \sum (\frac{\partial \rho}{\partial q_i} \dot{q}_i + \frac{\partial \rho}{\partial p_i} \dot{p}_i)] dt$
- $\frac{d\rho}{dt} = \frac{\partial \rho}{\partial t} + \sum (\frac{\partial \rho}{\partial q_i} \dot{q}_i + \frac{\partial \rho}{\partial p_i} \dot{p}_i).$

Phase space and Liouville's theorem

- $\rho(q + dq, p + dp, t + dt)$
 $= \rho(t) + (\frac{\partial \rho}{\partial t} + \frac{\partial \rho}{\partial q_1} \dot{q}_1 + \dots + \frac{\partial \rho}{\partial p_f} \dot{p}_f) dt$
 $= \rho(t) + [\frac{\partial \rho}{\partial t} + \sum (\frac{\partial \rho}{\partial q_i} \dot{q}_i + \frac{\partial \rho}{\partial p_i} \dot{p}_i)] dt$
- $\frac{d\rho}{dt} = \frac{\partial \rho}{\partial t} + \sum (\frac{\partial \rho}{\partial q_i} \dot{q}_i + \frac{\partial \rho}{\partial p_i} \dot{p}_i).$
- Provable: $\frac{d\rho}{dt} = 0.$

Phase space and Liouville's theorem

- $\rho(q + dq, p + dp, t + dt)$
 $= \rho(t) + (\frac{\partial \rho}{\partial t} + \frac{\partial \rho}{\partial q_1} \dot{q}_1 + \dots + \frac{\partial \rho}{\partial p_f} \dot{p}_f) dt$
 $= \rho(t) + [\frac{\partial \rho}{\partial t} + \sum (\frac{\partial \rho}{\partial q_i} \dot{q}_i + \frac{\partial \rho}{\partial p_i} \dot{p}_i)] dt$
- $\frac{d\rho}{dt} = \frac{\partial \rho}{\partial t} + \sum (\frac{\partial \rho}{\partial q_i} \dot{q}_i + \frac{\partial \rho}{\partial p_i} \dot{p}_i).$
- Provable: $\frac{d\rho}{dt} = 0.$
- Consider a fixed volume element:
 $d\Omega = dq_1 \dots dq_f dp_1 \dots dp_f.$

Phase space and Liouville's theorem

- $\rho(q + dq, p + dp, t + dt)$
 $= \rho(t) + \left(\frac{\partial \rho}{\partial t} + \frac{\partial \rho}{\partial q_1} \dot{q}_1 + \dots + \frac{\partial \rho}{\partial p_f} \dot{p}_f \right) dt$
 $= \rho(t) + \left[\frac{\partial \rho}{\partial t} + \sum \left(\frac{\partial \rho}{\partial q_i} \dot{q}_i + \frac{\partial \rho}{\partial p_i} \dot{p}_i \right) \right] dt$
- $\frac{d\rho}{dt} = \frac{\partial \rho}{\partial t} + \sum \left(\frac{\partial \rho}{\partial q_i} \dot{q}_i + \frac{\partial \rho}{\partial p_i} \dot{p}_i \right).$
- Provable: $\frac{d\rho}{dt} = 0$.
- Consider a fixed volume element:
 $d\Omega = dq_1 \dots dq_f dp_1 \dots dp_f.$
 At time t , the number of phase-space points: $\rho d\Omega$;

Phase space and Liouville's theorem

- $\rho(q + dq, p + dp, t + dt)$
 $= \rho(t) + \left(\frac{\partial \rho}{\partial t} + \frac{\partial \rho}{\partial q_1} \dot{q}_1 + \dots + \frac{\partial \rho}{\partial p_f} \dot{p}_f \right) dt$
 $= \rho(t) + \left[\frac{\partial \rho}{\partial t} + \sum \left(\frac{\partial \rho}{\partial q_i} \dot{q}_i + \frac{\partial \rho}{\partial p_i} \dot{p}_i \right) \right] dt$
- $\frac{d\rho}{dt} = \frac{\partial \rho}{\partial t} + \sum \left(\frac{\partial \rho}{\partial q_i} \dot{q}_i + \frac{\partial \rho}{\partial p_i} \dot{p}_i \right).$
- Provable: $\frac{d\rho}{dt} = 0$.
- Consider a fixed volume element:
 $d\Omega = dq_1 \dots dq_f dp_1 \dots dp_f.$
 At time t , the number of phase-space points: $\rho d\Omega$;
 at time $t + dt$, the number: $(\rho + \frac{\partial \rho}{\partial t} dt) d\Omega$.

Phase space and Liouville's theorem

- $\rho(q + dq, p + dp, t + dt)$
 $= \rho(t) + \left(\frac{\partial \rho}{\partial t} + \frac{\partial \rho}{\partial q_1} \dot{q}_1 + \dots + \frac{\partial \rho}{\partial p_f} \dot{p}_f \right) dt$
 $= \rho(t) + \left[\frac{\partial \rho}{\partial t} + \sum \left(\frac{\partial \rho}{\partial q_i} \dot{q}_i + \frac{\partial \rho}{\partial p_i} \dot{p}_i \right) \right] dt$
- $\frac{d\rho}{dt} = \frac{\partial \rho}{\partial t} + \sum \left(\frac{\partial \rho}{\partial q_i} \dot{q}_i + \frac{\partial \rho}{\partial p_i} \dot{p}_i \right).$
- Provable: $\frac{d\rho}{dt} = 0$.
- Consider a fixed volume element:
 $d\Omega = dq_1 \dots dq_f dp_1 \dots dp_f.$
 At time t , the number of phase-space points: $\rho d\Omega$;
 at time $t + dt$, the number: $(\rho + \frac{d\rho}{dt} dt) d\Omega.$
- The variance of the number comes from the flowing in and out through the surfaces of the volume element.

Phase space and Liouville's theorem

- Consider a pair of surfaces $q_i = q_i$ and $q_i = q_i + dq_i$ as an example.

Phase space and Liouville's theorem

- Consider a pair of surfaces $q_i = q_i$ and $q_i = q_i + dq_i$ as an example.
- \dot{q}_i is the velocity of flowing in; the surface $dA = dq_1 \dots dq_{i-1} dq_{i+1} \dots dq_f dp_1 \dots dp_f$.

Phase space and Liouville's theorem

- Consider a pair of surfaces $q_i = q_i$ and $q_i = q_i + dq_i$ as an example.
- \dot{q}_i is the velocity of flowing in; the surface $dA = dq_1 \dots dq_{i-1} dq_{i+1} \dots dq_f dp_1 \dots dp_f$.
- The number of points flowing in through the surface: $\rho \dot{q}_i dt dA$;

Phase space and Liouville's theorem

- Consider a pair of surfaces $q_i = q_i$ and $q_i = q_i + dq_i$ as an example.
- \dot{q}_i is the velocity of flowing in; the surface $dA = dq_1 \dots dq_{i-1} dq_{i+1} \dots dq_f dp_1 \dots dp_f$.
- The number of points flowing in through the surface: $\rho \dot{q}_i dt dA$;
similar, the number flowing out through the contrary surface: $(\rho \dot{q}_i)_{q_i + dq_i} dt dA$

Phase space and Liouville's theorem

- Consider a pair of surfaces $q_i = q_i$ and $q_i = q_i + dq_i$ as an example.
- \dot{q}_i is the velocity of flowing in; the surface $dA = dq_1 \dots dq_{i-1} dq_{i+1} \dots dq_f dp_1 \dots dp_f$.
- The number of points flowing in through the surface: $\rho \dot{q}_i dt dA$;
similar, the number flowing out through the contrary surface: $(\rho \dot{q}_i)_{q_i + dq_i} dt dA = [(\rho \dot{q}_i)_{q_i} + \frac{\partial(\rho \dot{q}_i)}{\partial q_i} dq_i] dt dA$.

Phase space and Liouville's theorem

- Consider a pair of surfaces $q_i = q_i$ and $q_i = q_i + dq_i$ as an example.
- \dot{q}_i is the velocity of flowing in; the surface $dA = dq_1 \dots dq_{i-1} dq_{i+1} \dots dq_f dp_1 \dots dp_f$.
- The number of points flowing in through the surface: $\rho \dot{q}_i dt dA$;
similar, the number flowing out through the contrary surface: $(\rho \dot{q}_i)_{q_i + dq_i} dt dA = [(\rho \dot{q}_i)_{q_i} + \frac{\partial(\rho \dot{q}_i)}{\partial q_i} dq_i] dt dA$.
- The net flowed in: $-\frac{\partial(\rho \dot{q}_i)}{\partial q_i} dq_i dt dA$

Phase space and Liouville's theorem

- Consider a pair of surfaces $q_i = q_i$ and $q_i = q_i + dq_i$ as an example.
- \dot{q}_i is the velocity of flowing in; the surface $dA = dq_1 \dots dq_{i-1} dq_{i+1} \dots dq_f dp_1 \dots dp_f$.
- The number of points flowing in through the surface: $\rho \dot{q}_i dt dA$;
similar, the number flowing out through the contrary surface: $(\rho \dot{q}_i)_{q_i + dq_i} dt dA = [(\rho \dot{q}_i)_{q_i} + \frac{\partial(\rho \dot{q}_i)}{\partial q_i} dq_i] dt dA$.
- The net flowed in: $-\frac{\partial(\rho \dot{q}_i)}{\partial q_i} dq_i dt dA = -\frac{\partial(\rho \dot{q}_i)}{\partial q_i} dt d\Omega$.

Phase space and Liouville's theorem

- Consider a pair of surfaces $q_i = q_i$ and $q_i = q_i + dq_i$ as an example.
- \dot{q}_i is the velocity of flowing in; the surface $dA = dq_1 \dots dq_{i-1} dq_{i+1} \dots dq_f dp_1 \dots dp_f$.
- The number of points flowing in through the surface: $\rho \dot{q}_i dt dA$;
similar, the number flowing out through the contrary surface: $(\rho \dot{q}_i)_{q_i + dq_i} dt dA = [(\rho \dot{q}_i)_{q_i} + \frac{\partial(\rho \dot{q}_i)}{\partial q_i} dq_i] dt dA$.
- The net flowed in: $-\frac{\partial(\rho \dot{q}_i)}{\partial q_i} dq_i dt dA = -\frac{\partial(\rho \dot{q}_i)}{\partial q_i} dt d\Omega$.
- Similar for p_i , the net in: $-\frac{\partial(\rho \dot{p}_i)}{\partial p_i} dt d\Omega$.

Phase space and Liouville's theorem

- Consider a pair of surfaces $q_i = q_i$ and $q_i = q_i + dq_i$ as an example.
- \dot{q}_i is the velocity of flowing in; the surface $dA = dq_1 \dots dq_{i-1} dq_{i+1} \dots dq_f dp_1 \dots dp_f$.
- The number of points flowing in through the surface: $\rho \dot{q}_i dt dA$;
similar, the number flowing out through the contrary surface: $(\rho \dot{q}_i)_{q_i + dq_i} dt dA = [(\rho \dot{q}_i)_{q_i} + \frac{\partial(\rho \dot{q}_i)}{\partial q_i} dq_i] dt dA$.
- The net flowed in: $-\frac{\partial(\rho \dot{q}_i)}{\partial q_i} dq_i dt dA = -\frac{\partial(\rho \dot{q}_i)}{\partial q_i} dt d\Omega$.
- Similar for p_i , the net in: $-\frac{\partial(\rho \dot{p}_i)}{\partial p_i} dt d\Omega$.
- Sum on all surfaces: $-\sum [\frac{\partial(\rho \dot{q}_i)}{\partial q_i} + \frac{\partial(\rho \dot{p}_i)}{\partial p_i}] dt d\Omega$.

Phase space and Liouville's theorem

- On the other hand, the net gain in the fixed volume:
 $\frac{\partial \rho}{\partial t} dt d\Omega$.

Phase space and Liouville's theorem

- On the other hand, the net gain in the fixed volume:
 $\frac{\partial \rho}{\partial t} dt d\Omega$.
- $\therefore - \sum \left[\frac{\partial(\rho \dot{q}_i)}{\partial q_i} + \frac{\partial(\rho \dot{p}_i)}{\partial p_i} \right] dt d\Omega = \frac{\partial \rho}{\partial t} dt d\Omega$

Phase space and Liouville's theorem

- On the other hand, the net gain in the fixed volume:
 $\frac{\partial \rho}{\partial t} dt d\Omega$.
- $\therefore - \sum \left[\frac{\partial(\rho \dot{q}_i)}{\partial q_i} + \frac{\partial(\rho \dot{p}_i)}{\partial p_i} \right] dt d\Omega = \frac{\partial \rho}{\partial t} dt d\Omega$
 $\Rightarrow \frac{\partial \rho}{\partial t} = - \sum \left[\frac{\partial(\rho \dot{q}_i)}{\partial q_i} + \frac{\partial(\rho \dot{p}_i)}{\partial p_i} \right]$

Phase space and Liouville's theorem

- On the other hand, the net gain in the fixed volume:
 $\frac{\partial \rho}{\partial t} dt d\Omega$.
- $\therefore - \sum \left[\frac{\partial(\rho \dot{q}_i)}{\partial q_i} + \frac{\partial(\rho \dot{p}_i)}{\partial p_i} \right] dt d\Omega = \frac{\partial \rho}{\partial t} dt d\Omega$
 $\Rightarrow \frac{\partial \rho}{\partial t} = - \sum \left[\frac{\partial(\rho \dot{q}_i)}{\partial q_i} + \frac{\partial(\rho \dot{p}_i)}{\partial p_i} \right]$
 $= - \sum \left[\frac{\partial \rho}{\partial q_i} \dot{q}_i + \frac{\partial \rho}{\partial p_i} \dot{p}_i + \rho \left(\frac{\partial \dot{q}_i}{\partial q_i} + \frac{\partial \dot{p}_i}{\partial p_i} \right) \right].$

Phase space and Liouville's theorem

- On the other hand, the net gain in the fixed volume:
 $\frac{\partial \rho}{\partial t} dt d\Omega$.
- $$\therefore - \sum \left[\frac{\partial(\rho \dot{q}_i)}{\partial q_i} + \frac{\partial(\rho \dot{p}_i)}{\partial p_i} \right] dt d\Omega = \frac{\partial \rho}{\partial t} dt d\Omega$$

$$\Rightarrow \frac{\partial \rho}{\partial t} = - \sum \left[\frac{\partial(\rho \dot{q}_i)}{\partial q_i} + \frac{\partial(\rho \dot{p}_i)}{\partial p_i} \right]$$

$$= - \sum \left[\frac{\partial \rho}{\partial q_i} \dot{q}_i + \frac{\partial \rho}{\partial p_i} \dot{p}_i + \rho \left(\frac{\partial \dot{q}_i}{\partial q_i} + \frac{\partial \dot{p}_i}{\partial p_i} \right) \right].$$
- According to $\dot{q}_i = \frac{\partial H}{\partial p_i}$, $\dot{p}_i = -\frac{\partial H}{\partial q_i}$,

Phase space and Liouville's theorem

- On the other hand, the net gain in the fixed volume:
 $\frac{\partial \rho}{\partial t} dt d\Omega$.
- $\therefore - \sum \left[\frac{\partial(\rho \dot{q}_i)}{\partial q_i} + \frac{\partial(\rho \dot{p}_i)}{\partial p_i} \right] dt d\Omega = \frac{\partial \rho}{\partial t} dt d\Omega$
 $\Rightarrow \frac{\partial \rho}{\partial t} = - \sum \left[\frac{\partial(\rho \dot{q}_i)}{\partial q_i} + \frac{\partial(\rho \dot{p}_i)}{\partial p_i} \right]$
 $= - \sum \left[\frac{\partial \rho}{\partial q_i} \dot{q}_i + \frac{\partial \rho}{\partial p_i} \dot{p}_i + \rho \left(\frac{\partial \dot{q}_i}{\partial q_i} + \frac{\partial \dot{p}_i}{\partial p_i} \right) \right].$
- According to $\dot{q}_i = \frac{\partial H}{\partial p_i}$, $\dot{p}_i = -\frac{\partial H}{\partial q_i}$, we have
 $\frac{\partial^2 H}{\partial p_i \partial q_i} = \frac{\partial^2 H}{\partial q_i \partial p_i} \Rightarrow -\frac{\partial \dot{p}_i}{\partial p_i} = \frac{\partial \dot{q}_i}{\partial q_i}.$

Phase space and Liouville's theorem

- On the other hand, the net gain in the fixed volume:

$$\frac{\partial \rho}{\partial t} dt d\Omega.$$

- $$\therefore - \sum \left[\frac{\partial(\rho \dot{q}_i)}{\partial q_i} + \frac{\partial(\rho \dot{p}_i)}{\partial p_i} \right] dt d\Omega = \frac{\partial \rho}{\partial t} dt d\Omega$$

$$\Rightarrow \frac{\partial \rho}{\partial t} = - \sum \left[\frac{\partial(\rho \dot{q}_i)}{\partial q_i} + \frac{\partial(\rho \dot{p}_i)}{\partial p_i} \right]$$

$$= - \sum \left[\frac{\partial \rho}{\partial q_i} \dot{q}_i + \frac{\partial \rho}{\partial p_i} \dot{p}_i + \rho \left(\frac{\partial \dot{q}_i}{\partial q_i} + \frac{\partial \dot{p}_i}{\partial p_i} \right) \right].$$

- According to $\dot{q}_i = \frac{\partial H}{\partial p_i}$, $\dot{p}_i = -\frac{\partial H}{\partial q_i}$, we have

$$\frac{\partial^2 H}{\partial p_i \partial q_i} = \frac{\partial^2 H}{\partial q_i \partial p_i} \Rightarrow -\frac{\partial \dot{p}_i}{\partial p_i} = \frac{\partial \dot{q}_i}{\partial q_i}.$$

$$\therefore \frac{\partial \rho}{\partial t} + \sum \left(\frac{\partial \rho}{\partial q_i} \dot{q}_i + \frac{\partial \rho}{\partial p_i} \dot{p}_i \right) = 0,$$

Phase space and Liouville's theorem

- On the other hand, the net gain in the fixed volume:

$$\frac{\partial \rho}{\partial t} dt d\Omega.$$

- $$\therefore - \sum \left[\frac{\partial(\rho \dot{q}_i)}{\partial q_i} + \frac{\partial(\rho \dot{p}_i)}{\partial p_i} \right] dt d\Omega = \frac{\partial \rho}{\partial t} dt d\Omega$$

$$\Rightarrow \frac{\partial \rho}{\partial t} = - \sum \left[\frac{\partial(\rho \dot{q}_i)}{\partial q_i} + \frac{\partial(\rho \dot{p}_i)}{\partial p_i} \right]$$

$$= - \sum \left[\frac{\partial \rho}{\partial q_i} \dot{q}_i + \frac{\partial \rho}{\partial p_i} \dot{p}_i + \rho \left(\frac{\partial \dot{q}_i}{\partial q_i} + \frac{\partial \dot{p}_i}{\partial p_i} \right) \right].$$

- According to $\dot{q}_i = \frac{\partial H}{\partial p_i}$, $\dot{p}_i = -\frac{\partial H}{\partial q_i}$, we have

$$\frac{\partial^2 H}{\partial p_i \partial q_i} = \frac{\partial^2 H}{\partial q_i \partial p_i} \Rightarrow -\frac{\partial \dot{p}_i}{\partial p_i} = \frac{\partial \dot{q}_i}{\partial q_i}.$$

$$\therefore \frac{\partial \rho}{\partial t} + \sum \left(\frac{\partial \rho}{\partial q_i} \dot{q}_i + \frac{\partial \rho}{\partial p_i} \dot{p}_i \right) = 0, \text{ or } \frac{d\rho}{dt} = 0.$$

Liouville's theorem.

Phase space and Liouville's theorem

- On the other hand, the net gain in the fixed volume:

$$\frac{\partial \rho}{\partial t} dt d\Omega.$$

- $$\therefore - \sum \left[\frac{\partial(\rho \dot{q}_i)}{\partial q_i} + \frac{\partial(\rho \dot{p}_i)}{\partial p_i} \right] dt d\Omega = \frac{\partial \rho}{\partial t} dt d\Omega$$

$$\Rightarrow \frac{\partial \rho}{\partial t} = - \sum \left[\frac{\partial(\rho \dot{q}_i)}{\partial q_i} + \frac{\partial(\rho \dot{p}_i)}{\partial p_i} \right]$$

$$= - \sum \left[\frac{\partial \rho}{\partial q_i} \dot{q}_i + \frac{\partial \rho}{\partial p_i} \dot{p}_i + \rho \left(\frac{\partial \dot{q}_i}{\partial q_i} + \frac{\partial \dot{p}_i}{\partial p_i} \right) \right].$$

- According to $\dot{q}_i = \frac{\partial H}{\partial p_i}$, $\dot{p}_i = -\frac{\partial H}{\partial q_i}$, we have

$$\frac{\partial^2 H}{\partial p_i \partial q_i} = \frac{\partial^2 H}{\partial q_i \partial p_i} \Rightarrow -\frac{\partial \dot{p}_i}{\partial p_i} = \frac{\partial \dot{q}_i}{\partial q_i}.$$

$$\therefore \frac{\partial \rho}{\partial t} + \sum \left(\frac{\partial \rho}{\partial q_i} \dot{q}_i + \frac{\partial \rho}{\partial p_i} \dot{p}_i \right) = 0, \text{ or } \frac{d\rho}{dt} = 0.$$

Liouville's theorem.

$$\text{Or } \frac{\partial \rho}{\partial t} = - \sum \left(\frac{\partial \rho}{\partial q_i} \frac{\partial H}{\partial p_i} - \frac{\partial \rho}{\partial p_i} \frac{\partial H}{\partial q_i} \right).$$

Table of contents

1 §9. Ensemble theory

- 9.1 Phase space and Liouville's theorem
- 9.2 The microcanonical ensemble
- 9.3 Thermodynamics in microcanonical ensemble theory
- 9.4 Canonical ensemble
- 9.5 Thermodynamics in canonical ensemble
- 9.6 EOS of real gas
- 9.7 Thermal capacity of solid
- 9.9 Ising model
- 9.10 Grand canonical ensemble
- 9.11 Thermodynamics in grand canonical ensemble
- 9.12 Applications of the grand canonical ensemble

9.2 The microcanonical ensemble

9.2 The microcanonical ensemble

- Redefine ρ (normalized): $\int \rho(q, p, t) d\Omega = 1$.

9.2 The microcanonical ensemble

- Redefine ρ (normalized): $\int \rho(q, p, t) d\Omega = 1$.
- Meaning: distribution function of the micro-states; probability to find the system in $d\Omega$ is: $\rho(q, p, t) d\Omega$.

9.2 The microcanonical ensemble

- Redefine ρ (normalized): $\int \rho(q, p, t) d\Omega = 1$.
- Meaning: distribution function of the micro-states; probability to find the system in $d\Omega$ is: $\rho(q, p, t) d\Omega$.
- For any variable $B(q, p)$, the average value:
$$\bar{B}(t) = \int B(q, p) \rho(q, p, t) d\Omega.$$

9.2 The microcanonical ensemble

- Redefine ρ (normalized): $\int \rho(q, p, t) d\Omega = 1$.
- Meaning: distribution function of the micro-states; probability to find the system in $d\Omega$ is: $\rho(q, p, t) d\Omega$.
- For any variable $B(q, p)$, the average value:
$$\bar{B}(t) = \int B(q, p) \rho(q, p, t) d\Omega.$$

Meaning...

9.2 The microcanonical ensemble

- Redefine ρ (normalized): $\int \rho(q, p, t) d\Omega = 1$.
- Meaning: distribution function of the micro-states; probability to find the system in $d\Omega$ is: $\rho(q, p, t) d\Omega$.
- For any variable $B(q, p)$, the average value:
$$\bar{B}(t) = \int B(q, p) \rho(q, p, t) d\Omega.$$

Meaning...
- For quantum system, $\rho_s(t)$ represents the probability for system being state s at time t .

9.2 The microcanonical ensemble

- Redefine ρ (normalized): $\int \rho(q, p, t) d\Omega = 1$.
- Meaning: distribution function of the micro-states; probability to find the system in $d\Omega$ is: $\rho(q, p, t) d\Omega$.
- For any variable $B(q, p)$, the average value:
$$\bar{B}(t) = \int B(q, p) \rho(q, p, t) d\Omega.$$

Meaning...
- For quantum system, $\rho_s(t)$ represents the probability for system being state s at time t .
$$\sum \rho_s(t) = 1;$$

9.2 The microcanonical ensemble

- Redefine ρ (normalized): $\int \rho(q, p, t) d\Omega = 1$.
- Meaning: distribution function of the micro-states; probability to find the system in $d\Omega$ is: $\rho(q, p, t) d\Omega$.
- For any variable $B(q, p)$, the average value:
$$\bar{B}(t) = \int B(q, p) \rho(q, p, t) d\Omega.$$

Meaning...
- For quantum system, $\rho_s(t)$ represents the probability for system being state s at time t .
$$\sum \rho_s(t) = 1; \quad \bar{B}(t) = \sum \rho_s(t) B.$$

9.2 The microcanonical ensemble

- Redefine ρ (normalized): $\int \rho(q, p, t) d\Omega = 1$.
- Meaning: distribution function of the micro-states; probability to find the system in $d\Omega$ is: $\rho(q, p, t) d\Omega$.
- For any variable $B(q, p)$, the average value:
$$\bar{B}(t) = \int B(q, p) \rho(q, p, t) d\Omega.$$

Meaning...

- For quantum system, $\rho_s(t)$ represents the probability for system being state s at time t .
$$\sum \rho_s(t) = 1; \bar{B}(t) = \sum \rho_s(t) B.$$
- Core task: distribution function ρ .

The microcanonical ensemble

- Consider isolated system in equilibrium, with distribution function ρ .

The microcanonical ensemble

- Consider isolated system in equilibrium, with distribution function ρ .
- Equilibrium $\rightarrow \frac{\partial \rho}{\partial t} = 0$;

The microcanonical ensemble

- Consider isolated system in equilibrium, with distribution function ρ .
- Equilibrium $\rightarrow \frac{\partial \rho}{\partial t} = 0$;
Liouville's theorem $\rightarrow \frac{d\rho}{dt} = 0$.

The microcanonical ensemble

- Consider isolated system in equilibrium, with distribution function ρ .
- Equilibrium $\rightarrow \frac{\partial \rho}{\partial t} = 0$;
Liouville's theorem $\rightarrow \frac{d\rho}{dt} = 0$.
- A natural **assumption**: $\rho(q, p) = \text{Const.}$

The microcanonical ensemble

- Consider isolated system in equilibrium, with distribution function ρ .
- Equilibrium $\rightarrow \frac{\partial \rho}{\partial t} = 0$;
Liouville's theorem $\rightarrow \frac{d\rho}{dt} = 0$.
- A natural **assumption**: $\rho(q, p) = \text{Const.}$
- As there is no absolute isolated system, it has more or less energy change with the environment.

The microcanonical ensemble

- Consider isolated system in equilibrium, with distribution function ρ .
- Equilibrium $\rightarrow \frac{\partial \rho}{\partial t} = 0$;
Liouville's theorem $\rightarrow \frac{d\rho}{dt} = 0$.
- A natural **assumption**: $\rho(q, p) = \text{Const.}$
- As there is no absolute isolated system, it has more or less energy change with the environment. Energy of the system is not an absolutely fixed value, but a range: $(E, E + \Delta E)$.

The microcanonical ensemble

- Consider isolated system in equilibrium, with distribution function ρ .
- Equilibrium $\rightarrow \frac{\partial \rho}{\partial t} = 0$;
Liouville's theorem $\rightarrow \frac{d\rho}{dt} = 0$.
- A natural **assumption**: $\rho(q, p) = \text{Const.}$
- As there is no absolute isolated system, it has more or less energy change with the environment. Energy of the system is not an absolutely fixed value, but a range: $(E, E + \Delta E)$.

The assumption goes to (equal probability principle):

$$\rho(q, p) = \begin{cases} \text{const.}, & E \leq H \leq E + \Delta E, \\ 0, & \text{others.} \end{cases}$$

The microcanonical ensemble

- For quantum system, $\rho_s = \frac{1}{\Omega}$, where Ω is the total number of micro-states of the system.

The microcanonical ensemble

- For quantum system, $\rho_s = \frac{1}{\Omega}$, where Ω is the total number of micro-states of the system.
- Consider the classical system as the limit of quantum system, the number of micro-state of the classical system: $\Omega = \frac{1}{N!} \frac{1}{h^{Nr}} \int_{E \leq H(q,p) \leq E + \Delta E} d\Omega$.

The microcanonical ensemble

- For quantum system, $\rho_s = \frac{1}{\Omega}$, where Ω is the total number of micro-states of the system.
- Consider the classical system as the limit of quantum system, the number of micro-state of the classical system: $\Omega = \frac{1}{N!} \frac{1}{h^{Nr}} \int_{E \leq H(q,p) \leq E + \Delta E} d\Omega$.
- For multiple kinds of particles:

$$\Omega = \frac{1}{\prod N_i!} \frac{1}{h^{Nr}} \int_{E \leq H(q,p) \leq E + \Delta E} d\Omega.$$

The microcanonical ensemble

- For quantum system, $\rho_s = \frac{1}{\Omega}$, where Ω is the total number of micro-states of the system.
- Consider the classical system as the limit of quantum system, the number of micro-state of the classical system: $\Omega = \frac{1}{N!} \frac{1}{h^{Nr}} \int_{E \leq H(q,p) \leq E + \Delta E} d\Omega$.
- For multiple kinds of particles:

$$\Omega = \frac{1}{\prod N_i!} \frac{1}{h^{Nr}} \int_{E \leq H(q,p) \leq E + \Delta E} d\Omega.$$
- “The most probable distribution” is different from the ensemble average, in concept.

The microcanonical ensemble

- For quantum system, $\rho_s = \frac{1}{\Omega}$, where Ω is the total number of micro-states of the system.
- Consider the classical system as the limit of quantum system, the number of micro-state of the classical system: $\Omega = \frac{1}{N!} \frac{1}{h^{Nr}} \int_{E \leq H(q,p) \leq E + \Delta E} d\Omega$.
- For multiple kinds of particles:

$$\Omega = \frac{1}{\prod N_i!} \frac{1}{h^{Nr}} \int_{E \leq H(q,p) \leq E + \Delta E} d\Omega.$$
- “The most probable distribution” is different from the ensemble average, in concept. But if the fluctuation is small, $\frac{\bar{B}^2 - (\bar{B})^2}{(\bar{B})^2} \ll 1$, both are equal.

Table of contents

1 §9. Ensemble theory

- 9.1 Phase space and Liouville's theorem
- 9.2 The microcanonical ensemble
- 9.3 Thermodynamics in microcanonical ensemble theory
- 9.4 Canonical ensemble
- 9.5 Thermodynamics in canonical ensemble
- 9.6 EOS of real gas
- 9.7 Thermal capacity of solid
- 9.9 Ising model
- 9.10 Grand canonical ensemble
- 9.11 Thermodynamics in grand canonical ensemble
- 9.12 Applications of the grand canonical ensemble

9.3 Thermodynamics in microcanonical ensemble theory

9.3 Thermodynamics in microcanonical ensemble theory

- For a given system (N, E, V) , the number of states: $\Omega(N, E, V)$.

9.3 Thermodynamics in microcanonical ensemble theory

- For a given system (N, E, V) , the number of states: $\Omega(N, E, V)$.
- Consider an isolated system $A^{(0)}$: $(N^{(0)}, E^{(0)}, V^{(0)})$, the number of states: $\Omega^{(0)}(N^{(0)}, E^{(0)}, V^{(0)})$.

9.3 Thermodynamics in microcanonical ensemble theory

- For a given system (N, E, V) , the number of states: $\Omega(N, E, V)$.
- Consider an isolated system $A^{(0)}$: $(N^{(0)}, E^{(0)}, V^{(0)})$, the number of states: $\Omega^{(0)}(N^{(0)}, E^{(0)}, V^{(0)})$.

Divide it into two sub-systems: A_1 : (N_1, E_1, V_1) , A_2 : (N_2, E_2, V_2) , with $\Omega_1(N_1, E_1, V_1)$, $\Omega_2(N_2, E_2, V_2)$.

9.3 Thermodynamics in microcanonical ensemble theory

- For a given system (N, E, V) , the number of states: $\Omega(N, E, V)$.

- Consider an isolated system $A^{(0)}$: $(N^{(0)}, E^{(0)}, V^{(0)})$, the number of states: $\Omega^{(0)}(N^{(0)}, E^{(0)}, V^{(0)})$.

Divide it into two sub-systems: A_1 : (N_1, E_1, V_1) , A_2 : (N_2, E_2, V_2) , with $\Omega_1(N_1, E_1, V_1)$, $\Omega_2(N_2, E_2, V_2)$.

- Obviously, $N^{(0)} = N_1 + N_2$, $E^{(0)} = E_1 + E_2$,
 $V^{(0)} = V_1 + V_2$,

9.3 Thermodynamics in microcanonical ensemble theory

- For a given system (N, E, V) , the number of states: $\Omega(N, E, V)$.
- Consider an isolated system $A^{(0)}$: $(N^{(0)}, E^{(0)}, V^{(0)})$, the number of states: $\Omega^{(0)}(N^{(0)}, E^{(0)}, V^{(0)})$.

Divide it into two sub-systems: A_1 : (N_1, E_1, V_1) , A_2 : (N_2, E_2, V_2) , with $\Omega_1(N_1, E_1, V_1)$, $\Omega_2(N_2, E_2, V_2)$.

- Obviously, $N^{(0)} = N_1 + N_2$, $E^{(0)} = E_1 + E_2$, $V^{(0)} = V_1 + V_2$, and $\Omega^{(0)} = \Omega_1 \cdot \Omega_2$.

9.3 Thermodynamics in microcanonical ensemble theory

- For a given system (N, E, V) , the number of states: $\Omega(N, E, V)$.
- Consider an isolated system $A^{(0)}$: $(N^{(0)}, E^{(0)}, V^{(0)})$, the number of states: $\Omega^{(0)}(N^{(0)}, E^{(0)}, V^{(0)})$.
Divide it into two sub-systems: A_1 : (N_1, E_1, V_1) , A_2 : (N_2, E_2, V_2) , with $\Omega_1(N_1, E_1, V_1)$, $\Omega_2(N_2, E_2, V_2)$.
- Obviously, $N^{(0)} = N_1 + N_2$, $E^{(0)} = E_1 + E_2$, $V^{(0)} = V_1 + V_2$, and $\Omega^{(0)} = \Omega_1 \cdot \Omega_2$.
- Keep (N_1, N_2, V_1, V_2) , let (E_1, E_2) vary, to see the condition to get equilibrium:

9.3 Thermodynamics in microcanonical ensemble theory

- For a given system (N, E, V) , the number of states: $\Omega(N, E, V)$.
- Consider an isolated system $A^{(0)}$: $(N^{(0)}, E^{(0)}, V^{(0)})$, the number of states: $\Omega^{(0)}(N^{(0)}, E^{(0)}, V^{(0)})$.

Divide it into two sub-systems: A_1 : (N_1, E_1, V_1) , A_2 : (N_2, E_2, V_2) , with $\Omega_1(N_1, E_1, V_1)$, $\Omega_2(N_2, E_2, V_2)$.

- Obviously, $N^{(0)} = N_1 + N_2$, $E^{(0)} = E_1 + E_2$, $V^{(0)} = V_1 + V_2$, and $\Omega^{(0)} = \Omega_1 \cdot \Omega_2$.
- Keep (N_1, N_2, V_1, V_2) , let (E_1, E_2) vary, to see the condition to get equilibrium: $\frac{\partial \Omega^{(0)}}{\partial E_1} = 0$,

9.3 Thermodynamics in microcanonical ensemble theory

- For a given system (N, E, V) , the number of states: $\Omega(N, E, V)$.
- Consider an isolated system $A^{(0)}$: $(N^{(0)}, E^{(0)}, V^{(0)})$, the number of states: $\Omega^{(0)}(N^{(0)}, E^{(0)}, V^{(0)})$.
Divide it into two sub-systems: A_1 : (N_1, E_1, V_1) , A_2 : (N_2, E_2, V_2) , with $\Omega_1(N_1, E_1, V_1)$, $\Omega_2(N_2, E_2, V_2)$.
- Obviously, $N^{(0)} = N_1 + N_2$, $E^{(0)} = E_1 + E_2$, $V^{(0)} = V_1 + V_2$, and $\Omega^{(0)} = \Omega_1 \cdot \Omega_2$.
- Keep (N_1, N_2, V_1, V_2) , let (E_1, E_2) vary, to see the condition to get equilibrium: $\frac{\partial \Omega^{(0)}}{\partial E_1} = 0$, i.e., with the choosing of (E_1, E_2) , number of states has maximum because of the equal probability principle.

Thermodynamics in microcanonical ensemble theory

- $\frac{\partial \Omega^{(0)}}{\partial E_1}$

Thermodynamics in microcanonical ensemble theory

- $$\frac{\partial \Omega^{(0)}}{\partial E_1} = \frac{\partial(\Omega_1 \Omega_2)}{\partial E_1}$$

Thermodynamics in microcanonical ensemble theory

- $$\frac{\partial \Omega^{(0)}}{\partial E_1} = \frac{\partial(\Omega_1 \Omega_2)}{\partial E_1} = \frac{\partial \Omega_1}{\partial E_1} \Omega_2 + \Omega_1 \frac{\partial \Omega_2}{\partial E_1}$$

Thermodynamics in microcanonical ensemble theory

- $$\frac{\partial \Omega^{(0)}}{\partial E_1} = \frac{\partial(\Omega_1 \Omega_2)}{\partial E_1} = \frac{\partial \Omega_1}{\partial E_1} \Omega_2 + \Omega_1 \frac{\partial \Omega_2}{\partial E_1} = \frac{\partial \Omega_1}{\partial E_1} \Omega_2 + \Omega_1 \frac{\partial \Omega_2}{\partial E_2} \frac{\partial E_2}{\partial E_1}$$

Thermodynamics in microcanonical ensemble theory

- $$\begin{aligned}\frac{\partial \Omega^{(0)}}{\partial E_1} &= \frac{\partial(\Omega_1 \Omega_2)}{\partial E_1} = \frac{\partial \Omega_1}{\partial E_1} \Omega_2 + \Omega_1 \frac{\partial \Omega_2}{\partial E_1} = \frac{\partial \Omega_1}{\partial E_1} \Omega_2 + \Omega_1 \frac{\partial \Omega_2}{\partial E_2} \frac{\partial E_2}{\partial E_1} \\ &= \frac{\partial \Omega_1}{\partial E_1} \Omega_2 - \Omega_1 \frac{\partial \Omega_2}{\partial E_2}\end{aligned}$$

Thermodynamics in microcanonical ensemble theory

- $$\begin{aligned}\frac{\partial \Omega^{(0)}}{\partial E_1} &= \frac{\partial(\Omega_1 \Omega_2)}{\partial E_1} = \frac{\partial \Omega_1}{\partial E_1} \Omega_2 + \Omega_1 \frac{\partial \Omega_2}{\partial E_1} = \frac{\partial \Omega_1}{\partial E_1} \Omega_2 + \Omega_1 \frac{\partial \Omega_2}{\partial E_2} \frac{\partial E_2}{\partial E_1} \\ &= \frac{\partial \Omega_1}{\partial E_1} \Omega_2 - \Omega_1 \frac{\partial \Omega_2}{\partial E_2} = 0;\end{aligned}$$

Thermodynamics in microcanonical ensemble theory

- $$\begin{aligned}\frac{\partial \Omega^{(0)}}{\partial E_1} &= \frac{\partial(\Omega_1 \Omega_2)}{\partial E_1} = \frac{\partial \Omega_1}{\partial E_1} \Omega_2 + \Omega_1 \frac{\partial \Omega_2}{\partial E_1} = \frac{\partial \Omega_1}{\partial E_1} \Omega_2 + \Omega_1 \frac{\partial \Omega_2}{\partial E_2} \frac{\partial E_2}{\partial E_1} \\ &= \frac{\partial \Omega_1}{\partial E_1} \Omega_2 - \Omega_1 \frac{\partial \Omega_2}{\partial E_2} = 0; \Rightarrow \frac{\partial \ln \Omega_1}{\partial E_1} = \frac{\partial \ln \Omega_2}{\partial E_2}.\end{aligned}$$

Thermodynamics in microcanonical ensemble theory

- $\frac{\partial \Omega^{(0)}}{\partial E_1} = \frac{\partial(\Omega_1 \Omega_2)}{\partial E_1} = \frac{\partial \Omega_1}{\partial E_1} \Omega_2 + \Omega_1 \frac{\partial \Omega_2}{\partial E_1} = \frac{\partial \Omega_1}{\partial E_1} \Omega_2 + \Omega_1 \frac{\partial \Omega_2}{\partial E_2} \frac{\partial E_2}{\partial E_1}$
 $= \frac{\partial \Omega_1}{\partial E_1} \Omega_2 - \Omega_1 \frac{\partial \Omega_2}{\partial E_2} = 0; \Rightarrow \frac{\partial \ln \Omega_1}{\partial E_1} = \frac{\partial \ln \Omega_2}{\partial E_2}.$
- More complete:
$$\left(\frac{\partial \ln \Omega_1(N_1, E_1, V_1)}{\partial E_1} \right)_{N_1, V_1} = \left(\frac{\partial \ln \Omega_2(N_2, E_2, V_2)}{\partial E_2} \right)_{N_2, V_2}.$$

Thermodynamics in microcanonical ensemble theory

- $$\frac{\partial \Omega^{(0)}}{\partial E_1} = \frac{\partial(\Omega_1 \Omega_2)}{\partial E_1} = \frac{\partial \Omega_1}{\partial E_1} \Omega_2 + \Omega_1 \frac{\partial \Omega_2}{\partial E_1} = \frac{\partial \Omega_1}{\partial E_1} \Omega_2 + \Omega_1 \frac{\partial \Omega_2}{\partial E_2} \frac{\partial E_2}{\partial E_1}$$

$$= \frac{\partial \Omega_1}{\partial E_1} \Omega_2 - \Omega_1 \frac{\partial \Omega_2}{\partial E_2} = 0; \Rightarrow \frac{\partial \ln \Omega_1}{\partial E_1} = \frac{\partial \ln \Omega_2}{\partial E_2}.$$
- More complete:**

$$\left(\frac{\partial \ln \Omega_1(N_1, E_1, V_1)}{\partial E_1} \right)_{N_1, V_1} = \left(\frac{\partial \ln \Omega_2(N_2, E_2, V_2)}{\partial E_2} \right)_{N_2, V_2}.$$

Set $\beta = \left(\frac{\partial \ln \Omega(N, E, V)}{\partial E} \right)_{N, V}$, $\beta_1 = \beta_2$.

Thermodynamics in microcanonical ensemble theory

- $$\frac{\partial \Omega^{(0)}}{\partial E_1} = \frac{\partial(\Omega_1 \Omega_2)}{\partial E_1} = \frac{\partial \Omega_1}{\partial E_1} \Omega_2 + \Omega_1 \frac{\partial \Omega_2}{\partial E_1} = \frac{\partial \Omega_1}{\partial E_1} \Omega_2 + \Omega_1 \frac{\partial \Omega_2}{\partial E_2} \frac{\partial E_2}{\partial E_1}$$

$$= \frac{\partial \Omega_1}{\partial E_1} \Omega_2 - \Omega_1 \frac{\partial \Omega_2}{\partial E_2} = 0; \Rightarrow \frac{\partial \ln \Omega_1}{\partial E_1} = \frac{\partial \ln \Omega_2}{\partial E_2}.$$
- More complete:

$$\left(\frac{\partial \ln \Omega_1(N_1, E_1, V_1)}{\partial E_1} \right)_{N_1, V_1} = \left(\frac{\partial \ln \Omega_2(N_2, E_2, V_2)}{\partial E_2} \right)_{N_2, V_2}.$$

Set $\beta = \left(\frac{\partial \ln \Omega(N, E, V)}{\partial E} \right)_{N, V}$, $\beta_1 = \beta_2$.
- Basic function (3.2.7): $dU = TdS - pdV + \mu dN$,

Thermodynamics in microcanonical ensemble theory

- $$\frac{\partial \Omega^{(0)}}{\partial E_1} = \frac{\partial(\Omega_1 \Omega_2)}{\partial E_1} = \frac{\partial \Omega_1}{\partial E_1} \Omega_2 + \Omega_1 \frac{\partial \Omega_2}{\partial E_1} = \frac{\partial \Omega_1}{\partial E_1} \Omega_2 + \Omega_1 \frac{\partial \Omega_2}{\partial E_2} \frac{\partial E_2}{\partial E_1}$$

$$= \frac{\partial \Omega_1}{\partial E_1} \Omega_2 - \Omega_1 \frac{\partial \Omega_2}{\partial E_2} = 0; \Rightarrow \frac{\partial \ln \Omega_1}{\partial E_1} = \frac{\partial \ln \Omega_2}{\partial E_2}.$$
- More complete:

$$\left(\frac{\partial \ln \Omega_1(N_1, E_1, V_1)}{\partial E_1} \right)_{N_1, V_1} = \left(\frac{\partial \ln \Omega_2(N_2, E_2, V_2)}{\partial E_2} \right)_{N_2, V_2}.$$

Set $\beta = \left(\frac{\partial \ln \Omega(N, E, V)}{\partial E} \right)_{N, V}$, $\beta_1 = \beta_2$.
- Basic function (3.2.7): $dU = TdS - pdV + \mu dN$, for fixed (N, V) , the condition for equilibrium is $T_1 = T_2$, i.e., $\frac{1}{T_1} = \left(\frac{\partial S_1}{\partial U_1} \right)_{N_1, V_1} = \left(\frac{\partial S_2}{\partial U_2} \right)_{N_2, V_2} = \frac{1}{T_2}$.

Thermodynamics in microcanonical ensemble theory

- $$\frac{\partial \Omega^{(0)}}{\partial E_1} = \frac{\partial(\Omega_1 \Omega_2)}{\partial E_1} = \frac{\partial \Omega_1}{\partial E_1} \Omega_2 + \Omega_1 \frac{\partial \Omega_2}{\partial E_1} = \frac{\partial \Omega_1}{\partial E_1} \Omega_2 + \Omega_1 \frac{\partial \Omega_2}{\partial E_2} \frac{\partial E_2}{\partial E_1}$$

$$= \frac{\partial \Omega_1}{\partial E_1} \Omega_2 - \Omega_1 \frac{\partial \Omega_2}{\partial E_2} = 0; \Rightarrow \frac{\partial \ln \Omega_1}{\partial E_1} = \frac{\partial \ln \Omega_2}{\partial E_2}.$$
- More complete:

$$\left(\frac{\partial \ln \Omega_1(N_1, E_1, V_1)}{\partial E_1} \right)_{N_1, V_1} = \left(\frac{\partial \ln \Omega_2(N_2, E_2, V_2)}{\partial E_2} \right)_{N_2, V_2}.$$
 Set $\beta = \left(\frac{\partial \ln \Omega(N, E, V)}{\partial E} \right)_{N, V}$, $\beta_1 = \beta_2$.
- Basic function (3.2.7): $dU = TdS - pdV + \mu dN$, for fixed (N, V) , the condition for equilibrium is $T_1 = T_2$, i.e., $\frac{1}{T_1} = \left(\frac{\partial S_1}{\partial U_1} \right)_{N_1, V_1} = \left(\frac{\partial S_2}{\partial U_2} \right)_{N_2, V_2} = \frac{1}{T_2}$.
- $\therefore \beta = \frac{1}{kT}$, where k is the Boltzmann constant, which can be found by taking into an example, like ideal gas.

Thermodynamics in microcanonical ensemble theory

- $$\frac{\partial \Omega^{(0)}}{\partial E_1} = \frac{\partial(\Omega_1 \Omega_2)}{\partial E_1} = \frac{\partial \Omega_1}{\partial E_1} \Omega_2 + \Omega_1 \frac{\partial \Omega_2}{\partial E_1} = \frac{\partial \Omega_1}{\partial E_1} \Omega_2 + \Omega_1 \frac{\partial \Omega_2}{\partial E_2} \frac{\partial E_2}{\partial E_1}$$

$$= \frac{\partial \Omega_1}{\partial E_1} \Omega_2 - \Omega_1 \frac{\partial \Omega_2}{\partial E_2} = 0; \Rightarrow \frac{\partial \ln \Omega_1}{\partial E_1} = \frac{\partial \ln \Omega_2}{\partial E_2}.$$
- More complete:

$$\left(\frac{\partial \ln \Omega_1(N_1, E_1, V_1)}{\partial E_1} \right)_{N_1, V_1} = \left(\frac{\partial \ln \Omega_2(N_2, E_2, V_2)}{\partial E_2} \right)_{N_2, V_2}.$$
 Set $\beta = \left(\frac{\partial \ln \Omega(N, E, V)}{\partial E} \right)_{N, V}$, $\beta_1 = \beta_2$.
- Basic function (3.2.7): $dU = TdS - pdV + \mu dN$, for fixed (N, V) , the condition for equilibrium is $T_1 = T_2$, i.e., $\frac{1}{T_1} = \left(\frac{\partial S_1}{\partial U_1} \right)_{N_1, V_1} = \left(\frac{\partial S_2}{\partial U_2} \right)_{N_2, V_2} = \frac{1}{T_2}$.
- $\therefore \beta = \frac{1}{kT}$, where k is the Boltzmann constant, which can be found by taking into an example, like ideal gas.
- As $\frac{1}{T} = k\beta$, $S = k \ln \Omega$.

Thermodynamics in microcanonical ensemble theory

- If between sub-systems A_1, A_2 , not only energy, but also particles and volume can exchange, we have:

Thermodynamics in microcanonical ensemble theory

- If between sub-systems A_1, A_2 , not only energy, but also particles and volume can exchange, we have:

$$\left(\frac{\partial \ln \Omega_1}{\partial E_1}\right)_{N_1, V_1} = \left(\frac{\partial \ln \Omega_2}{\partial E_2}\right)_{N_2, V_2} \rightarrow \beta_1 = \beta_2,$$

$$\left(\frac{\partial \ln \Omega_1}{\partial V_1}\right)_{N_1, E_1} = \left(\frac{\partial \ln \Omega_2}{\partial V_2}\right)_{N_2, E_2} \rightarrow \gamma_1 = \gamma_2,$$

$$\left(\frac{\partial \ln \Omega_1}{\partial N_1}\right)_{E_1, V_1} = \left(\frac{\partial \ln \Omega_2}{\partial N_2}\right)_{E_2, V_2} \rightarrow \alpha_1 = \alpha_2.$$

Thermodynamics in microcanonical ensemble theory

- If between sub-systems A_1, A_2 , not only energy, but also particles and volume can exchange, we have:

$$\left(\frac{\partial \ln \Omega_1}{\partial E_1}\right)_{N_1, V_1} = \left(\frac{\partial \ln \Omega_2}{\partial E_2}\right)_{N_2, V_2} \rightarrow \beta_1 = \beta_2,$$

$$\left(\frac{\partial \ln \Omega_1}{\partial V_1}\right)_{N_1, E_1} = \left(\frac{\partial \ln \Omega_2}{\partial V_2}\right)_{N_2, E_2} \rightarrow \gamma_1 = \gamma_2,$$

$$\left(\frac{\partial \ln \Omega_1}{\partial N_1}\right)_{E_1, V_1} = \left(\frac{\partial \ln \Omega_2}{\partial N_2}\right)_{E_2, V_2} \rightarrow \alpha_1 = \alpha_2.$$

- Then the complete differential:

$$d \ln \Omega = \beta dE + \gamma dV + \alpha dN.$$

Thermodynamics in microcanonical ensemble theory

- If between sub-systems A_1, A_2 , not only energy, but also particles and volume can exchange, we have:

$$\left(\frac{\partial \ln \Omega_1}{\partial E_1}\right)_{N_1, V_1} = \left(\frac{\partial \ln \Omega_2}{\partial E_2}\right)_{N_2, V_2} \rightarrow \beta_1 = \beta_2,$$

$$\left(\frac{\partial \ln \Omega_1}{\partial V_1}\right)_{N_1, E_1} = \left(\frac{\partial \ln \Omega_2}{\partial V_2}\right)_{N_2, E_2} \rightarrow \gamma_1 = \gamma_2,$$

$$\left(\frac{\partial \ln \Omega_1}{\partial N_1}\right)_{E_1, V_1} = \left(\frac{\partial \ln \Omega_2}{\partial N_2}\right)_{E_2, V_2} \rightarrow \alpha_1 = \alpha_2.$$

- Then the complete differential:

$$d \ln \Omega = \beta dE + \gamma dV + \alpha dN.$$

$$\text{Compare with (3.2.7): } dS = \frac{1}{T}dU + \frac{p}{T}dV - \frac{\mu}{T}dN.$$

Thermodynamics in microcanonical ensemble theory

- If between sub-systems A_1, A_2 , not only energy, but also particles and volume can exchange, we have:

$$\left(\frac{\partial \ln \Omega_1}{\partial E_1}\right)_{N_1, V_1} = \left(\frac{\partial \ln \Omega_2}{\partial E_2}\right)_{N_2, V_2} \rightarrow \beta_1 = \beta_2,$$

$$\left(\frac{\partial \ln \Omega_1}{\partial V_1}\right)_{N_1, E_1} = \left(\frac{\partial \ln \Omega_2}{\partial V_2}\right)_{N_2, E_2} \rightarrow \gamma_1 = \gamma_2,$$

$$\left(\frac{\partial \ln \Omega_1}{\partial N_1}\right)_{E_1, V_1} = \left(\frac{\partial \ln \Omega_2}{\partial N_2}\right)_{E_2, V_2} \rightarrow \alpha_1 = \alpha_2.$$

- Then the complete differential:

$$d \ln \Omega = \beta dE + \gamma dV + \alpha dN.$$

Compare with (3.2.7): $dS = \frac{1}{T}dU + \frac{p}{T}dV - \frac{\mu}{T}dN.$

$$\Rightarrow \gamma = \frac{p}{kT}, \alpha = -\frac{\mu}{kT}.$$

Thermodynamics in microcanonical ensemble theory

- If between sub-systems A_1, A_2 , not only energy, but also particles and volume can exchange, we have:

$$\left(\frac{\partial \ln \Omega_1}{\partial E_1}\right)_{N_1, V_1} = \left(\frac{\partial \ln \Omega_2}{\partial E_2}\right)_{N_2, V_2} \rightarrow \beta_1 = \beta_2,$$

$$\left(\frac{\partial \ln \Omega_1}{\partial V_1}\right)_{N_1, E_1} = \left(\frac{\partial \ln \Omega_2}{\partial V_2}\right)_{N_2, E_2} \rightarrow \gamma_1 = \gamma_2,$$

$$\left(\frac{\partial \ln \Omega_1}{\partial N_1}\right)_{E_1, V_1} = \left(\frac{\partial \ln \Omega_2}{\partial N_2}\right)_{E_2, V_2} \rightarrow \alpha_1 = \alpha_2.$$

- Then the complete differential:

$$d \ln \Omega = \beta dE + \gamma dV + \alpha dN.$$

$$\text{Compare with (3.2.7): } dS = \frac{1}{T}dU + \frac{p}{T}dV - \frac{\mu}{T}dN.$$

$$\Rightarrow \gamma = \frac{p}{kT}, \alpha = -\frac{\mu}{kT}.$$

- The condition for equilibrium: $T_1 = T_2$, $p_1 = p_2$, $\mu_1 = \mu_2$; also showed the meaning of α , β , and γ .

Thermodynamics in microcanonical ensemble theory

- Apply to classical ideal gas:

Thermodynamics in microcanonical ensemble theory

- Apply to classical ideal gas:
- $d\Omega = dq_1 \dots dq_f dp_1 \dots dp_f$

Thermodynamics in microcanonical ensemble theory

- Apply to classical ideal gas:
- $d\Omega = dq_1 \dots dq_f dp_1 \dots dp_f$
 $= dq_{11} dq_{12} dq_{13} \dots dq_{N3} dp_1 \dots dp_f,$

Thermodynamics in microcanonical ensemble theory

- Apply to classical ideal gas:
- $d\Omega = dq_1 \dots dq_f dp_1 \dots dp_f$
 $= dq_{11} dq_{12} dq_{13} \dots dq_{N3} dp_1 \dots dp_f,$
 $\therefore \Omega = \frac{1}{N!} \frac{1}{h^{Nr}} \int d\Omega = CV^N$, where C does not depend on q .

Thermodynamics in microcanonical ensemble theory

- Apply to classical ideal gas:
- $d\Omega = dq_1 \dots dq_f dp_1 \dots dp_f$
 $= dq_{11} dq_{12} dq_{13} \dots dq_{N3} dp_1 \dots dp_f,$
 $\therefore \Omega = \frac{1}{N!} \frac{1}{h^{Nr}} \int d\Omega = CV^N$, where C does not depend on q .
- $\frac{p}{kT} = \gamma$

Thermodynamics in microcanonical ensemble theory

- Apply to classical ideal gas:
- $d\Omega = dq_1 \dots dq_f dp_1 \dots dp_f$
 $= dq_{11} dq_{12} dq_{13} \dots dq_{N3} dp_1 \dots dp_f,$
 $\therefore \Omega = \frac{1}{N!} \frac{1}{h^{Nr}} \int d\Omega = CV^N$, where C does not depend on q .
- $\frac{p}{kT} = \gamma = \frac{\partial \ln \Omega}{\partial V}$

Thermodynamics in microcanonical ensemble theory

- Apply to classical ideal gas:
- $d\Omega = dq_1 \dots dq_f dp_1 \dots dp_f$
 $= dq_{11} dq_{12} dq_{13} \dots dq_{N3} dp_1 \dots dp_f,$
 $\therefore \Omega = \frac{1}{N!} \frac{1}{h^{Nr}} \int d\Omega = CV^N$, where C does not depend on q .
- $\frac{p}{kT} = \gamma = \frac{\partial \ln \Omega}{\partial V} = \frac{\partial \ln V^N}{\partial V}$

Thermodynamics in microcanonical ensemble theory

- Apply to classical ideal gas:
- $d\Omega = dq_1 \dots dq_f dp_1 \dots dp_f$
 $= dq_{11} dq_{12} dq_{13} \dots dq_{N3} dp_1 \dots dp_f,$
 $\therefore \Omega = \frac{1}{N!} \frac{1}{h^{Nr}} \int d\Omega = CV^N$, where C does not depend on q .
- $\frac{p}{kT} = \gamma = \frac{\partial \ln \Omega}{\partial V} = \frac{\partial \ln V^N}{\partial V} = \frac{N}{V}.$

Thermodynamics in microcanonical ensemble theory

- Apply to classical ideal gas:
- $d\Omega = dq_1 \dots dq_f dp_1 \dots dp_f$
 $= dq_{11} dq_{12} dq_{13} \dots dq_{N3} dp_1 \dots dp_f,$
 $\therefore \Omega = \frac{1}{N!} \frac{1}{h^{Nr}} \int d\Omega = CV^N$, where C does not depend on q .
- $\frac{p}{kT} = \gamma = \frac{\partial \ln \Omega}{\partial V} = \frac{\partial \ln V^N}{\partial V} = \frac{N}{V}.$
 $\Rightarrow pV = NkT.$

Thermodynamics in microcanonical ensemble theory

- Apply to classical ideal gas:
- $d\Omega = dq_1 \dots dq_f dp_1 \dots dp_f$
 $= dq_{11} dq_{12} dq_{13} \dots dq_{N3} dp_1 \dots dp_f,$
 $\therefore \Omega = \frac{1}{N!} \frac{1}{h^{Nr}} \int d\Omega = CV^N$, where C does not depend on q .
- $\frac{p}{kT} = \gamma = \frac{\partial \ln \Omega}{\partial V} = \frac{\partial \ln V^N}{\partial V} = \frac{N}{V}.$
 $\Rightarrow pV = NkT.$
- Known EOS of ideal gas: $pV = nRT.$

Thermodynamics in microcanonical ensemble theory

- Apply to classical ideal gas:
- $d\Omega = dq_1 \dots dq_f dp_1 \dots dp_f$
 $= dq_{11} dq_{12} dq_{13} \dots dq_{N3} dp_1 \dots dp_f,$
 $\therefore \Omega = \frac{1}{N!} \frac{1}{h^{Nr}} \int d\Omega = CV^N$, where C does not depend on q .
- $\frac{p}{kT} = \gamma = \frac{\partial \ln \Omega}{\partial V} = \frac{\partial \ln V^N}{\partial V} = \frac{N}{V}.$
 $\Rightarrow pV = NkT.$
- Known EOS of ideal gas: $pV = nRT.$
- $\therefore k = \frac{R}{N_0}$ is just the Boltzmann constant.

Thermodynamics in microcanonical ensemble theory

- General procedure in microcanonical ensemble theory:

Thermodynamics in microcanonical ensemble theory

- General procedure in microcanonical ensemble theory:
- 1. $\Omega(N, E, V)$ (Ω as function of (N, E, V));

Thermodynamics in microcanonical ensemble theory

- General procedure in microcanonical ensemble theory:
- 1. $\Omega(N, E, V)$ (Ω as function of (N, E, V));
- 2. $S(N, E, V) = k \ln \Omega(N, E, V)$ (S as function of (N, E, V)),

Thermodynamics in microcanonical ensemble theory

- General procedure in microcanonical ensemble theory:
- 1. $\Omega(N, E, V)$ (Ω as function of (N, E, V));
- 2. $S(N, E, V) = k \ln \Omega(N, E, V)$ (S as function of (N, E, V)),
 $\Rightarrow E(S, V, N)$ (E as function of (S, V, N));

Thermodynamics in microcanonical ensemble theory

- General procedure in microcanonical ensemble theory:
- 1. $\Omega(N, E, V)$ (Ω as function of (N, E, V));
- 2. $S(N, E, V) = k \ln \Omega(N, E, V)$ (S as function of (N, E, V)),
 $\Rightarrow E(S, V, N)$ (E as function of (S, V, N));
- 3. $dE = TdS - pdV + \mu dN$
 $\Rightarrow T = (\frac{\partial E}{\partial S})_{V,N}, p = -(\frac{\partial E}{\partial V})_{S,N},$

Thermodynamics in microcanonical ensemble theory

- General procedure in microcanonical ensemble theory:
- 1. $\Omega(N, E, V)$ (Ω as function of (N, E, V));
- 2. $S(N, E, V) = k \ln \Omega(N, E, V)$ (S as function of (N, E, V)),
 $\Rightarrow E(S, V, N)$ (E as function of (S, V, N));
- 3. $dE = TdS - pdV + \mu dN$
 $\Rightarrow T = (\frac{\partial E}{\partial S})_{V,N}, p = -(\frac{\partial E}{\partial V})_{S,N},$
 $\rightarrow T(S, V, N) \rightarrow S(T, V, N) \rightarrow E(T, V, N)$

Thermodynamics in microcanonical ensemble theory

- General procedure in microcanonical ensemble theory:
 - 1. $\Omega(N, E, V)$ (Ω as function of (N, E, V));
 - 2. $S(N, E, V) = k \ln \Omega(N, E, V)$ (S as function of (N, E, V)),
 $\Rightarrow E(S, V, N)$ (E as function of (S, V, N));
 - 3. $dE = TdS - pdV + \mu dN$
 $\Rightarrow T = (\frac{\partial E}{\partial S})_{V, N}, p = -(\frac{\partial E}{\partial V})_{S, N},$
 $\rightarrow T(S, V, N) \rightarrow S(T, V, N) \rightarrow E(T, V, N)$
 $p(S, V, N) \rightarrow p(T, V, N).$

Application: monatomic ideal gas.

Application: monatomic ideal gas.

- Hamiltonian: $H = \sum_{i=1}^{3N} \frac{p_i^2}{2m}$.

Application: monatomic ideal gas.

- Hamiltonian: $H = \sum_{i=1}^{3N} \frac{p_i^2}{2m}$.
- $\Omega(N, E, V) = \frac{1}{N!h^{3N}} \int_{(E, E+\Delta E)} dq_1 \dots dq_{3N} dp_1 \dots dp_{3N}$

Application: monatomic ideal gas.

- Hamiltonian: $H = \sum_{i=1}^{3N} \frac{p_i^2}{2m}$.
- $\Omega(N, E, V) = \frac{1}{N!h^{3N}} \int_{(E, E+\Delta E)} dq_1 \dots dq_{3N} dp_1 \dots dp_{3N}$
$$= \frac{1}{N!h^{3N}} \left(\int_{H \leq E+\Delta E} d\Omega - \int_{H \leq E} d\Omega \right)$$

Application: monatomic ideal gas.

- Hamiltonian: $H = \sum_{i=1}^{3N} \frac{p_i^2}{2m}$.
- $\Omega(N, E, V) = \frac{1}{N!h^{3N}} \int_{(E, E+\Delta E)} dq_1 \dots dq_{3N} dp_1 \dots dp_{3N}$

$$= \frac{1}{N!h^{3N}} \left(\int_{H \leq E+\Delta E} d\Omega - \int_{H \leq E} d\Omega \right)$$

Provable:

$$\int_{H \leq E} d\Omega = V^N \int_{H \leq E} dp_1 \dots dp_{3N} = V^N \cdot \frac{(2\pi m E)^{\frac{3}{2}N}}{(\frac{3}{2}N)!};$$

Application: monatomic ideal gas.

- Hamiltonian: $H = \sum_{i=1}^{3N} \frac{p_i^2}{2m}$.
- $\Omega(N, E, V) = \frac{1}{N!h^{3N}} \int_{(E, E+\Delta E)} dq_1 \dots dq_{3N} dp_1 \dots dp_{3N}$

$$= \frac{1}{N!h^{3N}} \left(\int_{H \leq E+\Delta E} d\Omega - \int_{H \leq E} d\Omega \right)$$

Provable:

$$\int_{H \leq E} d\Omega = V^N \int_{H \leq E} dp_1 \dots dp_{3N} = V^N \cdot \frac{(2\pi m E)^{\frac{3}{2}N}}{(\frac{3}{2}N)!};$$

- $\Omega(N, E, V) = \left(\frac{V}{h^3}\right)^N \cdot \frac{(2\pi m)^{\frac{3}{2}N}}{N!(\frac{3}{2}N)!} \left[(E + \Delta E)^{\frac{3}{2}N} - E^{\frac{3}{2}N} \right]$

Application: monatomic ideal gas.

- Hamiltonian: $H = \sum_{i=1}^{3N} \frac{p_i^2}{2m}$.
- $\Omega(N, E, V) = \frac{1}{N!h^{3N}} \int_{(E, E+\Delta E)} dq_1 \dots dq_{3N} dp_1 \dots dp_{3N}$

$$= \frac{1}{N!h^{3N}} \left(\int_{H \leq E+\Delta E} d\Omega - \int_{H \leq E} d\Omega \right)$$

Provable:

$$\int_{H \leq E} d\Omega = V^N \int_{H \leq E} dp_1 \dots dp_{3N} = V^N \cdot \frac{(2\pi m E)^{\frac{3}{2}N}}{(\frac{3}{2}N)!};$$

- $\Omega(N, E, V) = \left(\frac{V}{h^3}\right)^N \cdot \frac{(2\pi m)^{\frac{3}{2}N}}{N!(\frac{3}{2}N)!} \left[(E + \Delta E)^{\frac{3}{2}N} - E^{\frac{3}{2}N} \right]$

$$\simeq \frac{3}{2}N \cdot \left(\frac{V}{h^3}\right)^N \cdot \frac{(2\pi m E)^{\frac{3}{2}N}}{N!(\frac{3}{2}N)!} \frac{\Delta E}{E}.$$

Application: monatomic ideal gas. Prove...

- Prove: $\sigma(E) \equiv \int_{H \leq E} dp_1 \dots dp_{3N} = \frac{(2\pi m E)^{\frac{3}{2}N}}{(\frac{3}{2}N)!}$.

Application: monatomic ideal gas. Prove...

- Prove: $\sigma(E) \equiv \int_{H \leq E} dp_1 \dots dp_{3N} = \frac{(2\pi m E)^{\frac{3}{2}N}}{(\frac{3}{2}N)!}$.
- First: $p_i = \sqrt{2mE}x_i \Rightarrow \sigma(E) = (2mE)^{\frac{3}{2}N} \int_{\sum x_i^2 \leq 1} dx_1 \dots dx_{3N} \equiv K(2mE)^{\frac{3}{2}N} \dots (a)$

Application: monatomic ideal gas. Prove...

- Prove: $\sigma(E) \equiv \int_{H \leq E} dp_1 \dots dp_{3N} = \frac{(2\pi m E)^{\frac{3}{2}N}}{(\frac{3}{2}N)!}$.
- First: $p_i = \sqrt{2mE}x_i \Rightarrow \sigma(E) = (2mE)^{\frac{3}{2}N} \int_{\sum x_i^2 \leq 1} dx_1 \dots dx_{3N} \equiv K(2mE)^{\frac{3}{2}N} \dots (a)$
- Second, calculate $A \equiv \int_{-\infty}^{+\infty} e^{-\beta H} dp_1 \dots dp_{3N}$.

Application: monatomic ideal gas. Prove...

- Prove: $\sigma(E) \equiv \int_{H \leq E} dp_1 \dots dp_{3N} = \frac{(2\pi m E)^{\frac{3}{2}N}}{(\frac{3}{2}N)!}$.
- First: $p_i = \sqrt{2mE}x_i \Rightarrow \sigma(E) = (2mE)^{\frac{3}{2}N} \int_{\sum x_i^2 \leq 1} dx_1 \dots dx_{3N} \equiv K(2mE)^{\frac{3}{2}N} \dots (a)$
- Second, calculate $A \equiv \int_{-\infty}^{+\infty} e^{-\beta H} dp_1 \dots dp_{3N}$.
- Method 1. $A = \int_{-\infty}^{+\infty} e^{-\beta \sum \frac{p_i^2}{2m}} dp_1 \dots dp_{3N}$

Application: monatomic ideal gas. Prove...

- Prove: $\sigma(E) \equiv \int_{H \leq E} dp_1 \dots dp_{3N} = \frac{(2\pi m E)^{\frac{3}{2}N}}{(\frac{3}{2}N)!}$.
- First: $p_i = \sqrt{2mE}x_i \Rightarrow \sigma(E) = (2mE)^{\frac{3}{2}N} \int_{\sum x_i^2 \leq 1} dx_1 \dots dx_{3N} \equiv K(2mE)^{\frac{3}{2}N} \dots (a)$
- Second, calculate $A \equiv \int_{-\infty}^{+\infty} e^{-\beta H} dp_1 \dots dp_{3N}$.
- Method 1. $A = \int_{-\infty}^{+\infty} e^{-\beta \sum \frac{p_i^2}{2m}} dp_1 \dots dp_{3N}$

$$= \prod_{i=1}^{3N} \int_{-\infty}^{+\infty} e^{-\beta \frac{p_i^2}{2m}} dp_i$$

Application: monatomic ideal gas. Prove...

- Prove: $\sigma(E) \equiv \int_{H \leq E} dp_1 \dots dp_{3N} = \frac{(2\pi m E)^{\frac{3}{2}N}}{(\frac{3}{2}N)!}$.
- First: $p_i = \sqrt{2mE}x_i \Rightarrow \sigma(E) = (2mE)^{\frac{3}{2}N} \int_{\sum x_i^2 \leq 1} dx_1 \dots dx_{3N} \equiv K(2mE)^{\frac{3}{2}N} \dots (a)$
- Second, calculate $A \equiv \int_{-\infty}^{+\infty} e^{-\beta H} dp_1 \dots dp_{3N}$.
- Method 1. $A = \int_{-\infty}^{+\infty} e^{-\beta \sum \frac{p_i^2}{2m}} dp_1 \dots dp_{3N}$

$$= \prod_{i=1}^{3N} \int_{-\infty}^{+\infty} e^{-\beta \frac{p_i^2}{2m}} dp_i = \prod_{i=1}^{3N} \left(\frac{2m}{\beta}\right)^{\frac{1}{2}} \int_{-\infty}^{+\infty} e^{-\beta \frac{p_i^2}{2m}} d\left(\frac{\beta}{2m}\right)^{\frac{1}{2}} p_i$$

Application: monatomic ideal gas. Prove...

- Prove: $\sigma(E) \equiv \int_{H \leq E} dp_1 \dots dp_{3N} = \frac{(2\pi m E)^{\frac{3}{2}N}}{(\frac{3}{2}N)!}$.
- First: $p_i = \sqrt{2mE}x_i \Rightarrow \sigma(E) = (2mE)^{\frac{3}{2}N} \int_{\sum x_i^2 \leq 1} dx_1 \dots dx_{3N} \equiv K(2mE)^{\frac{3}{2}N} \dots (a)$
- Second, calculate $A \equiv \int_{-\infty}^{+\infty} e^{-\beta H} dp_1 \dots dp_{3N}$.
- Method 1. $A = \int_{-\infty}^{+\infty} e^{-\beta \sum \frac{p_i^2}{2m}} dp_1 \dots dp_{3N}$

$$= \prod_{i=1}^{3N} \int_{-\infty}^{+\infty} e^{-\beta \frac{p_i^2}{2m}} dp_i = \prod_{i=1}^{3N} \left(\frac{2m}{\beta}\right)^{\frac{1}{2}} \int_{-\infty}^{+\infty} e^{-\beta \frac{p_i^2}{2m}} d\left(\frac{\beta}{2m}\right)^{\frac{1}{2}} p_i$$

$$= \left(\frac{2\pi m}{\beta}\right)^{\frac{3}{2}N} \dots (b)$$

Application: monatomic ideal gas.

$$A = \left(\frac{2\pi m}{\beta}\right)^{\frac{3}{2}N} \dots (b),$$

$$\sigma(E) \equiv \int_{H \leq E} dp_1 \dots dp_{3N} = K(2mE)^{\frac{3}{2}N} \dots (a)$$

- Method 2. $A \equiv \int_{-\infty}^{+\infty} e^{-\beta E} dp_1 \dots dp_{3N}$

Application: monatomic ideal gas.

$$A = \left(\frac{2\pi m}{\beta}\right)^{\frac{3}{2}N} \dots (b),$$

$$\sigma(E) \equiv \int_{H \leq E} dp_1 \dots dp_{3N} = K(2mE)^{\frac{3}{2}N} \dots (a)$$

- Method 2. $A \equiv \int_{-\infty}^{+\infty} e^{-\beta E} dp_1 \dots dp_{3N} = \int e^{-\beta E} d\sigma(E)$

Application: monatomic ideal gas.

$$A = \left(\frac{2\pi m}{\beta} \right)^{\frac{3}{2}N} \dots (b),$$

$$\sigma(E) \equiv \int_{H \leq E} dp_1 \dots dp_{3N} = K(2mE)^{\frac{3}{2}N} \dots (a)$$

- Method 2. $A \equiv \int_{-\infty}^{+\infty} e^{-\beta E} dp_1 \dots dp_{3N} = \int e^{-\beta E} d\sigma(E)$
 $= \int_0^{+\infty} e^{-\beta E} \frac{d\sigma(E)}{dE} dE$

Application: monatomic ideal gas.

$$A = \left(\frac{2\pi m}{\beta}\right)^{\frac{3}{2}N} \dots (b),$$

$$\sigma(E) \equiv \int_{H \leq E} dp_1 \dots dp_{3N} = K(2mE)^{\frac{3}{2}N} \dots (a)$$

- Method 2. $A \equiv \int_{-\infty}^{+\infty} e^{-\beta E} dp_1 \dots dp_{3N} = \int e^{-\beta E} d\sigma(E)$
 $= \int_0^{+\infty} e^{-\beta E} \frac{d\sigma(E)}{dE} dE$
 $= K \cdot (2m)^{\frac{3}{2}N} \cdot \frac{3}{2}N \cdot \int_0^{\infty} e^{-\beta E} E^{\frac{3}{2}N-1} dE$

Application: monatomic ideal gas.

$$A = \left(\frac{2\pi m}{\beta}\right)^{\frac{3}{2}N} \dots (b),$$

$$\sigma(E) \equiv \int_{H \leq E} dp_1 \dots dp_{3N} = K(2mE)^{\frac{3}{2}N} \dots (a)$$

- Method 2. $A \equiv \int_{-\infty}^{+\infty} e^{-\beta E} dp_1 \dots dp_{3N} = \int e^{-\beta E} d\sigma(E)$
 $= \int_0^{+\infty} e^{-\beta E} \frac{d\sigma(E)}{dE} dE$
 $= K \cdot (2m)^{\frac{3}{2}N} \cdot \frac{3}{2}N \cdot \int_0^{\infty} e^{-\beta E} E^{\frac{3}{2}N-1} dE$
 $= K \cdot (2m)^{\frac{3}{2}N} \cdot \frac{3}{2}N \cdot \frac{1}{\beta} \left(- \int_0^{\infty} E^{\frac{3}{2}N-1} d e^{-\beta E} \right)$

Application: monatomic ideal gas.

$$A = \left(\frac{2\pi m}{\beta}\right)^{\frac{3}{2}N} \dots (b),$$

$$\sigma(E) \equiv \int_{H \leq E} dp_1 \dots dp_{3N} = K(2mE)^{\frac{3}{2}N} \dots (a)$$

- Method 2. $A \equiv \int_{-\infty}^{+\infty} e^{-\beta E} dp_1 \dots dp_{3N} = \int e^{-\beta E} d\sigma(E)$
 $= \int_0^{+\infty} e^{-\beta E} \frac{d\sigma(E)}{dE} dE$
 $= K \cdot (2m)^{\frac{3}{2}N} \cdot \frac{3}{2}N \cdot \int_0^{\infty} e^{-\beta E} E^{\frac{3}{2}N-1} dE$
 $= K \cdot (2m)^{\frac{3}{2}N} \cdot \frac{3}{2}N \cdot \frac{1}{\beta} \left(- \int_0^{\infty} E^{\frac{3}{2}N-1} d e^{-\beta E} \right)$
 $= K \cdot (2m)^{\frac{3}{2}N} \cdot \frac{3}{2}N \cdot \left(\frac{3}{2}N - 1 \right) \cdot \frac{1}{\beta} \int_0^{\infty} e^{-\beta E} E^{\frac{3}{2}N-2} dE$

Application: monatomic ideal gas.

$$A = \left(\frac{2\pi m}{\beta}\right)^{\frac{3}{2}N} \dots (b),$$

$$\sigma(E) \equiv \int_{H \leq E} dp_1 \dots dp_{3N} = K(2mE)^{\frac{3}{2}N} \dots (a)$$

- Method 2. $A \equiv \int_{-\infty}^{+\infty} e^{-\beta E} dp_1 \dots dp_{3N} = \int e^{-\beta E} d\sigma(E)$
 $= \int_0^{+\infty} e^{-\beta E} \frac{d\sigma(E)}{dE} dE$
 $= K \cdot (2m)^{\frac{3}{2}N} \cdot \frac{3}{2}N \cdot \int_0^{\infty} e^{-\beta E} E^{\frac{3}{2}N-1} dE$
 $= K \cdot (2m)^{\frac{3}{2}N} \cdot \frac{3}{2}N \cdot \frac{1}{\beta} \left(- \int_0^{\infty} E^{\frac{3}{2}N-1} d e^{-\beta E} \right)$
 $= K \cdot (2m)^{\frac{3}{2}N} \cdot \frac{3}{2}N \cdot \left(\frac{3}{2}N - 1 \right) \cdot \frac{1}{\beta} \int_0^{\infty} e^{-\beta E} E^{\frac{3}{2}N-2} dE \dots$

Application: monatomic ideal gas.

$$A = \left(\frac{2\pi m}{\beta}\right)^{\frac{3}{2}N} \dots (b)$$

$$\sigma(E) \equiv \int_{H \leq E} dp_1 \dots dp_{3N} = K(2mE)^{\frac{3}{2}N} \dots (a)$$

- Method 2. $A \equiv \int_{-\infty}^{+\infty} e^{-\beta E} dp_1 \dots dp_{3N} = \int e^{-\beta E} d\sigma(E)$

$$\begin{aligned}
 &= \int_0^{+\infty} e^{-\beta E} \frac{d\sigma(E)}{dE} dE \\
 &= K \cdot (2m)^{\frac{3}{2}N} \cdot \frac{3}{2}N \cdot \int_0^{\infty} e^{-\beta E} E^{\frac{3}{2}N-1} dE \\
 &= K \cdot (2m)^{\frac{3}{2}N} \cdot \frac{3}{2}N \cdot \frac{1}{\beta} \left(- \int_0^{\infty} E^{\frac{3}{2}N-1} d e^{-\beta E} \right) \\
 &= K \cdot (2m)^{\frac{3}{2}N} \cdot \frac{3}{2}N \cdot \left(\frac{3}{2}N - 1 \right) \cdot \frac{1}{\beta} \int_0^{\infty} e^{-\beta E} E^{\frac{3}{2}N-2} dE \dots \\
 &= K \cdot \left(\frac{2m}{\beta}\right)^{\frac{3}{2}N} \left(\frac{3}{2}N\right)! \dots (c)
 \end{aligned}$$

Application: monatomic ideal gas.

$$A = \left(\frac{2\pi m}{\beta}\right)^{\frac{3}{2}N} \dots (b),$$

$$\sigma(E) \equiv \int_{H \leq E} dp_1 \dots dp_{3N} = K(2mE)^{\frac{3}{2}N} \dots (a)$$

- Method 2. $A \equiv \int_{-\infty}^{+\infty} e^{-\beta E} dp_1 \dots dp_{3N} = \int e^{-\beta E} d\sigma(E)$
 $= \int_0^{+\infty} e^{-\beta E} \frac{d\sigma(E)}{dE} dE$
 $= K \cdot (2m)^{\frac{3}{2}N} \cdot \frac{3}{2}N \cdot \int_0^{\infty} e^{-\beta E} E^{\frac{3}{2}N-1} dE$
 $= K \cdot (2m)^{\frac{3}{2}N} \cdot \frac{3}{2}N \cdot \frac{1}{\beta} \left(- \int_0^{\infty} E^{\frac{3}{2}N-1} d e^{-\beta E} \right)$
 $= K \cdot (2m)^{\frac{3}{2}N} \cdot \frac{3}{2}N \cdot \left(\frac{3}{2}N - 1 \right) \cdot \frac{1}{\beta} \int_0^{\infty} e^{-\beta E} E^{\frac{3}{2}N-2} dE \dots$
 $= K \cdot \left(\frac{2m}{\beta}\right)^{\frac{3}{2}N} \left(\frac{3}{2}N\right)! \dots (c)$
- $(b)=(c) \Rightarrow K = \frac{\pi^{\frac{3}{2}N}}{(\frac{3}{2}N)!},$

Application: monatomic ideal gas.

$$A = \left(\frac{2\pi m}{\beta}\right)^{\frac{3}{2}N} \dots (b),$$

$$\sigma(E) \equiv \int_{H \leq E} dp_1 \dots dp_{3N} = K(2mE)^{\frac{3}{2}N} \dots (a)$$

- Method 2. $A \equiv \int_{-\infty}^{+\infty} e^{-\beta E} dp_1 \dots dp_{3N} = \int e^{-\beta E} d\sigma(E)$
 $= \int_0^{+\infty} e^{-\beta E} \frac{d\sigma(E)}{dE} dE$
 $= K \cdot (2m)^{\frac{3}{2}N} \cdot \frac{3}{2}N \cdot \int_0^{\infty} e^{-\beta E} E^{\frac{3}{2}N-1} dE$
 $= K \cdot (2m)^{\frac{3}{2}N} \cdot \frac{3}{2}N \cdot \frac{1}{\beta} \left(- \int_0^{\infty} E^{\frac{3}{2}N-1} d e^{-\beta E} \right)$
 $= K \cdot (2m)^{\frac{3}{2}N} \cdot \frac{3}{2}N \cdot \left(\frac{3}{2}N - 1 \right) \cdot \frac{1}{\beta} \int_0^{\infty} e^{-\beta E} E^{\frac{3}{2}N-2} dE \dots$
 $= K \cdot \left(\frac{2m}{\beta}\right)^{\frac{3}{2}N} \left(\frac{3}{2}N\right)! \dots (c)$
- $(b)=(c) \Rightarrow K = \frac{\pi^{\frac{3}{2}N}}{(\frac{3}{2}N)!},$
- $\therefore \sigma(E) = \frac{(2\pi m E)^{\frac{3}{2}N}}{(\frac{3}{2}N)!} \cdot \square$

Application: monatomic ideal gas.

$$\Omega(N, E, V) \simeq \frac{3}{2}N \cdot \left(\frac{V}{h^3}\right)^N \cdot \frac{(2\pi mE)^{\frac{3}{2}N}}{N!(\frac{3}{2}N)!} \frac{\Delta E}{E}.$$

Application: monatomic ideal gas.

$$\Omega(N, E, V) \simeq \frac{3}{2}N \cdot \left(\frac{V}{h^3}\right)^N \cdot \frac{(2\pi mE)^{\frac{3}{2}N}}{N! \left(\frac{3}{2}N\right)!} \frac{\Delta E}{E}.$$

- Entropy: $S = k \ln \Omega$

Application: monatomic ideal gas.

$$\Omega(N, E, V) \simeq \frac{3}{2}N \cdot \left(\frac{V}{h^3}\right)^N \cdot \frac{(2\pi mE)^{\frac{3}{2}N}}{N!(\frac{3}{2}N)!} \frac{\Delta E}{E}.$$

- Entropy: $S = k \ln \Omega = k \ln \left[\left(\frac{V}{h^3}\right)^N \cdot \frac{(2\pi mE)^{\frac{3}{2}N}}{N!(\frac{3}{2}N-1)!} \frac{\Delta E}{E} \right]$

Application: monatomic ideal gas.

$$\Omega(N, E, V) \simeq \frac{3}{2}N \cdot \left(\frac{V}{h^3}\right)^N \cdot \frac{(2\pi mE)^{\frac{3}{2}N}}{N!(\frac{3}{2}N)!} \frac{\Delta E}{E}.$$

- Entropy: $S = k \ln \Omega = k \ln \left[\left(\frac{V}{h^3}\right)^N \cdot \frac{(2\pi mE)^{\frac{3}{2}N}}{N!(\frac{3}{2}N-1)!} \frac{\Delta E}{E} \right]$
 $\simeq Nk \ln \left[\left(\frac{V}{h^3}\right) (2\pi mE)^{\frac{3}{2}} \right] - k [\ln N! + \ln (\frac{3}{2}N)!] + k \ln \frac{\Delta E}{E}$

Application: monatomic ideal gas.

$$\Omega(N, E, V) \simeq \frac{3}{2}N \cdot \left(\frac{V}{h^3}\right)^N \cdot \frac{(2\pi mE)^{\frac{3}{2}N}}{N!(\frac{3}{2}N)!} \frac{\Delta E}{E}.$$

- Entropy: $S = k \ln \Omega = k \ln \left[\left(\frac{V}{h^3}\right)^N \cdot \frac{(2\pi mE)^{\frac{3}{2}N}}{N!(\frac{3}{2}N-1)!} \frac{\Delta E}{E} \right]$
 $\simeq Nk \ln \left[\left(\frac{V}{h^3}\right) (2\pi mE)^{\frac{3}{2}} \right] - k [\ln N! + \ln (\frac{3}{2}N)!] + k \ln \frac{\Delta E}{E}$
 $= Nk \ln \left[\left(\frac{V}{h^3}\right) (2\pi mE)^{\frac{3}{2}} \right] - k [N(\ln N - 1) +$
 $\frac{3}{2}N(\ln \frac{3}{2}N - 1)] + k \ln \frac{\Delta E}{E}$

Application: monatomic ideal gas.

$$\Omega(N, E, V) \simeq \frac{3}{2}N \cdot \left(\frac{V}{h^3}\right)^N \cdot \frac{(2\pi mE)^{\frac{3}{2}N}}{N!(\frac{3}{2}N)!} \frac{\Delta E}{E}.$$

- Entropy: $S = k \ln \Omega = k \ln \left[\left(\frac{V}{h^3}\right)^N \cdot \frac{(2\pi mE)^{\frac{3}{2}N}}{N!(\frac{3}{2}N)!} \frac{\Delta E}{E} \right]$

$$\simeq Nk \ln \left[\left(\frac{V}{h^3}\right) (2\pi mE)^{\frac{3}{2}} \right] - k [\ln N! + \ln (\frac{3}{2}N)!] + k \ln \frac{\Delta E}{E}$$

$$= Nk \ln \left[\left(\frac{V}{h^3}\right) (2\pi mE)^{\frac{3}{2}} \right] - k [N(\ln N - 1) + \frac{3}{2}N(\ln \frac{3}{2}N - 1)] + k \ln \frac{\Delta E}{E}$$

$$= Nk \ln \left[\left(\frac{V}{h^3}\right) (2\pi mE)^{\frac{3}{2}} \right] - Nk \ln N + Nk - \frac{3}{2}Nk \ln (\frac{3}{2}N) + \frac{3}{2}Nk + k \ln \frac{\Delta E}{E}$$

Application: monatomic ideal gas.

$$\Omega(N, E, V) \simeq \frac{3}{2}N \cdot \left(\frac{V}{h^3}\right)^N \cdot \frac{(2\pi mE)^{\frac{3}{2}N}}{N!(\frac{3}{2}N)!} \frac{\Delta E}{E}.$$

- Entropy: $S = k \ln \Omega = k \ln \left[\left(\frac{V}{h^3}\right)^N \cdot \frac{(2\pi mE)^{\frac{3}{2}N}}{N!(\frac{3}{2}N-1)!} \frac{\Delta E}{E} \right]$

$$\begin{aligned} &\simeq Nk \ln \left[\left(\frac{V}{h^3}\right) (2\pi mE)^{\frac{3}{2}} \right] - k [\ln N! + \ln (\frac{3}{2}N)!] + k \ln \frac{\Delta E}{E} \\ &= Nk \ln \left[\left(\frac{V}{h^3}\right) (2\pi mE)^{\frac{3}{2}} \right] - k [N(\ln N - 1) + \\ &\quad \frac{3}{2}N(\ln \frac{3}{2}N - 1)] + k \ln \frac{\Delta E}{E} \\ &= Nk \ln \left[\left(\frac{V}{h^3}\right) (2\pi mE)^{\frac{3}{2}} \right] - Nk \ln N + Nk - \\ &\quad \frac{3}{2}Nk \ln (\frac{3}{2}N) + \frac{3}{2}Nk + k \ln \frac{\Delta E}{E} \\ &= Nk \ln \left[\left(\frac{V}{h^3N}\right) \left(\frac{4\pi mE}{3N}\right)^{\frac{3}{2}} \right] + \frac{5}{2}Nk + k \ln \frac{\Delta E}{E} \end{aligned}$$

Application: monatomic ideal gas.

$$\Omega(N, E, V) \simeq \frac{3}{2}N \cdot \left(\frac{V}{h^3}\right)^N \cdot \frac{(2\pi mE)^{\frac{3}{2}N}}{N!(\frac{3}{2}N)!} \frac{\Delta E}{E}.$$

- Entropy: $S = k \ln \Omega = k \ln \left[\left(\frac{V}{h^3}\right)^N \cdot \frac{(2\pi mE)^{\frac{3}{2}N}}{N!(\frac{3}{2}N-1)!} \frac{\Delta E}{E} \right]$

$$\begin{aligned} &\simeq Nk \ln \left[\left(\frac{V}{h^3}\right) (2\pi mE)^{\frac{3}{2}} \right] - k [\ln N! + \ln (\frac{3}{2}N)!] + k \ln \frac{\Delta E}{E} \\ &= Nk \ln \left[\left(\frac{V}{h^3}\right) (2\pi mE)^{\frac{3}{2}} \right] - k [N(\ln N - 1) + \\ &\quad \frac{3}{2}N(\ln \frac{3}{2}N - 1)] + k \ln \frac{\Delta E}{E} \\ &= Nk \ln \left[\left(\frac{V}{h^3}\right) (2\pi mE)^{\frac{3}{2}} \right] - Nk \ln N + Nk - \\ &\quad \frac{3}{2}Nk \ln (\frac{3}{2}N) + \frac{3}{2}Nk + k \ln \frac{\Delta E}{E} \\ &= Nk \ln \left[\left(\frac{V}{h^3N}\right) \left(\frac{4\pi mE}{3N}\right)^{\frac{3}{2}} \right] + \frac{5}{2}Nk + k \ln \frac{\Delta E}{E} \\ &\simeq Nk \ln \left[\left(\frac{V}{h^3N}\right) \left(\frac{4\pi mE}{3N}\right)^{\frac{3}{2}} \right] + \frac{5}{2}Nk. \end{aligned}$$

Application: monatomic ideal gas.

$$S = Nk \ln \left[\left(\frac{V}{h^3 N} \right) \left(\frac{4\pi m E}{3N} \right)^{\frac{3}{2}} \right] + \frac{5}{2} Nk$$

$$\bullet \Rightarrow E(N, S, V) = \frac{3h^2 N^{\frac{5}{3}}}{4\pi m V^{\frac{2}{3}}} e^{\left(\frac{2S}{3Nk} - \frac{5}{3} \right)}.$$

Application: monatomic ideal gas.

$$S = Nk \ln \left[\left(\frac{V}{h^3 N} \right) \left(\frac{4\pi m E}{3N} \right)^{\frac{3}{2}} \right] + \frac{5}{2} Nk$$

$$\bullet \Rightarrow E(N, S, V) = \frac{3h^2 N^{\frac{5}{3}}}{4\pi m V^{\frac{2}{3}}} e^{\left(\frac{2S}{3Nk} - \frac{5}{3} \right)}.$$

$$T = \left(\frac{\partial E}{\partial S} \right)_{N,V} = \frac{2}{3} \frac{E}{Nk}$$

Application: monatomic ideal gas.

$$S = Nk \ln \left[\left(\frac{V}{h^3 N} \right) \left(\frac{4\pi m E}{3N} \right)^{\frac{3}{2}} \right] + \frac{5}{2} Nk$$

$$\bullet \Rightarrow E(N, S, V) = \frac{3h^2 N^{\frac{5}{3}}}{4\pi m V^{\frac{2}{3}}} e^{\left(\frac{2S}{3Nk} - \frac{5}{3}\right)}.$$
$$T = \left(\frac{\partial E}{\partial S} \right)_{N,V} = \frac{2}{3} \frac{E}{Nk} \Rightarrow E = \frac{3}{2} NkT$$

Application: monatomic ideal gas.

$$S = Nk \ln \left[\left(\frac{V}{h^3 N} \right) \left(\frac{4\pi m E}{3N} \right)^{\frac{3}{2}} \right] + \frac{5}{2} Nk$$

$$\bullet \Rightarrow E(N, S, V) = \frac{3h^2 N^{\frac{5}{3}}}{4\pi m V^{\frac{2}{3}}} e^{\left(\frac{2S}{3Nk} - \frac{5}{3} \right)}.$$

$$T = \left(\frac{\partial E}{\partial S} \right)_{N,V} = \frac{2}{3} \frac{E}{Nk} \Rightarrow E = \frac{3}{2} NkT$$

$$p = - \left(\frac{\partial E}{\partial V} \right)_{N,S} = \frac{2}{3} \frac{E}{V}$$

Application: monatomic ideal gas.

$$S = Nk \ln \left[\left(\frac{V}{h^3 N} \right) \left(\frac{4\pi m E}{3N} \right)^{\frac{3}{2}} \right] + \frac{5}{2} Nk$$

- $\Rightarrow E(N, S, V) = \frac{3h^2 N^{\frac{5}{3}}}{4\pi m V^{\frac{2}{3}}} e^{\left(\frac{2S}{3Nk} - \frac{5}{3}\right)}.$
- $T = \left(\frac{\partial E}{\partial S}\right)_{N,V} = \frac{2}{3} \frac{E}{Nk} \Rightarrow E = \frac{3}{2} NkT$
- $p = -\left(\frac{\partial E}{\partial V}\right)_{N,S} = \frac{2}{3} \frac{E}{V}$
- $\Rightarrow pV = NkT,$

Application: monatomic ideal gas.

$$S = Nk \ln \left[\left(\frac{V}{h^3 N} \right) \left(\frac{4\pi m E}{3N} \right)^{\frac{3}{2}} \right] + \frac{5}{2} Nk$$

$$\bullet \Rightarrow E(N, S, V) = \frac{3h^2 N^{\frac{5}{3}}}{4\pi m V^{\frac{2}{3}}} e^{\left(\frac{2S}{3Nk} - \frac{5}{3} \right)}.$$

$$T = \left(\frac{\partial E}{\partial S} \right)_{N,V} = \frac{2}{3} \frac{E}{Nk} \Rightarrow E = \frac{3}{2} NkT$$

$$p = - \left(\frac{\partial E}{\partial V} \right)_{N,S} = \frac{2}{3} \frac{E}{V}$$

$$\bullet \Rightarrow pV = NkT,$$

$$S = Nk \ln \left[\frac{V}{N} \left(\frac{2\pi mkT}{h^2} \right)^{\frac{3}{2}} \right] + \frac{5}{2} Nk. \text{ Same as eq. (7.6.2).}$$

Table of contents

1 §9. Ensemble theory

- 9.1 Phase space and Liouville's theorem
- 9.2 The microcanonical ensemble
- 9.3 Thermodynamics in microcanonical ensemble theory
- 9.4 Canonical ensemble
- 9.5 Thermodynamics in canonical ensemble
- 9.6 EOS of real gas
- 9.7 Thermal capacity of solid
- 9.9 Ising model
- 9.10 Grand canonical ensemble
- 9.11 Thermodynamics in grand canonical ensemble
- 9.12 Applications of the grand canonical ensemble

9.4 Canonical ensemble

9.4 Canonical ensemble

- Object: system contacting with a thermal reservoir.

9.4 Canonical ensemble

- Object: system contacting with a thermal reservoir. Exists energy exchanges. Keep temperature T .

9.4 Canonical ensemble

- Object: system contacting with a thermal reservoir. Exists energy exchanges. Keep temperature T .
- System has fixed (N, V, T) .

9.4 Canonical ensemble

- Object: system contacting with a thermal reservoir. Exists energy exchanges. Keep temperature T .
- System has fixed (N, V, T) .
- Consider the combined system (labelled '0') contains both the 'system' and the reservoir (labelled 'r'), which is an isolated system.

9.4 Canonical ensemble

- Object: system contacting with a thermal reservoir. Exists energy exchanges. Keep temperature T .
- System has fixed (N, V, T) .
- Consider the combined system (labelled '0') contains both the 'system' and the reservoir (labelled 'r'), which is an isolated system. Back to the microcanonical ensemble.

9.4 Canonical ensemble

- Object: system contacting with a thermal reservoir. Exists energy exchanges. Keep temperature T .
- System has fixed (N, V, T) .
- Consider the combined system (labelled '0') contains both the 'system' and the reservoir (labelled 'r'), which is an isolated system. Back to the microcanonical ensemble.
- Set the energy of system E , of reservoir E_r , and of the whole system $E^{(0)}$.

9.4 Canonical ensemble

- Object: system contacting with a thermal reservoir. Exists energy exchanges. Keep temperature T .
- System has fixed (N, V, T) .
- Consider the combined system (labelled '0') contains both the 'system' and the reservoir (labelled 'r'), which is an isolated system. Back to the microcanonical ensemble.
- Set the energy of system E , of reservoir E_r , and of the whole system $E^{(0)}$.
- $E + E_r = E^{(0)}$.

9.4 Canonical ensemble

- Object: system contacting with a thermal reservoir. Exists energy exchanges. Keep temperature T .
- System has fixed (N, V, T) .
- Consider the combined system (labelled '0') contains both the 'system' and the reservoir (labelled 'r'), which is an isolated system. Back to the microcanonical ensemble.
- Set the energy of system E , of reservoir E_r , and of the whole system $E^{(0)}$.
- $E + E_r = E^{(0)}$.
- Consider the probability for the system at state s .

Canonical ensemble

- As state s , there exists energy E_s . Reservoir can be any state with energy $E_r = E^{(0)} - E_s$. Total number of these states: $\Omega_r(E^{(0)} - E_s)$. For the combined system being isolated system, obeys microcanonical ensemble's the principle of equal a priori probabilities.

Canonical ensemble

- As state s , there exists energy E_s . Reservoir can be any state with energy $E_r = E^{(0)} - E_s$. Total number of these states: $\Omega_r(E^{(0)} - E_s)$. For the combined system being isolated system, obeys microcanonical ensemble's the principle of equal a priori probabilities. I.e., $\rho_s \propto \Omega_r(E^{(0)} - E_s)$.

Canonical ensemble

- As state s , there exists energy E_s . Reservoir can be any state with energy $E_r = E^{(0)} - E_s$. Total number of these states: $\Omega_r(E^{(0)} - E_s)$. For the combined system being isolated system, obeys microcanonical ensemble's the principle of equal a priori probabilities. I.e., $\rho_s \propto \Omega_r(E^{(0)} - E_s)$.
- Use $\ln \Omega$ for convenience.

Canonical ensemble

- As state s , there exists energy E_s . Reservoir can be any state with energy $E_r = E^{(0)} - E_s$. Total number of these states: $\Omega_r(E^{(0)} - E_s)$. For the combined system being isolated system, obeys microcanonical ensemble's the principle of equal a priori probabilities. I.e., $\rho_s \propto \Omega_r(E^{(0)} - E_s)$.
- Use $\ln \Omega$ for convenience.
- As $E_s \ll E^{(0)}$,

$$\ln \Omega_r(E^{(0)} - E_s) \simeq \ln \Omega_r(E^{(0)}) + \left(\frac{\partial \ln \Omega_r}{\partial E_r} \right)_{E_r=E^{(0)}} \cdot (-E_s)$$

Canonical ensemble

- As state s , there exists energy E_s . Reservoir can be any state with energy $E_r = E^{(0)} - E_s$. Total number of these states: $\Omega_r(E^{(0)} - E_s)$. For the combined system being isolated system, obeys microcanonical ensemble's the principle of equal a priori probabilities. I.e., $\rho_s \propto \Omega_r(E^{(0)} - E_s)$.
- Use $\ln \Omega$ for convenience.
- As $E_s \ll E^{(0)}$,

$$\ln \Omega_r(E^{(0)} - E_s) \simeq \ln \Omega_r(E^{(0)}) + \left(\frac{\partial \ln \Omega_r}{\partial E_r}\right)_{E_r=E^{(0)}} \cdot (-E_s)$$

$$= \ln \Omega_r(E^{(0)}) - \beta E_s. \text{ Notice (9.3.5).}$$

Canonical ensemble

- As state s , there exists energy E_s . Reservoir can be any state with energy $E_r = E^{(0)} - E_s$. Total number of these states: $\Omega_r(E^{(0)} - E_s)$. For the combined system being isolated system, obeys microcanonical ensemble's the principle of equal a priori probabilities. I.e., $\rho_s \propto \Omega_r(E^{(0)} - E_s)$.
- Use $\ln \Omega$ for convenience.
- As $E_s \ll E^{(0)}$,

$$\ln \Omega_r(E^{(0)} - E_s) \simeq \ln \Omega_r(E^{(0)}) + \left(\frac{\partial \ln \Omega_r}{\partial E_r}\right)_{E_r=E^{(0)}} \cdot (-E_s)$$

$$= \ln \Omega_r(E^{(0)}) - \beta E_s. \text{ Notice (9.3.5).}$$
- $\therefore \rho_s \propto e^{\ln \Omega_r(E^{(0)}) - \beta E_s} \propto e^{-\beta E_s}$,
 where $\ln \Omega_r(E^{(0)}) = \text{Const.}$

Canonical ensemble

- Introduce a normalization factor: $\rho_s = \frac{1}{Z} e^{-\beta E_s}$,

Canonical ensemble

- Introduce a normalization factor: $\rho_s = \frac{1}{Z}e^{-\beta E_s}$,
where $Z = \sum_s e^{-\beta E_s}$ is called partition function.

Canonical ensemble

- Introduce a normalization factor: $\rho_s = \frac{1}{Z}e^{-\beta E_s}$, where $Z = \sum_s e^{-\beta E_s}$ is called partition function.
- Considering the degeneracy Ω_l at energy level E_l , then the probability at E_l is:

$$\rho_l = \Omega_l \cdot \frac{1}{Z}e^{-\beta E_l},$$

the partition function can also be written as

$$Z = \sum_l \Omega_l e^{-\beta E_l}.$$

Canonical ensemble

- Introduce a normalization factor: $\rho_s = \frac{1}{Z}e^{-\beta E_s}$, where $Z = \sum_s e^{-\beta E_s}$ is called partition function.
- Considering the degeneracy Ω_l at energy level E_l , then the probability at E_l is:

$$\rho_l = \Omega_l \cdot \frac{1}{Z}e^{-\beta E_l},$$

the partition function can also be written as

$$Z = \sum_l \Omega_l e^{-\beta E_l}.$$

- For continuous (classical) case:

$$\rho(q, p)d\Omega = \frac{1}{N!h^{Nr}} \frac{e^{-\beta E(q, p)}}{Z} d\Omega,$$

$$\text{where } Z = \frac{1}{N!h^{Nr}} \int e^{-\beta E(q, p)} d\Omega.$$

Table of contents

1 §9. Ensemble theory

- 9.1 Phase space and Liouville's theorem
- 9.2 The microcanonical ensemble
- 9.3 Thermodynamics in microcanonical ensemble theory
- 9.4 Canonical ensemble
- **9.5 Thermodynamics in canonical ensemble**
- 9.6 EOS of real gas
- 9.7 Thermal capacity of solid
- 9.9 Ising model
- 9.10 Grand canonical ensemble
- 9.11 Thermodynamics in grand canonical ensemble
- 9.12 Applications of the grand canonical ensemble

9.5 Thermodynamics in canonical ensemble

9.5 Thermodynamics in canonical ensemble

- Internal energy is the average energy for all possible micro-states:

9.5 Thermodynamics in canonical ensemble

- Internal energy is the average energy for all possible micro-states: $U = \bar{E}$

9.5 Thermodynamics in canonical ensemble

- Internal energy is the average energy for all possible micro-states: $U = \bar{E} = \sum E_s \rho_s$

9.5 Thermodynamics in canonical ensemble

- Internal energy is the average energy for all possible micro-states: $U = \bar{E} = \sum E_s \rho_s = \sum \frac{1}{Z} E_s e^{-\beta E_s}$

9.5 Thermodynamics in canonical ensemble

- Internal energy is the average energy for all possible micro-states:
$$U = \bar{E} = \sum E_s \rho_s = \sum \frac{1}{Z} E_s e^{-\beta E_s}$$
$$= \frac{1}{Z} \left(-\frac{\partial}{\partial \beta} \right) \sum e^{-\beta E_s}$$

9.5 Thermodynamics in canonical ensemble

- Internal energy is the average energy for all possible micro-states:
$$U = \bar{E} = \sum E_s \rho_s = \sum \frac{1}{Z} E_s e^{-\beta E_s}$$
$$= \frac{1}{Z} \left(-\frac{\partial}{\partial \beta} \right) \sum e^{-\beta E_s} = \frac{1}{Z} \left(-\frac{\partial}{\partial \beta} \right) Z$$

9.5 Thermodynamics in canonical ensemble

- Internal energy is the average energy for all possible micro-states:
$$U = \bar{E} = \sum E_s \rho_s = \sum \frac{1}{Z} E_s e^{-\beta E_s}$$
$$= \frac{1}{Z} \left(-\frac{\partial}{\partial \beta} \right) \sum e^{-\beta E_s} = \frac{1}{Z} \left(-\frac{\partial}{\partial \beta} \right) Z = -\frac{\partial}{\partial \beta} \ln Z.$$

9.5 Thermodynamics in canonical ensemble

- Internal energy is the average energy for all possible micro-states:
$$U = \bar{E} = \sum E_s \rho_s = \sum \frac{1}{Z} E_s e^{-\beta E_s} \\ = \frac{1}{Z} \left(-\frac{\partial}{\partial \beta} \right) \sum e^{-\beta E_s} = \frac{1}{Z} \left(-\frac{\partial}{\partial \beta} \right) Z = -\frac{\partial}{\partial \beta} \ln Z.$$
- Generalized force is the average of $\frac{\partial E_s}{\partial y}$:

9.5 Thermodynamics in canonical ensemble

- Internal energy is the average energy for all possible micro-states: $U = \bar{E} = \sum E_s \rho_s = \sum \frac{1}{Z} E_s e^{-\beta E_s}$
 $= \frac{1}{Z} \left(-\frac{\partial}{\partial \beta} \right) \sum e^{-\beta E_s} = \frac{1}{Z} \left(-\frac{\partial}{\partial \beta} \right) Z = -\frac{\partial}{\partial \beta} \ln Z.$
- Generalized force is the average of $\frac{\partial E_s}{\partial y}$:

$$Y = \frac{1}{Z} \sum \frac{\partial E_s}{\partial y} e^{-\beta E_s}$$

9.5 Thermodynamics in canonical ensemble

- Internal energy is the average energy for all possible micro-states: $U = \bar{E} = \sum E_s \rho_s = \sum \frac{1}{Z} E_s e^{-\beta E_s}$
 $= \frac{1}{Z} \left(-\frac{\partial}{\partial \beta} \right) \sum e^{-\beta E_s} = \frac{1}{Z} \left(-\frac{\partial}{\partial \beta} \right) Z = -\frac{\partial}{\partial \beta} \ln Z.$
- Generalized force is the average of $\frac{\partial E_s}{\partial y}$:
 $Y = \frac{1}{Z} \sum \frac{\partial E_s}{\partial y} e^{-\beta E_s} = \frac{1}{Z} \left(-\frac{1}{\beta} \frac{\partial}{\partial y} \right) \sum e^{-\beta E_s}$

9.5 Thermodynamics in canonical ensemble

- Internal energy is the average energy for all possible micro-states: $U = \bar{E} = \sum E_s \rho_s = \sum \frac{1}{Z} E_s e^{-\beta E_s}$
 $= \frac{1}{Z} \left(-\frac{\partial}{\partial \beta} \right) \sum e^{-\beta E_s} = \frac{1}{Z} \left(-\frac{\partial}{\partial \beta} \right) Z = -\frac{\partial}{\partial \beta} \ln Z.$
- Generalized force is the average of $\frac{\partial E_s}{\partial y}$:
 $Y = \frac{1}{Z} \sum \frac{\partial E_s}{\partial y} e^{-\beta E_s} = \frac{1}{Z} \left(-\frac{1}{\beta} \frac{\partial}{\partial y} \right) \sum e^{-\beta E_s}$
 $= -\frac{1}{\beta} \frac{\partial}{\partial y} \ln Z.$

9.5 Thermodynamics in canonical ensemble

- Internal energy is the average energy for all possible micro-states: $U = \bar{E} = \sum E_s \rho_s = \sum \frac{1}{Z} E_s e^{-\beta E_s}$
 $= \frac{1}{Z} \left(-\frac{\partial}{\partial \beta} \right) \sum e^{-\beta E_s} = \frac{1}{Z} \left(-\frac{\partial}{\partial \beta} \right) Z = -\frac{\partial}{\partial \beta} \ln Z.$
- Generalized force is the average of $\frac{\partial E_s}{\partial y}$:
 $Y = \frac{1}{Z} \sum \frac{\partial E_s}{\partial y} e^{-\beta E_s} = \frac{1}{Z} \left(-\frac{1}{\beta} \frac{\partial}{\partial y} \right) \sum e^{-\beta E_s}$
 $= -\frac{1}{\beta} \frac{\partial}{\partial y} \ln Z.$
- Similar to §7.1 and §8.1, the entropy:
 $S = k(\ln Z - \beta \frac{\partial}{\partial \beta} \ln Z).$

9.5 Thermodynamics in canonical ensemble

- Internal energy is the average energy for all possible micro-states: $U = \bar{E} = \sum E_s \rho_s = \sum \frac{1}{Z} E_s e^{-\beta E_s}$
 $= \frac{1}{Z} \left(-\frac{\partial}{\partial \beta}\right) \sum e^{-\beta E_s} = \frac{1}{Z} \left(-\frac{\partial}{\partial \beta}\right) Z = -\frac{\partial}{\partial \beta} \ln Z.$
- Generalized force is the average of $\frac{\partial E_s}{\partial y}$:
 $Y = \frac{1}{Z} \sum \frac{\partial E_s}{\partial y} e^{-\beta E_s} = \frac{1}{Z} \left(-\frac{1}{\beta} \frac{\partial}{\partial y}\right) \sum e^{-\beta E_s}$
 $= -\frac{1}{\beta} \frac{\partial}{\partial y} \ln Z.$
- Similar to §7.1 and §8.1, the entropy:
 $S = k(\ln Z - \beta \frac{\partial}{\partial \beta} \ln Z).$
- Free energy: $F = U - TS =$
 $-\frac{\partial}{\partial \beta} \ln Z - kT(\ln Z - \beta \frac{\partial}{\partial \beta} \ln Z) = -kT \ln Z,$

9.5 Thermodynamics in canonical ensemble

- Internal energy is the average energy for all possible micro-states: $U = \bar{E} = \sum E_s \rho_s = \sum \frac{1}{Z} E_s e^{-\beta E_s}$
 $= \frac{1}{Z} \left(-\frac{\partial}{\partial \beta} \right) \sum e^{-\beta E_s} = \frac{1}{Z} \left(-\frac{\partial}{\partial \beta} \right) Z = -\frac{\partial}{\partial \beta} \ln Z.$
- Generalized force is the average of $\frac{\partial E_s}{\partial y}$:
 $Y = \frac{1}{Z} \sum \frac{\partial E_s}{\partial y} e^{-\beta E_s} = \frac{1}{Z} \left(-\frac{1}{\beta} \frac{\partial}{\partial y} \right) \sum e^{-\beta E_s}$
 $= -\frac{1}{\beta} \frac{\partial}{\partial y} \ln Z.$
- Similar to §7.1 and §8.1, the entropy:
 $S = k(\ln Z - \beta \frac{\partial}{\partial \beta} \ln Z).$
- Free energy: $F = U - TS =$
 $-\frac{\partial}{\partial \beta} \ln Z - kT(\ln Z - \beta \frac{\partial}{\partial \beta} \ln Z) = -kT \ln Z,$ is
 characteristic function for $(N, V, T).$

Fluctuation

Fluctuation

- $\overline{(E - \bar{E})^2}$

Fluctuation

- $\overline{(E - \bar{E})^2} = \sum \rho_s (E_s - \bar{E})^2$

Fluctuation

- $$\overline{(E - \bar{E})^2} = \sum \rho_s (E_s - \bar{E})^2$$
$$= \sum \rho_s (E_s^2 - 2\bar{E}E_s + \bar{E}^2)$$

Fluctuation

- $$\begin{aligned}\overline{(E - \bar{E})^2} &= \sum \rho_s (E_s - \bar{E})^2 \\ &= \sum \rho_s (E_s^2 - 2\bar{E}E_s + \bar{E}^2) \\ &= \sum E_s^2 \rho_s - 2\bar{E} \sum \rho_s E_s + \bar{E}^2\end{aligned}$$

Fluctuation

- $$\begin{aligned}\overline{(E - \bar{E})^2} &= \sum \rho_s (E_s - \bar{E})^2 \\ &= \sum \rho_s (E_s^2 - 2\bar{E}E_s + \bar{E}^2) \\ &= \sum E_s^2 \rho_s - 2\bar{E} \sum \rho_s E_s + \bar{E}^2 = \sum E_s^2 \rho_s - \bar{E}^2\end{aligned}$$

Fluctuation

- $$\begin{aligned}\overline{(E - \bar{E})^2} &= \sum \rho_s (E_s - \bar{E})^2 \\ &= \sum \rho_s (E_s^2 - 2\bar{E}E_s + \bar{E}^2) \\ &= \sum E_s^2 \rho_s - 2\bar{E} \sum \rho_s E_s + \bar{E}^2 = \sum E_s^2 \rho_s - \bar{E}^2 \\ &= \overline{E^2} - \bar{E}^2;\end{aligned}$$

Fluctuation

- $$\begin{aligned}\overline{(E - \bar{E})^2} &= \sum \rho_s (E_s - \bar{E})^2 \\ &= \sum \rho_s (E_s^2 - 2\bar{E}E_s + \bar{E}^2) \\ &= \sum E_s^2 \rho_s - 2\bar{E} \sum \rho_s E_s + \bar{E}^2 = \sum E_s^2 \rho_s - \bar{E}^2 \\ &= \overline{E^2} - \bar{E}^2;\end{aligned}$$
- $$\frac{\partial \bar{E}}{\partial \beta}$$

Fluctuation

- $$\begin{aligned}\overline{(E - \bar{E})^2} &= \sum \rho_s (E_s - \bar{E})^2 \\ &= \sum \rho_s (E_s^2 - 2\bar{E}E_s + \bar{E}^2) \\ &= \sum E_s^2 \rho_s - 2\bar{E} \sum \rho_s E_s + \bar{E}^2 = \sum E_s^2 \rho_s - \bar{E}^2 \\ &= \overline{E^2} - \bar{E}^2;\end{aligned}$$
- $$\frac{\partial \bar{E}}{\partial \beta} = \frac{\partial \sum E_s \rho_s}{\partial \beta}$$

Fluctuation

- $$\begin{aligned}\overline{(E - \bar{E})^2} &= \sum \rho_s (E_s - \bar{E})^2 \\&= \sum \rho_s (E_s^2 - 2\bar{E}E_s + \bar{E}^2) \\&= \sum E_s^2 \rho_s - 2\bar{E} \sum \rho_s E_s + \bar{E}^2 = \sum E_s^2 \rho_s - \bar{E}^2 \\&= \overline{E^2} - \bar{E}^2;\end{aligned}$$
- $$\frac{\partial \bar{E}}{\partial \beta} = \frac{\partial \sum E_s \rho_s}{\partial \beta} = \frac{\partial \sum E_s e^{-\beta E_s}}{Z}$$

Fluctuation

- $$\begin{aligned}\overline{(E - \bar{E})^2} &= \sum \rho_s (E_s - \bar{E})^2 \\&= \sum \rho_s (E_s^2 - 2\bar{E}E_s + \bar{E}^2) \\&= \sum E_s^2 \rho_s - 2\bar{E} \sum \rho_s E_s + \bar{E}^2 = \sum E_s^2 \rho_s - \bar{E}^2 \\&= \overline{E^2} - \bar{E}^2;\end{aligned}$$
- $$\frac{\partial \bar{E}}{\partial \beta} = \frac{\partial \sum E_s \rho_s}{\partial \beta} = \frac{\partial \sum E_s e^{-\beta E_s}}{\partial \beta Z} = \frac{\partial \sum E_s e^{-\beta E_s}}{\partial \beta \sum e^{-\beta E_s}}$$

Fluctuation

- $$\begin{aligned}
 \overline{(E - \bar{E})^2} &= \sum \rho_s (E_s - \bar{E})^2 \\
 &= \sum \rho_s (E_s^2 - 2\bar{E}E_s + \bar{E}^2) \\
 &= \sum E_s^2 \rho_s - 2\bar{E} \sum \rho_s E_s + \bar{E}^2 = \sum E_s^2 \rho_s - \bar{E}^2 \\
 &= \overline{E^2} - \bar{E}^2;
 \end{aligned}$$
- $$\begin{aligned}
 \frac{\partial \bar{E}}{\partial \beta} &= \frac{\partial \sum E_s \rho_s}{\partial \beta} = \frac{\partial}{\partial \beta} \frac{\sum E_s e^{-\beta E_s}}{Z} = \frac{\partial}{\partial \beta} \frac{\sum E_s e^{-\beta E_s}}{\sum e^{-\beta E_s}} \\
 &= -\frac{\sum E_s^2 e^{-\beta E_s}}{\sum e^{-\beta E_s}} + \frac{(\sum E_s e^{-\beta E_s})^2}{(\sum e^{-\beta E_s})^2}
 \end{aligned}$$

Fluctuation

- $$\begin{aligned}
 \overline{(E - \bar{E})^2} &= \sum \rho_s (E_s - \bar{E})^2 \\
 &= \sum \rho_s (E_s^2 - 2\bar{E}E_s + \bar{E}^2) \\
 &= \sum E_s^2 \rho_s - 2\bar{E} \sum \rho_s E_s + \bar{E}^2 = \sum E_s^2 \rho_s - \bar{E}^2 \\
 &= \overline{E^2} - \bar{E}^2;
 \end{aligned}$$
- $$\begin{aligned}
 \frac{\partial \bar{E}}{\partial \beta} &= \frac{\partial \sum E_s \rho_s}{\partial \beta} = \frac{\partial}{\partial \beta} \frac{\sum E_s e^{-\beta E_s}}{Z} = \frac{\partial}{\partial \beta} \frac{\sum E_s e^{-\beta E_s}}{\sum e^{-\beta E_s}} \\
 &= -\frac{\sum E_s^2 e^{-\beta E_s}}{\sum e^{-\beta E_s}} + \frac{(\sum E_s e^{-\beta E_s})^2}{(\sum e^{-\beta E_s})^2} = -(\overline{E^2} - \bar{E}^2);
 \end{aligned}$$

Fluctuation

- $$\begin{aligned}
 \bullet \quad \overline{(E - \bar{E})^2} &= \sum \rho_s (E_s - \bar{E})^2 \\
 &= \sum \rho_s (E_s^2 - 2\bar{E}E_s + \bar{E}^2) \\
 &= \sum E_s^2 \rho_s - 2\bar{E} \sum \rho_s E_s + \bar{E}^2 = \sum E_s^2 \rho_s - \bar{E}^2 \\
 &= \overline{E^2} - \bar{E}^2;
 \end{aligned}$$
- $$\begin{aligned}
 \bullet \quad \frac{\partial \bar{E}}{\partial \beta} &= \frac{\partial \sum E_s \rho_s}{\partial \beta} = \frac{\partial}{\partial \beta} \frac{\sum E_s e^{-\beta E_s}}{Z} = \frac{\partial}{\partial \beta} \frac{\sum E_s e^{-\beta E_s}}{\sum e^{-\beta E_s}} \\
 &= -\frac{\sum E_s^2 e^{-\beta E_s}}{\sum e^{-\beta E_s}} + \frac{(\sum E_s e^{-\beta E_s})^2}{(\sum e^{-\beta E_s})^2} = -(\overline{E^2} - \bar{E}^2);
 \end{aligned}$$
- $$\bullet \quad \therefore \overline{(E - \bar{E})^2} = -\frac{\partial \bar{E}}{\partial \beta}$$

Fluctuation

- $$\begin{aligned}
 \bullet \quad \overline{(E - \bar{E})^2} &= \sum \rho_s (E_s - \bar{E})^2 \\
 &= \sum \rho_s (E_s^2 - 2\bar{E}E_s + \bar{E}^2) \\
 &= \sum E_s^2 \rho_s - 2\bar{E} \sum \rho_s E_s + \bar{E}^2 = \sum E_s^2 \rho_s - \bar{E}^2 \\
 &= \overline{E^2} - \bar{E}^2;
 \end{aligned}$$
- $$\begin{aligned}
 \bullet \quad \frac{\partial \bar{E}}{\partial \beta} &= \frac{\partial \sum E_s \rho_s}{\partial \beta} = \frac{\partial}{\partial \beta} \frac{\sum E_s e^{-\beta E_s}}{Z} = \frac{\partial}{\partial \beta} \frac{\sum E_s e^{-\beta E_s}}{\sum e^{-\beta E_s}} \\
 &= -\frac{\sum E_s^2 e^{-\beta E_s}}{\sum e^{-\beta E_s}} + \frac{(\sum E_s e^{-\beta E_s})^2}{(\sum e^{-\beta E_s})^2} = -(\overline{E^2} - \bar{E}^2);
 \end{aligned}$$
- $$\bullet \quad \therefore \overline{(E - \bar{E})^2} = -\frac{\partial \bar{E}}{\partial \beta} = kT^2 \frac{\partial \bar{E}}{\partial T}$$

Fluctuation

- $$\begin{aligned}
 \bullet \quad \overline{(E - \bar{E})^2} &= \sum \rho_s (E_s - \bar{E})^2 \\
 &= \sum \rho_s (E_s^2 - 2\bar{E}E_s + \bar{E}^2) \\
 &= \sum E_s^2 \rho_s - 2\bar{E} \sum \rho_s E_s + \bar{E}^2 = \sum E_s^2 \rho_s - \bar{E}^2 \\
 &= \overline{E^2} - \bar{E}^2;
 \end{aligned}$$
- $$\begin{aligned}
 \bullet \quad \frac{\partial \bar{E}}{\partial \beta} &= \frac{\partial \sum E_s \rho_s}{\partial \beta} = \frac{\partial}{\partial \beta} \frac{\sum E_s e^{-\beta E_s}}{Z} = \frac{\partial}{\partial \beta} \frac{\sum E_s e^{-\beta E_s}}{\sum e^{-\beta E_s}} \\
 &= -\frac{\sum E_s^2 e^{-\beta E_s}}{\sum e^{-\beta E_s}} + \frac{(\sum E_s e^{-\beta E_s})^2}{(\sum e^{-\beta E_s})^2} = -(\overline{E^2} - \bar{E}^2);
 \end{aligned}$$
- $$\bullet \quad \therefore \overline{(E - \bar{E})^2} = -\frac{\partial \bar{E}}{\partial \beta} = kT^2 \frac{\partial \bar{E}}{\partial T} = kT^2 \frac{\partial U}{\partial T}$$

Fluctuation

- $$\begin{aligned}
 \bullet \quad \overline{(E - \bar{E})^2} &= \sum \rho_s (E_s - \bar{E})^2 \\
 &= \sum \rho_s (E_s^2 - 2\bar{E}E_s + \bar{E}^2) \\
 &= \sum E_s^2 \rho_s - 2\bar{E} \sum \rho_s E_s + \bar{E}^2 = \sum E_s^2 \rho_s - \bar{E}^2 \\
 &= \overline{E^2} - \bar{E}^2;
 \end{aligned}$$
- $$\begin{aligned}
 \bullet \quad \frac{\partial \bar{E}}{\partial \beta} &= \frac{\partial \sum E_s \rho_s}{\partial \beta} = \frac{\partial}{\partial \beta} \frac{\sum E_s e^{-\beta E_s}}{Z} = \frac{\partial}{\partial \beta} \frac{\sum E_s e^{-\beta E_s}}{\sum e^{-\beta E_s}} \\
 &= -\frac{\sum E_s^2 e^{-\beta E_s}}{\sum e^{-\beta E_s}} + \frac{(\sum E_s e^{-\beta E_s})^2}{(\sum e^{-\beta E_s})^2} = -(\overline{E^2} - \bar{E}^2);
 \end{aligned}$$
- $$\bullet \quad \therefore \overline{(E - \bar{E})^2} = -\frac{\partial \bar{E}}{\partial \beta} = kT^2 \frac{\partial \bar{E}}{\partial T} = kT^2 \frac{\partial U}{\partial T} = kT^2 C_V.$$

Fluctuation

- $$\begin{aligned} \overline{(E - \bar{E})^2} &= \sum \rho_s (E_s - \bar{E})^2 \\ &= \sum \rho_s (E_s^2 - 2\bar{E}E_s + \bar{E}^2) \\ &= \sum E_s^2 \rho_s - 2\bar{E} \sum \rho_s E_s + \bar{E}^2 = \sum E_s^2 \rho_s - \bar{E}^2 \\ &= \overline{E^2} - \bar{E}^2; \end{aligned}$$
- $$\begin{aligned} \frac{\partial \bar{E}}{\partial \beta} &= \frac{\partial \sum E_s \rho_s}{\partial \beta} = \frac{\partial}{\partial \beta} \frac{\sum E_s e^{-\beta E_s}}{Z} = \frac{\partial}{\partial \beta} \frac{\sum E_s e^{-\beta E_s}}{\sum e^{-\beta E_s}} \\ &= -\frac{\sum E_s^2 e^{-\beta E_s}}{\sum e^{-\beta E_s}} + \frac{(\sum E_s e^{-\beta E_s})^2}{(\sum e^{-\beta E_s})^2} = -(\overline{E^2} - \bar{E}^2); \end{aligned}$$
- $$\therefore \overline{(E - \bar{E})^2} = -\frac{\partial \bar{E}}{\partial \beta} = kT^2 \frac{\partial \bar{E}}{\partial T} = kT^2 \frac{\partial U}{\partial T} = kT^2 C_V.$$
- $$\text{Relative fluctuation: } \frac{\overline{(E - \bar{E})^2}}{\bar{E}^2} = \frac{kT^2 C_V}{\bar{E}^2}$$

Fluctuation

- $$\begin{aligned} \overline{(E - \bar{E})^2} &= \sum \rho_s (E_s - \bar{E})^2 \\ &= \sum \rho_s (E_s^2 - 2\bar{E}E_s + \bar{E}^2) \\ &= \sum E_s^2 \rho_s - 2\bar{E} \sum \rho_s E_s + \bar{E}^2 = \sum E_s^2 \rho_s - \bar{E}^2 \\ &= \overline{E^2} - \bar{E}^2; \end{aligned}$$
- $$\begin{aligned} \frac{\partial \bar{E}}{\partial \beta} &= \frac{\partial \sum E_s \rho_s}{\partial \beta} = \frac{\partial}{\partial \beta} \frac{\sum E_s e^{-\beta E_s}}{Z} = \frac{\partial}{\partial \beta} \frac{\sum E_s e^{-\beta E_s}}{\sum e^{-\beta E_s}} \\ &= -\frac{\sum E_s^2 e^{-\beta E_s}}{\sum e^{-\beta E_s}} + \frac{(\sum E_s e^{-\beta E_s})^2}{(\sum e^{-\beta E_s})^2} = -(\overline{E^2} - \bar{E}^2); \end{aligned}$$
- $$\therefore \overline{(E - \bar{E})^2} = -\frac{\partial \bar{E}}{\partial \beta} = kT^2 \frac{\partial \bar{E}}{\partial T} = kT^2 \frac{\partial U}{\partial T} = kT^2 C_V.$$
- $$\text{Relative fluctuation: } \frac{\overline{(E - \bar{E})^2}}{\bar{E}^2} = \frac{kT^2 C_V}{\bar{E}^2} \propto \frac{1}{N}.$$

Fluctuation

- $$\begin{aligned}
 \bullet \quad \overline{(E - \bar{E})^2} &= \sum \rho_s (E_s - \bar{E})^2 \\
 &= \sum \rho_s (E_s^2 - 2\bar{E}E_s + \bar{E}^2) \\
 &= \sum E_s^2 \rho_s - 2\bar{E} \sum \rho_s E_s + \bar{E}^2 = \sum E_s^2 \rho_s - \bar{E}^2 \\
 &= \overline{E^2} - \bar{E}^2;
 \end{aligned}$$
- $$\begin{aligned}
 \bullet \quad \frac{\partial \bar{E}}{\partial \beta} &= \frac{\partial \sum E_s \rho_s}{\partial \beta} = \frac{\partial}{\partial \beta} \frac{\sum E_s e^{-\beta E_s}}{Z} = \frac{\partial}{\partial \beta} \frac{\sum E_s e^{-\beta E_s}}{\sum e^{-\beta E_s}} \\
 &= -\frac{\sum E_s^2 e^{-\beta E_s}}{\sum e^{-\beta E_s}} + \frac{(\sum E_s e^{-\beta E_s})^2}{(\sum e^{-\beta E_s})^2} = -(\overline{E^2} - \bar{E}^2);
 \end{aligned}$$
- $$\bullet \quad \therefore \overline{(E - \bar{E})^2} = -\frac{\partial \bar{E}}{\partial \beta} = kT^2 \frac{\partial \bar{E}}{\partial T} = kT^2 \frac{\partial U}{\partial T} = kT^2 C_V.$$
- $$\bullet \quad \text{Relative fluctuation: } \frac{\overline{(E - \bar{E})^2}}{\bar{E}^2} = \frac{kT^2 C_V}{\bar{E}^2} \propto \frac{1}{N}.$$

For $N \gg 1$, the relative fluctuation $\rightarrow 0$;

Fluctuation

- $$\begin{aligned} \overline{(E - \bar{E})^2} &= \sum \rho_s (E_s - \bar{E})^2 \\ &= \sum \rho_s (E_s^2 - 2\bar{E}E_s + \bar{E}^2) \\ &= \sum E_s^2 \rho_s - 2\bar{E} \sum \rho_s E_s + \bar{E}^2 = \sum E_s^2 \rho_s - \bar{E}^2 \\ &= \overline{E^2} - \bar{E}^2; \end{aligned}$$
- $$\begin{aligned} \frac{\partial \bar{E}}{\partial \beta} &= \frac{\partial \sum E_s \rho_s}{\partial \beta} = \frac{\partial}{\partial \beta} \frac{\sum E_s e^{-\beta E_s}}{Z} = \frac{\partial}{\partial \beta} \frac{\sum E_s e^{-\beta E_s}}{\sum e^{-\beta E_s}} \\ &= -\frac{\sum E_s^2 e^{-\beta E_s}}{\sum e^{-\beta E_s}} + \frac{(\sum E_s e^{-\beta E_s})^2}{(\sum e^{-\beta E_s})^2} = -(\overline{E^2} - \bar{E}^2); \end{aligned}$$
- $$\therefore \overline{(E - \bar{E})^2} = -\frac{\partial \bar{E}}{\partial \beta} = kT^2 \frac{\partial \bar{E}}{\partial T} = kT^2 \frac{\partial U}{\partial T} = kT^2 C_V.$$
- Relative fluctuation: $\frac{\overline{(E - \bar{E})^2}}{\bar{E}^2} = \frac{kT^2 C_V}{\bar{E}^2} \propto \frac{1}{N}.$

For $N \gg 1$, the relative fluctuation $\rightarrow 0$; back to microcanonical ensemble (equivalent).

Table of contents

1 §9. Ensemble theory

- 9.1 Phase space and Liouville's theorem
- 9.2 The microcanonical ensemble
- 9.3 Thermodynamics in microcanonical ensemble theory
- 9.4 Canonical ensemble
- 9.5 Thermodynamics in canonical ensemble
- **9.6 EOS of real gas**
- 9.7 Thermal capacity of solid
- 9.9 Ising model
- 9.10 Grand canonical ensemble
- 9.11 Thermodynamics in grand canonical ensemble
- 9.12 Applications of the grand canonical ensemble

9.6 EOS of real gas – 1st application.

9.6 EOS of real gas – 1st application.

- Boltzmann, Bose, Fermi statistics can only deal with ideal gas (nearly no interaction).

9.6 EOS of real gas – 1st application.

- Boltzmann, Bose, Fermi statistics can only deal with ideal gas (nearly no interaction). Ensemble theory is suitable for real gas.

9.6 EOS of real gas – 1st application.

- Boltzmann, Bose, Fermi statistics can only deal with ideal gas (nearly no interaction). Ensemble theory is suitable for real gas.

- 1st step: partition function,

$$Z = \frac{1}{N!h^{3N}} \int e^{-\beta E} dq_1 \dots dq_{3N} dp_1 \dots dp_{3N};$$

9.6 EOS of real gas – 1st application.

- Boltzmann, Bose, Fermi statistics can only deal with ideal gas (nearly no interaction). Ensemble theory is suitable for real gas.
- 1st step: partition function,
$$Z = \frac{1}{N!h^{3N}} \int e^{-\beta E} dq_1 \dots dq_{3N} dp_1 \dots dp_{3N};$$
- 2nd step: thermal quantities, like $p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln Z$.

9.6 EOS of real gas – 1st application.

- Boltzmann, Bose, Fermi statistics can only deal with ideal gas (nearly no interaction). Ensemble theory is suitable for real gas.
- 1st step: partition function,
$$Z = \frac{1}{N!h^{3N}} \int e^{-\beta E} dq_1 \dots dq_{3N} dp_1 \dots dp_{3N};$$
- 2nd step: thermal quantities, like $p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln Z$.
- E (kinetic energy and potential energy):

9.6 EOS of real gas – 1st application.

- Boltzmann, Bose, Fermi statistics can only deal with ideal gas (nearly no interaction). Ensemble theory is suitable for real gas.

- 1st step: partition function,

$$Z = \frac{1}{N!h^{3N}} \int e^{-\beta E} dq_1 \dots dq_{3N} dp_1 \dots dp_{3N};$$

- 2nd step: thermal quantities, like $p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln Z$.

- E (kinetic energy and potential energy):

$$E = \sum_{i=1}^N \frac{\vec{p}_i^2}{2m} + \sum_{i < j} \phi(\vec{r}_{ij}).$$

EOS of real gas

$$Z = \frac{1}{N!h^{3N}} \int e^{-\beta E} dq_1 \dots dq_{3N} dp_1 \dots dp_{3N}$$

- Kinetic term of E :

$$\int e^{-\beta \sum_{i=1}^N \frac{\vec{p}_i^2}{2m}} dp_1 \dots dp_{3N}$$

EOS of real gas

$$Z = \frac{1}{N!h^{3N}} \int e^{-\beta E} dq_1 \dots dq_{3N} dp_1 \dots dp_{3N}$$

- Kinetic term of E :

$$\int e^{-\beta \sum_{i=1}^N \frac{\vec{p}_i^2}{2m}} dp_1 \dots dp_{3N} = \int e^{-\beta \sum_{i=1}^{3N} \frac{p_i^2}{2m}} dp_1 \dots dp_{3N}$$

EOS of real gas

$$Z = \frac{1}{N!h^{3N}} \int e^{-\beta E} dq_1 \dots dq_{3N} dp_1 \dots dp_{3N}$$

- Kinetic term of E :

$$\begin{aligned} \int e^{-\beta \sum_{i=1}^N \frac{\vec{p}_i^2}{2m}} dp_1 \dots dp_{3N} &= \int e^{-\beta \sum_{i=1}^{3N} \frac{p_i^2}{2m}} dp_1 \dots dp_{3N} \\ &= \prod_{i=1}^{3N} \int e^{-\beta \frac{p_i^2}{2m}} dp_i \end{aligned}$$

EOS of real gas

$$Z = \frac{1}{N!h^{3N}} \int e^{-\beta E} dq_1 \dots dq_{3N} dp_1 \dots dp_{3N}$$

- Kinetic term of E :

$$\begin{aligned} \int e^{-\beta \sum_{i=1}^N \frac{\vec{p}_i^2}{2m}} dp_1 \dots dp_{3N} &= \int e^{-\beta \sum_{i=1}^{3N} \frac{p_i^2}{2m}} dp_1 \dots dp_{3N} \\ &= \prod_{i=1}^{3N} \int e^{-\beta \frac{p_i^2}{2m}} dp_i = \left(\frac{2\pi m}{\beta} \right)^{\frac{3}{2}N}. \end{aligned}$$

EOS of real gas

$$Z = \frac{1}{N!h^{3N}} \int e^{-\beta E} dq_1 \dots dq_{3N} dp_1 \dots dp_{3N}$$

- Kinetic term of E :

$$\begin{aligned} \int e^{-\beta \sum_{i=1}^N \frac{\vec{p}_i^2}{2m}} dp_1 \dots dp_{3N} &= \int e^{-\beta \sum_{i=1}^{3N} \frac{p_i^2}{2m}} dp_1 \dots dp_{3N} \\ &= \prod_{i=1}^{3N} \int e^{-\beta \frac{p_i^2}{2m}} dp_i = \left(\frac{2\pi m}{\beta} \right)^{\frac{3}{2}N}. \end{aligned}$$

- Potential part: $Q \equiv \int e^{-\beta \sum_{i<j} \phi(\vec{r}_{ij})} dq_1 \dots dq_{3N}$

EOS of real gas

$$Z = \frac{1}{N!h^{3N}} \int e^{-\beta E} dq_1 \dots dq_{3N} dp_1 \dots dp_{3N}$$

- Kinetic term of E :

$$\begin{aligned} \int e^{-\beta \sum_{i=1}^N \frac{\vec{p}_i^2}{2m}} dp_1 \dots dp_{3N} &= \int e^{-\beta \sum_{i=1}^{3N} \frac{p_i^2}{2m}} dp_1 \dots dp_{3N} \\ &= \prod_{i=1}^{3N} \int e^{-\beta \frac{p_i^2}{2m}} dp_i = \left(\frac{2\pi m}{\beta} \right)^{\frac{3}{2}N}. \end{aligned}$$

- Potential part: $Q \equiv \int e^{-\beta \sum_{i<j} \phi(\vec{r}_{ij})} dq_1 \dots dq_{3N}$
 $= \int e^{-\beta \sum_{i<j} \phi(\vec{r}_{ij})} d\tau_1 \dots d\tau_N.$

EOS of real gas

$$Z = \frac{1}{N!h^{3N}} \int e^{-\beta E} dq_1 \dots dq_{3N} dp_1 \dots dp_{3N}$$

- Kinetic term of E :

$$\begin{aligned} \int e^{-\beta \sum_{i=1}^N \frac{\vec{p}_i^2}{2m}} dp_1 \dots dp_{3N} &= \int e^{-\beta \sum_{i=1}^{3N} \frac{p_i^2}{2m}} dp_1 \dots dp_{3N} \\ &= \prod_{i=1}^{3N} \int e^{-\beta \frac{p_i^2}{2m}} dp_i = \left(\frac{2\pi m}{\beta} \right)^{\frac{3}{2}N}. \end{aligned}$$

- Potential part: $Q \equiv \int e^{-\beta \sum_{i<j} \phi(\vec{r}_{ij})} dq_1 \dots dq_{3N}$
 $= \int e^{-\beta \sum_{i<j} \phi(\vec{r}_{ij})} d\tau_1 \dots d\tau_N.$
- Then the partition function: $Z = \frac{1}{N!} \left(\frac{2\pi m}{\beta h^2} \right)^{\frac{3}{2}N} Q.$

EOS of real gas

$$Z = \frac{1}{N!h^{3N}} \int e^{-\beta E} dq_1 \dots dq_{3N} dp_1 \dots dp_{3N}$$

- Kinetic term of E :

$$\begin{aligned} \int e^{-\beta \sum_{i=1}^N \frac{\vec{p}_i^2}{2m}} dp_1 \dots dp_{3N} &= \int e^{-\beta \sum_{i=1}^{3N} \frac{p_i^2}{2m}} dp_1 \dots dp_{3N} \\ &= \prod_{i=1}^{3N} \int e^{-\beta \frac{p_i^2}{2m}} dp_i = \left(\frac{2\pi m}{\beta} \right)^{\frac{3}{2}N}. \end{aligned}$$

- Potential part: $Q \equiv \int e^{-\beta \sum_{i<j} \phi(\vec{r}_{ij})} dq_1 \dots dq_{3N}$
 $= \int e^{-\beta \sum_{i<j} \phi(\vec{r}_{ij})} d\tau_1 \dots d\tau_N.$
- Then the partition function: $Z = \frac{1}{N!} \left(\frac{2\pi m}{\beta h^2} \right)^{\frac{3}{2}N} Q.$
- Key: $Q.$

EOS of real gas

$$Q = \int e^{-\beta \sum_{i < j} \phi(\vec{r}_{ij})} d\tau_1 \dots d\tau_N$$

- Define $f_{ij} = e^{-\beta \phi(\vec{r}_{ij})} - 1$,

EOS of real gas

$$Q = \int e^{-\beta \sum_{i < j} \phi(\vec{r}_{ij})} d\tau_1 \dots d\tau_N$$

- Define $f_{ij} = e^{-\beta \phi(\vec{r}_{ij})} - 1$,
- If \vec{r}_{ij} is out of the interacting range, $\phi(\vec{r}_{ij}) = 0$, then $f_{ij} = 0$.

EOS of real gas

$$Q = \int e^{-\beta \sum_{i < j} \phi(\vec{r}_{ij})} d\tau_1 \dots d\tau_N$$

- Define $f_{ij} = e^{-\beta \phi(\vec{r}_{ij})} - 1$,
- If \vec{r}_{ij} is out of the interacting range, $\phi(\vec{r}_{ij}) = 0$, then $f_{ij} = 0$. Only the nearby neighbours are interacting.

EOS of real gas

$$Q = \int e^{-\beta \sum_{i < j} \phi(\vec{r}_{ij})} d\tau_1 \dots d\tau_N$$

- Define $f_{ij} = e^{-\beta \phi(\vec{r}_{ij})} - 1$,
- If \vec{r}_{ij} is out of the interacting range, $\phi(\vec{r}_{ij}) = 0$, then $f_{ij} = 0$. Only the nearby neighbours are interacting.
- $Q = \int \prod_{i < j} (1 + f_{ij}) d\tau_1 \dots d\tau_N$

EOS of real gas

$$Q = \int e^{-\beta \sum_{i < j} \phi(\vec{r}_{ij})} d\tau_1 \dots d\tau_N$$

- Define $f_{ij} = e^{-\beta \phi(\vec{r}_{ij})} - 1$,
- If \vec{r}_{ij} is out of the interacting range, $\phi(\vec{r}_{ij}) = 0$, then $f_{ij} = 0$. Only the nearby neighbours are interacting.
- $$Q = \int \prod_{i < j} (1 + f_{ij}) d\tau_1 \dots d\tau_N$$

$$= \int \left(1 + \sum_{i < j} f_{ij} + \sum_{i < j} \sum_{i' < j'} f_{ij} f_{i'j'} + \dots \right) d\tau_1 \dots d\tau_N.$$

EOS of real gas

$$Q = \int e^{-\beta \sum_{i<j} \phi(\vec{r}_{ij})} d\tau_1 \dots d\tau_N$$

- Define $f_{ij} = e^{-\beta \phi(\vec{r}_{ij})} - 1$,
- If \vec{r}_{ij} is out of the interacting range, $\phi(\vec{r}_{ij}) = 0$, then $f_{ij} = 0$. Only the nearby neighbours are interacting.
- $Q = \int \prod_{i<j} (1 + f_{ij}) d\tau_1 \dots d\tau_N$
 $= \int (1 + \sum_{i<j} f_{ij} + \sum_{i<j} \sum_{i'<j'} f_{ij} f_{i'j'} + \dots) d\tau_1 \dots d\tau_N.$
- If just keep “1”, no potential, it goes back to ideal gas. $Q = V^N$.

EOS of real gas

$$Q = \int e^{-\beta \sum_{i < j} \phi(\vec{r}_{ij})} d\tau_1 \dots d\tau_N$$

- Define $f_{ij} = e^{-\beta \phi(\vec{r}_{ij})} - 1$,
- If \vec{r}_{ij} is out of the interacting range, $\phi(\vec{r}_{ij}) = 0$, then $f_{ij} = 0$. Only the nearby neighbours are interacting.
- $$Q = \int \prod_{i < j} (1 + f_{ij}) d\tau_1 \dots d\tau_N$$

$$= \int (1 + \sum_{i < j} f_{ij} + \sum_{i < j} \sum_{i' < j'} f_{ij} f_{i'j'} + \dots) d\tau_1 \dots d\tau_N.$$
- If just keep “1”, no potential, it goes back to ideal gas. $Q = V^N$.
Then $Z = \frac{1}{N!} \left(\frac{2\pi m}{\beta h^2} \right)^{\frac{3}{2}N} V^N$.

EOS of real gas

$$Q = \int e^{-\beta \sum_{i < j} \phi(\vec{r}_{ij})} d\tau_1 \dots d\tau_N$$

- Define $f_{ij} = e^{-\beta \phi(\vec{r}_{ij})} - 1$,
- If \vec{r}_{ij} is out of the interacting range, $\phi(\vec{r}_{ij}) = 0$, then $f_{ij} = 0$. Only the nearby neighbours are interacting.
- $Q = \int \prod_{i < j} (1 + f_{ij}) d\tau_1 \dots d\tau_N$
 $= \int (1 + \sum_{i < j} f_{ij} + \sum_{i < j} \sum_{i' < j'} f_{ij} f_{i'j'} + \dots) d\tau_1 \dots d\tau_N$.
- If just keep “1”, no potential, it goes back to ideal gas. $Q = V^N$.

$$\text{Then } Z = \frac{1}{N!} \left(\frac{2\pi m}{\beta h^2} \right)^{\frac{3}{2}N} V^N.$$

$$p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln Z$$

EOS of real gas

$$Q = \int e^{-\beta \sum_{i < j} \phi(\vec{r}_{ij})} d\tau_1 \dots d\tau_N$$

- Define $f_{ij} = e^{-\beta \phi(\vec{r}_{ij})} - 1$,
- If \vec{r}_{ij} is out of the interacting range, $\phi(\vec{r}_{ij}) = 0$, then $f_{ij} = 0$. Only the nearby neighbours are interacting.
- $Q = \int \prod_{i < j} (1 + f_{ij}) d\tau_1 \dots d\tau_N$
 $= \int (1 + \sum_{i < j} f_{ij} + \sum_{i < j} \sum_{i' < j'} f_{ij} f_{i'j'} + \dots) d\tau_1 \dots d\tau_N$.
- If just keep “1”, no potential, it goes back to ideal gas. $Q = V^N$.

$$\text{Then } Z = \frac{1}{N!} \left(\frac{2\pi m}{\beta h^2} \right)^{\frac{3}{2}N} V^N.$$

$$p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln Z = \frac{1}{\beta} \frac{\partial \ln(CV^N)}{\partial V}$$

EOS of real gas

$$Q = \int e^{-\beta \sum_{i < j} \phi(\vec{r}_{ij})} d\tau_1 \dots d\tau_N$$

- Define $f_{ij} = e^{-\beta \phi(\vec{r}_{ij})} - 1$,
- If \vec{r}_{ij} is out of the interacting range, $\phi(\vec{r}_{ij}) = 0$, then $f_{ij} = 0$. Only the nearby neighbours are interacting.
- $Q = \int \prod_{i < j} (1 + f_{ij}) d\tau_1 \dots d\tau_N$
 $= \int (1 + \sum_{i < j} f_{ij} + \sum_{i < j} \sum_{i' < j'} f_{ij} f_{i'j'} + \dots) d\tau_1 \dots d\tau_N$.
- If just keep “1”, no potential, it goes back to ideal gas. $Q = V^N$.

$$\text{Then } Z = \frac{1}{N!} \left(\frac{2\pi m}{\beta h^2} \right)^{\frac{3}{2}N} V^N.$$

$$p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln Z = \frac{1}{\beta} \frac{\partial \ln(CV^N)}{\partial V} = \frac{1}{\beta} \frac{N}{V}$$

EOS of real gas

$$Q = \int e^{-\beta \sum_{i < j} \phi(\vec{r}_{ij})} d\tau_1 \dots d\tau_N$$

- Define $f_{ij} = e^{-\beta \phi(\vec{r}_{ij})} - 1$,
- If \vec{r}_{ij} is out of the interacting range, $\phi(\vec{r}_{ij}) = 0$, then $f_{ij} = 0$. Only the nearby neighbours are interacting.
- $Q = \int \prod_{i < j} (1 + f_{ij}) d\tau_1 \dots d\tau_N$
 $= \int (1 + \sum_{i < j} f_{ij} + \sum_{i < j} \sum_{i' < j'} f_{ij} f_{i'j'} + \dots) d\tau_1 \dots d\tau_N$.
- If just keep “1”, no potential, it goes back to ideal gas. $Q = V^N$.

$$\text{Then } Z = \frac{1}{N!} \left(\frac{2\pi m}{\beta h^2} \right)^{\frac{3}{2}N} V^N.$$

$$p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln Z = \frac{1}{\beta} \frac{\partial \ln(CV^N)}{\partial V} = \frac{1}{\beta} \frac{N}{V} = \frac{NkT}{V}.$$

EOS of real gas

- Else if the interacting is weak, i.e., $\phi(\vec{r}_{ij}) \rightarrow 0$, then $f_{ij} \rightarrow 0$.

EOS of real gas

- Else if the interacting is weak, i.e., $\phi(\vec{r}_{ij}) \rightarrow 0$, then $f_{ij} \rightarrow 0$. Only keep the first two terms:

EOS of real gas

- Else if the interacting is weak, i.e., $\phi(\vec{r}_{ij}) \rightarrow 0$, then $f_{ij} \rightarrow 0$. Only keep the first two terms:

$$Q = \int (1 + \sum_{i < j} f_{ij}) d\tau_1 \dots d\tau_N.$$

EOS of real gas

- Else if the interacting is weak, i.e., $\phi(\vec{r}_{ij}) \rightarrow 0$, then $f_{ij} \rightarrow 0$. Only keep the first two terms:
$$Q = \int (1 + \sum_{i < j} f_{ij}) d\tau_1 \dots d\tau_N.$$
- As there is no difference for the $\frac{1}{2}N(N-1)$ summation (neglect the boundary effect),

EOS of real gas

- Else if the interacting is weak, i.e., $\phi(\vec{r}_{ij}) \rightarrow 0$, then $f_{ij} \rightarrow 0$. Only keep the first two terms:

$$Q = \int (1 + \sum_{i < j} f_{ij}) d\tau_1 \dots d\tau_N.$$

- As there is no difference for the $\frac{1}{2}N(N-1)$ summation (neglect the boundary effect),
$$\int \sum_{i < j} f_{ij} d\tau_1 \dots d\tau_N = \frac{1}{2}N(N-1) \int f_{12} d\tau_1 \dots d\tau_N$$

EOS of real gas

- Else if the interacting is weak, i.e., $\phi(\vec{r}_{ij}) \rightarrow 0$, then $f_{ij} \rightarrow 0$. Only keep the first two terms:

$$Q = \int (1 + \sum_{i < j} f_{ij}) d\tau_1 \dots d\tau_N.$$

- As there is no difference for the $\frac{1}{2}N(N-1)$ summation (neglect the boundary effect),

$$\int \sum_{i < j} f_{ij} d\tau_1 \dots d\tau_N = \frac{1}{2}N(N-1) \int f_{12} d\tau_1 \dots d\tau_N$$

$$= \frac{1}{2}N(N-1)V^{N-2} \int f_{12} d\tau_1 d\tau_2.$$

EOS of real gas

- Else if the interacting is weak, i.e., $\phi(\vec{r}_{ij}) \rightarrow 0$, then $f_{ij} \rightarrow 0$. Only keep the first two terms:

$$Q = \int (1 + \sum_{i < j} f_{ij}) d\tau_1 \dots d\tau_N.$$

- As there is no difference for the $\frac{1}{2}N(N-1)$ summation (neglect the boundary effect),

$$\int \sum_{i < j} f_{ij} d\tau_1 \dots d\tau_N = \frac{1}{2}N(N-1) \int f_{12} d\tau_1 \dots d\tau_N$$

$$= \frac{1}{2}N(N-1)V^{N-2} \int f_{12} d\tau_1 d\tau_2.$$
- Integration does not depend on molecule 1's position,

EOS of real gas

- Else if the interacting is weak, i.e., $\phi(\vec{r}_{ij}) \rightarrow 0$, then $f_{ij} \rightarrow 0$. Only keep the first two terms:

$$Q = \int (1 + \sum_{i < j} f_{ij}) d\tau_1 \dots d\tau_N.$$

- As there is no difference for the $\frac{1}{2}N(N-1)$ summation (neglect the boundary effect),

$$\begin{aligned} \int \sum_{i < j} f_{ij} d\tau_1 \dots d\tau_N &= \frac{1}{2}N(N-1) \int f_{12} d\tau_1 \dots d\tau_N \\ &= \frac{1}{2}N(N-1)V^{N-2} \int f_{12} d\tau_1 d\tau_2. \end{aligned}$$

- Integration does not depend on molecule 1's position, $\int f_{12} d\tau_1 d\tau_2 = V \int f_{12} d\tau_2 = V \int f_{12} d\vec{r}$.

EOS of real gas

- Else if the interacting is weak, i.e., $\phi(\vec{r}_{ij}) \rightarrow 0$, then $f_{ij} \rightarrow 0$. Only keep the first two terms:

$$Q = \int (1 + \sum_{i < j} f_{ij}) d\tau_1 \dots d\tau_N.$$

- As there is no difference for the $\frac{1}{2}N(N-1)$ summation (neglect the boundary effect),

$$\begin{aligned} \int \sum_{i < j} f_{ij} d\tau_1 \dots d\tau_N &= \frac{1}{2}N(N-1) \int f_{12} d\tau_1 \dots d\tau_N \\ &= \frac{1}{2}N(N-1)V^{N-2} \int f_{12} d\tau_1 d\tau_2. \end{aligned}$$

- Integration does not depend on molecule 1's position,

$$\int f_{12} d\tau_1 d\tau_2 = V \int f_{12} d\tau_2 = V \int f_{12} d\vec{r}.$$

- $Q = V^N + \frac{1}{2}N(N-1)V^{N-1} \int f_{12} d\vec{r}$

EOS of real gas

- Else if the interacting is weak, i.e., $\phi(\vec{r}_{ij}) \rightarrow 0$, then $f_{ij} \rightarrow 0$. Only keep the first two terms:

$$Q = \int (1 + \sum_{i < j} f_{ij}) d\tau_1 \dots d\tau_N.$$

- As there is no difference for the $\frac{1}{2}N(N-1)$ summation (neglect the boundary effect),

$$\begin{aligned} \int \sum_{i < j} f_{ij} d\tau_1 \dots d\tau_N &= \frac{1}{2}N(N-1) \int f_{12} d\tau_1 \dots d\tau_N \\ &= \frac{1}{2}N(N-1)V^{N-2} \int f_{12} d\tau_1 d\tau_2. \end{aligned}$$

- Integration does not depend on molecule 1's position,

$$\int f_{12} d\tau_1 d\tau_2 = V \int f_{12} d\tau_2 = V \int f_{12} d\vec{r}.$$

- $Q = V^N + \frac{1}{2}N(N-1)V^{N-1} \int f_{12} d\vec{r}$
 $\simeq V^N + \frac{1}{2}N^2V^{N-1} \int f_{12} d\vec{r}$

EOS of real gas

- Else if the interacting is weak, i.e., $\phi(\vec{r}_{ij}) \rightarrow 0$, then $f_{ij} \rightarrow 0$. Only keep the first two terms:

$$Q = \int (1 + \sum_{i < j} f_{ij}) d\tau_1 \dots d\tau_N.$$

- As there is no difference for the $\frac{1}{2}N(N-1)$ summation (neglect the boundary effect),

$$\begin{aligned} \int \sum_{i < j} f_{ij} d\tau_1 \dots d\tau_N &= \frac{1}{2}N(N-1) \int f_{12} d\tau_1 \dots d\tau_N \\ &= \frac{1}{2}N(N-1)V^{N-2} \int f_{12} d\tau_1 d\tau_2. \end{aligned}$$

- Integration does not depend on molecule 1's position,

$$\int f_{12} d\tau_1 d\tau_2 = V \int f_{12} d\tau_2 = V \int f_{12} d\vec{r}.$$

- $Q = V^N + \frac{1}{2}N(N-1)V^{N-1} \int f_{12} d\vec{r}$
 $\simeq V^N + \frac{1}{2}N^2V^{N-1} \int f_{12} d\vec{r} = V^N(1 + \frac{N^2}{2V} \int f_{12} d\vec{r}).$

EOS of real gas $Q = V^N \left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r} \right)$

- $\ln Q = N \ln V + \ln \left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r} \right).$

EOS of real gas $Q = V^N \left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r} \right)$

- $\ln Q = N \ln V + \ln \left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r} \right).$
- For weak interaction,
 $\ln \left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r} \right) \simeq \frac{N^2}{2V} \int f_{12} d\vec{r}.$

EOS of real gas $Q = V^N \left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r} \right)$

- $\ln Q = N \ln V + \ln \left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r} \right).$
- For weak interaction,
 $\ln \left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r} \right) \simeq \frac{N^2}{2V} \int f_{12} d\vec{r}.$
 $\therefore \ln Q \simeq N \ln V + \frac{N^2}{2V} \int f_{12} d\vec{r}.$

EOS of real gas $Q = V^N \left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r}\right)$

- $\ln Q = N \ln V + \ln\left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r}\right)$.
- For weak interaction,
 $\ln\left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r}\right) \simeq \frac{N^2}{2V} \int f_{12} d\vec{r}$.
 $\therefore \ln Q \simeq N \ln V + \frac{N^2}{2V} \int f_{12} d\vec{r}$.
- As in $Z = \frac{1}{N!} \left(\frac{2\pi m}{\beta h^2}\right)^{\frac{3}{2}N} Q$, only Q depends on V ,

EOS of real gas $Q = V^N \left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r}\right)$

- $\ln Q = N \ln V + \ln\left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r}\right).$
- For weak interaction,
 $\ln\left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r}\right) \simeq \frac{N^2}{2V} \int f_{12} d\vec{r}.$
 $\therefore \ln Q \simeq N \ln V + \frac{N^2}{2V} \int f_{12} d\vec{r}.$
- As in $Z = \frac{1}{N!} \left(\frac{2\pi m}{\beta h^2}\right)^{\frac{3}{2}N} Q$, only Q depends on V ,
 $p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln Z$

EOS of real gas $Q = V^N \left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r}\right)$

- $\ln Q = N \ln V + \ln\left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r}\right)$.
- For weak interaction,
 $\ln\left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r}\right) \simeq \frac{N^2}{2V} \int f_{12} d\vec{r}$.
 $\therefore \ln Q \simeq N \ln V + \frac{N^2}{2V} \int f_{12} d\vec{r}$.
- As in $Z = \frac{1}{N!} \left(\frac{2\pi m}{\beta h^2}\right)^{\frac{3}{2}N} Q$, only Q depends on V ,
 $p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln Z = \frac{1}{\beta} \frac{\partial}{\partial V} \ln Q$

EOS of real gas $Q = V^N \left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r}\right)$

- $\ln Q = N \ln V + \ln\left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r}\right)$.
- For weak interaction,
 $\ln\left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r}\right) \simeq \frac{N^2}{2V} \int f_{12} d\vec{r}$.
 $\therefore \ln Q \simeq N \ln V + \frac{N^2}{2V} \int f_{12} d\vec{r}$.
- As in $Z = \frac{1}{N!} \left(\frac{2\pi m}{\beta h^2}\right)^{\frac{3}{2}N} Q$, only Q depends on V ,
 $p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln Z = \frac{1}{\beta} \frac{\partial}{\partial V} \ln Q$
 $= \frac{1}{\beta} \frac{\partial}{\partial V} \left(N \ln V + \frac{N^2}{2V} \int f_{12} d\vec{r}\right)$

EOS of real gas $Q = V^N \left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r}\right)$

- $\ln Q = N \ln V + \ln\left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r}\right).$
- For weak interaction,
 $\ln\left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r}\right) \simeq \frac{N^2}{2V} \int f_{12} d\vec{r}.$
 $\therefore \ln Q \simeq N \ln V + \frac{N^2}{2V} \int f_{12} d\vec{r}.$
- As in $Z = \frac{1}{N!} \left(\frac{2\pi m}{\beta h^2}\right)^{\frac{3}{2}N} Q$, only Q depends on V ,
 $p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln Z = \frac{1}{\beta} \frac{\partial}{\partial V} \ln Q$
 $= \frac{1}{\beta} \frac{\partial}{\partial V} \left(N \ln V + \frac{N^2}{2V} \int f_{12} d\vec{r}\right) = \frac{1}{\beta} \left(\frac{N}{V} - \frac{N^2}{2V^2} \int f_{12} d\vec{r}\right)$

EOS of real gas $Q = V^N \left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r}\right)$

- $\ln Q = N \ln V + \ln\left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r}\right).$
- For weak interaction,
 $\ln\left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r}\right) \simeq \frac{N^2}{2V} \int f_{12} d\vec{r}.$
 $\therefore \ln Q \simeq N \ln V + \frac{N^2}{2V} \int f_{12} d\vec{r}.$
- As in $Z = \frac{1}{N!} \left(\frac{2\pi m}{\beta h^2}\right)^{\frac{3}{2}N} Q$, only Q depends on V ,

$$p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln Z = \frac{1}{\beta} \frac{\partial}{\partial V} \ln Q$$

$$= \frac{1}{\beta} \frac{\partial}{\partial V} \left(N \ln V + \frac{N^2}{2V} \int f_{12} d\vec{r}\right) = \frac{1}{\beta} \left(\frac{N}{V} - \frac{N^2}{2V^2} \int f_{12} d\vec{r}\right)$$

$$= kT \frac{N}{V} \left[1 - \frac{N}{2V} \int f_{12} d\vec{r}\right].$$

EOS of real gas $Q = V^N \left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r}\right)$

- $\ln Q = N \ln V + \ln\left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r}\right).$
- For weak interaction,
 $\ln\left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r}\right) \simeq \frac{N^2}{2V} \int f_{12} d\vec{r}.$
 $\therefore \ln Q \simeq N \ln V + \frac{N^2}{2V} \int f_{12} d\vec{r}.$
- As in $Z = \frac{1}{N!} \left(\frac{2\pi m}{\beta h^2}\right)^{\frac{3}{2}N} Q$, only Q depends on V ,

$$p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln Z = \frac{1}{\beta} \frac{\partial}{\partial V} \ln Q$$

$$= \frac{1}{\beta} \frac{\partial}{\partial V} \left(N \ln V + \frac{N^2}{2V} \int f_{12} d\vec{r}\right) = \frac{1}{\beta} \left(\frac{N}{V} - \frac{N^2}{2V^2} \int f_{12} d\vec{r}\right)$$

$$= kT \frac{N}{V} \left[1 - \frac{N}{2V} \int f_{12} d\vec{r}\right].$$
- Define $B \equiv -\frac{N_A}{2} \int f_{12} d\vec{r}$, then $pV = NkT\left(1 + \frac{nB}{V}\right).$

EOS of real gas $Q = V^N \left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r}\right)$

- $\ln Q = N \ln V + \ln\left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r}\right).$
- For weak interaction,
 $\ln\left(1 + \frac{N^2}{2V} \int f_{12} d\vec{r}\right) \simeq \frac{N^2}{2V} \int f_{12} d\vec{r}.$
 $\therefore \ln Q \simeq N \ln V + \frac{N^2}{2V} \int f_{12} d\vec{r}.$
- As in $Z = \frac{1}{N!} \left(\frac{2\pi m}{\beta h^2}\right)^{\frac{3}{2}N} Q$, only Q depends on V ,
 $p = \frac{1}{\beta} \frac{\partial}{\partial V} \ln Z = \frac{1}{\beta} \frac{\partial}{\partial V} \ln Q$
 $= \frac{1}{\beta} \frac{\partial}{\partial V} \left(N \ln V + \frac{N^2}{2V} \int f_{12} d\vec{r}\right) = \frac{1}{\beta} \left(\frac{N}{V} - \frac{N^2}{2V^2} \int f_{12} d\vec{r}\right)$
 $= kT \frac{N}{V} \left[1 - \frac{N}{2V} \int f_{12} d\vec{r}\right].$
- Define $B \equiv -\frac{N_A}{2} \int f_{12} d\vec{r}$, then $pV = NkT\left(1 + \frac{nB}{V}\right).$
- Key: B .

EOS of real gas $B \equiv -\frac{N_A}{2} \int f_{12} d\vec{r}$

- Semi-empirical formula: $\phi(r) = \phi_0[(\frac{r_0}{r})^{12} - 2(\frac{r_0}{r})^6]$, fig.

EOS of real gas $B \equiv -\frac{N_A}{2} \int f_{12} d\vec{r}$

- Semi-empirical formula: $\phi(r) = \phi_0[(\frac{r_0}{r})^{12} - 2(\frac{r_0}{r})^6]$, fig.
- Simplify:

$$\phi(r) = \infty, \quad r < r_0,$$

$$\phi(r) = -\phi_0(\frac{r_0}{r})^6, \quad r \geq r_0.$$

EOS of real gas $B \equiv -\frac{N_A}{2} \int f_{12} d\vec{r}$

- Semi-empirical formula: $\phi(r) = \phi_0[(\frac{r_0}{r})^{12} - 2(\frac{r_0}{r})^6]$, fig.
- Simplify:

$$\phi(r) = \infty, \quad r < r_0,$$

$$\phi(r) = -\phi_0(\frac{r_0}{r})^6, \quad r \geq r_0.$$
- $B = -\frac{N_A}{2} \int f_{12} d\vec{r}$

EOS of real gas $B \equiv -\frac{N_A}{2} \int f_{12} d\vec{r}$

- Semi-empirical formula: $\phi(r) = \phi_0[(\frac{r_0}{r})^{12} - 2(\frac{r_0}{r})^6]$, fig.
- Simplify:

$$\phi(r) = \infty, \quad r < r_0,$$

$$\phi(r) = -\phi_0(\frac{r_0}{r})^6, \quad r \geq r_0.$$

- $B = -\frac{N_A}{2} \int f_{12} d\vec{r} = -\frac{N_A}{2} \int (e^{-\beta\phi(\vec{r})} - 1) d\vec{r}$

EOS of real gas $B \equiv -\frac{N_A}{2} \int f_{12} d\vec{r}$

- Semi-empirical formula: $\phi(r) = \phi_0[(\frac{r_0}{r})^{12} - 2(\frac{r_0}{r})^6]$, fig.
- Simplify:

$$\phi(r) = \infty, \quad r < r_0,$$

$$\phi(r) = -\phi_0(\frac{r_0}{r})^6, \quad r \geq r_0.$$

$$\begin{aligned} \bullet B &= -\frac{N_A}{2} \int f_{12} d\vec{r} = -\frac{N_A}{2} \int (e^{-\beta\phi(\vec{r})} - 1) d\vec{r} \\ &= -2\pi N_A \int_0^\infty (e^{-\frac{\phi(r)}{kT}} - 1) r^2 dr \end{aligned}$$

EOS of real gas $B \equiv -\frac{N_A}{2} \int f_{12} d\vec{r}$

- Semi-empirical formula: $\phi(r) = \phi_0[(\frac{r_0}{r})^{12} - 2(\frac{r_0}{r})^6]$, fig.
- Simplify:

$$\phi(r) = \infty, \quad r < r_0,$$

$$\phi(r) = -\phi_0(\frac{r_0}{r})^6, \quad r \geq r_0.$$

$$\begin{aligned} \bullet B &= -\frac{N_A}{2} \int f_{12} d\vec{r} = -\frac{N_A}{2} \int (e^{-\beta\phi(\vec{r})} - 1) d\vec{r} \\ &= -2\pi N_A \int_0^\infty (e^{-\frac{\phi(r)}{kT}} - 1) r^2 dr \\ &= -2\pi N_A [\int_0^{r_0} (-r^2) dr + \int_{r_0}^\infty (e^{\frac{\phi_0}{kT}(\frac{r_0}{r})^6} - 1) r^2 dr] \end{aligned}$$

EOS of real gas $B \equiv -\frac{N_A}{2} \int f_{12} d\vec{r}$

- Semi-empirical formula: $\phi(r) = \phi_0[(\frac{r_0}{r})^{12} - 2(\frac{r_0}{r})^6]$, fig.
- Simplify:

$$\phi(r) = \infty, \quad r < r_0,$$

$$\phi(r) = -\phi_0(\frac{r_0}{r})^6, \quad r \geq r_0.$$

$$\begin{aligned} \bullet B &= -\frac{N_A}{2} \int f_{12} d\vec{r} = -\frac{N_A}{2} \int (e^{-\beta\phi(\vec{r})} - 1) d\vec{r} \\ &= -2\pi N_A \int_0^\infty (e^{-\frac{\phi(r)}{kT}} - 1) r^2 dr \\ &= -2\pi N_A [\int_0^{r_0} (-r^2) dr + \int_{r_0}^\infty (e^{\frac{\phi_0}{kT}(\frac{r_0}{r})^6} - 1) r^2 dr] \\ &= 2\pi N_A [\frac{r_0^3}{3} - \int_{r_0}^\infty (e^{\frac{\phi_0}{kT}(\frac{r_0}{r})^6} - 1) r^2 dr]. \end{aligned}$$

EOS of real gas $B \equiv -\frac{N_A}{2} \int f_{12} d\vec{r}$

- Semi-empirical formula: $\phi(r) = \phi_0[(\frac{r_0}{r})^{12} - 2(\frac{r_0}{r})^6]$, fig.
- Simplify:

$$\phi(r) = \infty, \quad r < r_0,$$

$$\phi(r) = -\phi_0(\frac{r_0}{r})^6, \quad r \geq r_0.$$

$$\begin{aligned} \bullet B &= -\frac{N_A}{2} \int f_{12} d\vec{r} = -\frac{N_A}{2} \int (e^{-\beta\phi(\vec{r})} - 1) d\vec{r} \\ &= -2\pi N_A \int_0^\infty (e^{-\frac{\phi(r)}{kT}} - 1) r^2 dr \\ &= -2\pi N_A [\int_0^{r_0} (-r^2) dr + \int_{r_0}^\infty (e^{\frac{\phi_0}{kT}(\frac{r_0}{r})^6} - 1) r^2 dr] \\ &= 2\pi N_A [\frac{r_0^3}{3} - \int_{r_0}^\infty (e^{\frac{\phi_0}{kT}(\frac{r_0}{r})^6} - 1) r^2 dr]. \end{aligned}$$

- Simplify, at high temperature, molecule's kinetic energy can be greater than the potential energy ϕ_0 , i.e., $\phi_0 \ll kT$,

EOS of real gas $B \equiv -\frac{N_A}{2} \int f_{12} d\vec{r}$

- Semi-empirical formula: $\phi(r) = \phi_0[(\frac{r_0}{r})^{12} - 2(\frac{r_0}{r})^6]$, fig.
- Simplify:

$$\phi(r) = \infty, \quad r < r_0,$$

$$\phi(r) = -\phi_0(\frac{r_0}{r})^6, \quad r \geq r_0.$$

$$\begin{aligned} \bullet B &= -\frac{N_A}{2} \int f_{12} d\vec{r} = -\frac{N_A}{2} \int (e^{-\beta\phi(\vec{r})} - 1) d\vec{r} \\ &= -2\pi N_A \int_0^\infty (e^{-\frac{\phi(r)}{kT}} - 1) r^2 dr \\ &= -2\pi N_A [\int_0^{r_0} (-r^2) dr + \int_{r_0}^\infty (e^{\frac{\phi_0}{kT}(\frac{r_0}{r})^6} - 1) r^2 dr] \\ &= 2\pi N_A [\frac{r_0^3}{3} - \int_{r_0}^\infty (e^{\frac{\phi_0}{kT}(\frac{r_0}{r})^6} - 1) r^2 dr]. \end{aligned}$$

- Simplify, at high temperature, molecule's kinetic energy can be greater than the potential energy ϕ_0 , i.e., $\phi_0 \ll kT$, then $e^{\frac{\phi_0}{kT}(\frac{r_0}{r})^6} \simeq 1 + \frac{\phi_0}{kT}(\frac{r_0}{r})^6$.

EOS of real gas

$$B = 2\pi N_A \left[\frac{r_0^3}{3} - \int_{r_0}^{\infty} \left(e^{\frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6} - 1 \right) r^2 dr \right]$$

- $\int_{r_0}^{\infty} \left(e^{\frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6} - 1 \right) r^2 dr \simeq \int_{r_0}^{\infty} \frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6 r^2 dr$

EOS of real gas

$$B = 2\pi N_A \left[\frac{r_0^3}{3} - \int_{r_0}^{\infty} \left(e^{\frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6} - 1 \right) r^2 dr \right]$$

- $$\begin{aligned} \int_{r_0}^{\infty} \left(e^{\frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6} - 1 \right) r^2 dr &\simeq \int_{r_0}^{\infty} \frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6 r^2 dr \\ &= \frac{\phi_0 r_0^6}{kT} \int_{r_0}^{\infty} r^{-4} dr \end{aligned}$$

EOS of real gas

$$B = 2\pi N_A \left[\frac{r_0^3}{3} - \int_{r_0}^{\infty} \left(e^{\frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6} - 1 \right) r^2 dr \right]$$

- $$\begin{aligned} \int_{r_0}^{\infty} \left(e^{\frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6} - 1 \right) r^2 dr &\simeq \int_{r_0}^{\infty} \frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6 r^2 dr \\ &= \frac{\phi_0 r_0^6}{kT} \int_{r_0}^{\infty} r^{-4} dr = \frac{\phi_0 r_0^3}{3kT}. \end{aligned}$$

EOS of real gas

$$B = 2\pi N_A \left[\frac{r_0^3}{3} - \int_{r_0}^{\infty} \left(e^{\frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6} - 1 \right) r^2 dr \right]$$

- $\int_{r_0}^{\infty} \left(e^{\frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6} - 1 \right) r^2 dr \simeq \int_{r_0}^{\infty} \frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6 r^2 dr$
 $= \frac{\phi_0 r_0^6}{kT} \int_{r_0}^{\infty} r^{-4} dr = \frac{\phi_0 r_0^3}{3kT}.$
- $\therefore B \simeq 2\pi N_A \left(\frac{r_0^3}{3} - \frac{\phi_0 r_0^3}{3kT} \right).$

EOS of real gas

$$B = 2\pi N_A \left[\frac{r_0^3}{3} - \int_{r_0}^{\infty} \left(e^{\frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6} - 1 \right) r^2 dr \right]$$

- $\int_{r_0}^{\infty} \left(e^{\frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6} - 1 \right) r^2 dr \simeq \int_{r_0}^{\infty} \frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6 r^2 dr$
 $= \frac{\phi_0 r_0^6}{kT} \int_{r_0}^{\infty} r^{-4} dr = \frac{\phi_0 r_0^3}{3kT}.$
- $\therefore B \simeq 2\pi N_A \left(\frac{r_0^3}{3} - \frac{\phi_0 r_0^3}{3kT} \right).$
- Define $b = \frac{2}{3}\pi N_A r_0^3$, $a = \frac{2}{3}\pi N_A^2 \phi_0 r_0^3$, then
 $B = b - \frac{a}{N_A kT}.$

EOS of real gas

$$B = 2\pi N_A \left[\frac{r_0^3}{3} - \int_{r_0}^{\infty} \left(e^{\frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6} - 1 \right) r^2 dr \right]$$

- $\int_{r_0}^{\infty} \left(e^{\frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6} - 1 \right) r^2 dr \simeq \int_{r_0}^{\infty} \frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6 r^2 dr$
 $= \frac{\phi_0 r_0^6}{kT} \int_{r_0}^{\infty} r^{-4} dr = \frac{\phi_0 r_0^3}{3kT}.$
- $\therefore B \simeq 2\pi N_A \left(\frac{r_0^3}{3} - \frac{\phi_0 r_0^3}{3kT} \right).$
- Define $b = \frac{2}{3}\pi N_A r_0^3$, $a = \frac{2}{3}\pi N_A^2 \phi_0 r_0^3$, then
 $B = b - \frac{a}{N_A kT}.$
- $pV = NkT \left(1 + \frac{nB}{V} \right) = NkT \left(1 + \frac{nb}{V} \right) - \frac{n^2 a}{V}.$

EOS of real gas

$$B = 2\pi N_A \left[\frac{r_0^3}{3} - \int_{r_0}^{\infty} \left(e^{\frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6} - 1 \right) r^2 dr \right]$$

- $\int_{r_0}^{\infty} \left(e^{\frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6} - 1 \right) r^2 dr \simeq \int_{r_0}^{\infty} \frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6 r^2 dr$
 $= \frac{\phi_0 r_0^6}{kT} \int_{r_0}^{\infty} r^{-4} dr = \frac{\phi_0 r_0^3}{3kT}.$
- $\therefore B \simeq 2\pi N_A \left(\frac{r_0^3}{3} - \frac{\phi_0 r_0^3}{3kT} \right).$
- Define $b = \frac{2}{3}\pi N_A r_0^3$, $a = \frac{2}{3}\pi N_A^2 \phi_0 r_0^3$, then
 $B = b - \frac{a}{N_A kT}.$
- $pV = NkT \left(1 + \frac{nB}{V} \right) = NkT \left(1 + \frac{nb}{V} \right) - \frac{n^2 a}{V}.$
- As $\frac{nb}{V} = \frac{n}{V} \frac{2}{3}\pi N_A r_0^3 = \frac{\frac{2}{3}\pi r_0^3}{V/N} < 1,$

EOS of real gas

$$B = 2\pi N_A \left[\frac{r_0^3}{3} - \int_{r_0}^{\infty} \left(e^{\frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6} - 1 \right) r^2 dr \right]$$

- $\int_{r_0}^{\infty} \left(e^{\frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6} - 1 \right) r^2 dr \simeq \int_{r_0}^{\infty} \frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6 r^2 dr$
 $= \frac{\phi_0 r_0^6}{kT} \int_{r_0}^{\infty} r^{-4} dr = \frac{\phi_0 r_0^3}{3kT}.$
- $\therefore B \simeq 2\pi N_A \left(\frac{r_0^3}{3} - \frac{\phi_0 r_0^3}{3kT} \right).$
- Define $b = \frac{2}{3}\pi N_A r_0^3$, $a = \frac{2}{3}\pi N_A^2 \phi_0 r_0^3$, then
 $B = b - \frac{a}{N_A kT}.$
- $pV = NkT \left(1 + \frac{nB}{V} \right) = NkT \left(1 + \frac{nb}{V} \right) - \frac{n^2 a}{V}.$
- As $\frac{nb}{V} = \frac{n}{V} \frac{2}{3}\pi N_A r_0^3 = \frac{\frac{2}{3}\pi r_0^3}{V/N} < 1$, $1 + \frac{nb}{V} \simeq \frac{1}{1 - \frac{nb}{V}}.$

EOS of real gas

$$B = 2\pi N_A \left[\frac{r_0^3}{3} - \int_{r_0}^{\infty} \left(e^{\frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6} - 1 \right) r^2 dr \right]$$

- $\int_{r_0}^{\infty} \left(e^{\frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6} - 1 \right) r^2 dr \simeq \int_{r_0}^{\infty} \frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6 r^2 dr$
 $= \frac{\phi_0 r_0^6}{kT} \int_{r_0}^{\infty} r^{-4} dr = \frac{\phi_0 r_0^3}{3kT}.$
- $\therefore B \simeq 2\pi N_A \left(\frac{r_0^3}{3} - \frac{\phi_0 r_0^3}{3kT} \right).$
- Define $b = \frac{2}{3}\pi N_A r_0^3$, $a = \frac{2}{3}\pi N_A^2 \phi_0 r_0^3$, then
 $B = b - \frac{a}{N_A kT}.$
- $pV = NkT \left(1 + \frac{nb}{V} \right) = NkT \left(1 + \frac{nb}{V} \right) - \frac{n^2 a}{V}.$
- As $\frac{nb}{V} = \frac{n}{V} \frac{2}{3}\pi N_A r_0^3 = \frac{\frac{2}{3}\pi r_0^3}{V/N} < 1$, $1 + \frac{nb}{V} \simeq \frac{1}{1 - \frac{nb}{V}}.$
- $\Rightarrow p = \frac{NkT}{V - nb} - \frac{n^2 a}{V^2},$

EOS of real gas

$$B = 2\pi N_A \left[\frac{r_0^3}{3} - \int_{r_0}^{\infty} \left(e^{\frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6} - 1 \right) r^2 dr \right]$$

- $$\int_{r_0}^{\infty} \left(e^{\frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6} - 1 \right) r^2 dr \simeq \int_{r_0}^{\infty} \frac{\phi_0}{kT} \left(\frac{r_0}{r} \right)^6 r^2 dr$$

$$= \frac{\phi_0 r_0^6}{kT} \int_{r_0}^{\infty} r^{-4} dr = \frac{\phi_0 r_0^3}{3kT}.$$
- $$\therefore B \simeq 2\pi N_A \left(\frac{r_0^3}{3} - \frac{\phi_0 r_0^3}{3kT} \right).$$
- Define $b = \frac{2}{3}\pi N_A r_0^3$, $a = \frac{2}{3}\pi N_A^2 \phi_0 r_0^3$, then

$$B = b - \frac{a}{N_A kT}.$$
- $$pV = NkT \left(1 + \frac{nb}{V} \right) = NkT \left(1 + \frac{nb}{V} \right) - \frac{n^2 a}{V}.$$
- As $\frac{nb}{V} = \frac{n}{V} \frac{2}{3}\pi N_A r_0^3 = \frac{\frac{2}{3}\pi r_0^3}{V/N} < 1$, $1 + \frac{nb}{V} \simeq \frac{1}{1 - \frac{nb}{V}}.$
- $$\Rightarrow p = \frac{NkT}{V - nb} - \frac{n^2 a}{V^2}, \text{ or } (p + \frac{n^2 a}{V^2})(V - nb) = NkT.$$

van der Waals equation (3.5.2).

Table of contents

1 §9. Ensemble theory

- 9.1 Phase space and Liouville's theorem
- 9.2 The microcanonical ensemble
- 9.3 Thermodynamics in microcanonical ensemble theory
- 9.4 Canonical ensemble
- 9.5 Thermodynamics in canonical ensemble
- 9.6 EOS of real gas
- **9.7 Thermal capacity of solid**
- 9.9 Ising model
- 9.10 Grand canonical ensemble
- 9.11 Thermodynamics in grand canonical ensemble
- 9.12 Applications of the grand canonical ensemble

9.7 Thermal capacity of solid

9.7 Thermal capacity of solid

- Object: solid, atoms vibrate around the equilibrium position.

9.7 Thermal capacity of solid

- Object: solid, atoms vibrate around the equilibrium position.
- System: N atoms, $3N$ degrees of freedom.

9.7 Thermal capacity of solid

- Object: solid, atoms vibrate around the equilibrium position.
- System: N atoms, $3N$ degrees of freedom. Each one has a displacement ξ_i , and momentum p_{ξ_i} .

9.7 Thermal capacity of solid

- Object: solid, atoms vibrate around the equilibrium position.
- System: N atoms, $3N$ degrees of freedom. Each one has a displacement ξ_i , and momentum p_{ξ_i} .
- Total kinetic energy: $\sum_{i=1}^{3N} \frac{p_{\xi_i}^2}{2m}$,

9.7 Thermal capacity of solid

- Object: solid, atoms vibrate around the equilibrium position.
- System: N atoms, $3N$ degrees of freedom. Each one has a displacement ξ_i , and momentum p_{ξ_i} .
- Total kinetic energy: $\sum_{i=1}^{3N} \frac{p_{\xi_i}^2}{2m}$,

potential energy:

$$\phi \simeq \phi_0 + \sum_i \left(\frac{\partial \phi}{\partial \xi_i} \right)_0 \xi_i + \frac{1}{2} \sum_{i,j} \left(\frac{\partial^2 \phi}{\partial \xi_i \partial \xi_j} \right)_0 \xi_i \xi_j.$$

(Only suitable for small displacement.)

9.7 Thermal capacity of solid

- Object: solid, atoms vibrate around the equilibrium position.
- System: N atoms, $3N$ degrees of freedom. Each one has a displacement ξ_i , and momentum p_{ξ_i} .

- Total kinetic energy: $\sum_{i=1}^{3N} \frac{p_{\xi_i}^2}{2m}$,

potential energy:

$$\phi \simeq \phi_0 + \sum_i \left(\frac{\partial \phi}{\partial \xi_i} \right)_0 \xi_i + \frac{1}{2} \sum_{i,j} \left(\frac{\partial^2 \phi}{\partial \xi_i \partial \xi_j} \right)_0 \xi_i \xi_j.$$

(Only suitable for small displacement.)

- $\left(\frac{\partial \phi}{\partial \xi_i} \right)_0$ is the force exerted on the atom i at the equilibrium position,

9.7 Thermal capacity of solid

- Object: solid, atoms vibrate around the equilibrium position.
- System: N atoms, $3N$ degrees of freedom. Each one has a displacement ξ_i , and momentum p_{ξ_i} .

- Total kinetic energy: $\sum_{i=1}^{3N} \frac{p_{\xi_i}^2}{2m}$,

potential energy:

$$\phi \simeq \phi_0 + \sum_i \left(\frac{\partial \phi}{\partial \xi_i} \right)_0 \xi_i + \frac{1}{2} \sum_{i,j} \left(\frac{\partial^2 \phi}{\partial \xi_i \partial \xi_j} \right)_0 \xi_i \xi_j.$$

(Only suitable for small displacement.)

- $\left(\frac{\partial \phi}{\partial \xi_i} \right)_0$ is the force exerted on the atom i at the equilibrium position, i.e., $\left(\frac{\partial \phi}{\partial \xi_i} \right)_0 = 0$.

Total energy of solid

- Total energy $E = \sum_{i=1}^{3N} \frac{p_{\xi_i}^2}{2m} + \frac{1}{2} \sum_{i,j} a_{i,j} \xi_i \xi_j + \phi_0$.

Total energy of solid

- Total energy $E = \sum_{i=1}^{3N} \frac{p_{\xi_i}^2}{2m} + \frac{1}{2} \sum_{i,j} a_{i,j} \xi_i \xi_j + \phi_0$.
- In principle, can be transferred to:

$$E = \frac{1}{2} \sum_{i=1}^{3N} (p_i^2 + \omega_i^2 q_i^2) + \phi_0.$$

Total energy of solid

- Total energy $E = \sum_{i=1}^{3N} \frac{p_{\xi_i}^2}{2m} + \frac{1}{2} \sum_{i,j} a_{i,j} \xi_i \xi_j + \phi_0$.
- In principle, can be transferred to:

$$E = \frac{1}{2} \sum_{i=1}^{3N} (p_i^2 + \omega_i^2 q_i^2) + \phi_0.$$

It can be taken as $3N$ harmonic vibrations.

Total energy of solid

- Total energy $E = \sum_{i=1}^{3N} \frac{p_{\xi_i}^2}{2m} + \frac{1}{2} \sum_{i,j} a_{i,j} \xi_i \xi_j + \phi_0$.
- In principle, can be transferred to:

$$E = \frac{1}{2} \sum_{i=1}^{3N} (p_i^2 + \omega_i^2 q_i^2) + \phi_0.$$

It can be taken as $3N$ harmonic vibrations.

- Einstein's model: suppose all the vibrations vibrate at the same frequency ω , and the energy for each one:

$$\varepsilon_n = \hbar\omega(n + \frac{1}{2}); \text{ total } E = \sum_{i=1}^{3N} \hbar\omega(n_i + \frac{1}{2}).$$

Total energy of solid

- Total energy $E = \sum_{i=1}^{3N} \frac{p_{\xi_i}^2}{2m} + \frac{1}{2} \sum_{i,j} a_{i,j} \xi_i \xi_j + \phi_0$.

- In principle, can be transferred to:

$$E = \frac{1}{2} \sum_{i=1}^{3N} (p_i^2 + \omega_i^2 q_i^2) + \phi_0.$$

It can be taken as $3N$ harmonic vibrations.

- Einstein's model: suppose all the vibrations vibrate at the same frequency ω , and the energy for each one:

$$\varepsilon_n = \hbar\omega(n + \frac{1}{2}); \text{ total } E = \sum_{i=1}^{3N} \hbar\omega(n_i + \frac{1}{2}).$$

- More generally: ω is not the same (not independent).

Total energy of solid

- Total energy $E = \sum_{i=1}^{3N} \frac{p_{\xi_i}^2}{2m} + \frac{1}{2} \sum_{i,j} a_{i,j} \xi_i \xi_j + \phi_0$.

- In principle, can be transferred to:

$$E = \frac{1}{2} \sum_{i=1}^{3N} (p_i^2 + \omega_i^2 q_i^2) + \phi_0.$$

It can be taken as $3N$ harmonic vibrations.

- Einstein's model: suppose all the vibrations vibrate at the same frequency ω , and the energy for each one:

$$\varepsilon_n = \hbar\omega(n + \frac{1}{2}); \text{ total } E = \sum_{i=1}^{3N} \hbar\omega(n_i + \frac{1}{2}).$$

- More generally: ω is not the same (not independent).

$$E = \phi_0 + \sum_{i=1}^{3N} \hbar\omega_i(n_i + \frac{1}{2}).$$

Partition function

$$E = \phi_0 + \sum_{i=1}^{3N} \hbar \omega_i (n_i + \frac{1}{2})$$

- Partition function: $Z = \sum_s e^{-\beta E_s}$

Partition function

$$E = \phi_0 + \sum_{i=1}^{3N} \hbar \omega_i (n_i + \frac{1}{2})$$

- Partition function: $Z = \sum_s e^{-\beta E_s}$
$$= e^{-\beta \phi_0} \sum_{\{n_i\}} e^{-\beta \sum_i^s \hbar \omega_i (n_i + \frac{1}{2})}$$

Partition function

$$E = \phi_0 + \sum_{i=1}^{3N} \hbar \omega_i (n_i + \frac{1}{2})$$

- Partition function: $Z = \sum_s e^{-\beta E_s}$
$$= e^{-\beta \phi_0} \sum_{\{n_i\}} e^{-\beta \sum_i \hbar \omega_i (n_i + \frac{1}{2})} = e^{-\beta \phi_0} \sum_{\{n_i\}} \prod_i e^{-\beta \hbar \omega_i (n_i + \frac{1}{2})}$$

Partition function

$$E = \phi_0 + \sum_{i=1}^{3N} \hbar \omega_i (n_i + \frac{1}{2})$$

- Partition function: $Z = \sum_s e^{-\beta E_s}$
$$= e^{-\beta \phi_0} \sum_{\{n_i\}} e^{-\beta \sum_i \hbar \omega_i (n_i + \frac{1}{2})} = e^{-\beta \phi_0} \sum_{\{n_i\}} \prod_i e^{-\beta \hbar \omega_i (n_i + \frac{1}{2})}$$
$$= e^{-\beta \phi_0} \prod_i \sum_{n_i=0}^{\infty} e^{-\beta \hbar \omega_i (n_i + \frac{1}{2})}$$

Partition function

$$E = \phi_0 + \sum_{i=1}^{3N} \hbar\omega_i(n_i + \frac{1}{2})$$

- Partition function: $Z = \sum_s e^{-\beta E_s}$

$$= e^{-\beta\phi_0} \sum_{\{n_i\}} e^{-\beta \sum_i \hbar\omega_i(n_i + \frac{1}{2})} = e^{-\beta\phi_0} \sum_{\{n_i\}} \prod_i e^{-\beta \hbar\omega_i(n_i + \frac{1}{2})}$$

$$= e^{-\beta\phi_0} \prod_i \sum_{n_i=0}^{\infty} e^{-\beta \hbar\omega_i(n_i + \frac{1}{2})} = e^{-\beta\phi_0} \prod_i \frac{e^{-\frac{\beta \hbar\omega_i}{2}}}{1 - e^{-\beta \hbar\omega_i}}.$$

Partition function

$$E = \phi_0 + \sum_{i=1}^{3N} \hbar\omega_i(n_i + \frac{1}{2})$$

- Partition function: $Z = \sum_s e^{-\beta E_s}$
$$= e^{-\beta\phi_0} \sum_{\{n_i\}} e^{-\beta \sum_i \hbar\omega_i(n_i + \frac{1}{2})} = e^{-\beta\phi_0} \sum_{\{n_i\}} \prod_i e^{-\beta \hbar\omega_i(n_i + \frac{1}{2})}$$
$$= e^{-\beta\phi_0} \prod_i \sum_{n_i=0}^{\infty} e^{-\beta \hbar\omega_i(n_i + \frac{1}{2})} = e^{-\beta\phi_0} \prod_i \frac{e^{-\frac{\beta \hbar\omega_i}{2}}}{1 - e^{-\beta \hbar\omega_i}}.$$
- Internal energy: $U = -\frac{\partial \ln Z}{\partial \beta}$

Partition function

$$E = \phi_0 + \sum_{i=1}^{3N} \hbar\omega_i(n_i + \frac{1}{2})$$

- Partition function: $Z = \sum_s e^{-\beta E_s}$

$$= e^{-\beta\phi_0} \sum_{\{n_i\}} e^{-\beta \sum_i \hbar\omega_i(n_i + \frac{1}{2})} = e^{-\beta\phi_0} \sum_{\{n_i\}} \prod_i e^{-\beta \hbar\omega_i(n_i + \frac{1}{2})}$$

$$= e^{-\beta\phi_0} \prod_i \sum_{n_i=0}^{\infty} e^{-\beta \hbar\omega_i(n_i + \frac{1}{2})} = e^{-\beta\phi_0} \prod_i \frac{e^{-\frac{\beta \hbar\omega_i}{2}}}{1 - e^{-\beta \hbar\omega_i}}.$$
- Internal energy: $U = -\frac{\partial \ln Z}{\partial \beta}$

$$= \phi_0 + \sum_{i=1}^{3N} \frac{1}{2} \hbar\omega_i + \sum_{i=1}^{3N} \frac{\hbar\omega_i e^{-\beta \hbar\omega_i}}{1 - e^{-\beta \hbar\omega_i}}$$

Partition function

$$E = \phi_0 + \sum_{i=1}^{3N} \hbar\omega_i(n_i + \frac{1}{2})$$

- Partition function: $Z = \sum_s e^{-\beta E_s}$

$$= e^{-\beta\phi_0} \sum_{\{n_i\}} e^{-\beta \sum_i \hbar\omega_i(n_i + \frac{1}{2})} = e^{-\beta\phi_0} \sum_{\{n_i\}} \prod_i e^{-\beta \hbar\omega_i(n_i + \frac{1}{2})}$$

$$= e^{-\beta\phi_0} \prod_i \sum_{n_i=0}^{\infty} e^{-\beta \hbar\omega_i(n_i + \frac{1}{2})} = e^{-\beta\phi_0} \prod_i \frac{e^{-\frac{\beta \hbar\omega_i}{2}}}{1 - e^{-\beta \hbar\omega_i}}.$$
- Internal energy: $U = -\frac{\partial \ln Z}{\partial \beta}$

$$= \phi_0 + \sum_{i=1}^{3N} \frac{1}{2} \hbar\omega_i + \sum_{i=1}^{3N} \frac{\hbar\omega_i e^{-\beta \hbar\omega_i}}{1 - e^{-\beta \hbar\omega_i}} \equiv U_0 + \sum_{i=1}^{3N} \frac{\hbar\omega_i}{e^{\beta \hbar\omega_i} - 1}.$$

Partition function

$$E = \phi_0 + \sum_{i=1}^{3N} \hbar\omega_i(n_i + \frac{1}{2})$$

- Partition function: $Z = \sum_s e^{-\beta E_s}$

$$= e^{-\beta\phi_0} \sum_{\{n_i\}} e^{-\beta \sum_i \hbar\omega_i(n_i + \frac{1}{2})} = e^{-\beta\phi_0} \sum_{\{n_i\}} \prod_i e^{-\beta \hbar\omega_i(n_i + \frac{1}{2})}$$

$$= e^{-\beta\phi_0} \prod_i \sum_{n_i=0}^{\infty} e^{-\beta \hbar\omega_i(n_i + \frac{1}{2})} = e^{-\beta\phi_0} \prod_i \frac{e^{-\frac{\beta \hbar\omega_i}{2}}}{1 - e^{-\beta \hbar\omega_i}}.$$
- Internal energy: $U = -\frac{\partial \ln Z}{\partial \beta}$

$$= \phi_0 + \sum_{i=1}^{3N} \frac{1}{2} \hbar\omega_i + \sum_{i=1}^{3N} \frac{\hbar\omega_i e^{-\beta \hbar\omega_i}}{1 - e^{-\beta \hbar\omega_i}} \equiv U_0 + \sum_{i=1}^{3N} \frac{\hbar\omega_i}{e^{\beta \hbar\omega_i} - 1}.$$
- Simplest case: $\omega_i = \omega$,

Partition function

$$E = \phi_0 + \sum_{i=1}^{3N} \hbar\omega_i(n_i + \frac{1}{2})$$

- Partition function: $Z = \sum_s e^{-\beta E_s}$

$$= e^{-\beta\phi_0} \sum_{\{n_i\}} e^{-\beta \sum_i \hbar\omega_i(n_i + \frac{1}{2})} = e^{-\beta\phi_0} \sum_{\{n_i\}} \prod_i e^{-\beta \hbar\omega_i(n_i + \frac{1}{2})}$$

$$= e^{-\beta\phi_0} \prod_i \sum_{n_i=0}^{\infty} e^{-\beta \hbar\omega_i(n_i + \frac{1}{2})} = e^{-\beta\phi_0} \prod_i \frac{e^{-\frac{\beta \hbar\omega_i}{2}}}{1 - e^{-\beta \hbar\omega_i}}.$$
- Internal energy: $U = -\frac{\partial \ln Z}{\partial \beta}$

$$= \phi_0 + \sum_{i=1}^{3N} \frac{1}{2} \hbar\omega_i + \sum_{i=1}^{3N} \frac{\hbar\omega_i e^{-\beta \hbar\omega_i}}{1 - e^{-\beta \hbar\omega_i}} \equiv U_0 + \sum_{i=1}^{3N} \frac{\hbar\omega_i}{e^{\beta \hbar\omega_i} - 1}.$$
- Simplest case: $\omega_i = \omega$, $U = \phi_0 + \frac{3}{2}N\hbar\omega + \frac{3N\hbar\omega}{e^{\beta \hbar\omega} - 1},$

Partition function

$$E = \phi_0 + \sum_{i=1}^{3N} \hbar\omega_i(n_i + \frac{1}{2})$$

- Partition function: $Z = \sum_s e^{-\beta E_s}$

$$= e^{-\beta\phi_0} \sum_{\{n_i\}} e^{-\beta \sum_i \hbar\omega_i(n_i + \frac{1}{2})} = e^{-\beta\phi_0} \sum_{\{n_i\}} \prod_i e^{-\beta \hbar\omega_i(n_i + \frac{1}{2})}$$

$$= e^{-\beta\phi_0} \prod_i \sum_{n_i=0}^{\infty} e^{-\beta \hbar\omega_i(n_i + \frac{1}{2})} = e^{-\beta\phi_0} \prod_i \frac{e^{-\frac{\beta \hbar\omega_i}{2}}}{1 - e^{-\beta \hbar\omega_i}}.$$
- Internal energy: $U = -\frac{\partial \ln Z}{\partial \beta}$

$$= \phi_0 + \sum_{i=1}^{3N} \frac{1}{2} \hbar\omega_i + \sum_{i=1}^{3N} \frac{\hbar\omega_i e^{-\beta \hbar\omega_i}}{1 - e^{-\beta \hbar\omega_i}} \equiv U_0 + \sum_{i=1}^{3N} \frac{\hbar\omega_i}{e^{\beta \hbar\omega_i} - 1}.$$
- Simplest case: $\omega_i = \omega$, $U = \phi_0 + \frac{3}{2}N\hbar\omega + \frac{3N\hbar\omega}{e^{\beta \hbar\omega} - 1}$,
back to Einstein's model (7.7.3).

Debye's model

$$U = U_0 + \sum_{i=1}^{3N} \frac{\hbar\omega_i}{e^{\beta\hbar\omega_i} - 1}$$

Debye's model

$$U = U_0 + \sum_{i=1}^{3N} \frac{\hbar\omega_i}{e^{\beta\hbar\omega_i} - 1}$$

- Take the solid as continuous elastic medium.

Debye's model

$$U = U_0 + \sum_{i=1}^{3N} \frac{\hbar\omega_i}{e^{\beta\hbar\omega_i} - 1}$$

- Take the solid as continuous elastic medium. $3N$ normal vibrations can produce $3N$ different fundamental waves.

Debye's model

$$U = U_0 + \sum_{i=1}^{3N} \frac{\hbar\omega_i}{e^{\beta\hbar\omega_i} - 1}$$

- Take the solid as continuous elastic medium. $3N$ normal vibrations can produce $3N$ different fundamental waves.
- In elastic medium, there exists longitudinal (l) wave and transversal (t) wave (two directions),

Debye's model

$$U = U_0 + \sum_{i=1}^{3N} \frac{\hbar\omega_i}{e^{\beta\hbar\omega_i} - 1}$$

- Take the solid as continuous elastic medium. $3N$ normal vibrations can produce $3N$ different fundamental waves.
- In elastic medium, there exists longitudinal (l) wave and transversal (t) wave (two directions), with dispersion relations: $\omega = c_l k$, $\omega = c_t k$, where c is the speed of sound.

Debye's model

$$U = U_0 + \sum_{i=1}^{3N} \frac{\hbar\omega_i}{e^{\beta\hbar\omega_i} - 1}$$

- Take the solid as continuous elastic medium. $3N$ normal vibrations can produce $3N$ different fundamental waves.
- In elastic medium, there exists longitudinal (l) wave and transversal (t) wave (two directions), with dispersion relations: $\omega = c_l k$, $\omega = c_t k$, where c is the speed of sound. For low frequencies, ω_i can be taken as continuous.

Debye's model

$$U = U_0 + \sum_{i=1}^{3N} \frac{\hbar\omega_i}{e^{\beta\hbar\omega_i} - 1}$$

- Take the solid as continuous elastic medium. $3N$ normal vibrations can produce $3N$ different fundamental waves.
- In elastic medium, there exists longitudinal (l) wave and transversal (t) wave (two directions), with dispersion relations: $\omega = c_l k$, $\omega = c_t k$, where c is the speed of sound. For low frequencies, ω_i can be taken as continuous.
- The number of longitudinal normal vibrations (number of states):

Debye's model

$$U = U_0 + \sum_{i=1}^{3N} \frac{\hbar\omega_i}{e^{\beta\hbar\omega_i} - 1}$$

- Take the solid as continuous elastic medium. $3N$ normal vibrations can produce $3N$ different fundamental waves.
- In elastic medium, there exists longitudinal (l) wave and transversal (t) wave (two directions), with dispersion relations: $\omega = c_l k$, $\omega = c_t k$, where c is the speed of sound. For low frequencies, ω_i can be taken as continuous.
- The number of longitudinal normal vibrations (number of states): $D(p)dp = \frac{V}{h^3} 4\pi p^2 dp$

Debye's model

$$U = U_0 + \sum_{i=1}^{3N} \frac{\hbar\omega_i}{e^{\beta\hbar\omega_i} - 1}$$

- Take the solid as continuous elastic medium. $3N$ normal vibrations can produce $3N$ different fundamental waves.
- In elastic medium, there exists longitudinal (l) wave and transversal (t) wave (two directions), with dispersion relations: $\omega = c_l k$, $\omega = c_t k$, where c is the speed of sound. For low frequencies, ω_i can be taken as continuous.
- The number of longitudinal normal vibrations (number of states): $D(p)dp = \frac{V}{h^3} 4\pi p^2 dp = \frac{V}{h^3} 4\pi \hbar^3 k^2 dk$

Debye's model

$$U = U_0 + \sum_{i=1}^{3N} \frac{\hbar\omega_i}{e^{\beta\hbar\omega_i} - 1}$$

- Take the solid as continuous elastic medium. $3N$ normal vibrations can produce $3N$ different fundamental waves.
- In elastic medium, there exists longitudinal (l) wave and transversal (t) wave (two directions), with dispersion relations: $\omega = c_l k$, $\omega = c_t k$, where c is the speed of sound. For low frequencies, ω_i can be taken as continuous.
- The number of longitudinal normal vibrations (number of states): $D(p)dp = \frac{V}{h^3} 4\pi p^2 dp = \frac{V}{h^3} 4\pi \hbar^3 k^2 dk$
 $= \frac{V}{2\pi^2} k^2 dk$

Debye's model $U = U_0 + \sum_{i=1}^{3N} \frac{\hbar\omega_i}{e^{\beta\hbar\omega_i} - 1}$

- Take the solid as continuous elastic medium. $3N$ normal vibrations can produce $3N$ different fundamental waves.
- In elastic medium, there exists longitudinal (l) wave and transversal (t) wave (two directions), with dispersion relations: $\omega = c_l k$, $\omega = c_t k$, where c is the speed of sound. For low frequencies, ω_i can be taken as continuous.
- The number of longitudinal normal vibrations (number of states): $D(p)dp = \frac{V}{h^3} 4\pi p^2 dp = \frac{V}{h^3} 4\pi \hbar^3 k^2 dk$
 $= \frac{V}{2\pi^2} k^2 dk = \frac{V}{2\pi^2} \frac{1}{c_l^3} \omega^2 d\omega$

Debye's model

$$U = U_0 + \sum_{i=1}^{3N} \frac{\hbar\omega_i}{e^{\beta\hbar\omega_i} - 1}$$

- Take the solid as continuous elastic medium. $3N$ normal vibrations can produce $3N$ different fundamental waves.
- In elastic medium, there exists longitudinal (l) wave and transversal (t) wave (two directions), with dispersion relations: $\omega = c_l k$, $\omega = c_t k$, where c is the speed of sound. For low frequencies, ω_i can be taken as continuous.
- The number of longitudinal normal vibrations (number of states): $D(p)dp = \frac{V}{h^3} 4\pi p^2 dp = \frac{V}{h^3} 4\pi \hbar^3 k^2 dk$
 $= \frac{V}{2\pi^2} k^2 dk = \frac{V}{2\pi^2} \frac{1}{c_l^3} \omega^2 d\omega \equiv D_l(\omega) d\omega$.

Debye's model

$$U = U_0 + \sum_{i=1}^{3N} \frac{\hbar\omega_i}{e^{\beta\hbar\omega_i} - 1}$$

- Considering the two directions of transversal vibrations: $D_t(\omega)d\omega = 2\frac{V}{2\pi^2}\frac{1}{c_t^3}\omega^2d\omega$.

Debye's model $U = U_0 + \sum_{i=1}^{3N} \frac{\hbar\omega_i}{e^{\beta\hbar\omega_i} - 1}$

- Considering the two directions of transversal vibrations: $D_t(\omega)d\omega = 2\frac{V}{2\pi^2} \frac{1}{c_t^3} \omega^2 d\omega$.
- $D(\omega)d\omega = \frac{V}{2\pi^2} \left(\frac{1}{c_l^3} + \frac{2}{c_t^3} \right) \omega^2 d\omega$

Debye's model

$$U = U_0 + \sum_{i=1}^{3N} \frac{\hbar\omega_i}{e^{\beta\hbar\omega_i} - 1}$$

- Considering the two directions of transversal vibrations: $D_t(\omega)d\omega = 2\frac{V}{2\pi^2} \frac{1}{c_t^3} \omega^2 d\omega$.
- $D(\omega)d\omega = \frac{V}{2\pi^2} \left(\frac{1}{c_l^3} + \frac{2}{c_t^3} \right) \omega^2 d\omega \equiv B\omega^2 d\omega$, where $B = \frac{V}{2\pi^2} \left(\frac{1}{c_l^3} + \frac{2}{c_t^3} \right)$.

Debye's model

$$U = U_0 + \sum_{i=1}^{3N} \frac{\hbar\omega_i}{e^{\beta\hbar\omega_i} - 1}$$

- Considering the two directions of transversal vibrations: $D_t(\omega)d\omega = 2\frac{V}{2\pi^2}\frac{1}{c_t^3}\omega^2d\omega$.
- $D(\omega)d\omega = \frac{V}{2\pi^2}\left(\frac{1}{c_l^3} + \frac{2}{c_t^3}\right)\omega^2d\omega \equiv B\omega^2d\omega$, where $B = \frac{V}{2\pi^2}\left(\frac{1}{c_l^3} + \frac{2}{c_t^3}\right)$.
- Remember the total number of states: $3N$.

Debye's model

$$U = U_0 + \sum_{i=1}^{3N} \frac{\hbar\omega_i}{e^{\beta\hbar\omega_i} - 1}$$

- Considering the two directions of transversal vibrations: $D_t(\omega)d\omega = 2\frac{V}{2\pi^2}\frac{1}{c_t^3}\omega^2d\omega$.
- $D(\omega)d\omega = \frac{V}{2\pi^2}\left(\frac{1}{c_l^3} + \frac{2}{c_t^3}\right)\omega^2d\omega \equiv B\omega^2d\omega$, where $B = \frac{V}{2\pi^2}\left(\frac{1}{c_l^3} + \frac{2}{c_t^3}\right)$.
- Remember the total number of states: $3N$.

$$\int_0^{\omega_D} B\omega^2d\omega = 3N$$

Debye's model

$$U = U_0 + \sum_{i=1}^{3N} \frac{\hbar\omega_i}{e^{\beta\hbar\omega_i} - 1}$$

- Considering the two directions of transversal vibrations: $D_t(\omega)d\omega = 2\frac{V}{2\pi^2}\frac{1}{c_t^3}\omega^2d\omega$.
- $D(\omega)d\omega = \frac{V}{2\pi^2}\left(\frac{1}{c_l^3} + \frac{2}{c_t^3}\right)\omega^2d\omega \equiv B\omega^2d\omega$, where $B = \frac{V}{2\pi^2}\left(\frac{1}{c_l^3} + \frac{2}{c_t^3}\right)$.
- Remember the total number of states: $3N$.
 $\int_0^{\omega_D} B\omega^2d\omega = 3N \Rightarrow \omega_D^3 = \frac{9N}{B}$ is the maximum frequency, called Debye's frequency.

Debye's model

$$U = U_0 + \sum_{i=1}^{3N} \frac{\hbar\omega_i}{e^{\beta\hbar\omega_i} - 1}$$

- Considering the two directions of transversal vibrations: $D_t(\omega)d\omega = 2\frac{V}{2\pi^2} \frac{1}{c_t^3} \omega^2 d\omega$.
- $D(\omega)d\omega = \frac{V}{2\pi^2} \left(\frac{1}{c_l^3} + \frac{2}{c_t^3} \right) \omega^2 d\omega \equiv B\omega^2 d\omega$, where $B = \frac{V}{2\pi^2} \left(\frac{1}{c_l^3} + \frac{2}{c_t^3} \right)$.
- Remember the total number of states: $3N$.
 $\int_0^{\omega_D} B\omega^2 d\omega = 3N \Rightarrow \omega_D^3 = \frac{9N}{B}$ is the maximum frequency, called Debye's frequency.
- Internal energy ($\sum \rightarrow \int$):

$$U = U_0 + \int_0^{\omega_D} \frac{\hbar\omega}{e^{\beta\hbar\omega} - 1} D(\omega) d\omega$$

Debye's model

$$U = U_0 + \sum_{i=1}^{3N} \frac{\hbar\omega_i}{e^{\beta\hbar\omega_i} - 1}$$

- Considering the two directions of transversal vibrations: $D_t(\omega)d\omega = 2\frac{V}{2\pi^2} \frac{1}{c_t^3} \omega^2 d\omega$.
- $D(\omega)d\omega = \frac{V}{2\pi^2} \left(\frac{1}{c_l^3} + \frac{2}{c_t^3} \right) \omega^2 d\omega \equiv B\omega^2 d\omega$, where $B = \frac{V}{2\pi^2} \left(\frac{1}{c_l^3} + \frac{2}{c_t^3} \right)$.
- Remember the total number of states: $3N$.
 $\int_0^{\omega_D} B\omega^2 d\omega = 3N \Rightarrow \omega_D^3 = \frac{9N}{B}$ is the maximum frequency, called Debye's frequency.
- Internal energy ($\sum \rightarrow \int$):

$$U = U_0 + \int_0^{\omega_D} \frac{\hbar\omega}{e^{\beta\hbar\omega} - 1} D(\omega) d\omega = U_0 + B \int_0^{\omega_D} \frac{\hbar\omega^3}{e^{\beta\hbar\omega} - 1} d\omega.$$

Debye's model

$$U = U_0 + \sum_{i=1}^{3N} \frac{\hbar\omega_i}{e^{\beta\hbar\omega_i} - 1}$$

- Considering the two directions of transversal vibrations: $D_t(\omega)d\omega = 2\frac{V}{2\pi^2} \frac{1}{c_t^3} \omega^2 d\omega$.
- $D(\omega)d\omega = \frac{V}{2\pi^2} \left(\frac{1}{c_l^3} + \frac{2}{c_t^3} \right) \omega^2 d\omega \equiv B\omega^2 d\omega$, where $B = \frac{V}{2\pi^2} \left(\frac{1}{c_l^3} + \frac{2}{c_t^3} \right)$.
- Remember the total number of states: $3N$.
 $\int_0^{\omega_D} B\omega^2 d\omega = 3N \Rightarrow \omega_D^3 = \frac{9N}{B}$ is the maximum frequency, called Debye's frequency.
- Internal energy ($\sum \rightarrow \int$):

$$U = U_0 + \int_0^{\omega_D} \frac{\hbar\omega}{e^{\beta\hbar\omega} - 1} D(\omega) d\omega = U_0 + B \int_0^{\omega_D} \frac{\hbar\omega^3}{e^{\beta\hbar\omega} - 1} d\omega.$$
- Set $y = \frac{\hbar\omega}{kT}$,

Debye's model

$$U = U_0 + \sum_{i=1}^{3N} \frac{\hbar\omega_i}{e^{\beta\hbar\omega_i} - 1}$$

- Considering the two directions of transversal vibrations: $D_t(\omega)d\omega = 2\frac{V}{2\pi^2} \frac{1}{c_t^3} \omega^2 d\omega$.
- $D(\omega)d\omega = \frac{V}{2\pi^2} \left(\frac{1}{c_l^3} + \frac{2}{c_t^3} \right) \omega^2 d\omega \equiv B\omega^2 d\omega$, where $B = \frac{V}{2\pi^2} \left(\frac{1}{c_l^3} + \frac{2}{c_t^3} \right)$.
- Remember the total number of states: $3N$.
 $\int_0^{\omega_D} B\omega^2 d\omega = 3N \Rightarrow \omega_D^3 = \frac{9N}{B}$ is the maximum frequency, called Debye's frequency.
- Internal energy ($\sum \rightarrow \int$):

$$U = U_0 + \int_0^{\omega_D} \frac{\hbar\omega}{e^{\beta\hbar\omega} - 1} D(\omega) d\omega = U_0 + B \int_0^{\omega_D} \frac{\hbar\omega^3}{e^{\beta\hbar\omega} - 1} d\omega.$$
- Set $y = \frac{\hbar\omega}{kT}$, then $y_D = \frac{\hbar\omega_D}{kT} \equiv \frac{\theta_D}{T}$,

Debye's model

$$U = U_0 + \sum_{i=1}^{3N} \frac{\hbar\omega_i}{e^{\beta\hbar\omega_i} - 1}$$

- Considering the two directions of transversal vibrations: $D_t(\omega)d\omega = 2\frac{V}{2\pi^2} \frac{1}{c_t^3} \omega^2 d\omega$.
- $D(\omega)d\omega = \frac{V}{2\pi^2} \left(\frac{1}{c_l^3} + \frac{2}{c_t^3} \right) \omega^2 d\omega \equiv B\omega^2 d\omega$, where $B = \frac{V}{2\pi^2} \left(\frac{1}{c_l^3} + \frac{2}{c_t^3} \right)$.
- Remember the total number of states: $3N$.
 $\int_0^{\omega_D} B\omega^2 d\omega = 3N \Rightarrow \omega_D^3 = \frac{9N}{B}$ is the maximum frequency, called Debye's frequency.
- Internal energy ($\sum \rightarrow \int$):

$$U = U_0 + \int_0^{\omega_D} \frac{\hbar\omega}{e^{\beta\hbar\omega} - 1} D(\omega) d\omega = U_0 + B \int_0^{\omega_D} \frac{\hbar\omega^3}{e^{\beta\hbar\omega} - 1} d\omega.$$
- Set $y = \frac{\hbar\omega}{kT}$, then $y_D = \frac{\hbar\omega_D}{kT} \equiv \frac{\theta_D}{T}$, where $\theta_D = \frac{\hbar\omega_D}{k}$ is called Debye's temperature.

$$U = U_0 + B \int_0^{\omega_D} \frac{\hbar \omega^3}{e^{\beta \hbar \omega} - 1} d\omega, \quad \omega_D^3 = \frac{9N}{B}, \quad y_D = \frac{\hbar \omega_D}{kT}$$

$$\bullet U = U_0 + B \int_0^{\frac{\hbar \omega_D}{kT}} \frac{\hbar \cdot \left(\frac{\hbar \omega}{kT}\right)^3 \cdot \left(\frac{kT}{\hbar}\right)^3}{e^{\frac{\hbar \omega}{kT}} - 1} \frac{kT}{\hbar} d\frac{\hbar \omega}{kT}$$

$$U = U_0 + B \int_0^{\omega_D} \frac{\hbar \omega^3}{e^{\beta \hbar \omega} - 1} d\omega, \quad \omega_D^3 = \frac{9N}{B}, \quad y_D = \frac{\hbar \omega_D}{kT}$$

$$\begin{aligned} \bullet U &= U_0 + B \int_0^{\frac{\hbar \omega_D}{kT}} \frac{\hbar \cdot \left(\frac{\hbar \omega}{kT}\right)^3 \cdot \left(\frac{kT}{\hbar}\right)^3}{e^{\frac{\hbar \omega}{kT}} - 1} \frac{kT}{\hbar} d\frac{\hbar \omega}{kT} \\ &= U_0 + B \hbar \left(\frac{kT}{\hbar}\right)^4 \int_0^{y_D} \frac{y^3}{e^y - 1} dy \end{aligned}$$

$$U = U_0 + B \int_0^{\omega_D} \frac{\hbar \omega^3}{e^{\beta \hbar \omega} - 1} d\omega, \quad \omega_D^3 = \frac{9N}{B}, \quad y_D = \frac{\hbar \omega_D}{kT}$$

$$\begin{aligned} \bullet U &= U_0 + B \int_0^{\frac{\hbar \omega_D}{kT}} \frac{\hbar \cdot \left(\frac{\hbar \omega}{kT}\right)^3 \cdot \left(\frac{kT}{\hbar}\right)^3}{e^{\frac{\hbar \omega}{kT}} - 1} \frac{kT}{\hbar} d\frac{\hbar \omega}{kT} \\ &= U_0 + B \hbar \left(\frac{kT}{\hbar}\right)^4 \int_0^{y_D} \frac{y^3}{e^y - 1} dy \\ &= U_0 + \frac{9N}{\omega_D^3} \frac{(kT)^4}{\hbar^3} \int_0^{y_D} \frac{y^3}{e^y - 1} dy \end{aligned}$$

$$U = U_0 + B \int_0^{\omega_D} \frac{\hbar\omega^3}{e^{\beta\hbar\omega}-1} d\omega, \quad \omega_D^3 = \frac{9N}{B}, \quad y_D = \frac{\hbar\omega_D}{kT}$$

$$\begin{aligned} \bullet U &= U_0 + B \int_0^{\frac{\hbar\omega_D}{kT}} \frac{\hbar \cdot (\frac{\hbar\omega}{kT})^3 \cdot (\frac{kT}{\hbar})^3}{e^{\frac{\hbar\omega}{kT}} - 1} \frac{kT}{\hbar} d\frac{\hbar\omega}{kT} \\ &= U_0 + B\hbar \left(\frac{kT}{\hbar}\right)^4 \int_0^{y_D} \frac{y^3}{e^y - 1} dy \\ &= U_0 + \frac{9N}{\omega_D^3} \frac{(kT)^4}{\hbar^3} \int_0^{y_D} \frac{y^3}{e^y - 1} dy = U_0 + \frac{9N}{y_D^3} kT \int_0^{y_D} \frac{y^3}{e^y - 1} dy \end{aligned}$$

$$U = U_0 + B \int_0^{\omega_D} \frac{\hbar\omega^3}{e^{\beta\hbar\omega} - 1} d\omega, \quad \omega_D^3 = \frac{9N}{B}, \quad y_D = \frac{\hbar\omega_D}{kT}$$

$$\begin{aligned}
 \bullet \quad U &= U_0 + B \int_0^{\frac{\hbar\omega_D}{kT}} \frac{\hbar \cdot \left(\frac{\hbar\omega}{kT}\right)^3 \cdot \left(\frac{kT}{\hbar}\right)^3}{e^{\frac{\hbar\omega}{kT}} - 1} \frac{kT}{\hbar} d\frac{\hbar\omega}{kT} \\
 &= U_0 + B\hbar \left(\frac{kT}{\hbar}\right)^4 \int_0^{y_D} \frac{y^3}{e^y - 1} dy \\
 &= U_0 + \frac{9N}{\omega_D^3} \frac{(kT)^4}{\hbar^3} \int_0^{y_D} \frac{y^3}{e^y - 1} dy = U_0 + \frac{9N}{y_D^3} kT \int_0^{y_D} \frac{y^3}{e^y - 1} dy \\
 &\equiv U_0 + 3NkT \mathcal{D}(y_D), \text{ where } \mathcal{D}(y_D) = \frac{3}{y_D^3} \int_0^{y_D} \frac{y^3}{e^y - 1} dy.
 \end{aligned}$$

$$U = U_0 + B \int_0^{\omega_D} \frac{\hbar\omega^3}{e^{\beta\hbar\omega} - 1} d\omega, \quad \omega_D^3 = \frac{9N}{B}, \quad y_D = \frac{\hbar\omega_D}{kT}$$

- $$\begin{aligned}
 U &= U_0 + B \int_0^{\frac{\hbar\omega_D}{kT}} \frac{\hbar \cdot (\frac{\hbar\omega}{kT})^3 \cdot (\frac{kT}{\hbar})^3}{e^{\frac{\hbar\omega}{kT}} - 1} \frac{kT}{\hbar} d\frac{\hbar\omega}{kT} \\
 &= U_0 + B\hbar \left(\frac{kT}{\hbar}\right)^4 \int_0^{y_D} \frac{y^3}{e^y - 1} dy \\
 &= U_0 + \frac{9N}{\omega_D^3} \frac{(kT)^4}{\hbar^3} \int_0^{y_D} \frac{y^3}{e^y - 1} dy = U_0 + \frac{9N}{y_D^3} kT \int_0^{y_D} \frac{y^3}{e^y - 1} dy \\
 &\equiv U_0 + 3NkT \mathcal{D}(y_D), \text{ where } \mathcal{D}(y_D) = \frac{3}{y_D^3} \int_0^{y_D} \frac{y^3}{e^y - 1} dy.
 \end{aligned}$$
- High temperature, $y_D \ll 1$,

$$U = U_0 + B \int_0^{\omega_D} \frac{\hbar\omega^3}{e^{\beta\hbar\omega} - 1} d\omega, \quad \omega_D^3 = \frac{9N}{B}, \quad y_D = \frac{\hbar\omega_D}{kT}$$

- $$\begin{aligned}
 U &= U_0 + B \int_0^{\frac{\hbar\omega_D}{kT}} \frac{\hbar \cdot (\frac{\hbar\omega}{kT})^3 \cdot (\frac{kT}{\hbar})^3}{e^{\frac{\hbar\omega}{kT}} - 1} \frac{kT}{\hbar} d\frac{\hbar\omega}{kT} \\
 &= U_0 + B\hbar \left(\frac{kT}{\hbar}\right)^4 \int_0^{y_D} \frac{y^3}{e^y - 1} dy \\
 &= U_0 + \frac{9N}{\omega_D^3} \frac{(kT)^4}{\hbar^3} \int_0^{y_D} \frac{y^3}{e^y - 1} dy = U_0 + \frac{9N}{y_D^3} kT \int_0^{y_D} \frac{y^3}{e^y - 1} dy \\
 &\equiv U_0 + 3NkT \mathcal{D}(y_D), \text{ where } \mathcal{D}(y_D) = \frac{3}{y_D^3} \int_0^{y_D} \frac{y^3}{e^y - 1} dy.
 \end{aligned}$$
- High temperature, $y_D \ll 1$, $\mathcal{D}(y_D) \simeq \frac{3}{y_D^3} \int_0^{y_D} y^2 dy$

$$U = U_0 + B \int_0^{\omega_D} \frac{\hbar \omega^3}{e^{\beta \hbar \omega} - 1} d\omega, \quad \omega_D^3 = \frac{9N}{B}, \quad y_D = \frac{\hbar \omega_D}{kT}$$

- $$\begin{aligned}
 U &= U_0 + B \int_0^{\frac{\hbar \omega_D}{kT}} \frac{\hbar \cdot (\frac{\hbar \omega}{kT})^3 \cdot (\frac{kT}{\hbar})^3}{e^{\frac{\hbar \omega}{kT}} - 1} \frac{kT}{\hbar} d\frac{\hbar \omega}{kT} \\
 &= U_0 + B \hbar \left(\frac{kT}{\hbar}\right)^4 \int_0^{y_D} \frac{y^3}{e^y - 1} dy \\
 &= U_0 + \frac{9N}{\omega_D^3} \frac{(kT)^4}{\hbar^3} \int_0^{y_D} \frac{y^3}{e^y - 1} dy = U_0 + \frac{9N}{y_D^3} kT \int_0^{y_D} \frac{y^3}{e^y - 1} dy \\
 &\equiv U_0 + 3NkT \mathcal{D}(y_D), \text{ where } \mathcal{D}(y_D) = \frac{3}{y_D^3} \int_0^{y_D} \frac{y^3}{e^y - 1} dy.
 \end{aligned}$$
- High temperature, $y_D \ll 1$, $\mathcal{D}(y_D) \simeq \frac{3}{y_D^3} \int_0^{y_D} y^2 dy = 1$.

$$U = U_0 + B \int_0^{\omega_D} \frac{\hbar\omega^3}{e^{\beta\hbar\omega} - 1} d\omega, \quad \omega_D^3 = \frac{9N}{B}, \quad y_D = \frac{\hbar\omega_D}{kT}$$

- $$\begin{aligned}
 U &= U_0 + B \int_0^{\frac{\hbar\omega_D}{kT}} \frac{\hbar \cdot (\frac{\hbar\omega}{kT})^3 \cdot (\frac{kT}{\hbar})^3}{e^{\frac{\hbar\omega}{kT}} - 1} \frac{kT}{\hbar} d\frac{\hbar\omega}{kT} \\
 &= U_0 + B\hbar \left(\frac{kT}{\hbar}\right)^4 \int_0^{y_D} \frac{y^3}{e^y - 1} dy \\
 &= U_0 + \frac{9N}{\omega_D^3} \frac{(kT)^4}{\hbar^3} \int_0^{y_D} \frac{y^3}{e^y - 1} dy = U_0 + \frac{9N}{y_D^3} kT \int_0^{y_D} \frac{y^3}{e^y - 1} dy \\
 &\equiv U_0 + 3NkT \mathcal{D}(y_D), \text{ where } \mathcal{D}(y_D) = \frac{3}{y_D^3} \int_0^{y_D} \frac{y^3}{e^y - 1} dy.
 \end{aligned}$$
- High temperature, $y_D \ll 1$, $\mathcal{D}(y_D) \simeq \frac{3}{y_D^3} \int_0^{y_D} y^2 dy = 1$. $U = U_0 + 3NkT$, $C_V = 3Nk$.

$$U = U_0 + B \int_0^{\omega_D} \frac{\hbar\omega^3}{e^{\beta\hbar\omega} - 1} d\omega, \quad \omega_D^3 = \frac{9N}{B}, \quad y_D = \frac{\hbar\omega_D}{kT}$$

- $$\begin{aligned}
 U &= U_0 + B \int_0^{\frac{\hbar\omega_D}{kT}} \frac{\hbar \cdot (\frac{\hbar\omega}{kT})^3 \cdot (\frac{kT}{\hbar})^3}{e^{\frac{\hbar\omega}{kT}} - 1} \frac{kT}{\hbar} d\frac{\hbar\omega}{kT} \\
 &= U_0 + B\hbar \left(\frac{kT}{\hbar}\right)^4 \int_0^{y_D} \frac{y^3}{e^y - 1} dy \\
 &= U_0 + \frac{9N}{\omega_D^3} \frac{(kT)^4}{\hbar^3} \int_0^{y_D} \frac{y^3}{e^y - 1} dy = U_0 + \frac{9N}{y_D^3} kT \int_0^{y_D} \frac{y^3}{e^y - 1} dy \\
 &\equiv U_0 + 3NkT \mathcal{D}(y_D), \text{ where } \mathcal{D}(y_D) = \frac{3}{y_D^3} \int_0^{y_D} \frac{y^3}{e^y - 1} dy.
 \end{aligned}$$
- High temperature, $y_D \ll 1$, $\mathcal{D}(y_D) \simeq \frac{3}{y_D^3} \int_0^{y_D} y^2 dy = 1$. $U = U_0 + 3NkT$, $C_V = 3Nk$.
- Low temperature, $y_D \gg 1$,

$$U = U_0 + B \int_0^{\omega_D} \frac{\hbar \omega^3}{e^{\beta \hbar \omega} - 1} d\omega, \quad \omega_D^3 = \frac{9N}{B}, \quad y_D = \frac{\hbar \omega_D}{kT}$$

- $$\begin{aligned}
 U &= U_0 + B \int_0^{\frac{\hbar \omega_D}{kT}} \frac{\hbar \cdot (\frac{\hbar \omega}{kT})^3 \cdot (\frac{kT}{\hbar})^3}{e^{\frac{\hbar \omega}{kT}} - 1} \frac{kT}{\hbar} d\frac{\hbar \omega}{kT} \\
 &= U_0 + B \hbar \left(\frac{kT}{\hbar}\right)^4 \int_0^{y_D} \frac{y^3}{e^y - 1} dy \\
 &= U_0 + \frac{9N}{\omega_D^3} \frac{(kT)^4}{\hbar^3} \int_0^{y_D} \frac{y^3}{e^y - 1} dy = U_0 + \frac{9N}{y_D^3} kT \int_0^{y_D} \frac{y^3}{e^y - 1} dy \\
 &\equiv U_0 + 3NkT \mathcal{D}(y_D), \text{ where } \mathcal{D}(y_D) = \frac{3}{y_D^3} \int_0^{y_D} \frac{y^3}{e^y - 1} dy.
 \end{aligned}$$
- High temperature, $y_D \ll 1$, $\mathcal{D}(y_D) \simeq \frac{3}{y_D^3} \int_0^{y_D} y^2 dy = 1$. $U = U_0 + 3NkT$, $C_V = 3Nk$.
- Low temperature, $y_D \gg 1$, $\mathcal{D}(y_D) \simeq \frac{3}{y_D^3} \int_0^{\infty} \frac{y^3}{e^y - 1} dy$

$$U = U_0 + B \int_0^{\omega_D} \frac{\hbar\omega^3}{e^{\beta\hbar\omega} - 1} d\omega, \quad \omega_D^3 = \frac{9N}{B}, \quad y_D = \frac{\hbar\omega_D}{kT}$$

- $$U = U_0 + B \int_0^{\frac{\hbar\omega_D}{kT}} \frac{\hbar \cdot (\frac{\hbar\omega}{kT})^3 \cdot (\frac{kT}{\hbar})^3}{e^{\frac{\hbar\omega}{kT}} - 1} \frac{kT}{\hbar} d\frac{\hbar\omega}{kT}$$

$$= U_0 + B\hbar \left(\frac{kT}{\hbar}\right)^4 \int_0^{y_D} \frac{y^3}{e^y - 1} dy$$

$$= U_0 + \frac{9N}{\omega_D^3} \frac{(kT)^4}{\hbar^3} \int_0^{y_D} \frac{y^3}{e^y - 1} dy = U_0 + \frac{9N}{y_D^3} kT \int_0^{y_D} \frac{y^3}{e^y - 1} dy$$

$$\equiv U_0 + 3NkT \mathcal{D}(y_D), \text{ where } \mathcal{D}(y_D) = \frac{3}{y_D^3} \int_0^{y_D} \frac{y^3}{e^y - 1} dy.$$
- High temperature, $y_D \ll 1$, $\mathcal{D}(y_D) \simeq \frac{3}{y_D^3} \int_0^{y_D} y^2 dy$
 $= 1$. $U = U_0 + 3NkT$, $C_V = 3Nk$.
- Low temperature, $y_D \gg 1$, $\mathcal{D}(y_D) \simeq \frac{3}{y_D^3} \int_0^\infty \frac{y^3}{e^y - 1} dy$
 $= \frac{3}{y_D^3} \frac{\pi^4}{15}$

$$U = U_0 + B \int_0^{\omega_D} \frac{\hbar \omega^3}{e^{\beta \hbar \omega} - 1} d\omega, \quad \omega_D^3 = \frac{9N}{B}, \quad y_D = \frac{\hbar \omega_D}{kT}$$

- $$U = U_0 + B \int_0^{\frac{\hbar \omega_D}{kT}} \frac{\hbar \cdot (\frac{\hbar \omega}{kT})^3 \cdot (\frac{kT}{\hbar})^3}{e^{\frac{\hbar \omega}{kT}} - 1} \frac{kT}{\hbar} d\frac{\hbar \omega}{kT}$$

$$= U_0 + B \hbar \left(\frac{kT}{\hbar}\right)^4 \int_0^{y_D} \frac{y^3}{e^y - 1} dy$$

$$= U_0 + \frac{9N}{\omega_D^3} \frac{(kT)^4}{\hbar^3} \int_0^{y_D} \frac{y^3}{e^y - 1} dy = U_0 + \frac{9N}{y_D^3} kT \int_0^{y_D} \frac{y^3}{e^y - 1} dy$$

$$\equiv U_0 + 3NkT \mathcal{D}(y_D), \text{ where } \mathcal{D}(y_D) = \frac{3}{y_D^3} \int_0^{y_D} \frac{y^3}{e^y - 1} dy.$$
- High temperature, $y_D \ll 1$, $\mathcal{D}(y_D) \simeq \frac{3}{y_D^3} \int_0^{y_D} y^2 dy = 1$. $U = U_0 + 3NkT$, $C_V = 3Nk$.
- Low temperature, $y_D \gg 1$, $\mathcal{D}(y_D) \simeq \frac{3}{y_D^3} \int_0^\infty \frac{y^3}{e^y - 1} dy$

$$= \frac{3}{y_D^3} \frac{\pi^4}{15} = \frac{\pi^4}{5y_D^3}.$$

$$U = U_0 + B \int_0^{\omega_D} \frac{\hbar\omega^3}{e^{\beta\hbar\omega} - 1} d\omega, \quad \omega_D^3 = \frac{9N}{B}, \quad y_D = \frac{\hbar\omega_D}{kT}$$

- $$U = U_0 + B \int_0^{\frac{\hbar\omega_D}{kT}} \frac{\hbar \cdot (\frac{\hbar\omega}{kT})^3 \cdot (\frac{kT}{\hbar})^3}{e^{\frac{\hbar\omega}{kT}} - 1} \frac{kT}{\hbar} d\frac{\hbar\omega}{kT}$$

$$= U_0 + B\hbar \left(\frac{kT}{\hbar}\right)^4 \int_0^{y_D} \frac{y^3}{e^y - 1} dy$$

$$= U_0 + \frac{9N}{\omega_D^3} \frac{(kT)^4}{\hbar^3} \int_0^{y_D} \frac{y^3}{e^y - 1} dy = U_0 + \frac{9N}{y_D^3} kT \int_0^{y_D} \frac{y^3}{e^y - 1} dy$$

$$\equiv U_0 + 3NkT \mathcal{D}(y_D), \text{ where } \mathcal{D}(y_D) = \frac{3}{y_D^3} \int_0^{y_D} \frac{y^3}{e^y - 1} dy.$$
- High temperature, $y_D \ll 1$, $\mathcal{D}(y_D) \simeq \frac{3}{y_D^3} \int_0^{y_D} y^2 dy = 1$. $U = U_0 + 3NkT$, $C_V = 3Nk$.
- Low temperature, $y_D \gg 1$, $\mathcal{D}(y_D) \simeq \frac{3}{y_D^3} \int_0^\infty \frac{y^3}{e^y - 1} dy$

$$= \frac{3}{y_D^3} \frac{\pi^4}{15} = \frac{\pi^4}{5y_D^3}. \quad U = U_0 + 3Nk \cdot \frac{\pi^4}{5} \frac{T^4}{\theta_D^4},$$

$$C_V = 3nk \frac{4\pi^4}{5} \left(\frac{T}{\theta_D}\right)^3 \propto T^3. \text{ Source of (8.5.20).}$$

Table of contents

1 §9. Ensemble theory

- 9.1 Phase space and Liouville's theorem
- 9.2 The microcanonical ensemble
- 9.3 Thermodynamics in microcanonical ensemble theory
- 9.4 Canonical ensemble
- 9.5 Thermodynamics in canonical ensemble
- 9.6 EOS of real gas
- 9.7 Thermal capacity of solid
- 9.9 Ising model
- 9.10 Grand canonical ensemble
- 9.11 Thermodynamics in grand canonical ensemble
- 9.12 Applications of the grand canonical ensemble

9.9 Ising model

9.9 Ising model

- Problem: phase transition.

9.9 Ising model

- Problem: phase transition.
- Question: can statistical mechanics deal with phase transition?

9.9 Ising model

- Problem: phase transition.
- Question: can statistical mechanics deal with phase transition?
- Object: N magnetons at N lattice points.

9.9 Ising model

- Problem: phase transition.
- Question: can statistical mechanics deal with phase transition?
- Object: N magnetons at N lattice points. Analyze the phase transition between paramagnet and ferromagnet. (simple)

9.9 Ising model

- Problem: phase transition.
- Question: can statistical mechanics deal with phase transition?
- Object: N magnetons at N lattice points. Analyze the phase transition between paramagnet and ferromagnet. (simple)
- Suppose the total angular momentum quantum number $\frac{1}{2}$ (simple),

9.9 Ising model

- Problem: phase transition.
- Question: can statistical mechanics deal with phase transition?
- Object: N magnetons at N lattice points. Analyze the phase transition between paramagnet and ferromagnet. (simple)
- Suppose the total angular momentum quantum number $\frac{1}{2}$ (simple), the magnetic momentum: $\mu = \frac{e\hbar}{2m}$.

9.9 Ising model

- Problem: phase transition.
- Question: can statistical mechanics deal with phase transition?
- Object: N magnetons at N lattice points. Analyze the phase transition between paramagnet and ferromagnet. (simple)
- Suppose the total angular momentum quantum number $\frac{1}{2}$ (simple), the magnetic momentum: $\mu = \frac{e\hbar}{2m}$.
- In single-axis ferromagnet, only 2 choices for each magnetic momentum ($\sigma = \pm 1$).

Ferromagnet

- Potential energy between two magnetic momenta: $J_1 \vec{\mu}_1 \cdot \vec{\mu}_2$, where J_1 is a coefficient depending on position.

Ferromagnet

- Potential energy between two magnetic momenta: $J_1 \vec{\mu}_1 \cdot \vec{\mu}_2$, where J_1 is a coefficient depending on position.
- $J_1 \vec{\mu}_1 \cdot \vec{\mu}_2 = J_2 \sigma_1 \sigma_2$.

Ferromagnet

- Potential energy between two magnetic momenta: $J_1 \vec{\mu}_1 \cdot \vec{\mu}_2$, where J_1 is a coefficient depending on position.
- $J_1 \vec{\mu}_1 \cdot \vec{\mu}_2 = J_2 \sigma_1 \sigma_2$.
- Ising model: Similar form for the total interacting energy between two lattice points: $-J \sigma_1 \sigma_2$, where J is a constant for neighbors, 0 for others.

Ferromagnet

- Potential energy between two magnetic momenta: $J_1 \vec{\mu}_1 \cdot \vec{\mu}_2$, where J_1 is a coefficient depending on position.
- $J_1 \vec{\mu}_1 \cdot \vec{\mu}_2 = J_2 \sigma_1 \sigma_2$.
- Ising model: Similar form for the total interacting energy between two lattice points: $-J \sigma_1 \sigma_2$, where J is a constant for neighbors, 0 for others.
- Total interacting energy: $-J \sum'_{i,j} \sigma_i \sigma_j$, where $\sum'_{i,j}$ means the summation only exerts on the neighbors.

Ferromagnet

- Potential energy between two magnetic momenta: $J_1 \vec{\mu}_1 \cdot \vec{\mu}_2$, where J_1 is a coefficient depending on position.
- $J_1 \vec{\mu}_1 \cdot \vec{\mu}_2 = J_2 \sigma_1 \sigma_2$.
- Ising model: Similar form for the total interacting energy between two lattice points: $-J \sigma_1 \sigma_2$, where J is a constant for neighbors, 0 for others.
- Total interacting energy: $-J \sum'_{i,j} \sigma_i \sigma_j$, where $\sum'_{i,j}$ means the summation only exerts on the neighbors.
At absolute 0K, all directions being the same can get lowest total energy. Get macroscopic magnetic field, spontaneous magnetization.

Mean field approximation

- In external magnetic field, the total energy:

$$E\{\sigma_i\} = -J \sum_{i,j}' \sigma_i \sigma_j - \mu B \sum_i \sigma_i.$$

Mean field approximation

- In external magnetic field, the total energy:

$$E\{\sigma_i\} = -J \sum_{i,j}' \sigma_i \sigma_j - \mu B \sum_i \sigma_i.$$

- The partition function: $Z = \sum_s e^{-\beta E_s}$

Mean field approximation

- In external magnetic field, the total energy:

$$E\{\sigma_i\} = -J \sum_{i,j}' \sigma_i \sigma_j - \mu B \sum_i \sigma_i.$$

- The partition function: $Z = \sum_s e^{-\beta E_s} = \sum_{\{\sigma_i\}} e^{-\beta E\{\sigma_i\}}$

Mean field approximation

- In external magnetic field, the total energy:

$$E\{\sigma_i\} = -J \sum_{i,j}' \sigma_i \sigma_j - \mu B \sum_i \sigma_i.$$

- The partition function: $Z = \sum_s e^{-\beta E_s} = \sum_{\{\sigma_i\}} e^{-\beta E\{\sigma_i\}}$
$$= \sum_{\sigma_1=\pm 1} \sum_{\sigma_2=\pm 1} \cdots \sum_{\sigma_N=\pm 1} e^{-\beta E\{\sigma_i\}}.$$

Mean field approximation

- In external magnetic field, the total energy:

$$E\{\sigma_i\} = -J \sum_{i,j}' \sigma_i \sigma_j - \mu B \sum_i \sigma_i.$$

- The partition function: $Z = \sum_s e^{-\beta E_s} = \sum_{\{\sigma_i\}} e^{-\beta E\{\sigma_i\}}$

$$= \sum_{\sigma_1=\pm 1} \sum_{\sigma_2=\pm 1} \cdots \sum_{\sigma_N=\pm 1} e^{-\beta E\{\sigma_i\}}.$$

- Important task is to solve it.

Mean field approximation

- In external magnetic field, the total energy:

$$E\{\sigma_i\} = -J \sum_{i,j}' \sigma_i \sigma_j - \mu B \sum_i \sigma_i.$$

- The partition function: $Z = \sum_s e^{-\beta E_s} = \sum_{\{\sigma_i\}} e^{-\beta E\{\sigma_i\}}$

$$= \sum_{\sigma_1=\pm 1} \sum_{\sigma_2=\pm 1} \cdots \sum_{\sigma_N=\pm 1} e^{-\beta E\{\sigma_i\}}.$$

- Important task is to solve it.
- An approximation solution: mean field approximation.

Mean field approximation

- In external magnetic field, the total energy:

$$E\{\sigma_i\} = -J \sum'_{i,j} \sigma_i \sigma_j - \mu B \sum_i \sigma_i.$$

- The partition function: $Z = \sum_s e^{-\beta E_s} = \sum_{\{\sigma_i\}} e^{-\beta E\{\sigma_i\}}$

$$= \sum_{\sigma_1=\pm 1} \sum_{\sigma_2=\pm 1} \cdots \sum_{\sigma_N=\pm 1} e^{-\beta E\{\sigma_i\}}.$$

- Important task is to solve it.
- An approximation solution: mean field approximation.
- Rewrite the total energy:

$$E\{\sigma_i\} = -\frac{1}{2} \sum_{i,j} J_{ij} \sigma_i \sigma_j - \mu B \sum_i \sigma_i, \text{ where } J_{ij} = J \text{ for } i \text{ and } j \text{ being neighbors, } 0 \text{ for other.}$$

Mean field approximation

$$E = -\frac{1}{2} \sum_{i,j} J_{ij} \sigma_i \sigma_j - \mu B \sum_i \sigma_i$$

- $-\sigma_i$ acts as the general displacement,

Mean field approximation

$$E = -\frac{1}{2} \sum_{i,j} J_{ij} \sigma_i \sigma_j - \mu B \sum_i \sigma_i$$

- $-\sigma_i$ acts as the general displacement, the general force on lattice i : $-\frac{\partial E}{\partial \sigma_i} = \mu B + \sum_j J_{ij} \sigma_j$.

Mean field approximation

$$E = -\frac{1}{2} \sum_{i,j} J_{ij} \sigma_i \sigma_j - \mu B \sum_i \sigma_i$$

- $-\sigma_i$ acts as the general displacement, the general force on lattice i : $-\frac{\partial E}{\partial \sigma_i} = \mu B + \sum_j J_{ij} \sigma_j$.
- Like an effective magnetic field exerted on lattice i :
$$B_i = B + \frac{1}{\mu} \sum_j J_{ij} \sigma_j.$$

Mean field approximation

$$E = -\frac{1}{2} \sum_{i,j} J_{ij} \sigma_i \sigma_j - \mu B \sum_i \sigma_i$$

- $-\sigma_i$ acts as the general displacement, the general force on lattice i : $-\frac{\partial E}{\partial \sigma_i} = \mu B + \sum_j J_{ij} \sigma_j$.
- Like an effective magnetic field exerted on lattice i :
$$B_i = B + \frac{1}{\mu} \sum_j J_{ij} \sigma_j.$$
- As $\{\sigma_i\}$ changes for different states, the B_i is different for different states.

Mean field approximation

$$E = -\frac{1}{2} \sum_{i,j} J_{ij} \sigma_i \sigma_j - \mu B \sum_i \sigma_i$$

- $-\sigma_i$ acts as the general displacement, the general force on lattice i : $-\frac{\partial E}{\partial \sigma_i} = \mu B + \sum_j J_{ij} \sigma_j$.
- Like an effective magnetic field exerted on lattice i :
$$B_i = B + \frac{1}{\mu} \sum_j J_{ij} \sigma_j.$$
- As $\{\sigma_i\}$ changes for different states, the B_i is different for different states. However, choose an average value for the effective field: $\bar{B}_i = B + \frac{1}{\mu} \sum_j J_{ij} \bar{\sigma}_j,$

Mean field approximation

$$E = -\frac{1}{2} \sum_{i,j} J_{ij} \sigma_i \sigma_j - \mu B \sum_i \sigma_i$$

- $-\sigma_i$ acts as the general displacement, the general force on lattice i : $-\frac{\partial E}{\partial \sigma_i} = \mu B + \sum_j J_{ij} \sigma_j$.
- Like an effective magnetic field exerted on lattice i :
$$B_i = B + \frac{1}{\mu} \sum_j J_{ij} \sigma_j.$$
- As $\{\sigma_i\}$ changes for different states, the B_i is different for different states. However, choose an average value for the effective field: $\bar{B}_i = B + \frac{1}{\mu} \sum_j J_{ij} \bar{\sigma}_j$, called mean field approximation.

Mean field approximation

$$\bar{B}_i = B + \frac{1}{\mu} \sum_j J_{ij} \bar{\sigma}_j$$

- As $\bar{\sigma}_j$ is an average value, which doesn't depend on j ,
 $\bar{\sigma}_j = \bar{\sigma}_i$.

Mean field approximation

$$\bar{B}_i = B + \frac{1}{\mu} \sum_j J_{ij} \bar{\sigma}_j$$

- As $\bar{\sigma}_j$ is an average value, which doesn't depend on j ,
 $\bar{\sigma}_j = \bar{\sigma}_i \rightarrow \sum_j J_{ij} \bar{\sigma}_j = \bar{\sigma}_i \sum_j J_{ij}.$

Mean field approximation

$$\bar{B}_i = B + \frac{1}{\mu} \sum_j J_{ij} \bar{\sigma}_j$$

- As $\bar{\sigma}_j$ is an average value, which doesn't depend on j , $\bar{\sigma}_j = \bar{\sigma}_i$. $\rightarrow \sum_j J_{ij} \bar{\sigma}_j = \bar{\sigma}_i \sum_j J_{ij}$.
- $\sum_j J_{ij} = Jz$, where z is the number of neighbors.

Mean field approximation

$$\bar{B}_i = B + \frac{1}{\mu} \sum_j J_{ij} \bar{\sigma}_j$$

- As $\bar{\sigma}_j$ is an average value, which doesn't depend on j , $\bar{\sigma}_j = \bar{\sigma}_i$. $\rightarrow \sum_j J_{ij} \bar{\sigma}_j = \bar{\sigma}_i \sum_j J_{ij}$.
- $\sum_j J_{ij} = Jz$, where z is the number of neighbors.
- $\Rightarrow \bar{B}_i = B + \frac{1}{\mu} Jz \bar{\sigma}_i$.

Mean field approximation

$$\bar{B}_i = B + \frac{1}{\mu} \sum_j J_{ij} \bar{\sigma}_j$$

- As $\bar{\sigma}_j$ is an average value, which doesn't depend on j , $\bar{\sigma}_j = \bar{\sigma}_i$. $\rightarrow \sum_j J_{ij} \bar{\sigma}_j = \bar{\sigma}_i \sum_j J_{ij}$.
- $\sum_j J_{ij} = Jz$, where z is the number of neighbors.
- $\Rightarrow \bar{B}_i = B + \frac{1}{\mu} Jz \bar{\sigma}_i$.
- As $\bar{\sigma}_i$ doesn't depend on i ,

Mean field approximation

$$\bar{B}_i = B + \frac{1}{\mu} \sum_j J_{ij} \bar{\sigma}_j$$

- As $\bar{\sigma}_j$ is an average value, which doesn't depend on j , $\bar{\sigma}_j = \bar{\sigma}_i$. $\rightarrow \sum_j J_{ij} \bar{\sigma}_j = \bar{\sigma}_i \sum_j J_{ij}$.
- $\sum_j J_{ij} = Jz$, where z is the number of neighbors.
- $\Rightarrow \bar{B}_i = B + \frac{1}{\mu} Jz \bar{\sigma}_i$.
- As $\bar{\sigma}_i$ doesn't depend on i , $\bar{B} = B + \frac{1}{\mu} Jz \bar{\sigma}$.

Mean field approximation

$$\bar{B}_i = B + \frac{1}{\mu} \sum_j J_{ij} \bar{\sigma}_j$$

- As $\bar{\sigma}_j$ is an average value, which doesn't depend on j , $\bar{\sigma}_j = \bar{\sigma}_i$. $\rightarrow \sum_j J_{ij} \bar{\sigma}_j = \bar{\sigma}_i \sum_j J_{ij}$.
- $\sum_j J_{ij} = Jz$, where z is the number of neighbors.
- $\Rightarrow \bar{B}_i = B + \frac{1}{\mu} Jz \bar{\sigma}_i$.
- As $\bar{\sigma}_i$ doesn't depend on i , $\bar{B} = B + \frac{1}{\mu} Jz \bar{\sigma}$.
- The total energy is then: $E = -\mu \bar{B} \sum_i \sigma_i$.

Mean field approximation

$$\bar{B}_i = B + \frac{1}{\mu} \sum_j J_{ij} \bar{\sigma}_j$$

- As $\bar{\sigma}_j$ is an average value, which doesn't depend on j , $\bar{\sigma}_j = \bar{\sigma}_i$. $\rightarrow \sum_j J_{ij} \bar{\sigma}_j = \bar{\sigma}_i \sum_j J_{ij}$.
- $\sum_j J_{ij} = Jz$, where z is the number of neighbors.
- $\Rightarrow \bar{B}_i = B + \frac{1}{\mu} Jz \bar{\sigma}_i$.
- As $\bar{\sigma}_i$ doesn't depend on i , $\bar{B} = B + \frac{1}{\mu} Jz \bar{\sigma}$.
- The total energy is then: $E = -\mu \bar{B} \sum_i \sigma_i$.
- The partition function: $Z = \sum_{\sigma_1=\pm 1} \cdots \sum_{\sigma_N=\pm 1} e^{-\beta E\{\sigma_i\}}$

Mean field approximation

$$\bar{B}_i = B + \frac{1}{\mu} \sum_j J_{ij} \bar{\sigma}_j$$

- As $\bar{\sigma}_j$ is an average value, which doesn't depend on j , $\bar{\sigma}_j = \bar{\sigma}_i$. $\rightarrow \sum_j J_{ij} \bar{\sigma}_j = \bar{\sigma}_i \sum_j J_{ij}$.
- $\sum_j J_{ij} = Jz$, where z is the number of neighbors.
- $\Rightarrow \bar{B}_i = B + \frac{1}{\mu} Jz \bar{\sigma}_i$.
- As $\bar{\sigma}_i$ doesn't depend on i , $\bar{B} = B + \frac{1}{\mu} Jz \bar{\sigma}$.
- The total energy is then: $E = -\mu \bar{B} \sum_i \sigma_i$.
- The partition function:
$$Z = \sum_{\sigma_1=\pm 1} \cdots \sum_{\sigma_N=\pm 1} e^{-\beta E\{\sigma_i\}}$$
$$= \sum_{\sigma_1=\pm 1} \cdots \sum_{\sigma_N=\pm 1} e^{\beta \mu \bar{B} \sum_i \sigma_i}$$

Mean field approximation

$$\bar{B}_i = B + \frac{1}{\mu} \sum_j J_{ij} \bar{\sigma}_j$$

- As $\bar{\sigma}_j$ is an average value, which doesn't depend on j , $\bar{\sigma}_j = \bar{\sigma}_i$. $\rightarrow \sum_j J_{ij} \bar{\sigma}_j = \bar{\sigma}_i \sum_j J_{ij}$.
- $\sum_j J_{ij} = Jz$, where z is the number of neighbors.
- $\Rightarrow \bar{B}_i = B + \frac{1}{\mu} Jz \bar{\sigma}_i$.
- As $\bar{\sigma}_i$ doesn't depend on i , $\bar{B} = B + \frac{1}{\mu} Jz \bar{\sigma}$.
- The total energy is then: $E = -\mu \bar{B} \sum_i \sigma_i$.
- The partition function:
$$Z = \sum_{\sigma_1=\pm 1} \cdots \sum_{\sigma_N=\pm 1} e^{-\beta E\{\sigma_i\}}$$
$$= \sum_{\sigma_1=\pm 1} \cdots \sum_{\sigma_N=\pm 1} e^{\beta \mu \bar{B} \sum_i \sigma_i} = \prod_i \sum_{\sigma_i} e^{\beta \mu \bar{B} \sigma_i}$$

Mean field approximation

$$\bar{B} = B + \frac{1}{\mu} J z \bar{\sigma}$$

- $$Z = \prod_i \sum_{\sigma_i} e^{\beta \mu \bar{B} \sigma_i}$$

Mean field approximation

$$\bar{B} = B + \frac{1}{\mu} J z \bar{\sigma}$$

- $$Z = \prod_i \sum_{\sigma_i} e^{\beta \mu \bar{B} \sigma_i} = \prod_i (e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}})$$

Mean field approximation

$$\bar{B} = B + \frac{1}{\mu} J z \bar{\sigma}$$

- $$Z = \prod_i \sum_{\sigma_i} e^{\beta \mu \bar{B} \sigma_i} = \prod_i (e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}) = Z_1^N,$$

where $Z_1 = e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}$. (Comparing with (7.8.1))

Mean field approximation

$$\bar{B} = B + \frac{1}{\mu} J z \bar{\sigma}$$

- $Z = \prod_i \sum_{\sigma_i} e^{\beta \mu \bar{B} \sigma_i} = \prod_i (e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}) = Z_1^N$,
where $Z_1 = e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}$. (Comparing with (7.8.1))
- Magnetic momentum $-m$ and magnetic field B are a pair of general force and general displacement (2.7.19),

Mean field approximation

$$\bar{B} = B + \frac{1}{\mu} J z \bar{\sigma}$$

- $Z = \prod_i \sum_{\sigma_i} e^{\beta \mu \bar{B} \sigma_i} = \prod_i (e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}) = Z_1^N,$

where $Z_1 = e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}$. (Comparing with (7.8.1))

- Magnetic momentum $-m$ and magnetic field B are a pair of general force and general displacement (2.7.19), $\bar{m} = \frac{1}{\beta} \frac{\partial \ln Z}{\partial \bar{B}}$

Mean field approximation

$$\bar{B} = B + \frac{1}{\mu} J z \bar{\sigma}$$

- $Z = \prod_i \sum_{\sigma_i} e^{\beta \mu \bar{B} \sigma_i} = \prod_i (e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}) = Z_1^N$,
 where $Z_1 = e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}$. (Comparing with (7.8.1))
- Magnetic momentum $-m$ and magnetic field B are a pair of general force and general displacement (2.7.19), $\bar{m} = \frac{1}{\beta} \frac{\partial \ln Z}{\partial \bar{B}} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial \bar{B}}$

Mean field approximation

$$\bar{B} = B + \frac{1}{\mu} J z \bar{\sigma}$$

- $Z = \prod_i \sum_{\sigma_i} e^{\beta \mu \bar{B} \sigma_i} = \prod_i (e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}) = Z_1^N,$

where $Z_1 = e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}$. (Comparing with (7.8.1))

- Magnetic momentum $-m$ and magnetic field B are a pair of general force and general displacement (2.7.19), $\bar{m} = \frac{1}{\beta} \frac{\partial \ln Z}{\partial \bar{B}} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial \bar{B}} = N \mu \frac{e^{\beta \mu \bar{B}} - e^{-\beta \mu \bar{B}}}{e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}}.$

Mean field approximation

$$\bar{B} = B + \frac{1}{\mu} J z \bar{\sigma}$$

$$\bullet Z = \prod_i \sum_{\sigma_i} e^{\beta \mu \bar{B} \sigma_i} = \prod_i (e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}) = Z_1^N,$$

where $Z_1 = e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}$. (Comparing with (7.8.1))

- Magnetic momentum $-m$ and magnetic field B are a pair of general force and general displacement (2.7.19), $\bar{m} = \frac{1}{\beta} \frac{\partial \ln Z}{\partial \bar{B}} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial \bar{B}} = N \mu \frac{e^{\beta \mu \bar{B}} - e^{-\beta \mu \bar{B}}}{e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}}$.
- On the other hand, $\bar{m} = N \mu \bar{\sigma}$.

Mean field approximation

$$\bar{B} = B + \frac{1}{\mu} J z \bar{\sigma}$$

$$\bullet Z = \prod_i \sum_{\sigma_i} e^{\beta \mu \bar{B} \sigma_i} = \prod_i (e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}) = Z_1^N,$$

where $Z_1 = e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}$. (Comparing with (7.8.1))

- Magnetic momentum $-m$ and magnetic field B are a pair of general force and general displacement

$$(2.7.19), \bar{m} = \frac{1}{\beta} \frac{\partial \ln Z}{\partial \bar{B}} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial \bar{B}} = N \mu \frac{e^{\beta \mu \bar{B}} - e^{-\beta \mu \bar{B}}}{e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}}.$$

- On the other hand, $\bar{m} = N \mu \bar{\sigma}$.

$$\bullet \Rightarrow \bar{\sigma} = \frac{e^{\beta \mu \bar{B}} - e^{-\beta \mu \bar{B}}}{e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}}.$$

Mean field approximation

$$\bar{B} = B + \frac{1}{\mu} J z \bar{\sigma}$$

- $Z = \prod_i \sum_{\sigma_i} e^{\beta \mu \bar{B} \sigma_i} = \prod_i (e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}) = Z_1^N$,
where $Z_1 = e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}$. (Comparing with (7.8.1))
- Magnetic momentum $-m$ and magnetic field B are a pair of general force and general displacement (2.7.19), $\bar{m} = \frac{1}{\beta} \frac{\partial \ln Z}{\partial \bar{B}} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial \bar{B}} = N \mu \frac{e^{\beta \mu \bar{B}} - e^{-\beta \mu \bar{B}}}{e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}}$.
- On the other hand, $\bar{m} = N \mu \bar{\sigma}$.
- $\Rightarrow \bar{\sigma} = \frac{e^{\beta \mu \bar{B}} - e^{-\beta \mu \bar{B}}}{e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}}$.
- Without external field, $\bar{B} = \frac{1}{\mu} J z \bar{\sigma}$.

Mean field approximation

$$\bar{B} = B + \frac{1}{\mu} J z \bar{\sigma}$$

- $Z = \prod_i \sum_{\sigma_i} e^{\beta \mu \bar{B} \sigma_i} = \prod_i (e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}) = Z_1^N$,
where $Z_1 = e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}$. (Comparing with (7.8.1))
- Magnetic momentum $-m$ and magnetic field B are a pair of general force and general displacement (2.7.19), $\bar{m} = \frac{1}{\beta} \frac{\partial \ln Z}{\partial \bar{B}} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial \bar{B}} = N \mu \frac{e^{\beta \mu \bar{B}} - e^{-\beta \mu \bar{B}}}{e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}}$.
- On the other hand, $\bar{m} = N \mu \bar{\sigma}$.
- $\Rightarrow \bar{\sigma} = \frac{e^{\beta \mu \bar{B}} - e^{-\beta \mu \bar{B}}}{e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}}$.
- Without external field, $\bar{B} = \frac{1}{\mu} J z \bar{\sigma}$. $\rightarrow \beta \mu \bar{B} = \beta J z \bar{\sigma}$.

Mean field approximation

$$\bar{B} = B + \frac{1}{\mu} J z \bar{\sigma}$$

- $Z = \prod_i \sum_{\sigma_i} e^{\beta \mu \bar{B} \sigma_i} = \prod_i (e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}) = Z_1^N$,
 where $Z_1 = e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}$. (Comparing with (7.8.1))
- Magnetic momentum $-m$ and magnetic field B are a pair of general force and general displacement (2.7.19), $\bar{m} = \frac{1}{\beta} \frac{\partial \ln Z}{\partial \bar{B}} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial \bar{B}} = N \mu \frac{e^{\beta \mu \bar{B}} - e^{-\beta \mu \bar{B}}}{e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}}$.
- On the other hand, $\bar{m} = N \mu \bar{\sigma}$.
- $\Rightarrow \bar{\sigma} = \frac{e^{\beta \mu \bar{B}} - e^{-\beta \mu \bar{B}}}{e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}}$.
- Without external field, $\bar{B} = \frac{1}{\mu} J z \bar{\sigma}$. $\rightarrow \beta \mu \bar{B} = \beta J z \bar{\sigma}$.
 $\therefore Z_1 = e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}$;

Mean field approximation

$$\bar{B} = B + \frac{1}{\mu} J z \bar{\sigma}$$


- $Z = \prod_i \sum_{\sigma_i} e^{\beta \mu \bar{B} \sigma_i} = \prod_i (e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}) = Z_1^N$,
where $Z_1 = e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}$. (Comparing with (7.8.1))
- Magnetic momentum $-m$ and magnetic field B are a pair of general force and general displacement (2.7.19), $\bar{m} = \frac{1}{\beta} \frac{\partial \ln Z}{\partial \bar{B}} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial \bar{B}} = N \mu \frac{e^{\beta \mu \bar{B}} - e^{-\beta \mu \bar{B}}}{e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}}$.
- On the other hand, $\bar{m} = N \mu \bar{\sigma}$.
- $\Rightarrow \bar{\sigma} = \frac{e^{\beta \mu \bar{B}} - e^{-\beta \mu \bar{B}}}{e^{\beta \mu \bar{B}} + e^{-\beta \mu \bar{B}}}$.
- Without external field, $\bar{B} = \frac{1}{\mu} J z \bar{\sigma}$. $\rightarrow \beta \mu \bar{B} = \beta J z \bar{\sigma}$.
 $\therefore Z_1 = e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}};$
 $\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}}.$

Mean field approximation

$$\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}}$$

solution

The critical temperature ($\beta J z = 1$): $T_c = \frac{Jz}{k}$.



Thermal capacity $Z_1 = e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}, \quad T_c = \frac{Jz}{k},$

$$\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}}, \quad Z = Z_1^N$$

- Internal energy: $U = -\frac{\partial \ln Z}{\partial \beta}$

Thermal capacity $Z_1 = e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}$, $T_c = \frac{Jz}{k}$,

$$\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}}, \quad Z = Z_1^N$$

- Internal energy: $U = -\frac{\partial \ln Z}{\partial \beta} = -N \frac{\partial \ln Z_1}{\partial \beta}$

Thermal capacity $Z_1 = e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}, \quad T_c = \frac{Jz}{k},$

$$\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}}, \quad Z = Z_1^N$$

- Internal energy:
$$U = -\frac{\partial \ln Z}{\partial \beta} = -N \frac{\partial \ln Z_1}{\partial \beta}$$

$$= -N J z \bar{\sigma} \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}}$$

Thermal capacity $Z_1 = e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}, \quad T_c = \frac{Jz}{k},$

$$\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}}, \quad Z = Z_1^N$$

- Internal energy:
$$U = -\frac{\partial \ln Z}{\partial \beta} = -N \frac{\partial \ln Z_1}{\partial \beta}$$

$$= -N J z \bar{\sigma} \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}} = -N J z \bar{\sigma}^2$$

Thermal capacity $Z_1 = e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}, \quad T_c = \frac{Jz}{k},$

$$\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}}, \quad Z = Z_1^N$$

- Internal energy: $U = -\frac{\partial \ln Z}{\partial \beta} = -N \frac{\partial \ln Z_1}{\partial \beta}$
 $= -N J z \bar{\sigma} \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}} = -N J z \bar{\sigma}^2 = -N k T_c \bar{\sigma}^2.$

Thermal capacity

$$Z_1 = e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}$$

$$T_c = \frac{Jz}{k}$$

$$\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}},$$

$$Z = Z_1^N$$

- Internal energy: $U = -\frac{\partial \ln Z}{\partial \beta} = -N \frac{\partial \ln Z_1}{\partial \beta}$

$$= -N J z \bar{\sigma} \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}} = -N J z \bar{\sigma}^2 = -N k T_c \bar{\sigma}^2.$$
- $\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}}$

Thermal capacity

$$Z_1 = e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}$$

$$T_c = \frac{Jz}{k},$$

$$\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}},$$

$$Z = Z_1^N$$

- Internal energy: $U = -\frac{\partial \ln Z}{\partial \beta} = -N \frac{\partial \ln Z_1}{\partial \beta}$
 $= -N J z \bar{\sigma} \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}} = -N J z \bar{\sigma}^2 = -N k T_c \bar{\sigma}^2.$
- $\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}} = \frac{e^{\frac{T_c}{T} \bar{\sigma}} - e^{-\frac{T_c}{T} \bar{\sigma}}}{e^{\frac{T_c}{T} \bar{\sigma}} + e^{-\frac{T_c}{T} \bar{\sigma}}}.$

Thermal capacity

$$Z_1 = e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}$$

$$T_c = \frac{Jz}{k}$$

$$\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}},$$

$$Z = Z_1^N$$

- Internal energy: $U = -\frac{\partial \ln Z}{\partial \beta} = -N \frac{\partial \ln Z_1}{\partial \beta}$
 $= -N J z \bar{\sigma} \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}} = -N J z \bar{\sigma}^2 = -N k T_c \bar{\sigma}^2.$
- $\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}} = \frac{e^{\frac{T_c}{T} \bar{\sigma}} - e^{-\frac{T_c}{T} \bar{\sigma}}}{e^{\frac{T_c}{T} \bar{\sigma}} + e^{-\frac{T_c}{T} \bar{\sigma}}}.$
 As $\bar{\sigma}(T \rightarrow T_c^-) \rightarrow 0$, $x \equiv \frac{T_c}{T} \bar{\sigma} \rightarrow 0$.

Thermal capacity

$$Z_1 = e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}$$

$$T_c = \frac{Jz}{k},$$

$$\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}},$$

$$Z = Z_1^N$$

- Internal energy: $U = -\frac{\partial \ln Z}{\partial \beta} = -N \frac{\partial \ln Z_1}{\partial \beta}$
 $= -N J z \bar{\sigma} \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}} = -N J z \bar{\sigma}^2 = -N k T_c \bar{\sigma}^2.$

- $\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}} = \frac{e^{\frac{T_c}{T} \bar{\sigma}} - e^{-\frac{T_c}{T} \bar{\sigma}}}{e^{\frac{T_c}{T} \bar{\sigma}} + e^{-\frac{T_c}{T} \bar{\sigma}}}.$

As $\bar{\sigma}(T \rightarrow T_c^-) \rightarrow 0$, $x \equiv \frac{T_c}{T} \bar{\sigma} \rightarrow 0$.

$$\bar{\sigma} = \frac{e^x - e^{-x}}{e^x + e^{-x}}$$

Thermal capacity

$$Z_1 = e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}$$

$$T_c = \frac{Jz}{k},$$

$$\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}},$$

$$Z = Z_1^N$$

- Internal energy: $U = -\frac{\partial \ln Z}{\partial \beta} = -N \frac{\partial \ln Z_1}{\partial \beta}$
 $= -N J z \bar{\sigma} \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}} = -N J z \bar{\sigma}^2 = -N k T_c \bar{\sigma}^2.$

- $\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}} = \frac{e^{\frac{T_c}{T} \bar{\sigma}} - e^{-\frac{T_c}{T} \bar{\sigma}}}{e^{\frac{T_c}{T} \bar{\sigma}} + e^{-\frac{T_c}{T} \bar{\sigma}}}.$

As $\bar{\sigma}(T \rightarrow T_c^-) \rightarrow 0$, $x \equiv \frac{T_c}{T} \bar{\sigma} \rightarrow 0$.

$$\bar{\sigma} = \frac{e^x - e^{-x}}{e^x + e^{-x}} \simeq \frac{(1+x+\frac{1}{2}x^2+\frac{1}{6}x^3)-(1-x+\frac{1}{2}x^2-\frac{1}{6}x^3)}{(1+x+\frac{1}{2}x^2+\frac{1}{6}x^3)+(1-x+\frac{1}{2}x^2-\frac{1}{6}x^3)}$$

Thermal capacity

$$Z_1 = e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}$$

$$T_c = \frac{Jz}{k},$$

$$\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}},$$

$$Z = Z_1^N$$

- Internal energy: $U = -\frac{\partial \ln Z}{\partial \beta} = -N \frac{\partial \ln Z_1}{\partial \beta}$
 $= -N J z \bar{\sigma} \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}} = -N J z \bar{\sigma}^2 = -N k T_c \bar{\sigma}^2.$

- $\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}} = \frac{e^{\frac{T_c}{T} \bar{\sigma}} - e^{-\frac{T_c}{T} \bar{\sigma}}}{e^{\frac{T_c}{T} \bar{\sigma}} + e^{-\frac{T_c}{T} \bar{\sigma}}}.$

As $\bar{\sigma}(T \rightarrow T_c^-) \rightarrow 0$, $x \equiv \frac{T_c}{T} \bar{\sigma} \rightarrow 0$.

$$\bar{\sigma} = \frac{e^x - e^{-x}}{e^x + e^{-x}} \simeq \frac{(1+x+\frac{1}{2}x^2+\frac{1}{6}x^3)-(1-x+\frac{1}{2}x^2-\frac{1}{6}x^3)}{(1+x+\frac{1}{2}x^2+\frac{1}{6}x^3)+(1-x+\frac{1}{2}x^2-\frac{1}{6}x^3)} = \frac{2x+\frac{1}{3}x^3}{2+x^2}$$

Thermal capacity

$$Z_1 = e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}$$

$$T_c = \frac{Jz}{k},$$

$$\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}},$$

$$Z = Z_1^N$$

- Internal energy: $U = -\frac{\partial \ln Z}{\partial \beta} = -N \frac{\partial \ln Z_1}{\partial \beta}$
 $= -N J z \bar{\sigma} \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}} = -N J z \bar{\sigma}^2 = -N k T_c \bar{\sigma}^2.$

- $\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}} = \frac{e^{\frac{T_c}{T} \bar{\sigma}} - e^{-\frac{T_c}{T} \bar{\sigma}}}{e^{\frac{T_c}{T} \bar{\sigma}} + e^{-\frac{T_c}{T} \bar{\sigma}}}.$

As $\bar{\sigma}(T \rightarrow T_c^-) \rightarrow 0$, $x \equiv \frac{T_c}{T} \bar{\sigma} \rightarrow 0$.

$$\begin{aligned} \bar{\sigma} &= \frac{e^x - e^{-x}}{e^x + e^{-x}} \simeq \frac{(1+x+\frac{1}{2}x^2+\frac{1}{6}x^3)-(1-x+\frac{1}{2}x^2-\frac{1}{6}x^3)}{(1+x+\frac{1}{2}x^2+\frac{1}{6}x^3)+(1-x+\frac{1}{2}x^2-\frac{1}{6}x^3)} = \frac{2x+\frac{1}{3}x^3}{2+x^2} \\ &= \frac{x+\frac{1}{6}x^3}{1+\frac{1}{2}x^2} \end{aligned}$$

Thermal capacity

$$Z_1 = e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}$$

$$T_c = \frac{Jz}{k},$$

$$\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}},$$

$$Z = Z_1^N$$

- Internal energy: $U = -\frac{\partial \ln Z}{\partial \beta} = -N \frac{\partial \ln Z_1}{\partial \beta}$
 $= -N J z \bar{\sigma} \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}} = -N J z \bar{\sigma}^2 = -N k T_c \bar{\sigma}^2.$

- $\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}} = \frac{e^{\frac{T_c}{T} \bar{\sigma}} - e^{-\frac{T_c}{T} \bar{\sigma}}}{e^{\frac{T_c}{T} \bar{\sigma}} + e^{-\frac{T_c}{T} \bar{\sigma}}}.$

As $\bar{\sigma}(T \rightarrow T_c^-) \rightarrow 0$, $x \equiv \frac{T_c}{T} \bar{\sigma} \rightarrow 0$.

$$\begin{aligned} \bar{\sigma} &= \frac{e^x - e^{-x}}{e^x + e^{-x}} \simeq \frac{(1+x+\frac{1}{2}x^2+\frac{1}{6}x^3)-(1-x+\frac{1}{2}x^2-\frac{1}{6}x^3)}{(1+x+\frac{1}{2}x^2+\frac{1}{6}x^3)+(1-x+\frac{1}{2}x^2-\frac{1}{6}x^3)} = \frac{2x+\frac{1}{3}x^3}{2+x^2} \\ &= \frac{x+\frac{1}{6}x^3}{1+\frac{1}{2}x^2} \simeq x(1+\frac{1}{6}x^2)(1-\frac{1}{2}x^2) \end{aligned}$$

Thermal capacity

$$Z_1 = e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}$$

$$T_c = \frac{Jz}{k},$$

$$\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}},$$

$$Z = Z_1^N$$

- Internal energy: $U = -\frac{\partial \ln Z}{\partial \beta} = -N \frac{\partial \ln Z_1}{\partial \beta}$
 $= -N J z \bar{\sigma} \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}} = -N J z \bar{\sigma}^2 = -N k T_c \bar{\sigma}^2.$

- $\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}} = \frac{e^{\frac{T_c}{T} \bar{\sigma}} - e^{-\frac{T_c}{T} \bar{\sigma}}}{e^{\frac{T_c}{T} \bar{\sigma}} + e^{-\frac{T_c}{T} \bar{\sigma}}}.$

As $\bar{\sigma}(T \rightarrow T_c^-) \rightarrow 0$, $x \equiv \frac{T_c}{T} \bar{\sigma} \rightarrow 0$.

$$\begin{aligned} \bar{\sigma} &= \frac{e^x - e^{-x}}{e^x + e^{-x}} \simeq \frac{(1+x+\frac{1}{2}x^2+\frac{1}{6}x^3)-(1-x+\frac{1}{2}x^2-\frac{1}{6}x^3)}{(1+x+\frac{1}{2}x^2+\frac{1}{6}x^3)+(1-x+\frac{1}{2}x^2-\frac{1}{6}x^3)} = \frac{2x+\frac{1}{3}x^3}{2+x^2} \\ &= \frac{x+\frac{1}{6}x^3}{1+\frac{1}{2}x^2} \simeq x(1+\frac{1}{6}x^2)(1-\frac{1}{2}x^2) \simeq x(1-\frac{1}{3}x^2). \end{aligned}$$

Thermal capacity

$$Z_1 = e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}$$

$$T_c = \frac{Jz}{k},$$

$$\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}},$$

$$Z = Z_1^N$$

- Internal energy: $U = -\frac{\partial \ln Z}{\partial \beta} = -N \frac{\partial \ln Z_1}{\partial \beta}$
 $= -N J z \bar{\sigma} \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}} = -N J z \bar{\sigma}^2 = -N k T_c \bar{\sigma}^2.$

- $\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}} = \frac{e^{\frac{T_c}{T} \bar{\sigma}} - e^{-\frac{T_c}{T} \bar{\sigma}}}{e^{\frac{T_c}{T} \bar{\sigma}} + e^{-\frac{T_c}{T} \bar{\sigma}}}.$

As $\bar{\sigma}(T \rightarrow T_c^-) \rightarrow 0$, $x \equiv \frac{T_c}{T} \bar{\sigma} \rightarrow 0$.

$$\bar{\sigma} = \frac{e^x - e^{-x}}{e^x + e^{-x}} \simeq \frac{(1+x+\frac{1}{2}x^2+\frac{1}{6}x^3)-(1-x+\frac{1}{2}x^2-\frac{1}{6}x^3)}{(1+x+\frac{1}{2}x^2+\frac{1}{6}x^3)+(1-x+\frac{1}{2}x^2-\frac{1}{6}x^3)} = \frac{2x+\frac{1}{3}x^3}{2+x^2}$$

$$= \frac{x+\frac{1}{6}x^3}{1+\frac{1}{2}x^2} \simeq x(1+\frac{1}{6}x^2)(1-\frac{1}{2}x^2) \simeq x(1-\frac{1}{3}x^2).$$

I.e., $\bar{\sigma} = \frac{T_c}{T} \bar{\sigma} [1 - \frac{1}{3}(\frac{T_c}{T} \bar{\sigma})^2].$

Thermal capacity

$$Z_1 = e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}, \quad T_c = \frac{Jz}{k},$$

$$\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}}, \quad Z = Z_1^N$$

- Internal energy: $U = -\frac{\partial \ln Z}{\partial \beta} = -N \frac{\partial \ln Z_1}{\partial \beta}$
 $= -N J z \bar{\sigma} \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}} = -N J z \bar{\sigma}^2 = -N k T_c \bar{\sigma}^2.$

- $\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}} = \frac{e^{\frac{T_c}{T} \bar{\sigma}} - e^{-\frac{T_c}{T} \bar{\sigma}}}{e^{\frac{T_c}{T} \bar{\sigma}} + e^{-\frac{T_c}{T} \bar{\sigma}}}.$

As $\bar{\sigma}(T \rightarrow T_c^-) \rightarrow 0$, $x \equiv \frac{T_c}{T} \bar{\sigma} \rightarrow 0$.

$$\bar{\sigma} = \frac{e^x - e^{-x}}{e^x + e^{-x}} \simeq \frac{(1+x+\frac{1}{2}x^2+\frac{1}{6}x^3)-(1-x+\frac{1}{2}x^2-\frac{1}{6}x^3)}{(1+x+\frac{1}{2}x^2+\frac{1}{6}x^3)+(1-x+\frac{1}{2}x^2-\frac{1}{6}x^3)} = \frac{2x+\frac{1}{3}x^3}{2+x^2}$$

$$= \frac{x+\frac{1}{6}x^3}{1+\frac{1}{2}x^2} \simeq x(1+\frac{1}{6}x^2)(1-\frac{1}{2}x^2) \simeq x(1-\frac{1}{3}x^2).$$

I.e., $\bar{\sigma} = \frac{T_c}{T} \bar{\sigma} [1 - \frac{1}{3}(\frac{T_c}{T} \bar{\sigma})^2]. \Rightarrow \frac{T}{T_c} = 1 - \frac{1}{3}(\frac{T_c}{T} \bar{\sigma})^2$

Thermal capacity

$$Z_1 = e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}$$

$$T_c = \frac{Jz}{k}$$

$$\bar{\sigma} = \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}},$$

$$Z = Z_1^N$$

- Internal energy: $U = -\frac{\partial \ln Z}{\partial \beta} = -N \frac{\partial \ln Z_1}{\partial \beta}$
 $= -N J z \bar{\sigma} \frac{e^{\beta J z \bar{\sigma}} - e^{-\beta J z \bar{\sigma}}}{e^{\beta J z \bar{\sigma}} + e^{-\beta J z \bar{\sigma}}} = -N J z \bar{\sigma}^2 = -N k T_c \bar{\sigma}^2.$

- $\bar{\sigma} = \frac{e^{\frac{T_c}{T} \bar{\sigma}} - e^{-\frac{T_c}{T} \bar{\sigma}}}{e^{\frac{T_c}{T} \bar{\sigma}} + e^{-\frac{T_c}{T} \bar{\sigma}}} = \frac{e^{\frac{T_c}{T} \bar{\sigma}} - e^{-\frac{T_c}{T} \bar{\sigma}}}{e^{\frac{T_c}{T} \bar{\sigma}} + e^{-\frac{T_c}{T} \bar{\sigma}}}.$

As $\bar{\sigma}(T \rightarrow T_c^-) \rightarrow 0$, $x \equiv \frac{T_c}{T} \bar{\sigma} \rightarrow 0$.

$$\bar{\sigma} = \frac{e^x - e^{-x}}{e^x + e^{-x}} \simeq \frac{(1+x+\frac{1}{2}x^2+\frac{1}{6}x^3)-(1-x+\frac{1}{2}x^2-\frac{1}{6}x^3)}{(1+x+\frac{1}{2}x^2+\frac{1}{6}x^3)+(1-x+\frac{1}{2}x^2-\frac{1}{6}x^3)} = \frac{2x+\frac{1}{3}x^3}{2+x^2}$$

$$= \frac{x+\frac{1}{6}x^3}{1+\frac{1}{2}x^2} \simeq x(1+\frac{1}{6}x^2)(1-\frac{1}{2}x^2) \simeq x(1-\frac{1}{3}x^2).$$

I.e., $\bar{\sigma} = \frac{T_c}{T} \bar{\sigma} [1 - \frac{1}{3}(\frac{T_c}{T} \bar{\sigma})^2] \Rightarrow \frac{T}{T_c} = 1 - \frac{1}{3}(\frac{T_c}{T} \bar{\sigma})^2$

$$\Rightarrow \bar{\sigma}^2 = \frac{1 - \frac{T}{T_c}}{\frac{1}{3}(\frac{T_c}{T})^2} = 3(\frac{T}{T_c})^2 \frac{T_c - T}{T_c}.$$

Thermal capacity

$$U = -NkT_c\bar{\sigma}^2, \quad \bar{\sigma}^2(T \rightarrow T_c^-) = 3\left(\frac{T}{T_c}\right)^2\frac{T_c-T}{T_c}$$

- As $T > T_c$, $\bar{\sigma} = 0$ (from the figure),

Thermal capacity

$$U = -NkT_c\bar{\sigma}^2, \quad \bar{\sigma}^2(T \rightarrow T_c^-) = 3\left(\frac{T}{T_c}\right)^2\frac{T_c-T}{T_c}$$

- As $T > T_c$, $\bar{\sigma} = 0$ (from the figure), $U = 0$,
 $C_B(B = 0, T \rightarrow T_c^+) = 0$.

Thermal capacity

$$U = -NkT_c\bar{\sigma}^2, \quad \bar{\sigma}^2(T \rightarrow T_c^-) = 3\left(\frac{T}{T_c}\right)^2\frac{T_c-T}{T_c}$$

- As $T > T_c$, $\bar{\sigma} = 0$ (from the figure), $U = 0$,
 $C_B(B = 0, T \rightarrow T_c^+) = 0$.
- $T \rightarrow T_c^-$, $U(T \rightarrow T_c^-) = -NkT_c \cdot 3\left(\frac{T}{T_c}\right)^2\frac{T_c-T}{T_c}$

Thermal capacity

$$U = -NkT_c\bar{\sigma}^2, \quad \bar{\sigma}^2(T \rightarrow T_c^-) = 3\left(\frac{T}{T_c}\right)^2\frac{T_c-T}{T_c}$$

- As $T > T_c$, $\bar{\sigma} = 0$ (from the figure), $U = 0$,
 $C_B(B = 0, T \rightarrow T_c^+) = 0$.
- $T \rightarrow T_c^-$, $U(T \rightarrow T_c^-) = -NkT_c \cdot 3\left(\frac{T}{T_c}\right)^2\frac{T_c-T}{T_c}$
 $= 3Nk\frac{T^2(T-T_c)}{T_c^2}$

Thermal capacity

$$U = -NkT_c\bar{\sigma}^2, \quad \bar{\sigma}^2(T \rightarrow T_c^-) = 3\left(\frac{T}{T_c}\right)^2\frac{T_c-T}{T_c}$$

- As $T > T_c$, $\bar{\sigma} = 0$ (from the figure), $U = 0$,
 $C_B(B = 0, T \rightarrow T_c^+) = 0$.
- $T \rightarrow T_c^-$, $U(T \rightarrow T_c^-) = -NkT_c \cdot 3\left(\frac{T}{T_c}\right)^2\frac{T_c-T}{T_c}$
 $= 3Nk\frac{T^2(T-T_c)}{T_c^2}$ (notice not suitable for non $T \rightarrow T_c^-$).

Thermal capacity

$$U = -NkT_c\bar{\sigma}^2, \quad \bar{\sigma}^2(T \rightarrow T_c^-) = 3\left(\frac{T}{T_c}\right)^2\frac{T_c-T}{T_c}$$

- As $T > T_c$, $\bar{\sigma} = 0$ (from the figure), $U = 0$,
 $C_B(B=0, T \rightarrow T_c^+) = 0$.
- $T \rightarrow T_c^-$, $U(T \rightarrow T_c^-) = -NkT_c \cdot 3\left(\frac{T}{T_c}\right)^2\frac{T_c-T}{T_c}$
 $= 3Nk\frac{T^2(T-T_c)}{T_c^2}$ (notice not suitable for non $T \rightarrow T_c^-$).
 $C_B(B=0, T \rightarrow T_c^-) = 3Nk$.

Thermal capacity

$$U = -NkT_c\bar{\sigma}^2, \quad \bar{\sigma}^2(T \rightarrow T_c^-) = 3\left(\frac{T}{T_c}\right)^2\frac{T_c-T}{T_c}$$

- As $T > T_c$, $\bar{\sigma} = 0$ (from the figure), $U = 0$,
 $C_B(B=0, T \rightarrow T_c^+) = 0$.
- $T \rightarrow T_c^-$, $U(T \rightarrow T_c^-) = -NkT_c \cdot 3\left(\frac{T}{T_c}\right)^2\frac{T_c-T}{T_c}$
 $= 3Nk\frac{T^2(T-T_c)}{T_c^2}$ (notice not suitable for non $T \rightarrow T_c^-$).
 $C_B(B=0, T \rightarrow T_c^-) = 3Nk$.
- Phase transition.

Table of contents

1 §9. Ensemble theory

- 9.1 Phase space and Liouville's theorem
- 9.2 The microcanonical ensemble
- 9.3 Thermodynamics in microcanonical ensemble theory
- 9.4 Canonical ensemble
- 9.5 Thermodynamics in canonical ensemble
- 9.6 EOS of real gas
- 9.7 Thermal capacity of solid
- 9.9 Ising model
- **9.10 Grand canonical ensemble**
- 9.11 Thermodynamics in grand canonical ensemble
- 9.12 Applications of the grand canonical ensemble

9.10 Grand canonical ensemble

9.10 Grand canonical ensemble

- N is also not fixed.

9.10 Grand canonical ensemble

- N is also not fixed.
- System and the reservoir have particles and energy exchange.

9.10 Grand canonical ensemble

- N is also not fixed.
- System and the reservoir have particles and energy exchange.
- The combined system: $E + E_r = E^{(0)}$,
 $N + N_r = N^{(0)}$.

9.10 Grand canonical ensemble

- N is also not fixed.
- System and the reservoir have particles and energy exchange.
- The combined system: $E + E_r = E^{(0)}$,
 $N + N_r = N^{(0)}$.
- For a given micro-state of the system N_s , number of micro-states of the reservoir: $\Omega_r(N^{(0)} - N, E^{(0)} - E_s)$.

9.10 Grand canonical ensemble

- N is also not fixed.
- System and the reservoir have particles and energy exchange.
- The combined system: $E + E_r = E^{(0)}$,
 $N + N_r = N^{(0)}$.
- For a given micro-state of the system N_s , number of micro-states of the reservoir: $\Omega_r(N^{(0)} - N, E^{(0)} - E_s)$.
- Using the equal probability principle, $\rho_{N_s} \propto \Omega_r$.

9.10 Grand canonical ensemble

- N is also not fixed.
- System and the reservoir have particles and energy exchange.
- The combined system: $E + E_r = E^{(0)}$,
 $N + N_r = N^{(0)}$.
- For a given micro-state of the system N_s , number of micro-states of the reservoir: $\Omega_r(N^{(0)} - N, E^{(0)} - E_s)$.
- Using the equal probability principle, $\rho_{N_s} \propto \Omega_r$.
- Considering $N \ll N^{(0)}$, $E \ll E^{(0)}$,

9.10 Grand canonical ensemble

- N is also not fixed.
- System and the reservoir have particles and energy exchange.
- The combined system: $E + E_r = E^{(0)}$,
 $N + N_r = N^{(0)}$.
- For a given micro-state of the system N_s , number of micro-states of the reservoir: $\Omega_r(N^{(0)} - N, E^{(0)} - E_s)$.
- Using the equal probability principle, $\rho_{N_s} \propto \Omega_r$.
- Considering $N \ll N^{(0)}$, $E \ll E^{(0)}$,

$$\ln \Omega_r(N^{(0)} - N, E^{(0)} - E_s)$$

$$\simeq \ln \Omega_r(N^{(0)}, E^{(0)}) - \left. \frac{\partial \ln \Omega_r}{\partial N_r} \right|_{N_r=N^{(0)}} N - \left. \frac{\partial \ln \Omega_r}{\partial E_r} \right|_{E_r=E^{(0)}} E_s$$

9.10 Grand canonical ensemble

- N is also not fixed.
- System and the reservoir have particles and energy exchange.
- The combined system: $E + E_r = E^{(0)}$,
 $N + N_r = N^{(0)}$.
- For a given micro-state of the system N_s , number of micro-states of the reservoir: $\Omega_r(N^{(0)} - N, E^{(0)} - E_s)$.
- Using the equal probability principle, $\rho_{N_s} \propto \Omega_r$.
- Considering $N \ll N^{(0)}$, $E \ll E^{(0)}$,

$$\ln \Omega_r(N^{(0)} - N, E^{(0)} - E_s)$$

$$\simeq \ln \Omega_r(N^{(0)}, E^{(0)}) - \left. \frac{\partial \ln \Omega_r}{\partial N_r} \right|_{N_r=N^{(0)}} N - \left. \frac{\partial \ln \Omega_r}{\partial E_r} \right|_{E_r=E^{(0)}} E_s$$

$$= \ln \Omega_r(N^{(0)}, E^{(0)}) - \alpha N - \beta E_s.$$

Grand canonical ensemble

- $\alpha = \left. \frac{\partial \ln \Omega_r}{\partial N_r} \right|_{N_r=N^{(0)}} = -\frac{\mu}{kT},$

Grand canonical ensemble

- $\alpha = \left. \frac{\partial \ln \Omega_r}{\partial N_r} \right|_{N_r=N^{(0)}} = -\frac{\mu}{kT}, \quad \beta = \left. \frac{\partial \ln \Omega_r}{\partial E_r} \right|_{E_r=E^{(0)}} = \frac{1}{kT}.$

Grand canonical ensemble

- $\alpha = \left. \frac{\partial \ln \Omega_r}{\partial N_r} \right|_{N_r=N^{(0)}} = -\frac{\mu}{kT}, \quad \beta = \left. \frac{\partial \ln \Omega_r}{\partial E_r} \right|_{E_r=E^{(0)}} = \frac{1}{kT}.$
- $\rho_{N_s} \propto e^{\ln \Omega_r(N^{(0)}, E^{(0)}) - \alpha N - \beta E_s}$

Grand canonical ensemble

- $\alpha = \left. \frac{\partial \ln \Omega_r}{\partial N_r} \right|_{N_r=N^{(0)}} = -\frac{\mu}{kT}, \quad \beta = \left. \frac{\partial \ln \Omega_r}{\partial E_r} \right|_{E_r=E^{(0)}} = \frac{1}{kT}.$
- $\rho_{N_s} \propto e^{\ln \Omega_r(N^{(0)}, E^{(0)}) - \alpha N - \beta E_s} \propto e^{-\alpha N - \beta E_s}$

Grand canonical ensemble

- $\alpha = \left. \frac{\partial \ln \Omega_r}{\partial N_r} \right|_{N_r=N^{(0)}} = -\frac{\mu}{kT}, \quad \beta = \left. \frac{\partial \ln \Omega_r}{\partial E_r} \right|_{E_r=E^{(0)}} = \frac{1}{kT}.$
- $\rho_{N_s} \propto e^{\ln \Omega_r(N^{(0)}, E^{(0)}) - \alpha N - \beta E_s} \propto e^{-\alpha N - \beta E_s}$
 $= \frac{1}{\mathcal{Z}} e^{-\alpha N - \beta E_s}.$

Grand canonical ensemble

- $\alpha = \left. \frac{\partial \ln \Omega_r}{\partial N_r} \right|_{N_r=N^{(0)}} = -\frac{\mu}{kT}$, $\beta = \left. \frac{\partial \ln \Omega_r}{\partial E_r} \right|_{E_r=E^{(0)}} = \frac{1}{kT}$.
- $\rho_{Ns} \propto e^{\ln \Omega_r(N^{(0)}, E^{(0)}) - \alpha N - \beta E_s} \propto e^{-\alpha N - \beta E_s}$
 $= \frac{1}{\mathcal{Z}} e^{-\alpha N - \beta E_s}.$

The grand partition function: $\mathcal{Z} = \sum_{N=0}^{\infty} \sum_s e^{-\alpha N - \beta E_s}.$

Grand canonical ensemble

- $\alpha = \left. \frac{\partial \ln \Omega_r}{\partial N_r} \right|_{N_r=N^{(0)}} = -\frac{\mu}{kT}, \quad \beta = \left. \frac{\partial \ln \Omega_r}{\partial E_r} \right|_{E_r=E^{(0)}} = \frac{1}{kT}.$
- $\rho_{N_s} \propto e^{\ln \Omega_r(N^{(0)}, E^{(0)}) - \alpha N - \beta E_s} \propto e^{-\alpha N - \beta E_s}$
 $= \frac{1}{\mathcal{Z}} e^{-\alpha N - \beta E_s}.$

The grand partition function: $\mathcal{Z} = \sum_{N=0}^{\infty} \sum_s e^{-\alpha N - \beta E_s}.$

- Continuous case: $\rho_N(q, p) d\Omega = \frac{1}{N! h^{Nr}} \frac{e^{-\alpha N - \beta E(q, p)}}{\mathcal{Z}} d\Omega.$

Grand canonical ensemble

- $\alpha = \left. \frac{\partial \ln \Omega_r}{\partial N_r} \right|_{N_r=N^{(0)}} = -\frac{\mu}{kT}$, $\beta = \left. \frac{\partial \ln \Omega_r}{\partial E_r} \right|_{E_r=E^{(0)}} = \frac{1}{kT}$.
- $\rho_{N_s} \propto e^{\ln \Omega_r(N^{(0)}, E^{(0)}) - \alpha N - \beta E_s} \propto e^{-\alpha N - \beta E_s}$
 $= \frac{1}{\mathcal{Z}} e^{-\alpha N - \beta E_s}$.

The grand partition function: $\mathcal{Z} = \sum_{N=0}^{\infty} \sum_s e^{-\alpha N - \beta E_s}$.

- Continuous case: $\rho_N(q, p) d\Omega = \frac{1}{N! h^{Nr}} \frac{e^{-\alpha N - \beta E(q, p)}}{\mathcal{Z}} d\Omega$.
 Normalization: $\sum_N \int \rho_N d\Omega = 1$,

Grand canonical ensemble

- $\alpha = \left. \frac{\partial \ln \Omega_r}{\partial N_r} \right|_{N_r=N^{(0)}} = -\frac{\mu}{kT}$, $\beta = \left. \frac{\partial \ln \Omega_r}{\partial E_r} \right|_{E_r=E^{(0)}} = \frac{1}{kT}$.
- $\rho_{N_s} \propto e^{\ln \Omega_r(N^{(0)}, E^{(0)}) - \alpha N - \beta E_s} \propto e^{-\alpha N - \beta E_s}$
 $= \frac{1}{\mathcal{Z}} e^{-\alpha N - \beta E_s}$.

The grand partition function: $\mathcal{Z} = \sum_{N=0}^{\infty} \sum_s e^{-\alpha N - \beta E_s}$.

- Continuous case: $\rho_N(q, p) d\Omega = \frac{1}{N! h^{Nr}} \frac{e^{-\alpha N - \beta E(q, p)}}{\mathcal{Z}} d\Omega$.

Normalization: $\sum_N \int \rho_N d\Omega = 1$, i.e.,

$$\sum_N \frac{1}{N! h^{Nr}} \int \frac{e^{-\alpha N - \beta E(q, p)}}{\mathcal{Z}} d\Omega = 1,$$

Grand canonical ensemble

- $\alpha = \left. \frac{\partial \ln \Omega_r}{\partial N_r} \right|_{N_r=N^{(0)}} = -\frac{\mu}{kT}$, $\beta = \left. \frac{\partial \ln \Omega_r}{\partial E_r} \right|_{E_r=E^{(0)}} = \frac{1}{kT}$.
- $\rho_{N_s} \propto e^{\ln \Omega_r(N^{(0)}, E^{(0)}) - \alpha N - \beta E_s} \propto e^{-\alpha N - \beta E_s}$
 $= \frac{1}{\mathcal{Z}} e^{-\alpha N - \beta E_s}$.

The grand partition function: $\mathcal{Z} = \sum_{N=0}^{\infty} \sum_s e^{-\alpha N - \beta E_s}$.

- Continuous case: $\rho_N(q, p) d\Omega = \frac{1}{N! h^{Nr}} \frac{e^{-\alpha N - \beta E(q, p)}}{\mathcal{Z}} d\Omega$.

Normalization: $\sum_N \int \rho_N d\Omega = 1$, i.e.,

$$\sum_N \frac{1}{N! h^{Nr}} \int \frac{e^{-\alpha N - \beta E(q, p)}}{\mathcal{Z}} d\Omega = 1,$$

$$\Rightarrow \mathcal{Z} = \sum_N \frac{e^{-\alpha N}}{N! h^{Nr}} \int e^{-\beta E(q, p)} d\Omega.$$

Table of contents

1 §9. Ensemble theory

- 9.1 Phase space and Liouville's theorem
- 9.2 The microcanonical ensemble
- 9.3 Thermodynamics in microcanonical ensemble theory
- 9.4 Canonical ensemble
- 9.5 Thermodynamics in canonical ensemble
- 9.6 EOS of real gas
- 9.7 Thermal capacity of solid
- 9.9 Ising model
- 9.10 Grand canonical ensemble
- 9.11 Thermodynamics in grand canonical ensemble
- 9.12 Applications of the grand canonical ensemble

9.11 Thermodynamics in grand canonical ensemble

9.11 Thermodynamics in grand canonical ensemble

- Average number of particles: $\overline{N} = \sum_N \sum_s N \rho_{Ns}$

9.11 Thermodynamics in grand canonical ensemble

- Average number of particles: $\bar{N} = \sum_N \sum_s N \rho_{Ns}$
 $= \frac{1}{\mathcal{Z}} \sum_N \sum_s N e^{-\alpha N - \beta E_s}$

9.11 Thermodynamics in grand canonical ensemble

- Average number of particles: $\bar{N} = \sum_N \sum_s N \rho_{Ns}$
$$= \frac{1}{\mathcal{Z}} \sum_N \sum_s N e^{-\alpha N - \beta E_s} = \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \alpha} \right) \sum_N \sum_s e^{-\alpha N - \beta E_s}$$

9.11 Thermodynamics in grand canonical ensemble

- Average number of particles:
$$\begin{aligned}\bar{N} &= \sum_N \sum_s N \rho_{Ns} \\ &= \frac{1}{\mathcal{Z}} \sum_N \sum_s N e^{-\alpha N - \beta E_s} = \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \alpha} \right) \sum_N \sum_s e^{-\alpha N - \beta E_s} \\ &= \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \alpha} \right) \mathcal{Z}\end{aligned}$$

9.11 Thermodynamics in grand canonical ensemble

- Average number of particles: $\bar{N} = \sum_N \sum_s N \rho_{Ns}$
$$= \frac{1}{\mathcal{Z}} \sum_N \sum_s N e^{-\alpha N - \beta E_s} = \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \alpha} \right) \sum_N \sum_s e^{-\alpha N - \beta E_s}$$
$$= \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \alpha} \right) \mathcal{Z} = -\frac{\partial \ln \mathcal{Z}}{\partial \alpha}.$$

9.11 Thermodynamics in grand canonical ensemble

- Average number of particles: $\bar{N} = \sum_N \sum_s N \rho_{Ns}$
$$= \frac{1}{\mathcal{Z}} \sum_N \sum_s N e^{-\alpha N - \beta E_s} = \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \alpha} \right) \sum_N \sum_s e^{-\alpha N - \beta E_s}$$
$$= \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \alpha} \right) \mathcal{Z} = -\frac{\partial \ln \mathcal{Z}}{\partial \alpha}.$$
- Internal energy: $U = \bar{E}$

9.11 Thermodynamics in grand canonical ensemble

- Average number of particles: $\bar{N} = \sum_N \sum_s N \rho_{Ns}$
$$= \frac{1}{\mathcal{Z}} \sum_N \sum_s N e^{-\alpha N - \beta E_s} = \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \alpha} \right) \sum_N \sum_s e^{-\alpha N - \beta E_s}$$
$$= \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \alpha} \right) \mathcal{Z} = -\frac{\partial \ln \mathcal{Z}}{\partial \alpha}.$$
- Internal energy: $U = \bar{E} = \frac{1}{\mathcal{Z}} \sum_N \sum_s E_s e^{-\alpha N - \beta E_s}$

9.11 Thermodynamics in grand canonical ensemble

- Average number of particles: $\bar{N} = \sum_N \sum_s N \rho_{Ns}$
$$= \frac{1}{\mathcal{Z}} \sum_N \sum_s N e^{-\alpha N - \beta E_s} = \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \alpha} \right) \sum_N \sum_s e^{-\alpha N - \beta E_s}$$
$$= \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \alpha} \right) \mathcal{Z} = -\frac{\partial \ln \mathcal{Z}}{\partial \alpha}.$$
- Internal energy: $U = \bar{E} = \frac{1}{\mathcal{Z}} \sum_N \sum_s E_s e^{-\alpha N - \beta E_s}$
$$= \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \beta} \right) \sum_N \sum_s e^{-\alpha N - \beta E_s}$$

9.11 Thermodynamics in grand canonical ensemble

- Average number of particles: $\bar{N} = \sum_N \sum_s N \rho_{Ns}$

$$= \frac{1}{\mathcal{Z}} \sum_N \sum_s N e^{-\alpha N - \beta E_s} = \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \alpha} \right) \sum_N \sum_s e^{-\alpha N - \beta E_s}$$

$$= \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \alpha} \right) \mathcal{Z} = -\frac{\partial \ln \mathcal{Z}}{\partial \alpha}.$$
- Internal energy: $U = \bar{E} = \frac{1}{\mathcal{Z}} \sum_N \sum_s E_s e^{-\alpha N - \beta E_s}$

$$= \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \beta} \right) \sum_N \sum_s e^{-\alpha N - \beta E_s} = \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \beta} \right) \mathcal{Z}$$

9.11 Thermodynamics in grand canonical ensemble

- Average number of particles: $\bar{N} = \sum_N \sum_s N \rho_{Ns}$

$$= \frac{1}{\mathcal{Z}} \sum_N \sum_s N e^{-\alpha N - \beta E_s} = \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \alpha} \right) \sum_N \sum_s e^{-\alpha N - \beta E_s}$$

$$= \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \alpha} \right) \mathcal{Z} = -\frac{\partial \ln \mathcal{Z}}{\partial \alpha}.$$
- Internal energy: $U = \bar{E} = \frac{1}{\mathcal{Z}} \sum_N \sum_s E_s e^{-\alpha N - \beta E_s}$

$$= \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \beta} \right) \sum_N \sum_s e^{-\alpha N - \beta E_s} = \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \beta} \right) \mathcal{Z} = -\frac{\partial \ln \mathcal{Z}}{\partial \beta}.$$

9.11 Thermodynamics in grand canonical ensemble

- Average number of particles: $\bar{N} = \sum_N \sum_s N \rho_{Ns}$

$$= \frac{1}{\mathcal{Z}} \sum_N \sum_s N e^{-\alpha N - \beta E_s} = \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \alpha} \right) \sum_N \sum_s e^{-\alpha N - \beta E_s}$$

$$= \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \alpha} \right) \mathcal{Z} = -\frac{\partial \ln \mathcal{Z}}{\partial \alpha}.$$
- Internal energy: $U = \bar{E} = \frac{1}{\mathcal{Z}} \sum_N \sum_s E_s e^{-\alpha N - \beta E_s}$

$$= \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \beta} \right) \sum_N \sum_s e^{-\alpha N - \beta E_s} = \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \beta} \right) \mathcal{Z} = -\frac{\partial \ln \mathcal{Z}}{\partial \beta}.$$
- Generalized force: $Y = \frac{1}{\mathcal{Z}} \sum_N \sum_s \frac{\partial E_s}{\partial y} e^{-\alpha N - \beta E_s}$

9.11 Thermodynamics in grand canonical ensemble

- Average number of particles: $\bar{N} = \sum_N \sum_s N \rho_{Ns}$

$$= \frac{1}{\mathcal{Z}} \sum_N \sum_s N e^{-\alpha N - \beta E_s} = \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \alpha} \right) \sum_N \sum_s e^{-\alpha N - \beta E_s}$$

$$= \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \alpha} \right) \mathcal{Z} = -\frac{\partial \ln \mathcal{Z}}{\partial \alpha}.$$
- Internal energy: $U = \bar{E} = \frac{1}{\mathcal{Z}} \sum_N \sum_s E_s e^{-\alpha N - \beta E_s}$

$$= \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \beta} \right) \sum_N \sum_s e^{-\alpha N - \beta E_s} = \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \beta} \right) \mathcal{Z} = -\frac{\partial \ln \mathcal{Z}}{\partial \beta}.$$
- Generalized force: $Y = \frac{1}{\mathcal{Z}} \sum_N \sum_s \frac{\partial E_s}{\partial y} e^{-\alpha N - \beta E_s}$

$$= \frac{1}{\mathcal{Z}} \left(-\frac{1}{\beta} \frac{\partial}{\partial y} \right) \sum_N \sum_s e^{-\alpha N - \beta E_s}$$

9.11 Thermodynamics in grand canonical ensemble

- Average number of particles: $\bar{N} = \sum_N \sum_s N \rho_{Ns}$

$$= \frac{1}{\mathcal{Z}} \sum_N \sum_s N e^{-\alpha N - \beta E_s} = \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \alpha} \right) \sum_N \sum_s e^{-\alpha N - \beta E_s}$$

$$= \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \alpha} \right) \mathcal{Z} = -\frac{\partial \ln \mathcal{Z}}{\partial \alpha}.$$
- Internal energy: $U = \bar{E} = \frac{1}{\mathcal{Z}} \sum_N \sum_s E_s e^{-\alpha N - \beta E_s}$

$$= \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \beta} \right) \sum_N \sum_s e^{-\alpha N - \beta E_s} = \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \beta} \right) \mathcal{Z} = -\frac{\partial \ln \mathcal{Z}}{\partial \beta}.$$
- Generalized force: $Y = \frac{1}{\mathcal{Z}} \sum_N \sum_s \frac{\partial E_s}{\partial y} e^{-\alpha N - \beta E_s}$

$$= \frac{1}{\mathcal{Z}} \left(-\frac{1}{\beta} \frac{\partial}{\partial y} \right) \sum_N \sum_s e^{-\alpha N - \beta E_s} = \frac{1}{\mathcal{Z}} \left(-\frac{1}{\beta} \frac{\partial}{\partial y} \right) \mathcal{Z}$$

9.11 Thermodynamics in grand canonical ensemble

- Average number of particles: $\bar{N} = \sum_N \sum_s N \rho_{Ns}$

$$= \frac{1}{\mathcal{Z}} \sum_N \sum_s N e^{-\alpha N - \beta E_s} = \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \alpha} \right) \sum_N \sum_s e^{-\alpha N - \beta E_s}$$

$$= \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \alpha} \right) \mathcal{Z} = -\frac{\partial \ln \mathcal{Z}}{\partial \alpha}.$$
- Internal energy: $U = \bar{E} = \frac{1}{\mathcal{Z}} \sum_N \sum_s E_s e^{-\alpha N - \beta E_s}$

$$= \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \beta} \right) \sum_N \sum_s e^{-\alpha N - \beta E_s} = \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \beta} \right) \mathcal{Z} = -\frac{\partial \ln \mathcal{Z}}{\partial \beta}.$$
- Generalized force: $Y = \frac{1}{\mathcal{Z}} \sum_N \sum_s \frac{\partial E_s}{\partial y} e^{-\alpha N - \beta E_s}$

$$= \frac{1}{\mathcal{Z}} \left(-\frac{1}{\beta} \frac{\partial}{\partial y} \right) \sum_N \sum_s e^{-\alpha N - \beta E_s} = \frac{1}{\mathcal{Z}} \left(-\frac{1}{\beta} \frac{\partial}{\partial y} \right) \mathcal{Z} = -\frac{1}{\beta} \frac{\partial \ln \mathcal{Z}}{\partial y}.$$

9.11 Thermodynamics in grand canonical ensemble

- Average number of particles: $\bar{N} = \sum_N \sum_s N \rho_{Ns}$

$$= \frac{1}{\mathcal{Z}} \sum_N \sum_s N e^{-\alpha N - \beta E_s} = \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \alpha} \right) \sum_N \sum_s e^{-\alpha N - \beta E_s}$$

$$= \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \alpha} \right) \mathcal{Z} = -\frac{\partial \ln \mathcal{Z}}{\partial \alpha}.$$
- Internal energy: $U = \bar{E} = \frac{1}{\mathcal{Z}} \sum_N \sum_s E_s e^{-\alpha N - \beta E_s}$

$$= \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \beta} \right) \sum_N \sum_s e^{-\alpha N - \beta E_s} = \frac{1}{\mathcal{Z}} \left(-\frac{\partial}{\partial \beta} \right) \mathcal{Z} = -\frac{\partial \ln \mathcal{Z}}{\partial \beta}.$$
- Generalized force: $Y = \frac{1}{\mathcal{Z}} \sum_N \sum_s \frac{\partial E_s}{\partial y} e^{-\alpha N - \beta E_s}$

$$= \frac{1}{\mathcal{Z}} \left(-\frac{1}{\beta} \frac{\partial}{\partial y} \right) \sum_N \sum_s e^{-\alpha N - \beta E_s} = \frac{1}{\mathcal{Z}} \left(-\frac{1}{\beta} \frac{\partial}{\partial y} \right) \mathcal{Z} = -\frac{1}{\beta} \frac{\partial \ln \mathcal{Z}}{\partial y}.$$
- Entropy (same procedure as canonical ensemble):

$$S = k \left(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta} \right).$$

Thermodynamics in grand canonical ensemble

- Grand potential (characteristic function) (3.2.10):

$$J = F - \mu N$$

Thermodynamics in grand canonical ensemble

- Grand potential (characteristic function) (3.2.10):

$$\begin{aligned} J &= F - \mu N = U - TS - \mu N \\ &= -\frac{\partial \ln \mathcal{Z}}{\partial \beta} - kT \left(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta} \right) - \mu N \end{aligned}$$

Thermodynamics in grand canonical ensemble

- Grand potential (characteristic function) (3.2.10):

$$\begin{aligned} J &= F - \mu N = U - TS - \mu N \\ &= -\frac{\partial \ln \mathcal{Z}}{\partial \beta} - kT \left(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta} \right) - \mu N \\ &= -kT \left(\ln \mathcal{Z} - \frac{\mu}{kT} \overline{N} \right) - \mu N \end{aligned}$$

Thermodynamics in grand canonical ensemble

- Grand potential (characteristic function) (3.2.10):

$$\begin{aligned} J &= F - \mu N = U - TS - \mu N \\ &= -\frac{\partial \ln \mathcal{Z}}{\partial \beta} - kT \left(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta} \right) - \mu N \\ &= -kT \left(\ln \mathcal{Z} - \frac{\mu}{kT} \overline{N} \right) - \mu N = -kT \ln \mathcal{Z}. \end{aligned}$$

Thermodynamics in grand canonical ensemble

- Grand potential (characteristic function) (3.2.10):

$$\begin{aligned} J &= F - \mu N = U - TS - \mu N \\ &= -\frac{\partial \ln \mathcal{Z}}{\partial \beta} - kT(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}) - \mu N \\ &= -kT(\ln \mathcal{Z} - \frac{\mu}{kT} \overline{N}) - \mu N = -kT \ln \mathcal{Z}. \end{aligned}$$

- Fluctuation of particle number:

$$\overline{(N - \overline{N})^2}$$

Thermodynamics in grand canonical ensemble

- Grand potential (characteristic function) (3.2.10):

$$\begin{aligned} J &= F - \mu N = U - TS - \mu N \\ &= -\frac{\partial \ln \mathcal{Z}}{\partial \beta} - kT \left(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta} \right) - \mu N \\ &= -kT \left(\ln \mathcal{Z} - \frac{\mu}{kT} \bar{N} \right) - \mu N = -kT \ln \mathcal{Z}. \end{aligned}$$

- Fluctuation of particle number:

$$\overline{(N - \bar{N})^2} = \overline{N^2} - 2N\bar{N} + \bar{N}^2$$

Thermodynamics in grand canonical ensemble

- Grand potential (characteristic function) (3.2.10):

$$\begin{aligned} J &= F - \mu N = U - TS - \mu N \\ &= -\frac{\partial \ln \mathcal{Z}}{\partial \beta} - kT(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}) - \mu N \\ &= -kT(\ln \mathcal{Z} - \frac{\mu}{kT} \overline{N}) - \mu N = -kT \ln \mathcal{Z}. \end{aligned}$$

- Fluctuation of particle number:

$$\overline{(N - \overline{N})^2} = \overline{N^2 - 2N\overline{N} + \overline{N}^2} = \overline{N^2} - \overline{N}^2;$$

Thermodynamics in grand canonical ensemble

- Grand potential (characteristic function) (3.2.10):

$$\begin{aligned}
 J &= F - \mu N = U - TS - \mu N \\
 &= -\frac{\partial \ln \mathcal{Z}}{\partial \beta} - kT(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}) - \mu N \\
 &= -kT(\ln \mathcal{Z} - \frac{\mu}{kT} \overline{N}) - \mu N = -kT \ln \mathcal{Z}.
 \end{aligned}$$

- Fluctuation of particle number:

$$\overline{(N - \overline{N})^2} = \overline{N^2 - 2N\overline{N} + \overline{N}^2} = \overline{N^2} - \overline{N}^2;$$

- $\frac{\partial \overline{N}}{\partial \alpha}$

Thermodynamics in grand canonical ensemble

- Grand potential (characteristic function) (3.2.10):

$$\begin{aligned}
 J &= F - \mu N = U - TS - \mu N \\
 &= -\frac{\partial \ln \mathcal{Z}}{\partial \beta} - kT(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}) - \mu N \\
 &= -kT(\ln \mathcal{Z} - \frac{\mu}{kT} \bar{N}) - \mu N = -kT \ln \mathcal{Z}.
 \end{aligned}$$

- Fluctuation of particle number:

$$\overline{(N - \bar{N})^2} = \overline{N^2 - 2N\bar{N} + \bar{N}^2} = \overline{N^2} - \bar{N}^2;$$

- $$\frac{\partial \bar{N}}{\partial \alpha} = \frac{\partial}{\partial \alpha} \left(\frac{1}{\mathcal{Z}} \sum_N \sum_s N e^{-\alpha N - \beta E_s} \right)$$

Thermodynamics in grand canonical ensemble

- Grand potential (characteristic function) (3.2.10):

$$\begin{aligned}
 J &= F - \mu N = U - TS - \mu N \\
 &= -\frac{\partial \ln \mathcal{Z}}{\partial \beta} - kT(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}) - \mu N \\
 &= -kT(\ln \mathcal{Z} - \frac{\mu}{kT} \bar{N}) - \mu N = -kT \ln \mathcal{Z}.
 \end{aligned}$$

- Fluctuation of particle number:

$$\begin{aligned}
 \overline{(N - \bar{N})^2} &= \overline{N^2 - 2N\bar{N} + \bar{N}^2} = \overline{N^2} - \bar{N}^2; \\
 \frac{\partial \bar{N}}{\partial \alpha} &= \frac{\partial}{\partial \alpha} \left(\frac{1}{\mathcal{Z}} \sum_N \sum_s N e^{-\alpha N - \beta E_s} \right) = \frac{\partial}{\partial \alpha} \frac{\sum_N \sum_s N e^{-\alpha N - \beta E_s}}{\sum_N \sum_s e^{-\alpha N - \beta E_s}}
 \end{aligned}$$

Thermodynamics in grand canonical ensemble

- Grand potential (characteristic function) (3.2.10):

$$\begin{aligned}
 J &= F - \mu N = U - TS - \mu N \\
 &= -\frac{\partial \ln \mathcal{Z}}{\partial \beta} - kT(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}) - \mu N \\
 &= -kT(\ln \mathcal{Z} - \frac{\mu}{kT} \bar{N}) - \mu N = -kT \ln \mathcal{Z}.
 \end{aligned}$$

- Fluctuation of particle number:

$$\begin{aligned}
 \overline{(N - \bar{N})^2} &= \overline{N^2 - 2N\bar{N} + \bar{N}^2} = \overline{N^2} - \bar{N}^2; \\
 \bullet \quad \frac{\partial \bar{N}}{\partial \alpha} &= \frac{\partial}{\partial \alpha} \left(\frac{1}{\mathcal{Z}} \sum_N \sum_s N e^{-\alpha N - \beta E_s} \right) = \frac{\partial}{\partial \alpha} \frac{\sum_N \sum_s N e^{-\alpha N - \beta E_s}}{\sum_N \sum_s e^{-\alpha N - \beta E_s}} \\
 &= -\frac{\sum_N \sum_s N^2 e^{-\alpha N - \beta E_s}}{\sum_N \sum_s e^{-\alpha N - \beta E_s}} + \frac{(\sum_N \sum_s N e^{-\alpha N - \beta E_s})^2}{(\sum_N \sum_s e^{-\alpha N - \beta E_s})^2}
 \end{aligned}$$

Thermodynamics in grand canonical ensemble

- Grand potential (characteristic function) (3.2.10):

$$\begin{aligned}
 J &= F - \mu N = U - TS - \mu N \\
 &= -\frac{\partial \ln \mathcal{Z}}{\partial \beta} - kT(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}) - \mu N \\
 &= -kT(\ln \mathcal{Z} - \frac{\mu}{kT} \bar{N}) - \mu N = -kT \ln \mathcal{Z}.
 \end{aligned}$$

- Fluctuation of particle number:

$$\overline{(N - \bar{N})^2} = \overline{N^2 - 2N\bar{N} + \bar{N}^2} = \overline{N^2} - \bar{N}^2;$$

$$\begin{aligned}
 \bullet \quad \frac{\partial \bar{N}}{\partial \alpha} &= \frac{\partial}{\partial \alpha} \left(\frac{1}{\mathcal{Z}} \sum_N \sum_s N e^{-\alpha N - \beta E_s} \right) = \frac{\partial}{\partial \alpha} \frac{\sum_N \sum_s N e^{-\alpha N - \beta E_s}}{\sum_N \sum_s e^{-\alpha N - \beta E_s}} \\
 &= -\frac{\sum_N \sum_s N^2 e^{-\alpha N - \beta E_s}}{\sum_N \sum_s e^{-\alpha N - \beta E_s}} + \frac{(\sum_N \sum_s N e^{-\alpha N - \beta E_s})^2}{(\sum_N \sum_s e^{-\alpha N - \beta E_s})^2} = -\overline{N^2} + \bar{N}^2.
 \end{aligned}$$

Thermodynamics in grand canonical ensemble

- Grand potential (characteristic function) (3.2.10):

$$\begin{aligned}
 J &= F - \mu N = U - TS - \mu N \\
 &= -\frac{\partial \ln \mathcal{Z}}{\partial \beta} - kT(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}) - \mu N \\
 &= -kT(\ln \mathcal{Z} - \frac{\mu}{kT} \bar{N}) - \mu N = -kT \ln \mathcal{Z}.
 \end{aligned}$$

- Fluctuation of particle number:

$$\overline{(N - \bar{N})^2} = \overline{N^2 - 2N\bar{N} + \bar{N}^2} = \overline{N^2} - \bar{N}^2;$$

$$\begin{aligned}
 \bullet \quad \frac{\partial \bar{N}}{\partial \alpha} &= \frac{\partial}{\partial \alpha} \left(\frac{1}{\mathcal{Z}} \sum_N \sum_s N e^{-\alpha N - \beta E_s} \right) = \frac{\partial}{\partial \alpha} \frac{\sum_N \sum_s N e^{-\alpha N - \beta E_s}}{\sum_N \sum_s e^{-\alpha N - \beta E_s}} \\
 &= -\frac{\sum_N \sum_s N^2 e^{-\alpha N - \beta E_s}}{\sum_N \sum_s e^{-\alpha N - \beta E_s}} + \frac{(\sum_N \sum_s N e^{-\alpha N - \beta E_s})^2}{(\sum_N \sum_s e^{-\alpha N - \beta E_s})^2} = -\overline{N^2} + \bar{N}^2.
 \end{aligned}$$

$$\bullet \Rightarrow \overline{(N - \bar{N})^2} = -\left(\frac{\partial \bar{N}}{\partial \alpha}\right)_{\beta, \mu}$$

Thermodynamics in grand canonical ensemble

- Grand potential (characteristic function) (3.2.10):

$$\begin{aligned}
 J &= F - \mu N = U - TS - \mu N \\
 &= -\frac{\partial \ln \mathcal{Z}}{\partial \beta} - kT(\ln \mathcal{Z} - \alpha \frac{\partial \ln \mathcal{Z}}{\partial \alpha} - \beta \frac{\partial \ln \mathcal{Z}}{\partial \beta}) - \mu N \\
 &= -kT(\ln \mathcal{Z} - \frac{\mu}{kT} \bar{N}) - \mu N = -kT \ln \mathcal{Z}.
 \end{aligned}$$

- Fluctuation of particle number:

$$\overline{(N - \bar{N})^2} = \overline{N^2 - 2N\bar{N} + \bar{N}^2} = \overline{N^2} - \bar{N}^2;$$

$$\begin{aligned}
 \bullet \quad \frac{\partial \bar{N}}{\partial \alpha} &= \frac{\partial}{\partial \alpha} \left(\frac{1}{\mathcal{Z}} \sum_N \sum_s N e^{-\alpha N - \beta E_s} \right) = \frac{\partial}{\partial \alpha} \frac{\sum_N \sum_s N e^{-\alpha N - \beta E_s}}{\sum_N \sum_s e^{-\alpha N - \beta E_s}} \\
 &= -\frac{\sum_N \sum_s N^2 e^{-\alpha N - \beta E_s}}{\sum_N \sum_s e^{-\alpha N - \beta E_s}} + \frac{(\sum_N \sum_s N e^{-\alpha N - \beta E_s})^2}{(\sum_N \sum_s e^{-\alpha N - \beta E_s})^2} = -\overline{N^2} + \bar{N}^2.
 \end{aligned}$$

$$\bullet \Rightarrow \overline{(N - \bar{N})^2} = -\left(\frac{\partial \bar{N}}{\partial \alpha}\right)_{\beta, \gamma} = kT \left(\frac{\partial \bar{N}}{\partial \mu}\right)_{T, V}.$$

Fluctuation of particle number

$$\overline{(N - \bar{N})^2} = kT \left(\frac{\partial \bar{N}}{\partial \mu} \right)_{T,V}$$

- (3.2.2) $dG = -SdT + Vdp + \mu d\bar{N},$

Fluctuation of particle number

$$\overline{(N - \bar{N})^2} = kT \left(\frac{\partial \bar{N}}{\partial \mu} \right)_{T,V}$$

- (3.2.2) $dG = -SdT + Vdp + \mu d\bar{N}$,
i.e., $d(\bar{N}\mu) = -(\bar{N}s)dT + (\bar{N}v)dp + \mu d\bar{N}$

Fluctuation of particle number

$$\overline{(N - \bar{N})^2} = kT \left(\frac{\partial \bar{N}}{\partial \mu} \right)_{T,V}$$

- (3.2.2) $dG = -SdT + Vdp + \mu d\bar{N}$,
i.e., $d(\bar{N}\mu) = -(\bar{N}s)dT + (\bar{N}v)dp + \mu d\bar{N}$
 $\Rightarrow \bar{N}d\mu = -\bar{N}s dT + \bar{N}v dp$

Fluctuation of particle number

$$\overline{(N - \bar{N})^2} = kT \left(\frac{\partial \bar{N}}{\partial \mu} \right)_{T,V}$$

- (3.2.2) $dG = -SdT + Vdp + \mu d\bar{N}$,
i.e., $d(\bar{N}\mu) = -(\bar{N}s)dT + (\bar{N}v)dp + \mu d\bar{N}$
 $\Rightarrow \bar{N}d\mu = -\bar{N}s dT + \bar{N}v dp$
 $\Rightarrow d\mu = v dp - s dT$

Fluctuation of particle number

$$\overline{(N - \bar{N})^2} = kT \left(\frac{\partial \bar{N}}{\partial \mu} \right)_{T,V}$$

- (3.2.2) $dG = -SdT + Vdp + \mu d\bar{N}$,
i.e., $d(\bar{N}\mu) = -(\bar{N}s)dT + (\bar{N}v)dp + \mu d\bar{N}$
 $\Rightarrow \bar{N}d\mu = -\bar{N}s dT + \bar{N}v dp$
 $\Rightarrow d\mu = v dp - s dT$
 $\Rightarrow \left(\frac{\partial \mu}{\partial v} \right)_T = v \left(\frac{\partial p}{\partial v} \right)_T,$

Fluctuation of particle number

$$\overline{(N - \bar{N})^2} = kT \left(\frac{\partial \bar{N}}{\partial \mu} \right)_{T,V}$$

- (3.2.2) $dG = -SdT + Vdp + \mu d\bar{N}$,
i.e., $d(\bar{N}\mu) = -(\bar{N}s)dT + (\bar{N}v)dp + \mu d\bar{N}$
 $\Rightarrow \bar{N}d\mu = -\bar{N}s dT + \bar{N}v dp$
 $\Rightarrow d\mu = v dp - s dT$
 $\Rightarrow \left(\frac{\partial \mu}{\partial v} \right)_T = v \left(\frac{\partial p}{\partial v} \right)_T$, and
 $\left(\frac{\partial \mu}{\partial v} \right)_T = \left(\frac{\partial \mu}{\partial \frac{V}{\bar{N}}} \right)_T = -\frac{\bar{N}^2}{V} \left(\frac{\partial \mu}{\partial \bar{N}} \right)_{T,V}$

Fluctuation of particle number

$$\overline{(N - \bar{N})^2} = kT \left(\frac{\partial \bar{N}}{\partial \mu} \right)_{T,V}$$

- $(3.2.2) \quad dG = -SdT + Vdp + \mu d\bar{N},$
 i.e., $d(\bar{N}\mu) = -(\bar{N}s)dT + (\bar{N}v)dp + \mu d\bar{N}$
 $\Rightarrow \bar{N}d\mu = -\bar{N}s dT + \bar{N}v dp$
 $\Rightarrow d\mu = v dp - s dT$
 $\Rightarrow \left(\frac{\partial \mu}{\partial v} \right)_T = v \left(\frac{\partial p}{\partial v} \right)_T, \text{ and}$
 $\left(\frac{\partial \mu}{\partial v} \right)_T = \left(\frac{\partial \mu}{\partial \frac{V}{\bar{N}}} \right)_T = -\frac{\bar{N}^2}{V} \left(\frac{\partial \mu}{\partial \bar{N}} \right)_{T,V}$
 $\Rightarrow \left(\frac{\partial \bar{N}}{\partial \mu} \right)_{T,V} = -\frac{\bar{N}^2}{V} \frac{1}{v} \left(\frac{\partial v}{\partial p} \right)_T.$

Fluctuation of particle number

$$\overline{(N - \bar{N})^2} = kT \left(\frac{\partial \bar{N}}{\partial \mu} \right)_{T,V}$$

- $(3.2.2) \quad dG = -SdT + Vdp + \mu d\bar{N},$
 i.e., $d(\bar{N}\mu) = -(\bar{N}s)dT + (\bar{N}v)dp + \mu d\bar{N}$
 $\Rightarrow \bar{N}d\mu = -\bar{N}s dT + \bar{N}v dp$
 $\Rightarrow d\mu = v dp - s dT$
 $\Rightarrow \left(\frac{\partial \mu}{\partial v} \right)_T = v \left(\frac{\partial p}{\partial v} \right)_T, \text{ and}$
 $\left(\frac{\partial \mu}{\partial v} \right)_T = \left(\frac{\partial \mu}{\partial \frac{V}{\bar{N}}} \right)_T = -\frac{\bar{N}^2}{V} \left(\frac{\partial \mu}{\partial \bar{N}} \right)_{T,V}$
 $\Rightarrow \left(\frac{\partial \bar{N}}{\partial \mu} \right)_{T,V} = -\frac{\bar{N}^2}{V} \frac{1}{v} \left(\frac{\partial v}{\partial p} \right)_T.$
- $$\frac{\overline{(N - \bar{N})^2}}{(\bar{N})^2}$$

Fluctuation of particle number

$$\overline{(N - \bar{N})^2} = kT \left(\frac{\partial \bar{N}}{\partial \mu} \right)_{T,V}$$

- $(3.2.2) \quad dG = -SdT + Vdp + \mu d\bar{N},$
 i.e., $d(\bar{N}\mu) = -(\bar{N}s)dT + (\bar{N}v)dp + \mu d\bar{N}$
 $\Rightarrow \bar{N}d\mu = -\bar{N}s dT + \bar{N}v dp$
 $\Rightarrow d\mu = v dp - s dT$
 $\Rightarrow \left(\frac{\partial \mu}{\partial v} \right)_T = v \left(\frac{\partial p}{\partial v} \right)_T, \text{ and}$
 $\left(\frac{\partial \mu}{\partial v} \right)_T = \left(\frac{\partial \mu}{\partial \frac{V}{\bar{N}}} \right)_T = -\frac{\bar{N}^2}{V} \left(\frac{\partial \mu}{\partial \bar{N}} \right)_{T,V}$
 $\Rightarrow \left(\frac{\partial \bar{N}}{\partial \mu} \right)_{T,V} = -\frac{\bar{N}^2}{V} \frac{1}{v} \left(\frac{\partial v}{\partial p} \right)_T.$
- $\frac{\overline{(N - \bar{N})^2}}{(\bar{N})^2} = \frac{kT}{(\bar{N})^2} \cdot \left(-\frac{\bar{N}^2}{V} \right) \cdot \frac{1}{v} \left(\frac{\partial v}{\partial p} \right)_T$

Fluctuation of particle number

$$\overline{(N - \bar{N})^2} = kT \left(\frac{\partial \bar{N}}{\partial \mu} \right)_{T,V}$$

- $(3.2.2) \quad dG = -SdT + Vdp + \mu d\bar{N},$
 i.e., $d(\bar{N}\mu) = -(\bar{N}s)dT + (\bar{N}v)dp + \mu d\bar{N}$
 $\Rightarrow \bar{N}d\mu = -\bar{N}s dT + \bar{N}v dp$
 $\Rightarrow d\mu = v dp - s dT$
 $\Rightarrow \left(\frac{\partial \mu}{\partial v} \right)_T = v \left(\frac{\partial p}{\partial v} \right)_T, \text{ and}$
 $\left(\frac{\partial \mu}{\partial v} \right)_T = \left(\frac{\partial \mu}{\partial \frac{V}{\bar{N}}} \right)_T = -\frac{\bar{N}^2}{V} \left(\frac{\partial \mu}{\partial \bar{N}} \right)_{T,V}$
 $\Rightarrow \left(\frac{\partial \bar{N}}{\partial \mu} \right)_{T,V} = -\frac{\bar{N}^2}{V} \frac{1}{v} \left(\frac{\partial v}{\partial p} \right)_T.$
- $\frac{\overline{(N - \bar{N})^2}}{(\bar{N})^2} = \frac{kT}{(\bar{N})^2} \cdot \left(-\frac{\bar{N}^2}{V} \right) \cdot \frac{1}{v} \left(\frac{\partial v}{\partial p} \right)_T = \frac{kT}{V} \kappa_T$

Fluctuation of particle number

$$\overline{(N - \bar{N})^2} = kT \left(\frac{\partial \bar{N}}{\partial \mu} \right)_{T,V}$$

- $(3.2.2) \quad dG = -SdT + Vdp + \mu d\bar{N},$
 i.e., $d(\bar{N}\mu) = -(\bar{N}s)dT + (\bar{N}v)dp + \mu d\bar{N}$
 $\Rightarrow \bar{N}d\mu = -\bar{N}s dT + \bar{N}v dp$
 $\Rightarrow d\mu = v dp - s dT$
 $\Rightarrow \left(\frac{\partial \mu}{\partial v} \right)_T = v \left(\frac{\partial p}{\partial v} \right)_T, \text{ and}$
 $\left(\frac{\partial \mu}{\partial v} \right)_T = \left(\frac{\partial \mu}{\partial \frac{V}{\bar{N}}} \right)_T = -\frac{\bar{N}^2}{V} \left(\frac{\partial \mu}{\partial \bar{N}} \right)_{T,V}$
 $\Rightarrow \left(\frac{\partial \bar{N}}{\partial \mu} \right)_{T,V} = -\frac{\bar{N}^2}{V} \frac{1}{v} \left(\frac{\partial v}{\partial p} \right)_T.$
- $\frac{\overline{(N - \bar{N})^2}}{(\bar{N})^2} = \frac{kT}{(\bar{N})^2} \cdot \left(-\frac{\bar{N}^2}{V} \right) \cdot \frac{1}{v} \left(\frac{\partial v}{\partial p} \right)_T = \frac{kT}{V} \kappa_T$
 $\propto \frac{1}{\bar{N}}, \text{ in general;}$

Fluctuation of particle number

$$\overline{(N - \bar{N})^2} = kT \left(\frac{\partial \bar{N}}{\partial \mu} \right)_{T,V}$$

- $(3.2.2) \quad dG = -SdT + Vdp + \mu d\bar{N},$
 i.e., $d(\bar{N}\mu) = -(\bar{N}s)dT + (\bar{N}v)dp + \mu d\bar{N}$
 $\Rightarrow \bar{N}d\mu = -\bar{N}s dT + \bar{N}v dp$
 $\Rightarrow d\mu = v dp - s dT$
 $\Rightarrow \left(\frac{\partial \mu}{\partial v} \right)_T = v \left(\frac{\partial p}{\partial v} \right)_T, \text{ and}$
 $\left(\frac{\partial \mu}{\partial v} \right)_T = \left(\frac{\partial \mu}{\partial \frac{V}{\bar{N}}} \right)_T = -\frac{\bar{N}^2}{V} \left(\frac{\partial \mu}{\partial \bar{N}} \right)_{T,V}$
 $\Rightarrow \left(\frac{\partial \bar{N}}{\partial \mu} \right)_{T,V} = -\frac{\bar{N}^2}{V} \frac{1}{v} \left(\frac{\partial v}{\partial p} \right)_T.$
- $\frac{\overline{(N - \bar{N})^2}}{(\bar{N})^2} = \frac{kT}{(\bar{N})^2} \cdot \left(-\frac{\bar{N}^2}{V} \right) \cdot \frac{1}{v} \left(\frac{\partial v}{\partial p} \right)_T = \frac{kT}{V} \kappa_T$
 $\propto \frac{1}{\bar{N}}, \text{ in general;}$
 $\rightarrow 1, \text{ when } \kappa_T \text{ is very large (phase transition).}$

Table of contents

1 §9. Ensemble theory

- 9.1 Phase space and Liouville's theorem
- 9.2 The microcanonical ensemble
- 9.3 Thermodynamics in microcanonical ensemble theory
- 9.4 Canonical ensemble
- 9.5 Thermodynamics in canonical ensemble
- 9.6 EOS of real gas
- 9.7 Thermal capacity of solid
- 9.9 Ising model
- 9.10 Grand canonical ensemble
- 9.11 Thermodynamics in grand canonical ensemble
- 9.12 Applications of the grand canonical ensemble

9.12 Applications of the grand canonical ensemble

9.12 Applications of the grand canonical ensemble

- App 1. Absorption.

9.12 Applications of the grand canonical ensemble

- App 1. Absorption.
- N_0 absorbing points, each one with energy $-\varepsilon_0$, require $\frac{N}{N_0}$ at temperature T .

9.12 Applications of the grand canonical ensemble

- App 1. Absorption.
- N_0 absorbing points, each one with energy $-\varepsilon_0$, require $\frac{N}{N_0}$ at temperature T .
- The system of absorbing points obeys grand canonical distribution.

9.12 Applications of the grand canonical ensemble

- App 1. Absorption.
- N_0 absorbing points, each one with energy $-\varepsilon_0$, require $\frac{N}{N_0}$ at temperature T .
- The system of absorbing points obeys grand canonical distribution.
- $$\mathcal{Z} = \sum_{N=0}^{N_0} \sum_s e^{-\alpha N - \beta E_s}$$

9.12 Applications of the grand canonical ensemble

- App 1. Absorption.
- N_0 absorbing points, each one with energy $-\varepsilon_0$, require $\frac{N}{N_0}$ at temperature T .
- The system of absorbing points obeys grand canonical distribution.

$$\bullet \mathcal{Z} = \sum_{N=0}^{N_0} \sum_s e^{-\alpha N - \beta E_s} = \sum_{N=0}^{N_0} \frac{N_0!}{N!(N_0-N)!} e^{-\alpha N + \beta \varepsilon_0 N},$$

9.12 Applications of the grand canonical ensemble

- App 1. Absorption.
- N_0 absorbing points, each one with energy $-\varepsilon_0$, require $\frac{N}{N_0}$ at temperature T .
- The system of absorbing points obeys grand canonical distribution.
- $$\mathcal{Z} = \sum_{N=0}^{N_0} \sum_s e^{-\alpha N - \beta E_s} = \sum_{N=0}^{N_0} \frac{N_0!}{N!(N_0-N)!} e^{-\alpha N + \beta \varepsilon_0 N},$$
- As $-\alpha N + \beta \varepsilon_0 N = \beta(\mu + \varepsilon_0)N$,

9.12 Applications of the grand canonical ensemble

- App 1. Absorption.
- N_0 absorbing points, each one with energy $-\varepsilon_0$, require $\frac{N}{N_0}$ at temperature T .
- The system of absorbing points obeys grand canonical distribution.

$$\bullet \mathcal{Z} = \sum_{N=0}^{N_0} \sum_s e^{-\alpha N - \beta E_s} = \sum_{N=0}^{N_0} \frac{N_0!}{N!(N_0-N)!} e^{-\alpha N + \beta \varepsilon_0 N},$$

- As $-\alpha N + \beta \varepsilon_0 N = \beta(\mu + \varepsilon_0)N$, and

$$\sum_{N=0}^{N_0} \frac{N_0!}{N!(N_0-N)!} x^N = (1+x)^{N_0} \quad (9.12.15),$$

9.12 Applications of the grand canonical ensemble

- App 1. Absorption.
- N_0 absorbing points, each one with energy $-\varepsilon_0$, require $\frac{N}{N_0}$ at temperature T .
- The system of absorbing points obeys grand canonical distribution.

$$\bullet \mathcal{Z} = \sum_{N=0}^{N_0} \sum_s e^{-\alpha N - \beta E_s} = \sum_{N=0}^{N_0} \frac{N_0!}{N!(N_0-N)!} e^{-\alpha N + \beta \varepsilon_0 N},$$

- As $-\alpha N + \beta \varepsilon_0 N = \beta(\mu + \varepsilon_0)N$, and

$$\sum_{N=0}^{N_0} \frac{N_0!}{N!(N_0-N)!} x^N = (1+x)^{N_0} \quad (9.12.15),$$

$$\mathcal{Z} = [1 + e^{\beta(\mu + \varepsilon_0)}]^{N_0}.$$

Absorption

$$\mathcal{Z} = [1 + e^{\beta(\mu + \varepsilon_0)}]^{N_0}$$

- $\bar{N} = -\frac{\partial \ln \mathcal{Z}}{\partial \alpha}$

Absorption

$$\mathcal{Z} = [1 + e^{\beta(\mu + \varepsilon_0)}]^{N_0}$$

- $\bullet \quad \overline{N} = -\frac{\partial \ln \mathcal{Z}}{\partial \alpha} = kT \frac{\partial \ln \mathcal{Z}}{\partial \mu}$

Absorption

$$\mathcal{Z} = [1 + e^{\beta(\mu + \varepsilon_0)}]^{N_0}$$

- $$\bar{N} = -\frac{\partial \ln \mathcal{Z}}{\partial \alpha} = kT \frac{\partial \ln \mathcal{Z}}{\partial \mu} = kT \frac{\partial \{N_0 \ln[1 + e^{\beta(\mu + \varepsilon_0)}]\}}{\partial \mu}$$

Absorption

$$\mathcal{Z} = [1 + e^{\beta(\mu + \varepsilon_0)}]^{N_0}$$

- $$\begin{aligned}\overline{N} &= -\frac{\partial \ln \mathcal{Z}}{\partial \alpha} = kT \frac{\partial \ln \mathcal{Z}}{\partial \mu} = kT \frac{\partial \{N_0 \ln[1 + e^{\beta(\mu + \varepsilon_0)}]\}}{\partial \mu} \\ &= \frac{N_0 e^{\beta(\mu + \varepsilon_0)}}{1 + e^{\beta(\mu + \varepsilon_0)}}.\end{aligned}$$

Absorption

$$\mathcal{Z} = [1 + e^{\beta(\mu + \varepsilon_0)}]^{N_0}$$

- $$\begin{aligned} \overline{N} &= -\frac{\partial \ln \mathcal{Z}}{\partial \alpha} = kT \frac{\partial \ln \mathcal{Z}}{\partial \mu} = kT \frac{\partial \{N_0 \ln[1 + e^{\beta(\mu + \varepsilon_0)}]\}}{\partial \mu} \\ &= \frac{N_0 e^{\beta(\mu + \varepsilon_0)}}{1 + e^{\beta(\mu + \varepsilon_0)}}. \end{aligned}$$
- If ideal gas, $e^{\frac{\mu}{kT}} = \frac{p}{kT} \left(\frac{h^2}{2\pi m kT} \right)^{\frac{3}{2}}$ (7.6.8),

Absorption

$$\mathcal{Z} = [1 + e^{\beta(\mu + \varepsilon_0)}]^{N_0}$$

- $$\begin{aligned} \bar{N} &= -\frac{\partial \ln \mathcal{Z}}{\partial \alpha} = kT \frac{\partial \ln \mathcal{Z}}{\partial \mu} = kT \frac{\partial \{N_0 \ln[1 + e^{\beta(\mu + \varepsilon_0)}]\}}{\partial \mu} \\ &= \frac{N_0 e^{\beta(\mu + \varepsilon_0)}}{1 + e^{\beta(\mu + \varepsilon_0)}}. \end{aligned}$$
- If ideal gas, $e^{\frac{\mu}{kT}} = \frac{p}{kT} \left(\frac{h^2}{2\pi m kT} \right)^{\frac{3}{2}}$ (7.6.8),

$$\frac{\bar{N}}{N_0} = \left[1 + \frac{kT}{p} \left(\frac{2\pi m kT}{h^2} \right)^{\frac{3}{2}} e^{-\frac{\varepsilon_0}{kT}} \right]^{-1}.$$

Absorption

$$\mathcal{Z} = [1 + e^{\beta(\mu + \varepsilon_0)}]^{N_0}$$

- $$\begin{aligned}\bar{N} &= -\frac{\partial \ln \mathcal{Z}}{\partial \alpha} = kT \frac{\partial \ln \mathcal{Z}}{\partial \mu} = kT \frac{\partial \{N_0 \ln[1 + e^{\beta(\mu + \varepsilon_0)}]\}}{\partial \mu} \\ &= \frac{N_0 e^{\beta(\mu + \varepsilon_0)}}{1 + e^{\beta(\mu + \varepsilon_0)}}.\end{aligned}$$
- If ideal gas, $e^{\frac{\mu}{kT}} = \frac{p}{kT} \left(\frac{h^2}{2\pi m kT} \right)^{\frac{3}{2}}$ (7.6.8),

$$\frac{\bar{N}}{N_0} = \left[1 + \frac{kT}{p} \left(\frac{2\pi m kT}{h^2} \right)^{\frac{3}{2}} e^{-\frac{\varepsilon_0}{kT}} \right]^{-1}.$$
- App 2. \bar{a}_l of nearly independent particles.

Absorption

$$\mathcal{Z} = [1 + e^{\beta(\mu + \varepsilon_0)}]^{N_0}$$

- $$\overline{N} = -\frac{\partial \ln \mathcal{Z}}{\partial \alpha} = kT \frac{\partial \ln \mathcal{Z}}{\partial \mu} = kT \frac{\partial \{N_0 \ln[1 + e^{\beta(\mu + \varepsilon_0)}]\}}{\partial \mu}$$

$$= \frac{N_0 e^{\beta(\mu + \varepsilon_0)}}{1 + e^{\beta(\mu + \varepsilon_0)}}.$$
- If ideal gas, $e^{\frac{\mu}{kT}} = \frac{p}{kT} \left(\frac{h^2}{2\pi m kT} \right)^{\frac{3}{2}}$ (7.6.8),

$$\frac{\overline{N}}{N_0} = \left[1 + \frac{kT}{p} \left(\frac{2\pi m kT}{h^2} \right)^{\frac{3}{2}} e^{-\frac{\varepsilon_0}{kT}} \right]^{-1}.$$
- App 2. \overline{a}_l of nearly independent particles.
- Grand canonical distribution: $\rho_{NS} = \frac{1}{\mathcal{Z}} e^{-\alpha N - \beta E_S}$,
where S is the micro-state of the **system**.

Absorption

$$\mathcal{Z} = [1 + e^{\beta(\mu + \varepsilon_0)}]^{N_0}$$

- $\bar{N} = -\frac{\partial \ln \mathcal{Z}}{\partial \alpha} = kT \frac{\partial \ln \mathcal{Z}}{\partial \mu} = kT \frac{\partial \{N_0 \ln[1 + e^{\beta(\mu + \varepsilon_0)}]\}}{\partial \mu}$
 $= \frac{N_0 e^{\beta(\mu + \varepsilon_0)}}{1 + e^{\beta(\mu + \varepsilon_0)}}.$
- If ideal gas, $e^{\frac{\mu}{kT}} = \frac{p}{kT} \left(\frac{h^2}{2\pi m kT} \right)^{\frac{3}{2}}$ (7.6.8),
 $\frac{\bar{N}}{N_0} = \left[1 + \frac{kT}{p} \left(\frac{2\pi m kT}{h^2} \right)^{\frac{3}{2}} e^{-\frac{\varepsilon_0}{kT}} \right]^{-1}.$
- App 2. \bar{a}_l of nearly independent particles.
- Grand canonical distribution: $\rho_{NS} = \frac{1}{\mathcal{Z}} e^{-\alpha N - \beta E_S}$,
 where S is the micro-state of the **system**.
- a_s represents the number of particles at the particle's state s , then

Absorption

$$\mathcal{Z} = [1 + e^{\beta(\mu + \varepsilon_0)}]^{N_0}$$

- $\bar{N} = -\frac{\partial \ln \mathcal{Z}}{\partial \alpha} = kT \frac{\partial \ln \mathcal{Z}}{\partial \mu} = kT \frac{\partial \{N_0 \ln[1 + e^{\beta(\mu + \varepsilon_0)}]\}}{\partial \mu}$
 $= \frac{N_0 e^{\beta(\mu + \varepsilon_0)}}{1 + e^{\beta(\mu + \varepsilon_0)}}.$
- If ideal gas, $e^{\frac{\mu}{kT}} = \frac{p}{kT} \left(\frac{h^2}{2\pi m kT} \right)^{\frac{3}{2}}$ (7.6.8),
 $\frac{\bar{N}}{N_0} = [1 + \frac{kT}{p} \left(\frac{2\pi m kT}{h^2} \right)^{\frac{3}{2}} e^{-\frac{\varepsilon_0}{kT}}]^{-1}.$
- App 2. \bar{a}_l of nearly independent particles.
- Grand canonical distribution: $\rho_{NS} = \frac{1}{\mathcal{Z}} e^{-\alpha N - \beta E_S}$,
 where S is the micro-state of the **system**.
- a_s represents the number of particles at the particle's state s , then

$$N = \sum_s a_s, \quad E_S = \sum_s \varepsilon_s a_s.$$

\bar{a}_l of nearly independent particles

- Grand partition function: $\mathcal{Z} = \sum_N \sum_S e^{-\alpha N - \beta E_S}$

\bar{a}_l of nearly independent particles

- Grand partition function: $\mathcal{Z} = \sum_N \sum_S e^{-\alpha N - \beta E_S}$
$$= \sum_{\{a_s\}} e^{-\sum_s (\alpha + \beta \varepsilon_s) a_s}$$

\bar{a}_l of nearly independent particles

- Grand partition function:
$$\mathcal{Z} = \sum_N \sum_S e^{-\alpha N - \beta E_S}$$
$$= \sum_{\{a_s\}} e^{-\sum_s (\alpha + \beta \varepsilon_s) a_s} = \sum_{\{a_s\}} \prod_s e^{-(\alpha + \beta \varepsilon_s) a_s}$$

\bar{a}_l of nearly independent particles

- Grand partition function:
$$\begin{aligned}\mathcal{Z} &= \sum_N \sum_S e^{-\alpha N - \beta E_S} \\ &= \sum_{\{a_s\}} e^{-\sum_s (\alpha + \beta \varepsilon_s) a_s} = \sum_{\{a_s\}} \prod_s e^{-(\alpha + \beta \varepsilon_s) a_s} \\ &= \prod_s \sum_{a_s} e^{-(\alpha + \beta \varepsilon_s) a_s}.\end{aligned}$$

\bar{a}_l of nearly independent particles

- Grand partition function: $\mathcal{Z} = \sum_N \sum_S e^{-\alpha N - \beta E_S}$

$$= \sum_{\{a_s\}} e^{-\sum_s (\alpha + \beta \varepsilon_s) a_s} = \sum_{\{a_s\}} \prod_s e^{-(\alpha + \beta \varepsilon_s) a_s}$$

$$= \prod_s \sum_{a_s} e^{-(\alpha + \beta \varepsilon_s) a_s}.$$
- Set $\mathcal{Z}_s = \sum_{a_s} e^{-(\alpha + \beta \varepsilon_s) a_s}$, then $\mathcal{Z} = \prod_s \mathcal{Z}_s$.

\bar{a}_l of nearly independent particles

- Grand partition function: $\mathcal{Z} = \sum_N \sum_S e^{-\alpha N - \beta E_S}$

$$= \sum_{\{a_s\}} e^{-\sum_s (\alpha + \beta \varepsilon_s) a_s} = \sum_{\{a_s\}} \prod_s e^{-(\alpha + \beta \varepsilon_s) a_s}$$

$$= \prod_s \sum_{a_s} e^{-(\alpha + \beta \varepsilon_s) a_s}.$$
- Set $\mathcal{Z}_s = \sum_{a_s} e^{-(\alpha + \beta \varepsilon_s) a_s}$, then $\mathcal{Z} = \prod_s \mathcal{Z}_s$.
- For Bosons, $\mathcal{Z}_s = \sum_{a_s=0}^{\infty} e^{-(\alpha + \beta \varepsilon_s) a_s}$

\bar{a}_l of nearly independent particles

- Grand partition function: $\mathcal{Z} = \sum_N \sum_S e^{-\alpha N - \beta E_S}$

$$= \sum_{\{a_s\}} e^{-\sum_s (\alpha + \beta \varepsilon_s) a_s} = \sum_{\{a_s\}} \prod_s e^{-(\alpha + \beta \varepsilon_s) a_s}$$

$$= \prod_s \sum_{a_s} e^{-(\alpha + \beta \varepsilon_s) a_s}.$$
- Set $\mathcal{Z}_s = \sum_{a_s} e^{-(\alpha + \beta \varepsilon_s) a_s}$, then $\mathcal{Z} = \prod_s \mathcal{Z}_s$.
- For Bosons, $\mathcal{Z}_s = \sum_{a_s=0}^{\infty} e^{-(\alpha + \beta \varepsilon_s) a_s} = \frac{1}{1 - e^{-\alpha - \beta \varepsilon_s}};$

\bar{a}_l of nearly independent particles

- Grand partition function: $\mathcal{Z} = \sum_N \sum_S e^{-\alpha N - \beta E_S}$

$$= \sum_{\{a_s\}} e^{-\sum_s (\alpha + \beta \varepsilon_s) a_s} = \sum_{\{a_s\}} \prod_s e^{-(\alpha + \beta \varepsilon_s) a_s}$$

$$= \prod_s \sum_{a_s} e^{-(\alpha + \beta \varepsilon_s) a_s}.$$
- Set $\mathcal{Z}_s = \sum_{a_s} e^{-(\alpha + \beta \varepsilon_s) a_s}$, then $\mathcal{Z} = \prod_s \mathcal{Z}_s$.
- For Bosons, $\mathcal{Z}_s = \sum_{a_s=0}^{\infty} e^{-(\alpha + \beta \varepsilon_s) a_s} = \frac{1}{1 - e^{-\alpha - \beta \varepsilon_s}};$
- For Fermions, $\mathcal{Z}_s = \sum_{a_s=0}^1 e^{-(\alpha + \beta \varepsilon_s) a_s}$

\bar{a}_l of nearly independent particles

- Grand partition function: $\mathcal{Z} = \sum_N \sum_S e^{-\alpha N - \beta E_S}$

$$= \sum_{\{a_s\}} e^{-\sum_s (\alpha + \beta \varepsilon_s) a_s} = \sum_{\{a_s\}} \prod_s e^{-(\alpha + \beta \varepsilon_s) a_s}$$

$$= \prod_s \sum_{a_s} e^{-(\alpha + \beta \varepsilon_s) a_s}.$$
- Set $\mathcal{Z}_s = \sum_{a_s} e^{-(\alpha + \beta \varepsilon_s) a_s}$, then $\mathcal{Z} = \prod_s \mathcal{Z}_s$.
- For Bosons, $\mathcal{Z}_s = \sum_{a_s=0}^{\infty} e^{-(\alpha + \beta \varepsilon_s) a_s} = \frac{1}{1 - e^{-\alpha - \beta \varepsilon_s}};$
- For Fermions, $\mathcal{Z}_s = \sum_{a_s=0}^1 e^{-(\alpha + \beta \varepsilon_s) a_s} = 1 + e^{-\alpha - \beta \varepsilon_s}.$

$$\bar{a}_l \quad \mathcal{Z}_{s,\text{Bose}} = \frac{1}{1 - e^{-\alpha - \beta \varepsilon_s}}, \quad \mathcal{Z}_{s,\text{Fermi}} = 1 + e^{-\alpha - \beta \varepsilon_s}$$

- Average number of particles (6.7.10):

$$\bar{a}_l \quad \mathcal{Z}_{s,\text{Bose}} = \frac{1}{1 - e^{-\alpha - \beta \varepsilon_s}}, \quad \mathcal{Z}_{s,\text{Fermi}} = 1 + e^{-\alpha - \beta \varepsilon_s}$$

- Average number of particles (6.7.10):

$$f_s = \bar{a}_s$$

$$\bar{a}_l \quad \mathcal{Z}_{s,\text{Bose}} = \frac{1}{1 - e^{-\alpha - \beta \varepsilon_s}}, \quad \mathcal{Z}_{s,\text{Fermi}} = 1 + e^{-\alpha - \beta \varepsilon_s}$$

- Average number of particles (6.7.10):

$$f_s = \bar{a}_s = \frac{1}{\mathcal{Z}} \sum_N \sum_{S'} a_s e^{-\alpha N - \beta E_{S'}}$$

$$\bar{a}_l \quad \mathcal{Z}_{s,\text{Bose}} = \frac{1}{1 - e^{-\alpha - \beta \varepsilon_s}}, \quad \mathcal{Z}_{s,\text{Fermi}} = 1 + e^{-\alpha - \beta \varepsilon_s}$$

- Average number of particles (6.7.10):

$$\begin{aligned} f_s = \bar{a}_s &= \frac{1}{\mathcal{Z}} \sum_N \sum_{S'} a_s e^{-\alpha N - \beta E_{S'}} \\ &= \frac{1}{\mathcal{Z}} \sum_N \sum_{S'} a_s e^{-\sum (\alpha + \beta \varepsilon_{s'}) a_{s'}} \end{aligned}$$

$$\bar{a}_l \quad \mathcal{Z}_{s,\text{Bose}} = \frac{1}{1 - e^{-\alpha - \beta \varepsilon_s}}, \quad \mathcal{Z}_{s,\text{Fermi}} = 1 + e^{-\alpha - \beta \varepsilon_s}$$

- Average number of particles (6.7.10):

$$\begin{aligned} f_s = \bar{a}_s &= \frac{1}{\mathcal{Z}} \sum_N \sum_{S'} a_s e^{-\alpha N - \beta E_{S'}} \\ &= \frac{1}{\mathcal{Z}} \sum_N \sum_{S'} a_s e^{-\sum (\alpha + \beta \varepsilon_{s'}) a_{s'}} = \frac{1}{\mathcal{Z}} \sum_{\{a_{s'}\}} a_s e^{-\sum (\alpha + \beta \varepsilon_{s'}) a_{s'}} \end{aligned}$$

$$\bar{a}_l \quad \mathcal{Z}_{s,\text{Bose}} = \frac{1}{1 - e^{-\alpha - \beta \varepsilon_s}}, \quad \mathcal{Z}_{s,\text{Fermi}} = 1 + e^{-\alpha - \beta \varepsilon_s}$$

- Average number of particles (6.7.10):

$$\begin{aligned} f_s = \bar{a}_s &= \frac{1}{\mathcal{Z}} \sum_N \sum_{S'} a_s e^{-\alpha N - \beta E_{S'}} \\ &= \frac{1}{\mathcal{Z}} \sum_N \sum_{S'} a_s e^{-\sum (\alpha + \beta \varepsilon_{s'}) a_{s'}} = \frac{1}{\mathcal{Z}} \sum_{\{a_{s'}\}} a_s e^{-\sum (\alpha + \beta \varepsilon_{s'}) a_{s'}} \\ &= \frac{1}{\mathcal{Z}} \sum_{\{a_{s'}\}} a_s \prod_{s'} e^{-(\alpha + \beta \varepsilon_{s'}) a_{s'}} \end{aligned}$$

$$\bar{a}_l \quad \mathcal{Z}_{s,\text{Bose}} = \frac{1}{1 - e^{-\alpha - \beta \varepsilon_s}}, \quad \mathcal{Z}_{s,\text{Fermi}} = 1 + e^{-\alpha - \beta \varepsilon_s}$$

- Average number of particles (6.7.10):

$$\begin{aligned} f_s = \bar{a}_s &= \frac{1}{\mathcal{Z}} \sum_N \sum_{S'} a_s e^{-\alpha N - \beta E_{S'}} \\ &= \frac{1}{\mathcal{Z}} \sum_N \sum_{S'} a_s e^{-\sum (\alpha + \beta \varepsilon_{s'}) a_{s'}} = \frac{1}{\mathcal{Z}} \sum_{\{a_{s'}\}} a_s e^{-\sum (\alpha + \beta \varepsilon_{s'}) a_{s'}} \\ &= \frac{1}{\mathcal{Z}} \sum_{\{a_{s'}\}} a_s \prod_{s'} e^{-(\alpha + \beta \varepsilon_{s'}) a_{s'}} = \frac{1}{\mathcal{Z}} \prod_{s'} \sum_{a_{s'}} a_s^{\frac{1}{2}} e^{-(\alpha + \beta \varepsilon_{s'}) a_{s'}} \end{aligned}$$

$$\bar{a}_l \quad \mathcal{Z}_{s,\text{Bose}} = \frac{1}{1 - e^{-\alpha - \beta \varepsilon_s}}, \quad \mathcal{Z}_{s,\text{Fermi}} = 1 + e^{-\alpha - \beta \varepsilon_s}$$

- Average number of particles (6.7.10):

$$\begin{aligned} f_s = \bar{a}_s &= \frac{1}{\mathcal{Z}} \sum_N \sum_{S'} a_s e^{-\alpha N - \beta E_{S'}} \\ &= \frac{1}{\mathcal{Z}} \sum_N \sum_{S'} a_s e^{-\sum (\alpha + \beta \varepsilon_{s'}) a_{s'}} = \frac{1}{\mathcal{Z}} \sum_{\{a_{s'}\}} a_s e^{-\sum (\alpha + \beta \varepsilon_{s'}) a_{s'}} \\ &= \frac{1}{\mathcal{Z}} \sum_{\{a_{s'}\}} a_s \prod_{s'} e^{-(\alpha + \beta \varepsilon_{s'}) a_{s'}} = \frac{1}{\mathcal{Z}} \prod_{s'} \sum_{a_{s'}} a_s^{\frac{1}{2}} e^{-(\alpha + \beta \varepsilon_{s'}) a_{s'}} \\ &= \frac{1}{\mathcal{Z}} \sum_{a_s} a_s e^{-(\alpha + \beta \varepsilon_s) a_s} \cdot \prod_{s' \neq s} \sum_{a_{s'}} e^{-(\alpha + \beta \varepsilon_{s'}) a_{s'}} \end{aligned}$$

$$\bar{a}_l \quad \mathcal{Z}_{s,\text{Bose}} = \frac{1}{1 - e^{-\alpha - \beta \varepsilon_s}}, \quad \mathcal{Z}_{s,\text{Fermi}} = 1 + e^{-\alpha - \beta \varepsilon_s}$$

- Average number of particles (6.7.10):

$$\begin{aligned}
 f_s = \bar{a}_s &= \frac{1}{\mathcal{Z}} \sum_N \sum_{S'} a_s e^{-\alpha N - \beta E_{S'}} \\
 &= \frac{1}{\mathcal{Z}} \sum_N \sum_{S'} a_s e^{-\sum (\alpha + \beta \varepsilon_{s'}) a_{s'}} = \frac{1}{\mathcal{Z}} \sum_{\{a_{s'}\}} a_s e^{-\sum (\alpha + \beta \varepsilon_{s'}) a_{s'}} \\
 &= \frac{1}{\mathcal{Z}} \sum_{\{a_{s'}\}} a_s \prod_{s'} e^{-(\alpha + \beta \varepsilon_{s'}) a_{s'}} = \frac{1}{\mathcal{Z}} \prod_{s'} \sum_{a_{s'}} a_s^{\frac{1}{2}} e^{-(\alpha + \beta \varepsilon_{s'}) a_{s'}} \\
 &= \frac{1}{\mathcal{Z}} \sum_{a_s} a_s e^{-(\alpha + \beta \varepsilon_s) a_s} \cdot \prod_{s' \neq s} \sum_{a_{s'}} e^{-(\alpha + \beta \varepsilon_{s'}) a_{s'}} \\
 &= \frac{1}{\mathcal{Z}_s} \sum_{a_s} a_s e^{-(\alpha + \beta \varepsilon_s) a_s}
 \end{aligned}$$

$$\bar{a}_l \quad \mathcal{Z}_{s,\text{Bose}} = \frac{1}{1 - e^{-\alpha - \beta \varepsilon_s}}, \quad \mathcal{Z}_{s,\text{Fermi}} = 1 + e^{-\alpha - \beta \varepsilon_s}$$

- Average number of particles (6.7.10):

$$\begin{aligned}
 f_s = \bar{a}_s &= \frac{1}{\mathcal{Z}} \sum_N \sum_{S'} a_s e^{-\alpha N - \beta E_{S'}} \\
 &= \frac{1}{\mathcal{Z}} \sum_N \sum_{S'} a_s e^{-\sum (\alpha + \beta \varepsilon_{s'}) a_{s'}} = \frac{1}{\mathcal{Z}} \sum_{\{a_{s'}\}} a_s e^{-\sum (\alpha + \beta \varepsilon_{s'}) a_{s'}} \\
 &= \frac{1}{\mathcal{Z}} \sum_{\{a_{s'}\}} a_s \prod_{s'} e^{-(\alpha + \beta \varepsilon_{s'}) a_{s'}} = \frac{1}{\mathcal{Z}} \prod_{s'} \sum_{a_{s'}} a_s^{\frac{1}{2}} e^{-(\alpha + \beta \varepsilon_{s'}) a_{s'}} \\
 &= \frac{1}{\mathcal{Z}} \sum_{a_s} a_s e^{-(\alpha + \beta \varepsilon_s) a_s} \cdot \prod_{s' \neq s} \sum_{a_{s'}} e^{-(\alpha + \beta \varepsilon_{s'}) a_{s'}} \\
 &= \frac{1}{\mathcal{Z}_s} \sum_{a_s} a_s e^{-(\alpha + \beta \varepsilon_s) a_s} = \frac{1}{\mathcal{Z}_s} \left(-\frac{\partial}{\partial \alpha} \right) \mathcal{Z}_s
 \end{aligned}$$

\bar{a}_l	$\mathcal{Z}_{s,\text{Bose}} = \frac{1}{1 - e^{-\alpha - \beta \varepsilon_s}}, \quad \mathcal{Z}_{s,\text{Fermi}} = 1 + e^{-\alpha - \beta \varepsilon_s}$
-------------	---

- Average number of particles (6.7.10):

$$\begin{aligned}
 f_s = \bar{a}_s &= \frac{1}{\mathcal{Z}} \sum_N \sum_{S'} a_s e^{-\alpha N - \beta E_{S'}} \\
 &= \frac{1}{\mathcal{Z}} \sum_N \sum_{S'} a_s e^{-\sum (\alpha + \beta \varepsilon_{s'}) a_{s'}} = \frac{1}{\mathcal{Z}} \sum_{\{a_{s'}\}} a_s e^{-\sum (\alpha + \beta \varepsilon_{s'}) a_{s'}} \\
 &= \frac{1}{\mathcal{Z}} \sum_{\{a_{s'}\}} a_s \prod_{s'} e^{-(\alpha + \beta \varepsilon_{s'}) a_{s'}} = \frac{1}{\mathcal{Z}} \prod_{s'} \sum_{a_{s'}} a_s^{\frac{1}{2}} e^{-(\alpha + \beta \varepsilon_{s'}) a_{s'}} \\
 &= \frac{1}{\mathcal{Z}} \sum_{a_s} a_s e^{-(\alpha + \beta \varepsilon_s) a_s} \cdot \prod_{s' \neq s} \sum_{a_{s'}} e^{-(\alpha + \beta \varepsilon_{s'}) a_{s'}} \\
 &= \frac{1}{\mathcal{Z}_s} \sum_{a_s} a_s e^{-(\alpha + \beta \varepsilon_s) a_s} = \frac{1}{\mathcal{Z}_s} \left(-\frac{\partial}{\partial \alpha} \right) \mathcal{Z}_s = -\frac{\partial \ln \mathcal{Z}_s}{\partial \alpha}.
 \end{aligned}$$

$$\bar{a}_l \quad \mathcal{Z}_{s,\text{Bose}} = \frac{1}{1 - e^{-\alpha - \beta \varepsilon_s}}, \quad \mathcal{Z}_{s,\text{Fermi}} = 1 + e^{-\alpha - \beta \varepsilon_s}$$

- Average number of particles (6.7.10):

$$\begin{aligned} f_s = \bar{a}_s &= \frac{1}{\mathcal{Z}} \sum_N \sum_{S'} a_s e^{-\alpha N - \beta E_{S'}} \\ &= \frac{1}{\mathcal{Z}} \sum_N \sum_{S'} a_s e^{-\sum (\alpha + \beta \varepsilon_{s'}) a_{s'}} = \frac{1}{\mathcal{Z}} \sum_{\{a_{s'}\}} a_s e^{-\sum (\alpha + \beta \varepsilon_{s'}) a_{s'}} \\ &= \frac{1}{\mathcal{Z}} \sum_{\{a_{s'}\}} a_s \prod_{s'} e^{-(\alpha + \beta \varepsilon_{s'}) a_{s'}} = \frac{1}{\mathcal{Z}} \prod_{s'} \sum_{a_{s'}} a_s^{\frac{1}{2}} e^{-(\alpha + \beta \varepsilon_{s'}) a_{s'}} \\ &= \frac{1}{\mathcal{Z}} \sum_{a_s} a_s e^{-(\alpha + \beta \varepsilon_s) a_s} \cdot \prod_{s' \neq s} \sum_{a_{s'}} e^{-(\alpha + \beta \varepsilon_{s'}) a_{s'}} \\ &= \frac{1}{\mathcal{Z}_s} \sum_{a_s} a_s e^{-(\alpha + \beta \varepsilon_s) a_s} = \frac{1}{\mathcal{Z}_s} \left(-\frac{\partial}{\partial \alpha} \right) \mathcal{Z}_s = -\frac{\partial \ln \mathcal{Z}_s}{\partial \alpha}. \end{aligned}$$

- For Bosons: $\bar{a}_s = \frac{1}{e^{\alpha + \beta \varepsilon_s} - 1},$

$$\bar{a}_l \quad \mathcal{Z}_{s,\text{Bose}} = \frac{1}{1 - e^{-\alpha - \beta \varepsilon_s}}, \quad \mathcal{Z}_{s,\text{Fermi}} = 1 + e^{-\alpha - \beta \varepsilon_s}$$

- Average number of particles (6.7.10):

$$\begin{aligned} f_s = \bar{a}_s &= \frac{1}{\mathcal{Z}} \sum_N \sum_{S'} a_s e^{-\alpha N - \beta E_{S'}} \\ &= \frac{1}{\mathcal{Z}} \sum_N \sum_{S'} a_s e^{-\sum (\alpha + \beta \varepsilon_{s'}) a_{s'}} = \frac{1}{\mathcal{Z}} \sum_{\{a_{s'}\}} a_s e^{-\sum (\alpha + \beta \varepsilon_{s'}) a_{s'}} \\ &= \frac{1}{\mathcal{Z}} \sum_{\{a_{s'}\}} a_s \prod_{s'} e^{-(\alpha + \beta \varepsilon_{s'}) a_{s'}} = \frac{1}{\mathcal{Z}} \prod_{s'} \sum_{a_{s'}} a_s^{\frac{1}{2}} e^{-(\alpha + \beta \varepsilon_{s'}) a_{s'}} \\ &= \frac{1}{\mathcal{Z}} \sum_{a_s} a_s e^{-(\alpha + \beta \varepsilon_s) a_s} \cdot \prod_{s' \neq s} \sum_{a_{s'}} e^{-(\alpha + \beta \varepsilon_{s'}) a_{s'}} \\ &= \frac{1}{\mathcal{Z}_s} \sum_{a_s} a_s e^{-(\alpha + \beta \varepsilon_s) a_s} = \frac{1}{\mathcal{Z}_s} \left(-\frac{\partial}{\partial \alpha} \right) \mathcal{Z}_s = -\frac{\partial \ln \mathcal{Z}_s}{\partial \alpha}. \end{aligned}$$

- For Bosons: $\bar{a}_s = \frac{1}{e^{\alpha + \beta \varepsilon_s} - 1}$, for Fermions, $\bar{a}_s = \frac{1}{e^{\alpha + \beta \varepsilon_s} + 1}$.

$$\bar{a}_l \quad \mathcal{Z}_{s,\text{Bose}} = \frac{1}{1 - e^{-\alpha - \beta \varepsilon_s}}, \quad \mathcal{Z}_{s,\text{Fermi}} = 1 + e^{-\alpha - \beta \varepsilon_s}$$

- Average number of particles (6.7.10):

$$\begin{aligned} f_s &= \bar{a}_s = \frac{1}{\mathcal{Z}} \sum_N \sum_{s'} a_s e^{-\alpha N - \beta E_{s'}} \\ &= \frac{1}{\mathcal{Z}} \sum_N \sum_{s'} a_s e^{-\sum (\alpha + \beta \varepsilon_{s'}) a_{s'}} = \frac{1}{\mathcal{Z}} \sum_{\{a_{s'}\}} a_s e^{-\sum (\alpha + \beta \varepsilon_{s'}) a_{s'}} \\ &= \frac{1}{\mathcal{Z}} \sum_{\{a_{s'}\}} a_s \prod_{s'} e^{-(\alpha + \beta \varepsilon_{s'}) a_{s'}} = \frac{1}{\mathcal{Z}} \prod_{s'} \sum_{a_{s'}} a_s^{\frac{1}{2}} e^{-(\alpha + \beta \varepsilon_{s'}) a_{s'}} \\ &= \frac{1}{\mathcal{Z}} \sum_{a_s} a_s e^{-(\alpha + \beta \varepsilon_s) a_s} \cdot \prod_{s' \neq s} \sum_{a_{s'}} e^{-(\alpha + \beta \varepsilon_{s'}) a_{s'}} \\ &= \frac{1}{\mathcal{Z}_s} \sum_{a_s} a_s e^{-(\alpha + \beta \varepsilon_s) a_s} = \frac{1}{\mathcal{Z}_s} \left(-\frac{\partial}{\partial \alpha} \right) \mathcal{Z}_s = -\frac{\partial \ln \mathcal{Z}_s}{\partial \alpha}. \end{aligned}$$

- For Bosons: $\bar{a}_s = \frac{1}{e^{\alpha + \beta \varepsilon_s} - 1}$, for Fermions, $\bar{a}_s = \frac{1}{e^{\alpha + \beta \varepsilon_s} + 1}$.
- For degenerated states: $\bar{a}_l = \sum_{\omega_l} \bar{a}_s$

$$\bar{a}_l \quad \mathcal{Z}_{s,\text{Bose}} = \frac{1}{1 - e^{-\alpha - \beta \varepsilon_s}}, \quad \mathcal{Z}_{s,\text{Fermi}} = 1 + e^{-\alpha - \beta \varepsilon_s}$$

- Average number of particles (6.7.10):

$$\begin{aligned} f_s &= \bar{a}_s = \frac{1}{\mathcal{Z}} \sum_N \sum_{S'} a_s e^{-\alpha N - \beta E_{S'}} \\ &= \frac{1}{\mathcal{Z}} \sum_N \sum_{S'} a_s e^{-\sum (\alpha + \beta \varepsilon_{s'}) a_{s'}} = \frac{1}{\mathcal{Z}} \sum_{\{a_{s'}\}} a_s e^{-\sum (\alpha + \beta \varepsilon_{s'}) a_{s'}} \\ &= \frac{1}{\mathcal{Z}} \sum_{\{a_{s'}\}} a_s \prod_{s'} e^{-(\alpha + \beta \varepsilon_{s'}) a_{s'}} = \frac{1}{\mathcal{Z}} \prod_{s'} \sum_{a_{s'}} a_s^{\frac{1}{2}} e^{-(\alpha + \beta \varepsilon_{s'}) a_{s'}} \\ &= \frac{1}{\mathcal{Z}} \sum_{a_s} a_s e^{-(\alpha + \beta \varepsilon_s) a_s} \cdot \prod_{s' \neq s} \sum_{a_{s'}} e^{-(\alpha + \beta \varepsilon_{s'}) a_{s'}} \\ &= \frac{1}{\mathcal{Z}_s} \sum_{a_s} a_s e^{-(\alpha + \beta \varepsilon_s) a_s} = \frac{1}{\mathcal{Z}_s} \left(-\frac{\partial}{\partial \alpha} \right) \mathcal{Z}_s = -\frac{\partial \ln \mathcal{Z}_s}{\partial \alpha}. \end{aligned}$$

- For Bosons: $\bar{a}_s = \frac{1}{e^{\alpha + \beta \varepsilon_s} - 1}$, for Fermions, $\bar{a}_s = \frac{1}{e^{\alpha + \beta \varepsilon_s} + 1}$.
- For degenerated states: $\bar{a}_l = \sum_l \bar{a}_s = \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} \pm 1}$.

Fluctuation of Bose/Fermi distribution

$$\bar{a}_l = \frac{\omega_l}{e^{\alpha + \beta \epsilon_l} \pm 1}$$

- App 3. Fluctuation of Bose/Fermi distribution.

Fluctuation of Bose/Fermi distribution

$$\overline{a}_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l} \pm 1}$$

- App 3. Fluctuation of Bose/Fermi distribution.
- Take the particles at energy level ε_l as an open system.

Fluctuation of Bose/Fermi distribution

$$\overline{a_l} = \frac{\omega_l}{e^{\alpha + \beta \varepsilon_l} \pm 1}$$

- App 3. Fluctuation of Bose/Fermi distribution.
- Take the particles at energy level ε_l as an open system.
 a_l acts like N in (9.11.8).

Fluctuation of Bose/Fermi distribution

$$\bar{a}_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l} \pm 1}$$

- App 3. Fluctuation of Bose/Fermi distribution.
- Take the particles at energy level ε_l as an open system.
 a_l acts like N in (9.11.8).
- $\overline{(a_l - \bar{a}_l)^2} = -\frac{\partial \bar{a}_l}{\partial \alpha}$

Fluctuation of Bose/Fermi distribution

$$\bar{a}_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l} \pm 1}$$

- App 3. Fluctuation of Bose/Fermi distribution.
- Take the particles at energy level ε_l as an open system.
 a_l acts like N in (9.11.8).
- $\overline{(a_l - \bar{a}_l)^2} = -\frac{\partial \bar{a}_l}{\partial \alpha} = -\frac{\partial \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l} \pm 1}}{\partial \alpha}$

Fluctuation of Bose/Fermi distribution

$$\bar{a}_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}\pm 1}$$

- App 3. Fluctuation of Bose/Fermi distribution.
- Take the particles at energy level ε_l as an open system. a_l acts like N in (9.11.8).
- $$\overline{(a_l - \bar{a}_l)^2} = -\frac{\partial \bar{a}_l}{\partial \alpha} = -\frac{\partial \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}\pm 1}}{\partial \alpha} = \frac{\omega_l e^{\alpha+\beta\varepsilon_l}}{(e^{\alpha+\beta\varepsilon_l}\pm 1)^2}$$

Fluctuation of Bose/Fermi distribution

$$\bar{a}_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}\pm 1}$$

- App 3. Fluctuation of Bose/Fermi distribution.
- Take the particles at energy level ε_l as an open system.
 a_l acts like N in (9.11.8).

$$\begin{aligned} \overline{(a_l - \bar{a}_l)^2} &= -\frac{\partial \bar{a}_l}{\partial \alpha} = -\frac{\partial \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}\pm 1}}{\partial \alpha} = \frac{\omega_l e^{\alpha+\beta\varepsilon_l}}{(e^{\alpha+\beta\varepsilon_l}\pm 1)^2} \\ &= \bar{a}_l \frac{e^{\alpha+\beta\varepsilon_l}}{e^{\alpha+\beta\varepsilon_l}\pm 1} \end{aligned}$$

Fluctuation of Bose/Fermi distribution

$$\bar{a}_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l\pm 1}}$$

- App 3. Fluctuation of Bose/Fermi distribution.
- Take the particles at energy level ε_l as an open system.
 a_l acts like N in (9.11.8).

$$\begin{aligned} \overline{(a_l - \bar{a}_l)^2} &= -\frac{\partial \bar{a}_l}{\partial \alpha} = -\frac{\partial \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l\pm 1}}}{\partial \alpha} = \frac{\omega_l e^{\alpha+\beta\varepsilon_l}}{(e^{\alpha+\beta\varepsilon_l\pm 1})^2} \\ &= \bar{a}_l \frac{e^{\alpha+\beta\varepsilon_l}}{e^{\alpha+\beta\varepsilon_l\pm 1}} = \bar{a}_l \left(1 \mp \frac{1}{e^{\alpha+\beta\varepsilon_l\pm 1}}\right) \end{aligned}$$

Fluctuation of Bose/Fermi distribution

$$\bar{a}_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l\pm 1}}$$

- App 3. Fluctuation of Bose/Fermi distribution.
- Take the particles at energy level ε_l as an open system. a_l acts like N in (9.11.8).

$$\begin{aligned} \overline{(a_l - \bar{a}_l)^2} &= -\frac{\partial \bar{a}_l}{\partial \alpha} = -\frac{\partial \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l\pm 1}}}{\partial \alpha} = \frac{\omega_l e^{\alpha+\beta\varepsilon_l}}{(e^{\alpha+\beta\varepsilon_l\pm 1})^2} \\ &= \bar{a}_l \frac{e^{\alpha+\beta\varepsilon_l}}{e^{\alpha+\beta\varepsilon_l\pm 1}} = \bar{a}_l \left(1 \mp \frac{1}{e^{\alpha+\beta\varepsilon_l\pm 1}}\right) = \bar{a}_l \left(1 \mp \frac{\bar{a}_l}{\omega_l}\right). \end{aligned}$$

Fluctuation of Bose/Fermi distribution

$$\bar{a}_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l\pm 1}}$$

- App 3. Fluctuation of Bose/Fermi distribution.
- Take the particles at energy level ε_l as an open system. a_l acts like N in (9.11.8).
- $$\overline{(a_l - \bar{a}_l)^2} = -\frac{\partial \bar{a}_l}{\partial \alpha} = -\frac{\partial \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l\pm 1}}}{\partial \alpha} = \frac{\omega_l e^{\alpha+\beta\varepsilon_l}}{(e^{\alpha+\beta\varepsilon_l\pm 1})^2}$$
$$= \bar{a}_l \frac{e^{\alpha+\beta\varepsilon_l}}{e^{\alpha+\beta\varepsilon_l\pm 1}} = \bar{a}_l \left(1 \mp \frac{1}{e^{\alpha+\beta\varepsilon_l\pm 1}}\right) = \bar{a}_l \left(1 \mp \frac{\bar{a}_l}{\omega_l}\right).$$
- For Fermions, $\varepsilon < \mu$, $\frac{\bar{a}_l}{\omega_l} \simeq 1$; $\varepsilon > \mu$, $\bar{a}_l \simeq 0$, so the fluctuation is small.

Fluctuation of Bose/Fermi distribution

$$\bar{a}_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l\pm 1}}$$

- App 3. Fluctuation of Bose/Fermi distribution.
- Take the particles at energy level ε_l as an open system. a_l acts like N in (9.11.8).
- $$\overline{(a_l - \bar{a}_l)^2} = -\frac{\partial \bar{a}_l}{\partial \alpha} = -\frac{\partial \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l\pm 1}}}{\partial \alpha} = \frac{\omega_l e^{\alpha+\beta\varepsilon_l}}{(e^{\alpha+\beta\varepsilon_l\pm 1})^2}$$
$$= \bar{a}_l \frac{e^{\alpha+\beta\varepsilon_l}}{e^{\alpha+\beta\varepsilon_l\pm 1}} = \bar{a}_l \left(1 \mp \frac{1}{e^{\alpha+\beta\varepsilon_l\pm 1}}\right) = \bar{a}_l \left(1 \mp \frac{\bar{a}_l}{\omega_l}\right).$$
- For Fermions, $\varepsilon < \mu$, $\frac{\bar{a}_l}{\omega_l} \simeq 1$; $\varepsilon > \mu$, $\bar{a}_l \simeq 0$, so the fluctuation is small. For Bosons, no such restrict.

Fluctuation of Bose/Fermi distribution

$$\bar{a}_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}\pm 1}$$

- App 3. Fluctuation of Bose/Fermi distribution.
- Take the particles at energy level ε_l as an open system. a_l acts like N in (9.11.8).
- $$\overline{(a_l - \bar{a}_l)^2} = -\frac{\partial \bar{a}_l}{\partial \alpha} = -\frac{\partial \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}\pm 1}}{\partial \alpha} = \frac{\omega_l e^{\alpha+\beta\varepsilon_l}}{(e^{\alpha+\beta\varepsilon_l}\pm 1)^2}$$
$$= \bar{a}_l \frac{e^{\alpha+\beta\varepsilon_l}}{e^{\alpha+\beta\varepsilon_l}\pm 1} = \bar{a}_l \left(1 \mp \frac{1}{e^{\alpha+\beta\varepsilon_l}\pm 1}\right) = \bar{a}_l \left(1 \mp \frac{\bar{a}_l}{\omega_l}\right).$$
- For Fermions, $\varepsilon < \mu$, $\frac{\bar{a}_l}{\omega_l} \simeq 1$; $\varepsilon > \mu$, $\bar{a}_l \simeq 0$, so the fluctuation is small. For Bosons, no such restrict.
- The correlation: $\overline{a_l a_m}$

Fluctuation of Bose/Fermi distribution

$$\bar{a}_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l\pm 1}}$$

- App 3. Fluctuation of Bose/Fermi distribution.
- Take the particles at energy level ε_l as an open system. a_l acts like N in (9.11.8).
- $\overline{(a_l - \bar{a}_l)^2} = -\frac{\partial \bar{a}_l}{\partial \alpha} = -\frac{\partial \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l\pm 1}}}{\partial \alpha} = \frac{\omega_l e^{\alpha+\beta\varepsilon_l}}{(e^{\alpha+\beta\varepsilon_l\pm 1})^2}$
 $= \bar{a}_l \frac{e^{\alpha+\beta\varepsilon_l}}{e^{\alpha+\beta\varepsilon_l\pm 1}} = \bar{a}_l (1 \mp \frac{1}{e^{\alpha+\beta\varepsilon_l\pm 1}}) = \bar{a}_l (1 \mp \frac{\bar{a}_l}{\omega_l}).$
- For Fermions, $\varepsilon < \mu$, $\frac{\bar{a}_l}{\omega_l} \simeq 1$; $\varepsilon > \mu$, $\bar{a}_l \simeq 0$, so the fluctuation is small. For Bosons, no such restrict.
- The correlation: $\overline{a_l a_m} = \frac{1}{\mathcal{Z}} \sum_N \sum_S a_l a_m e^{-(\alpha N + \beta E_S)}$

Fluctuation of Bose/Fermi distribution

$$\bar{a}_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}\pm 1}$$

- App 3. Fluctuation of Bose/Fermi distribution.
- Take the particles at energy level ε_l as an open system. a_l acts like N in (9.11.8).
- $\overline{(a_l - \bar{a}_l)^2} = -\frac{\partial \bar{a}_l}{\partial \alpha} = -\frac{\partial \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}\pm 1}}{\partial \alpha} = \frac{\omega_l e^{\alpha+\beta\varepsilon_l}}{(e^{\alpha+\beta\varepsilon_l}\pm 1)^2}$
 $= \bar{a}_l \frac{e^{\alpha+\beta\varepsilon_l}}{e^{\alpha+\beta\varepsilon_l}\pm 1} = \bar{a}_l (1 \mp \frac{1}{e^{\alpha+\beta\varepsilon_l}\pm 1}) = \bar{a}_l (1 \mp \frac{\bar{a}_l}{\omega_l}).$
- For Fermions, $\varepsilon < \mu$, $\frac{\bar{a}_l}{\omega_l} \simeq 1$; $\varepsilon > \mu$, $\bar{a}_l \simeq 0$, so the fluctuation is small. For Bosons, no such restrict.
- The correlation: $\overline{a_l a_m} = \frac{1}{\mathcal{Z}} \sum_N \sum_S a_l a_m e^{-(\alpha N + \beta E_S)}$
 $= \frac{1}{\mathcal{Z}_l} \left[\sum_{a_l} a_l e^{-(\alpha + \beta \varepsilon_l) a_l} \right] \cdot \frac{1}{\mathcal{Z}_m} \left[\sum_{a_m} a_m e^{-(\alpha + \beta \varepsilon_m) a_m} \right]$

Fluctuation of Bose/Fermi distribution

$$\bar{a}_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}\pm 1}$$

- App 3. Fluctuation of Bose/Fermi distribution.
- Take the particles at energy level ε_l as an open system. a_l acts like N in (9.11.8).
- $\overline{(a_l - \bar{a}_l)^2} = -\frac{\partial \bar{a}_l}{\partial \alpha} = -\frac{\partial \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}\pm 1}}{\partial \alpha} = \frac{\omega_l e^{\alpha+\beta\varepsilon_l}}{(e^{\alpha+\beta\varepsilon_l}\pm 1)^2}$
 $= \bar{a}_l \frac{e^{\alpha+\beta\varepsilon_l}}{e^{\alpha+\beta\varepsilon_l}\pm 1} = \bar{a}_l \left(1 \mp \frac{1}{e^{\alpha+\beta\varepsilon_l}\pm 1}\right) = \bar{a}_l \left(1 \mp \frac{\bar{a}_l}{\omega_l}\right).$
- For Fermions, $\varepsilon < \mu$, $\frac{\bar{a}_l}{\omega_l} \simeq 1$; $\varepsilon > \mu$, $\bar{a}_l \simeq 0$, so the fluctuation is small. For Bosons, no such restrict.
- The correlation: $\overline{a_l a_m} = \frac{1}{\mathcal{Z}} \sum_N \sum_S a_l a_m e^{-(\alpha N + \beta E_S)}$
 $= \frac{1}{\mathcal{Z}_l} \left[\sum_{a_l} a_l e^{-(\alpha + \beta\varepsilon_l)a_l} \right] \cdot \frac{1}{\mathcal{Z}_m} \left[\sum_{a_m} a_m e^{-(\alpha + \beta\varepsilon_m)a_m} \right] = \bar{a}_l \bar{a}_m.$

Fluctuation of Bose/Fermi distribution

$$\bar{a}_l = \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}\pm 1}$$

- App 3. Fluctuation of Bose/Fermi distribution.
- Take the particles at energy level ε_l as an open system. a_l acts like N in (9.11.8).
- $\overline{(a_l - \bar{a}_l)^2} = -\frac{\partial \bar{a}_l}{\partial \alpha} = -\frac{\partial \frac{\omega_l}{e^{\alpha+\beta\varepsilon_l}\pm 1}}{\partial \alpha} = \frac{\omega_l e^{\alpha+\beta\varepsilon_l}}{(e^{\alpha+\beta\varepsilon_l}\pm 1)^2}$
 $= \bar{a}_l \frac{e^{\alpha+\beta\varepsilon_l}}{e^{\alpha+\beta\varepsilon_l}\pm 1} = \bar{a}_l (1 \mp \frac{1}{e^{\alpha+\beta\varepsilon_l}\pm 1}) = \bar{a}_l (1 \mp \frac{\bar{a}_l}{\omega_l}).$
- For Fermions, $\varepsilon < \mu$, $\frac{\bar{a}_l}{\omega_l} \simeq 1$; $\varepsilon > \mu$, $\bar{a}_l \simeq 0$, so the fluctuation is small. For Bosons, no such restrict.
- The correlation: $\overline{a_l a_m} = \frac{1}{\mathcal{Z}} \sum_N \sum_S a_l a_m e^{-(\alpha N + \beta E_S)}$
 $= \frac{1}{\mathcal{Z}_l} [\sum_{a_l} a_l e^{-(\alpha+\beta\varepsilon_l)a_l}] \cdot \frac{1}{\mathcal{Z}_m} [\sum_{a_m} a_m e^{-(\alpha+\beta\varepsilon_m)a_m}] = \bar{a}_l \bar{a}_m.$
- $\overline{(a_l - \bar{a}_l)(a_m - \bar{a}_m)} = \overline{a_l a_m} - \bar{a}_l \bar{a}_m = 0$, no correlation.

Table of contents

1 §9. Ensemble theory

- 9.1 Phase space and Liouville's theorem
- 9.2 The microcanonical ensemble
- 9.3 Thermodynamics in microcanonical ensemble theory
- 9.4 Canonical ensemble
- 9.5 Thermodynamics in canonical ensemble
- 9.6 EOS of real gas
- 9.7 Thermal capacity of solid
- 9.9 Ising model
- 9.10 Grand canonical ensemble
- 9.11 Thermodynamics in grand canonical ensemble
- 9.12 Applications of the grand canonical ensemble