

# Thermodynamics & Statistical Physics

## Chapter 7. Boltzmann Statistics

Yuan-Chuan Zou  
zouyc@hust.edu.cn

School of Physics, Huazhong University of Science and Technology

December 30, 2013

# Table of contents

## 1 §7 Boltzmann Statistics

- 7.1 Thermal quantities in statistics
- 7.2 Equation of state of ideal gas
- 7.3 Maxwell speed distribution
- 7.4 Energy equipartition theorem
- 7.5 Internal energy and heat capacity of the ideal gas
- 7.6 Entropy of the ideal gas
- 7.7 Einstein's theory on heat capacity of solid
- 7.8 Paramagnetic solid
- 7.9 Negative temperature

# 7.1 Thermal quantities in statistics

## 7.1 Thermal quantities in statistics

- Internal energy:  $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}.$

## 7.1 Thermal quantities in statistics

- Internal energy:  $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}.$
- Introduce partition function:  $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}.$

## 7.1 Thermal quantities in statistics

- Internal energy:  $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$ .
- Introduce partition function:  $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$ .  
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)

## 7.1 Thermal quantities in statistics

- Internal energy:  $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$ .
- Introduce partition function:  $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$ .  
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)
- $N = \sum a_l$

## 7.1 Thermal quantities in statistics

- Internal energy:  $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$ .
- Introduce partition function:  $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$ .  
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)
- $N = \sum a_l = \sum \omega_l e^{-\alpha - \beta \varepsilon_l}$



## 7.1 Thermal quantities in statistics

- Internal energy:  $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$ .
- Introduce partition function:  $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$ .  
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)
- $N = \sum a_l = \sum \omega_l e^{-\alpha - \beta \varepsilon_l} = e^{-\alpha} \sum \omega_l e^{-\beta \varepsilon_l}$

## 7.1 Thermal quantities in statistics

- Internal energy:  $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$ .
- Introduce partition function:  $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$ .  
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)
- $N = \sum a_l = \sum \omega_l e^{-\alpha - \beta \varepsilon_l} = e^{-\alpha} \sum \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} Z_1$ .

## 7.1 Thermal quantities in statistics

- Internal energy:  $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$ .
- Introduce partition function:  $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$ .  
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)
- $N = \sum a_l = \sum \omega_l e^{-\alpha - \beta \varepsilon_l} = e^{-\alpha} \sum \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} Z_1$ .
- $\therefore e^{\alpha} = \frac{Z_1}{N}$ .

## 7.1 Thermal quantities in statistics

- Internal energy:  $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$ .
- Introduce partition function:  $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$ .  
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)
- $N = \sum a_l = \sum \omega_l e^{-\alpha - \beta \varepsilon_l} = e^{-\alpha} \sum \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} Z_1$ .
- $\therefore e^{\alpha} = \frac{Z_1}{N}$ .
- $U = e^{-\alpha} \sum \varepsilon_l \omega_l e^{-\beta \varepsilon_l}$

## 7.1 Thermal quantities in statistics

- Internal energy:  $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$ .
- Introduce partition function:  $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$ .  
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)
- $N = \sum a_l = \sum \omega_l e^{-\alpha - \beta \varepsilon_l} = e^{-\alpha} \sum \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} Z_1$ .
- $\therefore e^{\alpha} = \frac{Z_1}{N}$ .
- $U = e^{-\alpha} \sum \varepsilon_l \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} \sum -\frac{\partial}{\partial \beta} (\omega_l e^{-\beta \varepsilon_l})$

## 7.1 Thermal quantities in statistics

- Internal energy:  $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$ .
- Introduce partition function:  $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$ .  
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)
- $N = \sum a_l = \sum \omega_l e^{-\alpha - \beta \varepsilon_l} = e^{-\alpha} \sum \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} Z_1$ .
- $\therefore e^{\alpha} = \frac{Z_1}{N}$ .
- $U = e^{-\alpha} \sum \varepsilon_l \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} \sum -\frac{\partial}{\partial \beta} (\omega_l e^{-\beta \varepsilon_l})$   
 $= e^{-\alpha} \left(-\frac{\partial}{\partial \beta}\right) \sum (\omega_l e^{-\beta \varepsilon_l})$

# 7.1 Thermal quantities in statistics

- Internal energy:  $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$ .
- Introduce partition function:  $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$ .  
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)
- $N = \sum a_l = \sum \omega_l e^{-\alpha - \beta \varepsilon_l} = e^{-\alpha} \sum \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} Z_1$ .
- $\therefore e^{\alpha} = \frac{Z_1}{N}$ .
- $U = e^{-\alpha} \sum \varepsilon_l \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} \sum -\frac{\partial}{\partial \beta} (\omega_l e^{-\beta \varepsilon_l})$   
 $= e^{-\alpha} \left(-\frac{\partial}{\partial \beta}\right) \sum (\omega_l e^{-\beta \varepsilon_l}) = \frac{N}{Z_1} \left(-\frac{\partial}{\partial \beta}\right) Z_1$

# 7.1 Thermal quantities in statistics

- Internal energy:  $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$ .
- Introduce partition function:  $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$ .  
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)
- $N = \sum a_l = \sum \omega_l e^{-\alpha - \beta \varepsilon_l} = e^{-\alpha} \sum \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} Z_1$ .
- $\therefore e^{\alpha} = \frac{Z_1}{N}$ .
- $U = e^{-\alpha} \sum \varepsilon_l \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} \sum -\frac{\partial}{\partial \beta} (\omega_l e^{-\beta \varepsilon_l})$   
 $= e^{-\alpha} \left(-\frac{\partial}{\partial \beta}\right) \sum (\omega_l e^{-\beta \varepsilon_l}) = \frac{N}{Z_1} \left(-\frac{\partial}{\partial \beta}\right) Z_1 = -N \frac{\partial}{\partial \beta} \ln Z_1$ .



# 7.1 Thermal quantities in statistics

- Internal energy:  $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$ .
- Introduce partition function:  $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$ .  
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)
- $N = \sum a_l = \sum \omega_l e^{-\alpha - \beta \varepsilon_l} = e^{-\alpha} \sum \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} Z_1$ .
- $\therefore e^{\alpha} = \frac{Z_1}{N}$ .
- $U = e^{-\alpha} \sum \varepsilon_l \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} \sum -\frac{\partial}{\partial \beta} (\omega_l e^{-\beta \varepsilon_l})$   
 $= e^{-\alpha} \left(-\frac{\partial}{\partial \beta}\right) \sum (\omega_l e^{-\beta \varepsilon_l}) = \frac{N}{Z_1} \left(-\frac{\partial}{\partial \beta}\right) Z_1 = -N \frac{\partial}{\partial \beta} \ln Z_1$ .
- 1st law:  $dU = \delta W + \delta Q$ .

# 7.1 Thermal quantities in statistics

- Internal energy:  $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$ .
- Introduce partition function:  $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$ .  
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)
- $N = \sum a_l = \sum \omega_l e^{-\alpha - \beta \varepsilon_l} = e^{-\alpha} \sum \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} Z_1$ .
- $\therefore e^{\alpha} = \frac{Z_1}{N}$ .
- $U = e^{-\alpha} \sum \varepsilon_l \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} \sum -\frac{\partial}{\partial \beta} (\omega_l e^{-\beta \varepsilon_l})$   
 $= e^{-\alpha} \left(-\frac{\partial}{\partial \beta}\right) \sum (\omega_l e^{-\beta \varepsilon_l}) = \frac{N}{Z_1} \left(-\frac{\partial}{\partial \beta}\right) Z_1 = -N \frac{\partial}{\partial \beta} \ln Z_1$ .
- 1st law:  $dU = \delta W + \delta Q$ .  $\delta W = \sum_i Y_i dy_i$  ( $Y_i$  generalized force,  $dy_i$  generalized displacement)

# 7.1 Thermal quantities in statistics

- Internal energy:  $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$ .
- Introduce partition function:  $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$ .  
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)
- $N = \sum a_l = \sum \omega_l e^{-\alpha - \beta \varepsilon_l} = e^{-\alpha} \sum \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} Z_1$ .
- $\therefore e^{\alpha} = \frac{Z_1}{N}$ .
- $U = e^{-\alpha} \sum \varepsilon_l \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} \sum -\frac{\partial}{\partial \beta} (\omega_l e^{-\beta \varepsilon_l})$   
 $= e^{-\alpha} \left(-\frac{\partial}{\partial \beta}\right) \sum (\omega_l e^{-\beta \varepsilon_l}) = \frac{N}{Z_1} \left(-\frac{\partial}{\partial \beta}\right) Z_1 = -N \frac{\partial}{\partial \beta} \ln Z_1$ .
- 1st law:  $dU = \delta W + \delta Q$ .  $\delta W = \sum_i Y_i dy_i$  ( $Y_i$  generalized force,  $dy_i$  generalized displacement)  
 A force exerted on the system,  $U$  increases. But  $\{a_l\}$  does not change. What changes?

# 7.1 Thermal quantities in statistics

- Internal energy:  $U = \sum a_l \varepsilon_l = \sum \varepsilon_l \omega_l e^{-\alpha - \beta \varepsilon_l}$ .
- Introduce partition function:  $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$ .  
(Most thermal variables can be expressed in terms of the partition function or its derivatives.)
- $N = \sum a_l = \sum \omega_l e^{-\alpha - \beta \varepsilon_l} = e^{-\alpha} \sum \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} Z_1$ .
- $\therefore e^{\alpha} = \frac{Z_1}{N}$ .
- $U = e^{-\alpha} \sum \varepsilon_l \omega_l e^{-\beta \varepsilon_l} = e^{-\alpha} \sum -\frac{\partial}{\partial \beta} (\omega_l e^{-\beta \varepsilon_l})$   
 $= e^{-\alpha} \left(-\frac{\partial}{\partial \beta}\right) \sum (\omega_l e^{-\beta \varepsilon_l}) = \frac{N}{Z_1} \left(-\frac{\partial}{\partial \beta}\right) Z_1 = -N \frac{\partial}{\partial \beta} \ln Z_1$ .
- 1st law:  $dU = \delta W + \delta Q$ .  $\delta W = \sum_i Y_i dy_i$  ( $Y_i$  generalized force,  $dy_i$  generalized displacement)  
 A force exerted on the system,  $U$  increases. But  $\{a_l\}$  does not change. What changes?  $\{\varepsilon_l\}$ .

# Thermal quantities in statistics

- For each particle, the generalized force is  $\frac{\partial \varepsilon_l}{\partial y}$ ,

# Thermal quantities in statistics

- For each particle, the generalized force is  $\frac{\partial \varepsilon_l}{\partial y}$ , the whole force ( $Y_i$ ):  $Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l$

# Thermal quantities in statistics

- For each particle, the generalized force is  $\frac{\partial \varepsilon_l}{\partial y}$ , the whole force ( $Y_i$ ):  $Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$

# Thermal quantities in statistics

- For each particle, the generalized force is  $\frac{\partial \varepsilon_l}{\partial y}$ , the whole force ( $Y_i$ ):  
$$Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$$
$$= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l}$$



# Thermal quantities in statistics

- For each particle, the generalized force is  $\frac{\partial \varepsilon_l}{\partial y}$ , the whole force ( $Y_i$ ):  
$$Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$$
$$= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l} = e^{-\alpha} \sum \omega_l \left(-\frac{1}{\beta}\right) \frac{\partial e^{-\beta \varepsilon_l}}{\partial y}$$

# Thermal quantities in statistics

- For each particle, the generalized force is  $\frac{\partial \varepsilon_l}{\partial y}$ , the whole force ( $Y_i$ ):
$$Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$$
$$= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l} = e^{-\alpha} \sum \omega_l \left(-\frac{1}{\beta}\right) \frac{\partial e^{-\beta \varepsilon_l}}{\partial y}$$
$$= e^{-\alpha} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} \sum \omega_l e^{-\beta \varepsilon_l}$$

# Thermal quantities in statistics

- For each particle, the generalized force is  $\frac{\partial \varepsilon_l}{\partial y}$ , the whole force ( $Y_i$ ):
$$Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$$
$$= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l} = e^{-\alpha} \sum \omega_l \left(-\frac{1}{\beta}\right) \frac{\partial e^{-\beta \varepsilon_l}}{\partial y}$$
$$= e^{-\alpha} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} \sum \omega_l e^{-\beta \varepsilon_l} = \frac{N}{Z_1} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} Z_1$$

# Thermal quantities in statistics

- For each particle, the generalized force is  $\frac{\partial \varepsilon_l}{\partial y}$ , the whole force ( $Y_i$ ):
$$Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$$
$$= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l} = e^{-\alpha} \sum \omega_l \left(-\frac{1}{\beta}\right) \frac{\partial e^{-\beta \varepsilon_l}}{\partial y}$$
$$= e^{-\alpha} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} \sum \omega_l e^{-\beta \varepsilon_l} = \frac{N}{Z_1} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} Z_1 = -\frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y}.$$

# Thermal quantities in statistics

- For each particle, the generalized force is  $\frac{\partial \varepsilon_l}{\partial y}$ , the whole force ( $Y_i$ ): 
$$Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$$
$$= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l} = e^{-\alpha} \sum \omega_l \left(-\frac{1}{\beta}\right) \frac{\partial e^{-\beta \varepsilon_l}}{\partial y}$$
$$= e^{-\alpha} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} \sum \omega_l e^{-\beta \varepsilon_l} = \frac{N}{Z_1} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} Z_1 = -\frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y}.$$
- For  $p$  and  $V$ ,  $\delta W = -p dV$ , so  $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V}$ .

# Thermal quantities in statistics

- For each particle, the generalized force is  $\frac{\partial \varepsilon_l}{\partial y}$ , the whole force ( $Y_i$ ): 
$$Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$$
$$= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l} = e^{-\alpha} \sum \omega_l \left(-\frac{1}{\beta}\right) \frac{\partial e^{-\beta \varepsilon_l}}{\partial y}$$
$$= e^{-\alpha} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} \sum \omega_l e^{-\beta \varepsilon_l} = \frac{N}{Z_1} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} Z_1 = -\frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y}.$$
- For  $p$  and  $V$ ,  $\delta W = -p dV$ , so  $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V}$ .
- As  $d\varepsilon_l = \sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i$ ,

# Thermal quantities in statistics

- For each particle, the generalized force is  $\frac{\partial \varepsilon_l}{\partial y}$ , the whole force ( $Y_i$ ): 
$$Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$$
$$= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l} = e^{-\alpha} \sum \omega_l \left(-\frac{1}{\beta}\right) \frac{\partial e^{-\beta \varepsilon_l}}{\partial y}$$
$$= e^{-\alpha} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} \sum \omega_l e^{-\beta \varepsilon_l} = \frac{N}{Z_1} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} Z_1 = -\frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y}.$$
- For  $p$  and  $V$ ,  $\delta W = -p dV$ , so  $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V}$ .
- As  $d\varepsilon_l = \sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i$ , then  $\delta W = \sum_i Y_i dy_i$

# Thermal quantities in statistics

- For each particle, the generalized force is  $\frac{\partial \varepsilon_l}{\partial y}$ , the whole force ( $Y_i$ ):  $Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$   
 $= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l} = e^{-\alpha} \sum \omega_l \left(-\frac{1}{\beta}\right) \frac{\partial e^{-\beta \varepsilon_l}}{\partial y}$   
 $= e^{-\alpha} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} \sum \omega_l e^{-\beta \varepsilon_l} = \frac{N}{Z_1} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} Z_1 = -\frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y}.$
- For  $p$  and  $V$ ,  $\delta W = -p dV$ , so  $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V}.$
- As  $d\varepsilon_l = \sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i$ , then  $\delta W = \sum_i Y_i dy_i$   
 $= \sum_i dy_i \sum \frac{\partial \varepsilon_l}{\partial y_i} a_l$



# Thermal quantities in statistics

- For each particle, the generalized force is  $\frac{\partial \varepsilon_l}{\partial y}$ , the whole force ( $Y_i$ ): 
$$Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$$
$$= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l} = e^{-\alpha} \sum \omega_l \left(-\frac{1}{\beta}\right) \frac{\partial e^{-\beta \varepsilon_l}}{\partial y}$$
$$= e^{-\alpha} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} \sum \omega_l e^{-\beta \varepsilon_l} = \frac{N}{Z_1} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} Z_1 = -\frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y}.$$
- For  $p$  and  $V$ ,  $\delta W = -p dV$ , so  $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V}$ .
- As  $d\varepsilon_l = \sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i$ , then  $\delta W = \sum_i Y_i dy_i$ 
$$= \sum_i dy_i \sum \frac{\partial \varepsilon_l}{\partial y_i} a_l = \sum \left(\sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i\right) a_l$$

# Thermal quantities in statistics

- For each particle, the generalized force is  $\frac{\partial \varepsilon_l}{\partial y}$ , the whole force ( $Y_i$ ): 
$$Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$$
$$= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l} = e^{-\alpha} \sum \omega_l \left(-\frac{1}{\beta}\right) \frac{\partial e^{-\beta \varepsilon_l}}{\partial y}$$
$$= e^{-\alpha} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} \sum \omega_l e^{-\beta \varepsilon_l} = \frac{N}{Z_1} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} Z_1 = -\frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y}.$$
- For  $p$  and  $V$ ,  $\delta W = -p dV$ , so  $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V}$ .
- As  $d\varepsilon_l = \sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i$ , then  $\delta W = \sum_i Y_i dy_i$ 
$$= \sum_i dy_i \sum \frac{\partial \varepsilon_l}{\partial y_i} a_l = \sum \left(\sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i\right) a_l = \sum a_l d\varepsilon_l.$$

# Thermal quantities in statistics

- For each particle, the generalized force is  $\frac{\partial \varepsilon_l}{\partial y}$ , the whole force ( $Y_i$ ):  $Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$   
 $= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l} = e^{-\alpha} \sum \omega_l \left(-\frac{1}{\beta}\right) \frac{\partial e^{-\beta \varepsilon_l}}{\partial y}$   
 $= e^{-\alpha} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} \sum \omega_l e^{-\beta \varepsilon_l} = \frac{N}{Z_1} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} Z_1 = -\frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y}.$
- For  $p$  and  $V$ ,  $\delta W = -p dV$ , so  $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V}.$
- As  $d\varepsilon_l = \sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i$ , then  $\delta W = \sum_i Y_i dy_i$   
 $= \sum_i dy_i \sum \frac{\partial \varepsilon_l}{\partial y_i} a_l = \sum \left(\sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i\right) a_l = \sum a_l d\varepsilon_l.$
- $\therefore U = \sum \varepsilon_l a_l,$

# Thermal quantities in statistics

- For each particle, the generalized force is  $\frac{\partial \varepsilon_l}{\partial y}$ , the whole force ( $Y_i$ ): 
$$Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$$
$$= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l} = e^{-\alpha} \sum \omega_l \left(-\frac{1}{\beta}\right) \frac{\partial e^{-\beta \varepsilon_l}}{\partial y}$$
$$= e^{-\alpha} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} \sum \omega_l e^{-\beta \varepsilon_l} = \frac{N}{Z_1} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} Z_1 = -\frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y}.$$
- For  $p$  and  $V$ ,  $\delta W = -p dV$ , so  $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V}$ .
- As  $d\varepsilon_l = \sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i$ , then  $\delta W = \sum_i Y_i dy_i$ 
$$= \sum_i dy_i \sum \frac{\partial \varepsilon_l}{\partial y_i} a_l = \sum \left(\sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i\right) a_l = \sum a_l d\varepsilon_l.$$
- $\because U = \sum \varepsilon_l a_l, \therefore dU = \sum a_l d\varepsilon_l + \sum \varepsilon_l da_l$

# Thermal quantities in statistics

- For each particle, the generalized force is  $\frac{\partial \varepsilon_l}{\partial y}$ , the whole force ( $Y_i$ ): 
$$Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$$
$$= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l} = e^{-\alpha} \sum \omega_l \left(-\frac{1}{\beta}\right) \frac{\partial e^{-\beta \varepsilon_l}}{\partial y}$$
$$= e^{-\alpha} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} \sum \omega_l e^{-\beta \varepsilon_l} = \frac{N}{Z_1} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} Z_1 = -\frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y}.$$
- For  $p$  and  $V$ ,  $\delta W = -p dV$ , so  $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V}$ .
- As  $d\varepsilon_l = \sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i$ , then  $\delta W = \sum_i Y_i dy_i$ 
$$= \sum_i dy_i \sum \frac{\partial \varepsilon_l}{\partial y_i} a_l = \sum \left(\sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i\right) a_l = \sum a_l d\varepsilon_l.$$
- $\because U = \sum \varepsilon_l a_l, \therefore dU = \sum a_l d\varepsilon_l + \sum \varepsilon_l da_l$ 
$$= \delta W + \delta Q.$$

# Thermal quantities in statistics

- For each particle, the generalized force is  $\frac{\partial \varepsilon_l}{\partial y}$ , the whole force ( $Y_i$ ): 
$$Y = \sum \frac{\partial \varepsilon_l}{\partial y} a_l = \sum \frac{\partial \varepsilon_l}{\partial y} \omega_l e^{-\alpha - \beta \varepsilon_l}$$
$$= e^{-\alpha} \sum \omega_l \frac{\partial \varepsilon_l}{\partial y} e^{-\beta \varepsilon_l} = e^{-\alpha} \sum \omega_l \left(-\frac{1}{\beta}\right) \frac{\partial e^{-\beta \varepsilon_l}}{\partial y}$$
$$= e^{-\alpha} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} \sum \omega_l e^{-\beta \varepsilon_l} = \frac{N}{Z_1} \left(-\frac{1}{\beta}\right) \frac{\partial}{\partial y} Z_1 = -\frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y}.$$
  - For  $p$  and  $V$ ,  $\delta W = -p dV$ , so  $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V}$ .
  - As  $d\varepsilon_l = \sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i$ , then  $\delta W = \sum_i Y_i dy_i$ 
$$= \sum_i dy_i \sum \frac{\partial \varepsilon_l}{\partial y_i} a_l = \sum \left(\sum_i \frac{\partial \varepsilon_l}{\partial y_i} dy_i\right) a_l = \sum a_l d\varepsilon_l.$$
  - $\because U = \sum \varepsilon_l a_l, \therefore dU = \sum a_l d\varepsilon_l + \sum \varepsilon_l da_l$ 
$$= \delta W + \delta Q.$$
- $\delta W$  changes the energy levels;  $\delta Q$  changes  $\{a_l\}$ .

# Thermal quantities in statistics — $\beta = \frac{1}{kT}$

- Using the partition function,

# Thermal quantities in statistics — $\beta = \frac{1}{kT}$

- Using the partition function,

$$\delta Q = dU - Y dy = -N d\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} dy$$



# Thermal quantities in statistics — $\beta = \frac{1}{kT}$

- Using the partition function,

$$\mathfrak{d}Q = dU - Ydy = -Nd\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} dy$$

$$\beta \mathfrak{d}Q = \beta(dU - Ydy) = -N\beta d\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + N \frac{\partial \ln Z_1}{\partial y} dy$$

# Thermal quantities in statistics — $\beta = \frac{1}{kT}$

- Using the partition function,

$$\mathfrak{d}Q = dU - Ydy = -Nd\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} dy$$

$$\begin{aligned}\beta \mathfrak{d}Q &= \beta(dU - Ydy) = -N\beta d\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + N \frac{\partial \ln Z_1}{\partial y} dy \\ &= \left[N \frac{\partial \ln Z_1}{\partial \beta} d\beta - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right)\right] + N \frac{\partial \ln Z_1}{\partial y} dy\end{aligned}$$

# Thermal quantities in statistics — $\beta = \frac{1}{kT}$

- Using the partition function,

$$\mathfrak{d}Q = dU - Ydy = -Nd\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} dy$$

$$\beta \mathfrak{d}Q = \beta(dU - Ydy) = -N\beta d\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[ N \frac{\partial \ln Z_1}{\partial \beta} d\beta - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right) \right] + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[ N \frac{\partial \ln Z_1}{\partial \beta} d\beta + N \frac{\partial \ln Z_1}{\partial y} dy \right] - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right),$$

# Thermal quantities in statistics — $\beta = \frac{1}{kT}$

- Using the partition function,

$$\mathfrak{d}Q = dU - Ydy = -Nd\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} dy$$

$$\beta \mathfrak{d}Q = \beta(dU - Ydy) = -N\beta d\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[ N \frac{\partial \ln Z_1}{\partial \beta} d\beta - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right) \right] + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[ N \frac{\partial \ln Z_1}{\partial \beta} d\beta + N \frac{\partial \ln Z_1}{\partial y} dy \right] - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right),$$

$$Z_1(\beta, \varepsilon_l), \text{ and } \varepsilon(y), \text{ so, } d \ln Z_1 = \frac{\partial \ln Z_1}{\partial \beta} d\beta + \frac{\partial \ln Z_1}{\partial y} dy.$$

# Thermal quantities in statistics — $\beta = \frac{1}{kT}$

- Using the partition function,

$$\mathfrak{d}Q = dU - Ydy = -Nd\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} dy$$

$$\beta \mathfrak{d}Q = \beta(dU - Ydy) = -N\beta d\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[ N \frac{\partial \ln Z_1}{\partial \beta} d\beta - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right) \right] + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[ N \frac{\partial \ln Z_1}{\partial \beta} d\beta + N \frac{\partial \ln Z_1}{\partial y} dy \right] - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right),$$

$$Z_1(\beta, \varepsilon_l), \text{ and } \varepsilon(y), \text{ so, } d \ln Z_1 = \frac{\partial \ln Z_1}{\partial \beta} d\beta + \frac{\partial \ln Z_1}{\partial y} dy.$$

$$\beta \mathfrak{d}Q = Nd \ln Z_1 - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right)$$

# Thermal quantities in statistics — $\beta = \frac{1}{kT}$

- Using the partition function,

$$\mathfrak{d}Q = dU - Ydy = -Nd\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} dy$$

$$\beta \mathfrak{d}Q = \beta(dU - Ydy) = -N\beta d\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[ N \frac{\partial \ln Z_1}{\partial \beta} d\beta - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right) \right] + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[ N \frac{\partial \ln Z_1}{\partial \beta} d\beta + N \frac{\partial \ln Z_1}{\partial y} dy \right] - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right),$$

$$Z_1(\beta, \varepsilon_l), \text{ and } \varepsilon(y), \text{ so, } d \ln Z_1 = \frac{\partial \ln Z_1}{\partial \beta} d\beta + \frac{\partial \ln Z_1}{\partial y} dy.$$

$$\beta \mathfrak{d}Q = Nd \ln Z_1 - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right)$$

$$= Nd\left(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta}\right),$$

# Thermal quantities in statistics — $\beta = \frac{1}{kT}$

- Using the partition function,

$$\mathfrak{d}Q = dU - Ydy = -Nd\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} dy$$

$$\beta \mathfrak{d}Q = \beta(dU - Ydy) = -N\beta d\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[ N \frac{\partial \ln Z_1}{\partial \beta} d\beta - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right) \right] + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[ N \frac{\partial \ln Z_1}{\partial \beta} d\beta + N \frac{\partial \ln Z_1}{\partial y} dy \right] - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right),$$

$$Z_1(\beta, \varepsilon_l), \text{ and } \varepsilon(y), \text{ so, } d \ln Z_1 = \frac{\partial \ln Z_1}{\partial \beta} d\beta + \frac{\partial \ln Z_1}{\partial y} dy.$$

$$\beta \mathfrak{d}Q = Nd \ln Z_1 - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right)$$

$$= Nd\left(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta}\right), \text{ exact differential.}$$

# Thermal quantities in statistics — $\beta = \frac{1}{kT}$

- Using the partition function,

$$\mathfrak{d}Q = dU - Y dy = -N d\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} dy$$

$$\beta \mathfrak{d}Q = \beta(dU - Y dy) = -N \beta d\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[ N \frac{\partial \ln Z_1}{\partial \beta} d\beta - N d\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right) \right] + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[ N \frac{\partial \ln Z_1}{\partial \beta} d\beta + N \frac{\partial \ln Z_1}{\partial y} dy \right] - N d\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right),$$

$$Z_1(\beta, \varepsilon_l), \text{ and } \varepsilon(y), \text{ so, } d \ln Z_1 = \frac{\partial \ln Z_1}{\partial \beta} d\beta + \frac{\partial \ln Z_1}{\partial y} dy.$$

$$\beta \mathfrak{d}Q = N d \ln Z_1 - N d\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right)$$

$$= N d\left(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta}\right), \text{ exact differential.}$$

- $\frac{1}{T} \mathfrak{d}Q = dS$  is also an exact differential.



# Thermal quantities in statistics — $\beta = \frac{1}{kT}$

- Using the partition function,

$$\mathfrak{d}Q = dU - Ydy = -Nd\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} dy$$

$$\beta \mathfrak{d}Q = \beta(dU - Ydy) = -N\beta d\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[ N \frac{\partial \ln Z_1}{\partial \beta} d\beta - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right) \right] + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[ N \frac{\partial \ln Z_1}{\partial \beta} d\beta + N \frac{\partial \ln Z_1}{\partial y} dy \right] - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right),$$

$$Z_1(\beta, \varepsilon_l), \text{ and } \varepsilon(y), \text{ so, } d \ln Z_1 = \frac{\partial \ln Z_1}{\partial \beta} d\beta + \frac{\partial \ln Z_1}{\partial y} dy.$$

$$\beta \mathfrak{d}Q = Nd \ln Z_1 - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right)$$

$$= Nd\left(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta}\right), \text{ exact differential.}$$

- $\frac{1}{T} \mathfrak{d}Q = dS$  is also an exact differential.
- $\therefore \beta = \frac{1}{kT}$ , where  $k$  is a constant.

# Thermal quantities in statistics — $\beta = \frac{1}{kT}$

- Using the partition function,

$$\mathfrak{d}Q = dU - Ydy = -Nd\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} dy$$

$$\beta \mathfrak{d}Q = \beta(dU - Ydy) = -N\beta d\left(\frac{\partial \ln Z_1}{\partial \beta}\right) + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[ N \frac{\partial \ln Z_1}{\partial \beta} d\beta - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right) \right] + N \frac{\partial \ln Z_1}{\partial y} dy$$

$$= \left[ N \frac{\partial \ln Z_1}{\partial \beta} d\beta + N \frac{\partial \ln Z_1}{\partial y} dy \right] - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right),$$

$$Z_1(\beta, \varepsilon_l), \text{ and } \varepsilon(y), \text{ so, } d \ln Z_1 = \frac{\partial \ln Z_1}{\partial \beta} d\beta + \frac{\partial \ln Z_1}{\partial y} dy.$$

$$\beta \mathfrak{d}Q = Nd \ln Z_1 - Nd\left(\beta \frac{\partial \ln Z_1}{\partial \beta}\right)$$

$$= Nd\left(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta}\right), \text{ exact differential.}$$

- $\frac{1}{T} \mathfrak{d}Q = dS$  is also an exact differential.

- $\therefore \beta = \frac{1}{kT}$ , where  $k$  is a constant.

$k = 1.381 \times 10^{-23} \text{ J} \cdot \text{K}^{-1}$  can be measured.

# Thermal quantities in statistics — entropy

- $dS = \frac{1}{T} dQ$

# Thermal quantities in statistics — entropy

- $dS = \frac{1}{T}dQ = k\beta dQ$

# Thermal quantities in statistics — entropy

- $dS = \frac{1}{T}dQ = k\beta dQ = kN d(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$ .

# Thermal quantities in statistics — entropy

- $dS = \frac{1}{T} \delta Q = k\beta \delta Q = kN d(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta}).$   
 $\Rightarrow S = Nk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta}).$

# Thermal quantities in statistics — entropy

- $dS = \frac{1}{T} \delta Q = k\beta \delta Q = kN d(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$ .  
 $\Rightarrow S = Nk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$ .
- $N = e^{-\alpha} Z_1, \Rightarrow \ln Z_1 = \ln N + \alpha$ .

# Thermal quantities in statistics — entropy

- $dS = \frac{1}{T}dQ = k\beta dQ = kN d(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$ .  
 $\Rightarrow S = Nk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$ .
- $N = e^{-\alpha} Z_1, \Rightarrow \ln Z_1 = \ln N + \alpha$ .
- $S = Nk[\ln N + \alpha - \beta \frac{\partial \ln Z_1}{\partial \beta}]$



# Thermal quantities in statistics — entropy

- $dS = \frac{1}{T} \delta Q = k\beta \delta Q = kN d(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$ .  
 $\Rightarrow S = Nk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$ .
- $N = e^{-\alpha} Z_1, \Rightarrow \ln Z_1 = \ln N + \alpha$ .
- $S = Nk[\ln N + \alpha - \beta \frac{\partial \ln Z_1}{\partial \beta}]$   
 $= k[N \ln N + \alpha N + \beta U]$

# Thermal quantities in statistics — entropy

- $dS = \frac{1}{T} \delta Q = k\beta \delta Q = kN d(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$ .  
 $\Rightarrow S = Nk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$ .
- $N = e^{-\alpha} Z_1, \Rightarrow \ln Z_1 = \ln N + \alpha$ .
- $S = Nk[\ln N + \alpha - \beta \frac{\partial \ln Z_1}{\partial \beta}]$   
 $= k[N \ln N + \alpha N + \beta U] = k[N \ln N + \sum (\alpha + \beta \varepsilon_l) a_l]$

# Thermal quantities in statistics — entropy

- $dS = \frac{1}{T}dQ = k\beta dQ = kN d(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$ .  
 $\Rightarrow S = Nk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$ .
- $N = e^{-\alpha} Z_1, \Rightarrow \ln Z_1 = \ln N + \alpha$ .
- $S = Nk[\ln N + \alpha - \beta \frac{\partial \ln Z_1}{\partial \beta}]$   
 $= k[N \ln N + \alpha N + \beta U] = k[N \ln N + \sum(\alpha + \beta \varepsilon_l) a_l]$
- Boltzmann distribution:  $\ln \frac{\omega_l}{a_l} = \alpha + \beta \varepsilon_l$

# Thermal quantities in statistics — entropy

- $dS = \frac{1}{T}dQ = k\beta dQ = kN d(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$ .  
 $\Rightarrow S = Nk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$ .
- $N = e^{-\alpha} Z_1, \Rightarrow \ln Z_1 = \ln N + \alpha$ .
- $S = Nk[\ln N + \alpha - \beta \frac{\partial \ln Z_1}{\partial \beta}]$   
 $= k[N \ln N + \alpha N + \beta U] = k[N \ln N + \sum (\alpha + \beta \varepsilon_l) a_l]$
- Boltzmann distribution:  $\ln \frac{\omega_l}{a_l} = \alpha + \beta \varepsilon_l$
- $\therefore S = k[N \ln N + \sum \ln \frac{\omega_l}{a_l} a_l]$

# Thermal quantities in statistics — entropy

- $dS = \frac{1}{T}dQ = k\beta dQ = kNd(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$ .  
 $\Rightarrow S = Nk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$ .
- $N = e^{-\alpha} Z_1, \Rightarrow \ln Z_1 = \ln N + \alpha$ .
- $S = Nk[\ln N + \alpha - \beta \frac{\partial \ln Z_1}{\partial \beta}]$   
 $= k[N \ln N + \alpha N + \beta U] = k[N \ln N + \sum(\alpha + \beta \varepsilon_l) a_l]$
- Boltzmann distribution:  $\ln \frac{\omega_l}{a_l} = \alpha + \beta \varepsilon_l$
- $\therefore S = k[N \ln N + \sum \ln \frac{\omega_l}{a_l} a_l] = k \ln \Omega_M$ .

# Thermal quantities in statistics — entropy

- $dS = \frac{1}{T} \delta Q = k\beta \delta Q = kN d(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$ .  
 $\Rightarrow S = Nk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$ .
- $N = e^{-\alpha} Z_1, \Rightarrow \ln Z_1 = \ln N + \alpha$ .
- $S = Nk[\ln N + \alpha - \beta \frac{\partial \ln Z_1}{\partial \beta}]$   
 $= k[N \ln N + \alpha N + \beta U] = k[N \ln N + \sum (\alpha + \beta \varepsilon_l) a_l]$
- Boltzmann distribution:  $\ln \frac{\omega_l}{a_l} = \alpha + \beta \varepsilon_l$
- $\therefore S = k[N \ln N + \sum \ln \frac{\omega_l}{a_l} a_l] = k \ln \Omega_M$ .
- Entropy statistical meaning: number of micro-states.

# Thermal quantities in statistics — entropy

- $dS = \frac{1}{T} \delta Q = k\beta \delta Q = kN d(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$ .  
 $\Rightarrow S = Nk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$ .
- $N = e^{-\alpha} Z_1, \Rightarrow \ln Z_1 = \ln N + \alpha$ .
- $S = Nk[\ln N + \alpha - \beta \frac{\partial \ln Z_1}{\partial \beta}]$   
 $= k[N \ln N + \alpha N + \beta U] = k[N \ln N + \sum (\alpha + \beta \varepsilon_l) a_l]$
- Boltzmann distribution:  $\ln \frac{\omega_l}{a_l} = \alpha + \beta \varepsilon_l$
- $\therefore S = k[N \ln N + \sum \ln \frac{\omega_l}{a_l} a_l] = k \ln \Omega_M$ .
- Entropy statistical meaning: number of micro-states.
- $F = U - TS$

# Thermal quantities in statistics — entropy

- $dS = \frac{1}{T}dQ = k\beta dQ = kN d(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$ .  
 $\Rightarrow S = Nk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$ .
- $N = e^{-\alpha} Z_1, \Rightarrow \ln Z_1 = \ln N + \alpha$ .
- $S = Nk[\ln N + \alpha - \beta \frac{\partial \ln Z_1}{\partial \beta}]$   
 $= k[N \ln N + \alpha N + \beta U] = k[N \ln N + \sum(\alpha + \beta \varepsilon_l) a_l]$
- Boltzmann distribution:  $\ln \frac{\omega_l}{a_l} = \alpha + \beta \varepsilon_l$
- $\therefore S = k[N \ln N + \sum \ln \frac{\omega_l}{a_l} a_l] = k \ln \Omega_M$ .
- Entropy statistical meaning: number of micro-states.
- $F = U - TS = -N \frac{\partial \ln Z_1}{\partial \beta} - TNk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$



# Thermal quantities in statistics — entropy

- $dS = \frac{1}{T} \delta Q = k\beta \delta Q = kN d(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$ .  
 $\Rightarrow S = Nk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$ .
- $N = e^{-\alpha} Z_1, \Rightarrow \ln Z_1 = \ln N + \alpha$ .
- $S = Nk[\ln N + \alpha - \beta \frac{\partial \ln Z_1}{\partial \beta}]$   
 $= k[N \ln N + \alpha N + \beta U] = k[N \ln N + \sum (\alpha + \beta \varepsilon_l) a_l]$
- Boltzmann distribution:  $\ln \frac{\omega_l}{a_l} = \alpha + \beta \varepsilon_l$
- $\therefore S = k[N \ln N + \sum \ln \frac{\omega_l}{a_l} a_l] = k \ln \Omega_M$ .
- Entropy statistical meaning: number of micro-states.
- $F = U - TS = -N \frac{\partial \ln Z_1}{\partial \beta} - TNk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$   
 $= -NkT \ln Z_1$ .

# Thermal quantities in statistics — entropy

- $dS = \frac{1}{T}dQ = k\beta dQ = kN d(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$ .  
 $\Rightarrow S = Nk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$ .
- $N = e^{-\alpha} Z_1, \Rightarrow \ln Z_1 = \ln N + \alpha$ .
- $S = Nk[\ln N + \alpha - \beta \frac{\partial \ln Z_1}{\partial \beta}]$   
 $= k[N \ln N + \alpha N + \beta U] = k[N \ln N + \sum(\alpha + \beta \varepsilon_l) a_l]$
- Boltzmann distribution:  $\ln \frac{\omega_l}{a_l} = \alpha + \beta \varepsilon_l$
- $\therefore S = k[N \ln N + \sum \ln \frac{\omega_l}{a_l} a_l] = k \ln \Omega_M$ .
- Entropy statistical meaning: number of micro-states.
- $F = U - TS = -N \frac{\partial \ln Z_1}{\partial \beta} - TNk(\ln Z_1 - \beta \frac{\partial \ln Z_1}{\partial \beta})$   
 $= -NkT \ln Z_1$ .
- Conclusion: Given the partition function  
 $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$ , thermal variables are determined.

# Thermal quantities in statistics

- For classical particle system,  $\omega_l \rightarrow \frac{\Delta\omega_l}{h_0^r}$ ,

# Thermal quantities in statistics

- For classical particle system,  $\omega_l \rightarrow \frac{\Delta\omega_l}{h_0^r}$ ,  
the partition function:  $Z_1 = \sum e^{-\beta\epsilon_l} \frac{\Delta\omega_l}{h_0^r}$ .

# Thermal quantities in statistics

- For classical particle system,  $\omega_l \rightarrow \frac{\Delta\omega_l}{h_0^r}$ ,  
the partition function:  $Z_1 = \sum e^{-\beta\epsilon_l} \frac{\Delta\omega_l}{h_0^r}$ .
- For continuous  $(q, p)$ ,  $\Delta\omega_l \rightarrow d\omega = dq_1 \dots dq_r dp_1 \dots dp_r$ ,

# Thermal quantities in statistics

- For classical particle system,  $\omega_l \rightarrow \frac{\Delta\omega_l}{h_0^r}$ ,  
the partition function:  $Z_1 = \sum e^{-\beta\epsilon_l} \frac{\Delta\omega_l}{h_0^r}$ .
- For continuous  $(q, p)$ ,  $\Delta\omega_l \rightarrow d\omega = dq_1 \dots dq_r dp_1 \dots dp_r$ ,  
the partition function:  $Z_1 = \int e^{-\beta\epsilon} \frac{d\omega}{h_0^r}$

# Thermal quantities in statistics

- For classical particle system,  $\omega_l \rightarrow \frac{\Delta\omega_l}{h_0^r}$ ,  
the partition function:  $Z_1 = \sum e^{-\beta\epsilon_l} \frac{\Delta\omega_l}{h_0^r}$ .
- For continuous  $(q, p)$ ,  $\Delta\omega_l \rightarrow d\omega = dq_1 \dots dq_r dp_1 \dots dp_r$ ,  
the partition function:  $Z_1 = \int e^{-\beta\epsilon} \frac{d\omega}{h_0^r}$   
 $= \int \dots \int e^{-\beta\epsilon(p,q)} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r}$ .

# Table of contents

## 1 §7 Boltzmann Statistics

- 7.1 Thermal quantities in statistics
- 7.2 Equation of state of ideal gas
- 7.3 Maxwell speed distribution
- 7.4 Energy equipartition theorem
- 7.5 Internal energy and heat capacity of the ideal gas
- 7.6 Entropy of the ideal gas
- 7.7 Einstein's theory on heat capacity of solid
- 7.8 Paramagnetic solid
- 7.9 Negative temperature



## 7.2 Equation of state of ideal gas

- A simple application of Boltzmann statistics.

## 7.2 Equation of state of ideal gas

- A simple application of Boltzmann statistics.
- Object: ideal gas with monatomic molecule, nearly independent.

## 7.2 Equation of state of ideal gas

- A simple application of Boltzmann statistics.
- Object: ideal gas with monatomic molecule, nearly independent.
- Energy of each particle:  $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$ .

## 7.2 Equation of state of ideal gas

- A simple application of Boltzmann statistics.
- Object: ideal gas with monatomic molecule, nearly independent.
- Energy of each particle:  $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$ .
- Micro-states number:  $\frac{dx dy dz dp_x dp_y dp_z}{h^3}$ .

## 7.2 Equation of state of ideal gas

- A simple application of Boltzmann statistics.
- Object: ideal gas with monatomic molecule, nearly independent.
- Energy of each particle:  $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$ .
- Micro-states number:  $\frac{dx dy dz dp_x dp_y dp_z}{h^3}$ .
- Partition function:

$$Z_1 = \frac{1}{h^3} \int \dots \int e^{-\beta \varepsilon_l} dx dy dz dp_x dp_y dp_z$$

## 7.2 Equation of state of ideal gas

- A simple application of Boltzmann statistics.
- Object: ideal gas with monatomic molecule, nearly independent.
- Energy of each particle:  $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$ .
- Micro-states number:  $\frac{dx dy dz dp_x dp_y dp_z}{h^3}$ .
- Partition function:

$$\begin{aligned} Z_1 &= \frac{1}{h^3} \int \dots \int e^{-\beta \varepsilon_l} dx dy dz dp_x dp_y dp_z \\ &= \frac{1}{h^3} \iiint dx dy dz \iiint e^{-\frac{\beta}{2m}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z \end{aligned}$$

## 7.2 Equation of state of ideal gas

- A simple application of Boltzmann statistics.
- Object: ideal gas with monatomic molecule, nearly independent.
- Energy of each particle:  $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$ .
- Micro-states number:  $\frac{dx dy dz dp_x dp_y dp_z}{h^3}$ .
- Partition function:

$$\begin{aligned}
 Z_1 &= \frac{1}{h^3} \int \dots \int e^{-\beta \varepsilon_l} dx dy dz dp_x dp_y dp_z \\
 &= \frac{1}{h^3} \iiint dx dy dz \iiint e^{-\frac{\beta}{2m}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z \\
 &= \frac{V}{h^3} \int_{-\infty}^{+\infty} e^{-\frac{\beta}{2m}p_x^2} dp_x \int_{-\infty}^{+\infty} e^{-\frac{\beta}{2m}p_y^2} dp_y \int_{-\infty}^{+\infty} e^{-\frac{\beta}{2m}p_z^2} dp_z
 \end{aligned}$$

## 7.2 Equation of state of ideal gas

- A simple application of Boltzmann statistics.
- Object: ideal gas with monatomic molecule, nearly independent.
- Energy of each particle:  $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$ .
- Micro-states number:  $\frac{dx dy dz dp_x dp_y dp_z}{h^3}$ .
- Partition function:

$$\begin{aligned}
 Z_1 &= \frac{1}{h^3} \int \dots \int e^{-\beta \varepsilon_l} dx dy dz dp_x dp_y dp_z \\
 &= \frac{1}{h^3} \iiint dx dy dz \iiint e^{-\frac{\beta}{2m}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z \\
 &= \frac{V}{h^3} \int_{-\infty}^{+\infty} e^{-\frac{\beta}{2m}p_x^2} dp_x \int_{-\infty}^{+\infty} e^{-\frac{\beta}{2m}p_y^2} dp_y \int_{-\infty}^{+\infty} e^{-\frac{\beta}{2m}p_z^2} dp_z \\
 &= \left(\frac{2\pi m}{h^2 \beta}\right)^{\frac{3}{2}} V.
 \end{aligned}$$



# Equation of state of ideal gas

- $$p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y}$$

# Equation of state of ideal gas

- $$p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V}$$

# Equation of state of ideal gas

$$\bullet p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln \left[ \left( \frac{2\pi m}{h^2 \beta} \right)^{\frac{3}{2}} V \right]}{\partial V}$$

# Equation of state of ideal gas

$$\bullet p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V}$$

# Equation of state of ideal gas

$$\bullet p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V} = \frac{NkT}{V}.$$

# Equation of state of ideal gas

- $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V} = \frac{NkT}{V}.$
- Comparing with  $pV = nRT$ ,  $k = \frac{n}{N}R$  is determined.

# Equation of state of ideal gas

- $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V} = \frac{NkT}{V}.$
- Comparing with  $pV = nRT$ ,  $k = \frac{n}{N}R$  is determined.
- Classical limit condition:  $\frac{\omega_l}{a_l} \gg 1$ , or  $e^\alpha \gg 1$ .

# Equation of state of ideal gas

- $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V} = \frac{NkT}{V}.$
- Comparing with  $pV = nRT$ ,  $k = \frac{n}{N}R$  is determined.
- Classical limit condition:  $\frac{\omega_l}{a_l} \gg 1$ , or  $e^\alpha \gg 1$ .  
$$e^\alpha = \frac{Z_1}{N}$$



# Equation of state of ideal gas

- $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V} = \frac{NkT}{V}.$
- Comparing with  $pV = nRT$ ,  $k = \frac{n}{N}R$  is determined.
- Classical limit condition:  $\frac{\omega_l}{a_l} \gg 1$ , or  $e^\alpha \gg 1$ .  
$$e^\alpha = \frac{Z_1}{N} = \frac{V}{N} \left( \frac{2\pi m}{h^2 \beta} \right)^{\frac{3}{2}}$$

# Equation of state of ideal gas

- $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V} = \frac{NkT}{V}.$
- Comparing with  $pV = nRT$ ,  $k = \frac{n}{N}R$  is determined.
- Classical limit condition:  $\frac{\omega_l}{a_l} \gg 1$ , or  $e^\alpha \gg 1$ .  
$$e^\alpha = \frac{Z_1}{N} = \frac{V}{N} \left( \frac{2\pi m}{h^2 \beta} \right)^{\frac{3}{2}} = \frac{V}{N} \left( \frac{2\pi m kT}{h^2} \right)^{\frac{3}{2}} \gg 1.$$

# Equation of state of ideal gas

- $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V} = \frac{NkT}{V}.$
- Comparing with  $pV = nRT$ ,  $k = \frac{n}{N}R$  is determined.
- Classical limit condition:  $\frac{\omega_l}{a_l} \gg 1$ , or  $e^\alpha \gg 1$ .  

$$e^\alpha = \frac{Z_1}{N} = \frac{V}{N} \left( \frac{2\pi m}{h^2 \beta} \right)^{\frac{3}{2}} = \frac{V}{N} \left( \frac{2\pi m kT}{h^2} \right)^{\frac{3}{2}} \gg 1.$$
 I.e., 1.  $N/V \rightarrow 0$ , rare;

# Equation of state of ideal gas

- $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V} = \frac{NkT}{V}.$
- Comparing with  $pV = nRT$ ,  $k = \frac{n}{N}R$  is determined.
- Classical limit condition:  $\frac{\omega_l}{a_l} \gg 1$ , or  $e^\alpha \gg 1$ .  

$$e^\alpha = \frac{Z_1}{N} = \frac{V}{N} \left( \frac{2\pi m}{h^2 \beta} \right)^{\frac{3}{2}} = \frac{V}{N} \left( \frac{2\pi m kT}{h^2} \right)^{\frac{3}{2}} \gg 1.$$
 I.e., 1.  $N/V \rightarrow 0$ , rare; 2. high temperature;

# Equation of state of ideal gas

- $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V} = \frac{NkT}{V}.$
- Comparing with  $pV = nRT$ ,  $k = \frac{n}{N}R$  is determined.
- Classical limit condition:  $\frac{\omega_l}{a_l} \gg 1$ , or  $e^\alpha \gg 1$ .  

$$e^\alpha = \frac{Z_1}{N} = \frac{V}{N} \left( \frac{2\pi m}{h^2 \beta} \right)^{\frac{3}{2}} = \frac{V}{N} \left( \frac{2\pi m kT}{h^2} \right)^{\frac{3}{2}} \gg 1.$$

i.e., 1.  $N/V \rightarrow 0$ , rare; 2. high temperature; 3. high mass of the molecule.

# Equation of state of ideal gas

- $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V} = \frac{NkT}{V}.$
- Comparing with  $pV = nRT$ ,  $k = \frac{n}{N}R$  is determined.
- Classical limit condition:  $\frac{\omega_l}{a_l} \gg 1$ , or  $e^\alpha \gg 1$ .  

$$e^\alpha = \frac{Z_1}{N} = \frac{V}{N} \left( \frac{2\pi m}{h^2 \beta} \right)^{\frac{3}{2}} = \frac{V}{N} \left( \frac{2\pi m kT}{h^2} \right)^{\frac{3}{2}} \gg 1.$$

i.e., 1.  $N/V \rightarrow 0$ , rare; 2. high temperature; 3. high mass of the molecule.
- de Broglie wavelength:  $\lambda = \frac{h}{p}$

# Equation of state of ideal gas

- $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V} = \frac{NkT}{V}.$
- Comparing with  $pV = nRT$ ,  $k = \frac{n}{N}R$  is determined.
- Classical limit condition:  $\frac{\omega_l}{a_l} \gg 1$ , or  $e^\alpha \gg 1$ .  

$$e^\alpha = \frac{Z_1}{N} = \frac{V}{N} \left(\frac{2\pi m}{h^2 \beta}\right)^{\frac{3}{2}} = \frac{V}{N} \left(\frac{2\pi m kT}{h^2}\right)^{\frac{3}{2}} \gg 1.$$

i.e., 1.  $N/V \rightarrow 0$ , rare; 2. high temperature; 3. high mass of the molecule.
- de Broglie wavelength:  $\lambda = \frac{h}{p} = \frac{h}{\sqrt{2m\varepsilon}}$

# Equation of state of ideal gas

- $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V} = \frac{NkT}{V}.$
- Comparing with  $pV = nRT$ ,  $k = \frac{n}{N}R$  is determined.
- Classical limit condition:  $\frac{\omega_l}{a_l} \gg 1$ , or  $e^\alpha \gg 1$ .  

$$e^\alpha = \frac{Z_1}{N} = \frac{V}{N} \left( \frac{2\pi m}{h^2 \beta} \right)^{\frac{3}{2}} = \frac{V}{N} \left( \frac{2\pi mkT}{h^2} \right)^{\frac{3}{2}} \gg 1.$$

i.e., 1.  $N/V \rightarrow 0$ , rare; 2. high temperature; 3. high mass of the molecule.
- de Broglie wavelength:  $\lambda = \frac{h}{p} = \frac{h}{\sqrt{2m\varepsilon}} \simeq \frac{h}{\sqrt{2\pi mkT}}.$



# Equation of state of ideal gas

- $p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V} = \frac{NkT}{V}$ .
- Comparing with  $pV = nRT$ ,  $k = \frac{n}{N}R$  is determined.
- Classical limit condition:  $\frac{\omega_l}{a_l} \gg 1$ , or  $e^\alpha \gg 1$ .  

$$e^\alpha = \frac{Z_1}{N} = \frac{V}{N} \left(\frac{2\pi m}{h^2 \beta}\right)^{\frac{3}{2}} = \frac{V}{N} \left(\frac{2\pi m kT}{h^2}\right)^{\frac{3}{2}} \gg 1.$$

I.e., 1.  $N/V \rightarrow 0$ , rare; 2. high temperature; 3. high mass of the molecule.
- de Broglie wavelength:  $\lambda = \frac{h}{p} = \frac{h}{\sqrt{2m\varepsilon}} \simeq \frac{h}{\sqrt{2\pi m kT}}$ .  
 So,  $e^\alpha \gg 1 \Rightarrow \frac{1}{n} \lambda^{-3} \gg 1$ , or  $n\lambda^3 \ll 1$ , or  $\lambda \ll \left(\frac{1}{n}\right)^{\frac{1}{3}}$ .

# Equation of state of ideal gas

$$\bullet p = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial y} = \frac{N}{\beta} \frac{\partial \ln Z_1}{\partial V} = \frac{N}{\beta} \frac{\partial \ln[(\frac{2\pi m}{h^2 \beta})^{\frac{3}{2}} V]}{\partial V} = \frac{N}{\beta} \frac{1}{V} = \frac{NkT}{V}.$$

$$\bullet \text{Comparing with } pV = nRT, k = \frac{n}{N}R \text{ is determined.}$$

$$\bullet \text{Classical limit condition: } \frac{\omega_l}{a_l} \gg 1, \text{ or } e^\alpha \gg 1.$$

$$e^\alpha = \frac{Z_1}{N} = \frac{V}{N} \left( \frac{2\pi m}{h^2 \beta} \right)^{\frac{3}{2}} = \frac{V}{N} \left( \frac{2\pi mkT}{h^2} \right)^{\frac{3}{2}} \gg 1.$$

I.e., 1.  $N/V \rightarrow 0$ , rare; 2. high temperature; 3. high mass of the molecule.

$$\bullet \text{de Broglie wavelength: } \lambda = \frac{h}{p} = \frac{h}{\sqrt{2m\varepsilon}} \simeq \frac{h}{\sqrt{2\pi mkT}}.$$

$$\text{So, } e^\alpha \gg 1 \Rightarrow \frac{1}{n} \lambda^{-3} \gg 1, \text{ or } n\lambda^3 \ll 1, \text{ or } \lambda \ll \left(\frac{1}{n}\right)^{\frac{1}{3}}.$$

Means, the wavelength should be much less than the distance of any two molecules. (Can be distinguished.)

# Table of contents

## 1 §7 Boltzmann Statistics

- 7.1 Thermal quantities in statistics
- 7.2 Equation of state of ideal gas
- 7.3 Maxwell speed distribution
- 7.4 Energy equipartition theorem
- 7.5 Internal energy and heat capacity of the ideal gas
- 7.6 Entropy of the ideal gas
- 7.7 Einstein's theory on heat capacity of solid
- 7.8 Paramagnetic solid
- 7.9 Negative temperature

## 7.3 Maxwell speed distribution

## 7.3 Maxwell speed distribution

- Object: gas. Obeys classical limit,

## 7.3 Maxwell speed distribution

- Object: gas. Obeys classical limit,  
so Boltzmann distribution:  $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$ .

## 7.3 Maxwell speed distribution

- Object: gas. Obeys classical limit,  
so Boltzmann distribution:  $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$ .
- Simple case: monatomic molecules.

## 7.3 Maxwell speed distribution

- Object: gas. Obeys classical limit,  
so Boltzmann distribution:  $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$ .
- Simple case: monatomic molecules.
- Energy:  $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$ ;



## 7.3 Maxwell speed distribution

- Object: gas. Obeys classical limit,  
so Boltzmann distribution:  $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$ .
- Simple case: monatomic molecules.
- Energy:  $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$ ;  
Degeneracy:  $\omega = \frac{V}{h^3} dp_x dp_y dp_z$ .

## 7.3 Maxwell speed distribution

- Object: gas. Obeys classical limit,  
so Boltzmann distribution:  $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$ .
- Simple case: monatomic molecules.
- Energy:  $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$ ;  
Degeneracy:  $\omega = \frac{V}{h^3} dp_x dp_y dp_z$ .
- $a = \frac{V}{h^3} e^{-\alpha - \frac{1}{2mkT}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z$ .

## 7.3 Maxwell speed distribution

- Object: gas. Obeys classical limit,  
so Boltzmann distribution:  $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$ .
- Simple case: monatomic molecules.
- Energy:  $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$ ;  
Degeneracy:  $\omega = \frac{V}{h^3} dp_x dp_y dp_z$ .
- $a = \frac{V}{h^3} e^{-\alpha - \frac{1}{2mkT}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z$ .
- $N = \sum a_l$

## 7.3 Maxwell speed distribution

- Object: gas. Obeys classical limit,  
so Boltzmann distribution:  $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$ .
- Simple case: monatomic molecules.
- Energy:  $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$ ;  
Degeneracy:  $\omega = \frac{V}{h^3} dp_x dp_y dp_z$ .
- $a = \frac{V}{h^3} e^{-\alpha - \frac{1}{2mkT}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z$ .
- $N = \sum a_l = \frac{V}{h^3} \iiint e^{-\alpha - \frac{1}{2mkT}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z$

## 7.3 Maxwell speed distribution

- Object: gas. Obeys classical limit,  
so Boltzmann distribution:  $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$ .

- Simple case: monatomic molecules.

- Energy:  $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$ ;

Degeneracy:  $\omega = \frac{V}{h^3} dp_x dp_y dp_z$ .

- $a = \frac{V}{h^3} e^{-\alpha - \frac{1}{2mkT}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z$ .

- $N = \sum a_l = \frac{V}{h^3} \iiint e^{-\alpha - \frac{1}{2mkT}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z =$   
 $e^{-\alpha} \frac{V}{h^3} (2mkT)^{\frac{3}{2}} \int e^{-\frac{p_x^2}{2mkT}} d\frac{p_x}{\sqrt{2mkT}} \int e^{-\frac{p_y^2}{2mkT}} d\frac{p_y}{\sqrt{2mkT}} \int e^{-\frac{p_z^2}{2mkT}} d\frac{p_z}{\sqrt{2mkT}}$

## 7.3 Maxwell speed distribution

- Object: gas. Obeys classical limit,  
so Boltzmann distribution:  $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$ .

- Simple case: monatomic molecules.

- Energy:  $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$ ;

Degeneracy:  $\omega = \frac{V}{h^3} dp_x dp_y dp_z$ .

- $a = \frac{V}{h^3} e^{-\alpha - \frac{1}{2mkT}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z$ .

- $$N = \sum a_l = \frac{V}{h^3} \iiint e^{-\alpha - \frac{1}{2mkT}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z =$$

$$e^{-\alpha} \frac{V}{h^3} (2mkT)^{\frac{3}{2}} \int e^{-\frac{p_x^2}{2mkT}} d\frac{p_x}{\sqrt{2mkT}} \int e^{-\frac{p_y^2}{2mkT}} d\frac{p_y}{\sqrt{2mkT}} \int e^{-\frac{p_z^2}{2mkT}} d\frac{p_z}{\sqrt{2mkT}}$$

$$= e^{-\alpha} \frac{V}{h^3} (2mkT)^{\frac{3}{2}} (\sqrt{\pi})^3$$

## 7.3 Maxwell speed distribution

- Object: gas. Obeys classical limit,  
so Boltzmann distribution:  $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$ .

- Simple case: monatomic molecules.

- Energy:  $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$ ;

Degeneracy:  $\omega = \frac{V}{h^3} dp_x dp_y dp_z$ .

- $a = \frac{V}{h^3} e^{-\alpha - \frac{1}{2mkT}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z$ .

- $$N = \sum a_l = \frac{V}{h^3} \iiint e^{-\alpha - \frac{1}{2mkT}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z =$$

$$e^{-\alpha} \frac{V}{h^3} (2mkT)^{\frac{3}{2}} \int e^{-\frac{p_x^2}{2mkT}} d\frac{p_x}{\sqrt{2mkT}} \int e^{-\frac{p_y^2}{2mkT}} d\frac{p_y}{\sqrt{2mkT}} \int e^{-\frac{p_z^2}{2mkT}} d\frac{p_z}{\sqrt{2mkT}}$$

$$= e^{-\alpha} \frac{V}{h^3} (2mkT)^{\frac{3}{2}} (\sqrt{\pi})^3 \Rightarrow e^{-\alpha} = \frac{N}{V} \left( \frac{h^2}{2\pi mkT} \right)^{\frac{3}{2}}.$$

## 7.3 Maxwell speed distribution

- Object: gas. Obeys classical limit,  
so Boltzmann distribution:  $a_l = \omega_l e^{-\alpha - \beta \varepsilon_l}$ .
- Simple case: monatomic molecules.
- Energy:  $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$ ;  
Degeneracy:  $\omega = \frac{V}{h^3} dp_x dp_y dp_z$ .
- $a = \frac{V}{h^3} e^{-\alpha - \frac{1}{2mkT}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z$ .
- $N = \sum a_l = \frac{V}{h^3} \iiint e^{-\alpha - \frac{1}{2mkT}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z =$   

$$e^{-\alpha} \frac{V}{h^3} (2mkT)^{\frac{3}{2}} \int e^{-\frac{p_x^2}{2mkT}} d\frac{p_x}{\sqrt{2mkT}} \int e^{-\frac{p_y^2}{2mkT}} d\frac{p_y}{\sqrt{2mkT}} \int e^{-\frac{p_z^2}{2mkT}} d\frac{p_z}{\sqrt{2mkT}}$$

$$= e^{-\alpha} \frac{V}{h^3} (2mkT)^{\frac{3}{2}} (\sqrt{\pi})^3 \Rightarrow e^{-\alpha} = \frac{N}{V} \left( \frac{h^2}{2\pi mkT} \right)^{\frac{3}{2}}.$$
- $\therefore a = N \left( \frac{1}{2\pi mkT} \right)^{\frac{3}{2}} e^{-\frac{1}{2mkT}(p_x^2 + p_y^2 + p_z^2)} dp_x dp_y dp_z$ .



# Maxwell speed distribution

- Turn to velocity space:  $v_x = \frac{p_x}{m}$ ,

# Maxwell speed distribution

- Turn to velocity space:  $v_x = \frac{p_x}{m}$ ,  
 $a \equiv f(v_x, v_y, v_z) dv_x dv_y dv_z$   
 $= n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z.$

# Maxwell speed distribution

- Turn to velocity space:  $v_x = \frac{p_x}{m}$ ,  
 $a \equiv f(v_x, v_y, v_z) dv_x dv_y dv_z$   
 $= n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z.$   
 $f(v_x, v_y, v_z)$  is the Maxwell velocity distribution.

# Maxwell speed distribution

- Turn to velocity space:  $v_x = \frac{p_x}{m}$ ,

$$\begin{aligned} a &\equiv \int f(v_x, v_y, v_z) dv_x dv_y dv_z \\ &= n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z. \end{aligned}$$

$f(v_x, v_y, v_z)$  is the Maxwell velocity distribution.

Obeys  $\iiint f(v_x, v_y, v_z) dv_x dv_y dv_z = n$ .

# Maxwell speed distribution

- Turn to velocity space:  $v_x = \frac{p_x}{m}$ ,  
 $a \equiv f(v_x, v_y, v_z)dv_x dv_y dv_z$   
 $= n\left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z.$   
 $f(v_x, v_y, v_z)$  is the Maxwell velocity distribution.  
Obeys  $\iiint f(v_x, v_y, v_z)dv_x dv_y dv_z = n.$
- Convert to spheric polar coordinates  $(v, \theta, \varphi)$ ,  
 $dv_x dv_y dv_z = v^2 \sin \theta dv d\theta d\varphi.$

# Maxwell speed distribution

- Turn to velocity space:  $v_x = \frac{p_x}{m}$ ,  
 $a \equiv f(v_x, v_y, v_z)dv_x dv_y dv_z$   
 $= n\left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z.$   
 $f(v_x, v_y, v_z)$  is the Maxwell velocity distribution.  
 Obeys  $\iiint f(v_x, v_y, v_z)dv_x dv_y dv_z = n.$
- Convert to spheric polar coordinates  $(v, \theta, \varphi)$ ,  
 $dv_x dv_y dv_z = v^2 \sin \theta dv d\theta d\varphi.$   
 For isotropic velocity,  $\int_0^\pi \sin \theta d\theta \int_0^{2\pi} d\varphi = 4\pi,$

# Maxwell speed distribution

- Turn to velocity space:  $v_x = \frac{p_x}{m}$ ,

$$a \equiv \int f(v_x, v_y, v_z) dv_x dv_y dv_z$$

$$= n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z.$$

$f(v_x, v_y, v_z)$  is the Maxwell velocity distribution.

Obeys  $\iiint f(v_x, v_y, v_z) dv_x dv_y dv_z = n$ .

- Convert to spheric polar coordinates  $(v, \theta, \varphi)$ ,

$$dv_x dv_y dv_z = v^2 \sin \theta dv d\theta d\varphi.$$

For isotropic velocity,  $\int_0^\pi \sin \theta d\theta \int_0^{2\pi} d\varphi = 4\pi$ , the number of molecules:

$$f(v) dv = 4\pi n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2 dv.$$

# Maxwell speed distribution

- Turn to velocity space:  $v_x = \frac{p_x}{m}$ ,

$$a \equiv \int f(v_x, v_y, v_z) dv_x dv_y dv_z$$

$$= n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z.$$

$f(v_x, v_y, v_z)$  is the Maxwell velocity distribution.

Obeys  $\iiint f(v_x, v_y, v_z) dv_x dv_y dv_z = n$ .

- Convert to spheric polar coordinates  $(v, \theta, \varphi)$ ,

$$dv_x dv_y dv_z = v^2 \sin \theta dv d\theta d\varphi.$$

For isotropic velocity,  $\int_0^\pi \sin \theta d\theta \int_0^{2\pi} d\varphi = 4\pi$ , the number of molecules:

$$f(v) dv = 4\pi n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2 dv.$$

Maxwell speed distribution. Obeys:

$$\int 4\pi n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2 dv = n.$$



## Maxwell speed distribution

$$f(v) = 4\pi n \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2$$

- The most probable speed  $v_m$ :  $\frac{df(v)}{dv} = 0$ .

## Maxwell speed distribution

$$f(v) = 4\pi n \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2$$

- The most probable speed  $v_m$ :  $\frac{df(v)}{dv} = 0$ .

$$\frac{d(e^{-\frac{m}{2kT}v^2} v^2)}{dv} = e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{kT}\right) \cdot 2v \cdot v^2 + e^{-\frac{m}{2kT}v^2} \cdot 2v$$

## Maxwell speed distribution

$$f(v) = 4\pi n \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2$$

- The most probable speed  $v_m$ :  $\frac{df(v)}{dv} = 0$ .

$$\begin{aligned}\frac{d(e^{-\frac{m}{2kT}v^2} v^2)}{dv} &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{kT}\right) \cdot 2v \cdot v^2 + e^{-\frac{m}{2kT}v^2} \cdot 2v \\ &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{kT}v^2 + 1\right) \cdot 2v\end{aligned}$$

## Maxwell speed distribution

$$f(v) = 4\pi n \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2$$

- The most probable speed  $v_m$ :  $\frac{df(v)}{dv} = 0$ .

$$\begin{aligned}\frac{d(e^{-\frac{m}{2kT}v^2} v^2)}{dv} &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{kT}\right) \cdot 2v \cdot v^2 + e^{-\frac{m}{2kT}v^2} \cdot 2v \\ &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{kT}v^2 + 1\right) \cdot 2v = 0,\end{aligned}$$

## Maxwell speed distribution

$$f(v) = 4\pi n \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2$$

- The most probable speed  $v_m$ :  $\frac{df(v)}{dv} = 0$ .

$$\begin{aligned}\frac{d(e^{-\frac{m}{2kT}v^2} v^2)}{dv} &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{kT}\right) \cdot 2v \cdot v^2 + e^{-\frac{m}{2kT}v^2} \cdot 2v \\ &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{kT}v^2 + 1\right) \cdot 2v = 0, \Rightarrow v_m^2 = \frac{2kT}{m}.\end{aligned}$$

## Maxwell speed distribution

$$f(v) = 4\pi n \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2$$

- The most probable speed  $v_m$ :  $\frac{df(v)}{dv} = 0$ .

$$\begin{aligned}\frac{d(e^{-\frac{m}{2kT}v^2} v^2)}{dv} &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2kT}\right) \cdot 2v \cdot v^2 + e^{-\frac{m}{2kT}v^2} \cdot 2v \\ &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2kT}v^2 + 1\right) \cdot 2v = 0, \Rightarrow v_m^2 = \frac{2kT}{m}.\end{aligned}$$

- Average speed  $\bar{v} = \int v f(v) / n dv$

## Maxwell speed distribution

$$f(v) = 4\pi n \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2$$

- The most probable speed  $v_m$ :  $\frac{df(v)}{dv} = 0$ .

$$\begin{aligned} \frac{d(e^{-\frac{m}{2kT}v^2} v^2)}{dv} &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2kT}\right) \cdot 2v \cdot v^2 + e^{-\frac{m}{2kT}v^2} \cdot 2v \\ &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2\pi kT}v^2 + 1\right) \cdot 2v = 0, \Rightarrow v_m^2 = \frac{2kT}{m}. \end{aligned}$$

- Average speed  $\bar{v} = \int v f(v) / n dv$
- $$= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \int_0^\infty v e^{-\frac{m}{2kT}v^2} v^2 dv$$

## Maxwell speed distribution

$$f(v) = 4\pi n \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2$$

- The most probable speed  $v_m$ :  $\frac{df(v)}{dv} = 0$ .

$$\begin{aligned}\frac{d(e^{-\frac{m}{2kT}v^2} v^2)}{dv} &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2kT}\right) \cdot 2v \cdot v^2 + e^{-\frac{m}{2kT}v^2} \cdot 2v \\ &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2\pi kT}v^2 + 1\right) \cdot 2v = 0, \Rightarrow v_m^2 = \frac{2kT}{m}.\end{aligned}$$

- Average speed  $\bar{v} = \int v f(v) / n dv$

$$\begin{aligned}&= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \int_0^\infty v e^{-\frac{m}{2kT}v^2} v^2 dv \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \int_0^\infty e^{-\frac{m}{2kT}v^2} v^2 dv^2\end{aligned}$$



## Maxwell speed distribution

$$f(v) = 4\pi n \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2$$

- The most probable speed  $v_m$ :  $\frac{df(v)}{dv} = 0$ .

$$\begin{aligned} \frac{d(e^{-\frac{m}{2kT}v^2} v^2)}{dv} &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2kT}\right) \cdot 2v \cdot v^2 + e^{-\frac{m}{2kT}v^2} \cdot 2v \\ &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2\pi kT}v^2 + 1\right) \cdot 2v = 0, \Rightarrow v_m^2 = \frac{2kT}{m}. \end{aligned}$$

- Average speed  $\bar{v} = \int v f(v) / n dv$

$$\begin{aligned} &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \int_0^\infty v e^{-\frac{m}{2kT}v^2} v^2 dv \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \int_0^\infty e^{-\frac{m}{2kT}v^2} v^2 dv^2 \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(-\frac{2kT}{m}\right) \int_0^\infty v^2 de^{-\frac{m}{2kT}v^2} \end{aligned}$$

## Maxwell speed distribution

$$f(v) = 4\pi n \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2$$

- The most probable speed  $v_m$ :  $\frac{df(v)}{dv} = 0$ .

$$\begin{aligned} \frac{d(e^{-\frac{m}{2kT}v^2} v^2)}{dv} &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2kT}\right) \cdot 2v \cdot v^2 + e^{-\frac{m}{2kT}v^2} \cdot 2v \\ &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2kT}v^2 + 1\right) \cdot 2v = 0, \Rightarrow v_m^2 = \frac{2kT}{m}. \end{aligned}$$

- Average speed  $\bar{v} = \int v f(v) / n dv$

$$\begin{aligned} &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \int_0^\infty v e^{-\frac{m}{2kT}v^2} v^2 dv \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \int_0^\infty e^{-\frac{m}{2kT}v^2} v^2 dv^2 \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(-\frac{2kT}{m}\right) \int_0^\infty v^2 de^{-\frac{m}{2kT}v^2} \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(-\frac{2kT}{m}\right) [(v^2 e^{-\frac{m}{2kT}v^2})|_0^\infty - \int_0^\infty e^{-\frac{m}{2kT}v^2} dv^2] \end{aligned}$$

## Maxwell speed distribution

$$f(v) = 4\pi n \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2$$

- The most probable speed  $v_m$ :  $\frac{df(v)}{dv} = 0$ .

$$\begin{aligned} \frac{d(e^{-\frac{m}{2kT}v^2} v^2)}{dv} &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2kT}\right) \cdot 2v \cdot v^2 + e^{-\frac{m}{2kT}v^2} \cdot 2v \\ &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2\pi kT}v^2 + 1\right) \cdot 2v = 0, \Rightarrow v_m^2 = \frac{2kT}{m}. \end{aligned}$$

- Average speed  $\bar{v} = \int v f(v) / n dv$

$$\begin{aligned} &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \int_0^\infty v e^{-\frac{m}{2kT}v^2} v^2 dv \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \int_0^\infty e^{-\frac{m}{2kT}v^2} v^2 dv^2 \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(-\frac{2kT}{m}\right) \int_0^\infty v^2 de^{-\frac{m}{2kT}v^2} \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(-\frac{2kT}{m}\right) [(v^2 e^{-\frac{m}{2kT}v^2})|_0^\infty - \int_0^\infty e^{-\frac{m}{2kT}v^2} dv^2] \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(\frac{2kT}{m}\right) \int_0^\infty e^{-\frac{m}{2kT}v^2} dv^2 \end{aligned}$$

## Maxwell speed distribution

$$f(v) = 4\pi n \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2$$

- The most probable speed  $v_m$ :  $\frac{df(v)}{dv} = 0$ .

$$\begin{aligned} \frac{d(e^{-\frac{m}{2kT}v^2} v^2)}{dv} &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2kT}\right) \cdot 2v \cdot v^2 + e^{-\frac{m}{2kT}v^2} \cdot 2v \\ &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2\pi kT}v^2 + 1\right) \cdot 2v = 0, \Rightarrow v_m^2 = \frac{2kT}{m}. \end{aligned}$$

- Average speed  $\bar{v} = \int v f(v) / n dv$

$$\begin{aligned} &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \int_0^\infty v e^{-\frac{m}{2kT}v^2} v^2 dv \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \int_0^\infty e^{-\frac{m}{2kT}v^2} v^2 dv^2 \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(-\frac{2kT}{m}\right) \int_0^\infty v^2 de^{-\frac{m}{2kT}v^2} \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(-\frac{2kT}{m}\right) [(v^2 e^{-\frac{m}{2kT}v^2})|_0^\infty - \int_0^\infty e^{-\frac{m}{2kT}v^2} dv^2] \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(\frac{2kT}{m}\right) \int_0^\infty e^{-\frac{m}{2kT}v^2} dv^2 \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(\frac{2kT}{m}\right)^2 \int_0^\infty e^{-\frac{m}{2kT}v^2} d\left(\frac{m}{2kT}v^2\right) \end{aligned}$$

## Maxwell speed distribution

$$f(v) = 4\pi n \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2$$

- The most probable speed  $v_m$ :  $\frac{df(v)}{dv} = 0$ .

$$\begin{aligned} \frac{d(e^{-\frac{m}{2kT}v^2} v^2)}{dv} &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2kT}\right) \cdot 2v \cdot v^2 + e^{-\frac{m}{2kT}v^2} \cdot 2v \\ &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2\pi kT}v^2 + 1\right) \cdot 2v = 0, \Rightarrow v_m^2 = \frac{2kT}{m}. \end{aligned}$$

- Average speed  $\bar{v} = \int v f(v) / n dv$

$$\begin{aligned} &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \int_0^\infty v e^{-\frac{m}{2kT}v^2} v^2 dv \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \int_0^\infty e^{-\frac{m}{2kT}v^2} v^2 dv^2 \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(-\frac{2kT}{m}\right) \int_0^\infty v^2 de^{-\frac{m}{2kT}v^2} \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(-\frac{2kT}{m}\right) [(v^2 e^{-\frac{m}{2kT}v^2})|_0^\infty - \int_0^\infty e^{-\frac{m}{2kT}v^2} dv^2] \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(\frac{2kT}{m}\right) \int_0^\infty e^{-\frac{m}{2kT}v^2} dv^2 \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(\frac{2kT}{m}\right)^2 \int_0^\infty e^{-\frac{m}{2kT}v^2} d\left(\frac{m}{2kT}v^2\right) \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(\frac{2kT}{m}\right)^2 \int_0^\infty e^{-u} du \end{aligned}$$

## Maxwell speed distribution

$$f(v) = 4\pi n \left(\frac{m}{2kT}\right)^{\frac{3}{2}} e^{-\frac{m}{2kT}v^2} v^2$$

- The most probable speed  $v_m$ :  $\frac{df(v)}{dv} = 0$ .

$$\begin{aligned} \frac{d(e^{-\frac{m}{2kT}v^2} v^2)}{dv} &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2kT}\right) \cdot 2v \cdot v^2 + e^{-\frac{m}{2kT}v^2} \cdot 2v \\ &= e^{-\frac{m}{2kT}v^2} \left(-\frac{m}{2\pi kT}v^2 + 1\right) \cdot 2v = 0, \Rightarrow v_m^2 = \frac{2kT}{m}. \end{aligned}$$

- Average speed  $\bar{v} = \int v f(v) / n dv$

$$\begin{aligned} &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \int_0^\infty v e^{-\frac{m}{2kT}v^2} v^2 dv \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \int_0^\infty e^{-\frac{m}{2kT}v^2} v^2 dv^2 \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(-\frac{2kT}{m}\right) \int_0^\infty v^2 de^{-\frac{m}{2kT}v^2} \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(-\frac{2kT}{m}\right) [(v^2 e^{-\frac{m}{2kT}v^2})|_0^\infty - \int_0^\infty e^{-\frac{m}{2kT}v^2} dv^2] \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(\frac{2kT}{m}\right) \int_0^\infty e^{-\frac{m}{2kT}v^2} dv^2 \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(\frac{2kT}{m}\right)^2 \int_0^\infty e^{-\frac{m}{2kT}v^2} d\left(\frac{m}{2kT}v^2\right) \\ &= 4\pi \left(\frac{m}{2\pi kT}\right)^{\frac{3}{2}} \cdot \frac{1}{2} \left(\frac{2kT}{m}\right)^2 = \sqrt{\frac{8kT}{\pi m}}. \end{aligned}$$

# Maxwell speed distribution

- Root-mean-square speed:

# Maxwell speed distribution

- Root-mean-square speed:

$$v_s^2 \equiv \overline{v^2}$$



# Maxwell speed distribution

- Root-mean-square speed:

$$v_s^2 \equiv \bar{v^2} = \int v^2 f(v) / n dv$$

# Maxwell speed distribution

- Root-mean-square speed:

$$v_s^2 \equiv \bar{v^2} = \int v^2 f(v) / n dv = \frac{3kT}{m}.$$

# Maxwell speed distribution

- Root-mean-square speed:

$$v_s^2 \equiv \bar{v^2} = \int v^2 f(v) / n dv = \frac{3kT}{m}.$$

Useful in calculating average energy  $\bar{\varepsilon} = \frac{1}{2} m \bar{v^2}$ .

# Maxwell speed distribution

- Root-mean-square speed:

$$v_s^2 \equiv \bar{v^2} = \int v^2 f(v) / n dv = \frac{3kT}{m}.$$

Useful in calculating average energy  $\bar{\varepsilon} = \frac{1}{2} m \bar{v^2}$ .

Example:  $N_2$  at  $0^\circ\text{C}$ .

# Maxwell speed distribution

- Root-mean-square speed:

$$v_s^2 \equiv \bar{v^2} = \int v^2 f(v) / n dv = \frac{3kT}{m}.$$

Useful in calculating average energy  $\bar{\varepsilon} = \frac{1}{2} m \bar{v^2}$ .

Example:  $N_2$  at  $0^\circ\text{C}$ .  $m = 28 \times 1.67 \times 10^{-27} \text{kg}$ ,

# Maxwell speed distribution

- Root-mean-square speed:

$$v_s^2 \equiv \bar{v^2} = \int v^2 f(v) n dv = \frac{3kT}{m}.$$

Useful in calculating average energy  $\bar{\varepsilon} = \frac{1}{2} m \bar{v^2}$ .

Example:  $N_2$  at  $0^\circ\text{C}$ .  $m = 28 \times 1.67 \times 10^{-27} \text{kg}$ ,

$$v_s = \left( \frac{3kT}{m} \right)^{\frac{1}{2}}$$

# Maxwell speed distribution

- Root-mean-square speed:

$$v_s^2 \equiv \bar{v^2} = \int v^2 f(v) / n dv = \frac{3kT}{m}.$$

Useful in calculating average energy  $\bar{\epsilon} = \frac{1}{2} m \bar{v^2}$ .

Example:  $N_2$  at  $0^\circ\text{C}$ .  $m = 28 \times 1.67 \times 10^{-27} \text{kg}$ ,

$$v_s = \left( \frac{3kT}{m} \right)^{\frac{1}{2}} = \left( \frac{3 \cdot 1.38 \times 10^{-23} \text{JK}^{-1} \cdot 273 \text{K}}{28 \times 1.67 \times 10^{-27} \text{kg}} \right)^{\frac{1}{2}}$$

# Maxwell speed distribution

- Root-mean-square speed:

$$v_s^2 \equiv \bar{v^2} = \int v^2 f(v) / n dv = \frac{3kT}{m}.$$

Useful in calculating average energy  $\bar{\epsilon} = \frac{1}{2} m \bar{v^2}$ .

Example:  $N_2$  at  $0^\circ\text{C}$ .  $m = 28 \times 1.67 \times 10^{-27} \text{kg}$ ,

$$v_s = \left( \frac{3kT}{m} \right)^{\frac{1}{2}} = \left( \frac{3 \cdot 1.38 \times 10^{-23} \text{JK}^{-1} \cdot 273 \text{K}}{28 \times 1.67 \times 10^{-27} \text{kg}} \right)^{\frac{1}{2}} \simeq 493 \text{m/s}.$$



# Maxwell speed distribution

- Root-mean-square speed:

$$v_s^2 \equiv \bar{v^2} = \int v^2 f(v) / n dv = \frac{3kT}{m}.$$


Useful in calculating average energy  $\bar{\epsilon} = \frac{1}{2} m \bar{v^2}$ .

Example:  $N_2$  at  $0^\circ\text{C}$ .  $m = 28 \times 1.67 \times 10^{-27} \text{kg}$ ,

$$v_s = \left( \frac{3kT}{m} \right)^{\frac{1}{2}} = \left( \frac{3 \cdot 1.38 \times 10^{-23} \text{JK}^{-1} \cdot 273 \text{K}}{28 \times 1.67 \times 10^{-27} \text{kg}} \right)^{\frac{1}{2}} \simeq 493 \text{m/s}.$$

Application 1: frequency of colliding.

$\Gamma$ : Number of colliding molecules on the surface in unit time unit area.



# Maxwell speed distribution

- Root-mean-square speed:

$$v_s^2 \equiv \bar{v^2} = \int v^2 f(v) / n dv = \frac{3kT}{m}.$$


Useful in calculating average energy  $\bar{\varepsilon} = \frac{1}{2} m \bar{v^2}$ .

Example:  $N_2$  at  $0^\circ\text{C}$ .  $m = 28 \times 1.67 \times 10^{-27} \text{kg}$ ,

$$v_s = \left( \frac{3kT}{m} \right)^{\frac{1}{2}} = \left( \frac{3 \cdot 1.38 \times 10^{-23} \text{JK}^{-1} \cdot 273 \text{K}}{28 \times 1.67 \times 10^{-27} \text{kg}} \right)^{\frac{1}{2}} \simeq 493 \text{m/s}.$$

Application 1: frequency of colliding.

$\Gamma$ : Number of colliding molecules on the surface in unit time unit area.



For a given velocity,  $(v_x, v_y, v_z)$ , all particles inside the oblique cylinder can collide.

# Maxwell speed distribution

- Root-mean-square speed:

$$v_s^2 \equiv \bar{v^2} = \int v^2 f(v) / n dv = \frac{3kT}{m}.$$


Useful in calculating average energy  $\bar{\varepsilon} = \frac{1}{2} m \bar{v^2}$ .

Example:  $N_2$  at  $0^\circ\text{C}$ .  $m = 28 \times 1.67 \times 10^{-27} \text{kg}$ ,

$$v_s = \left( \frac{3kT}{m} \right)^{\frac{1}{2}} = \left( \frac{3 \cdot 1.38 \times 10^{-23} \text{JK}^{-1} \cdot 273 \text{K}}{28 \times 1.67 \times 10^{-27} \text{kg}} \right)^{\frac{1}{2}} \simeq 493 \text{m/s}.$$

Application 1: frequency of colliding.

$\Gamma$ : Number of colliding molecules on the surface in unit time unit area.



For a given velocity,  $(v_x, v_y, v_z)$ , all particles inside the oblique cylinder can collide. The number is

$$f(v_x, v_y, v_z) dv_x dv_y dv_z \cdot v_x dt dA.$$

# Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- In unit time unit area, all velocity:

$$\Gamma = \iiint f(v_x, v_y, v_z) v_x dv_x dv_y dv_z$$

# Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- In unit time unit area, all velocity:

$$\begin{aligned}\Gamma &= \iiint f(v_x, v_y, v_z) v_x dv_x dv_y dv_z \\ &= n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} \iiint v_x e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z\end{aligned}$$

# Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- In unit time unit area, all velocity:

$$\begin{aligned} \Gamma &= \iiint f(v_x, v_y, v_z) v_x dv_x dv_y dv_z \\ &= n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} \iiint v_x e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z \\ &= n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} \int_0^\infty v_x e^{-\frac{m}{2kT}v_x^2} dv_x \iint_{-\infty}^\infty e^{-\frac{m}{2kT}(v_y^2 + v_z^2)} dv_y dv_z \end{aligned}$$

# Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- In unit time unit area, all velocity:

$$\begin{aligned} \Gamma &= \iiint f(v_x, v_y, v_z) v_x dv_x dv_y dv_z \\ &= n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} \iiint v_x e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z \\ &= n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} \int_0^\infty v_x e^{-\frac{m}{2kT}v_x^2} dv_x \iint_{-\infty}^\infty e^{-\frac{m}{2kT}(v_y^2 + v_z^2)} dv_y dv_z \\ &= n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} \cdot \left( \frac{1}{2} \frac{2kT}{m} \right) \cdot \left( \frac{2kT}{m} \pi \right) \end{aligned}$$

# Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- In unit time unit area, all velocity:

$$\begin{aligned} \Gamma &= \iiint f(v_x, v_y, v_z) v_x dv_x dv_y dv_z \\ &= n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} \iiint v_x e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z \\ &= n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} \int_0^\infty v_x e^{-\frac{m}{2kT}v_x^2} dv_x \iint_{-\infty}^\infty e^{-\frac{m}{2kT}(v_y^2 + v_z^2)} dv_y dv_z \\ &= n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} \cdot \left( \frac{1}{2} \frac{2kT}{m} \right) \cdot \left( \frac{2kT}{m} \pi \right) \\ &= n \sqrt{\frac{kT}{2\pi m}} \end{aligned}$$



# Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- In unit time unit area, all velocity:

$$\begin{aligned} \Gamma &= \iiint f(v_x, v_y, v_z) v_x dv_x dv_y dv_z \\ &= n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} \iiint v_x e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z \\ &= n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} \int_0^\infty v_x e^{-\frac{m}{2kT}v_x^2} dv_x \iint_{-\infty}^\infty e^{-\frac{m}{2kT}(v_y^2 + v_z^2)} dv_y dv_z \\ &= n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} \cdot \left( \frac{1}{2} \frac{2kT}{m} \right) \cdot \left( \frac{2kT}{m} \pi \right) \\ &= n \sqrt{\frac{kT}{2\pi m}} \\ &= \frac{1}{4} n \bar{v}. \end{aligned}$$

# Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- Application 2: Pressure.

# Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- Application 2: Pressure.

$$p = \frac{F}{A}$$

# Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- Application 2: Pressure.

$$p = \frac{F}{A} = \frac{\Delta p_x}{\Delta t A}$$

# Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- Application 2: Pressure.

$$p = \frac{F}{A} = \frac{\Delta p_x}{\Delta t A} = \frac{\sum m v_x \cdot 2}{dA dt}$$

# Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- Application 2: Pressure.

$$p = \frac{F}{A} = \frac{\Delta p_x}{\Delta t A} = \frac{\sum m v_x \cdot 2}{dA dt}$$

Number of particles  $f(v_x, v_y, v_z) v_x dv_x dv_y dv_z dA dt$ .

# Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- Application 2: Pressure.

$$p = \frac{F}{A} = \frac{\Delta p_x}{\Delta t A} = \frac{\sum m v_x \cdot 2}{dA dt}$$

Number of particles  $f(v_x, v_y, v_z) v_x dv_x dv_y dv_z dA dt$ .

- $\therefore p = \iiint f(v_x, v_y, v_z) v_x \cdot 2m v_x dv_x dv_y dv_z$

# Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- Application 2: Pressure.

$$p = \frac{F}{A} = \frac{\Delta p_x}{\Delta t A} = \frac{\sum m v_x \cdot 2}{dA dt}$$

Number of particles  $f(v_x, v_y, v_z) v_x dv_x dv_y dv_z dA dt$ .

- $\therefore p = \iiint f(v_x, v_y, v_z) v_x \cdot 2m v_x dv_x dv_y dv_z$   
 $= n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} \iiint 2m v_x^2 e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z$



# Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- Application 2: Pressure.

$$p = \frac{F}{A} = \frac{\Delta p_x}{\Delta t A} = \frac{\sum m v_x \cdot 2}{dA dt}$$

Number of particles  $f(v_x, v_y, v_z) v_x dv_x dv_y dv_z dA dt$ .

$$\begin{aligned} \bullet \therefore p &= \iiint f(v_x, v_y, v_z) v_x \cdot 2m v_x dv_x dv_y dv_z \\ &= n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} \iiint 2m v_x^2 e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z \\ &= 2mn \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} \left( \frac{2kT}{m} \right)^{\frac{3}{2}} \int_0^\infty \left( \frac{m}{2kT} v_x^2 \right) e^{-\frac{m}{2kT} v_x^2} d\left( \sqrt{\frac{m}{2kT}} v_x \right) \cdot \\ &\quad \frac{2kT}{m} \iint_{-\infty}^\infty e^{-\frac{m}{2kT}(v_y^2 + v_z^2)} d\left( \sqrt{\frac{m}{2kT}} v_y \right) d\left( \sqrt{\frac{m}{2kT}} v_z \right) \end{aligned}$$

# Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- Application 2: Pressure.

$$p = \frac{F}{A} = \frac{\Delta p_x}{\Delta t A} = \frac{\sum m v_x \cdot 2}{dA dt}$$

Number of particles  $f(v_x, v_y, v_z) v_x dv_x dv_y dv_z dA dt$ .

- $\therefore p = \iiint f(v_x, v_y, v_z) v_x \cdot 2m v_x dv_x dv_y dv_z$   
 $= n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} \iiint 2m v_x^2 e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z$   
 $= 2mn \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} \left( \frac{2kT}{m} \right)^{\frac{3}{2}} \int_0^\infty \left( \frac{m}{2kT} v_x^2 \right) e^{-\frac{m}{2kT} v_x^2} d\left( \sqrt{\frac{m}{2kT}} v_x \right) \cdot$   
 $\frac{2kT}{m} \int_{-\infty}^\infty e^{-\frac{m}{2kT}(v_y^2 + v_z^2)} d\left( \sqrt{\frac{m}{2kT}} v_y \right) d\left( \sqrt{\frac{m}{2kT}} v_z \right)$   
 $= 2mn \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} \left( \frac{2kT}{m} \right)^{\frac{3}{2}} \cdot \frac{\sqrt{\pi}}{4} \cdot \frac{2kT}{m} \pi$

# Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- Application 2: Pressure.

$$p = \frac{F}{A} = \frac{\Delta p_x}{\Delta t A} = \frac{\sum m v_x \cdot 2}{dA dt}$$

Number of particles  $f(v_x, v_y, v_z) v_x dv_x dv_y dv_z dA dt$ .

- $\therefore p = \iiint f(v_x, v_y, v_z) v_x \cdot 2m v_x dv_x dv_y dv_z$   
 $= n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} \iiint 2m v_x^2 e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z$   
 $= 2mn \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} \left( \frac{2kT}{m} \right)^{\frac{3}{2}} \int_0^\infty \left( \frac{m}{2kT} v_x^2 \right) e^{-\frac{m}{2kT} v_x^2} d\left( \sqrt{\frac{m}{2kT}} v_x \right) \cdot$   
 $\frac{2kT}{m} \iint_{-\infty}^\infty e^{-\frac{m}{2kT}(v_y^2 + v_z^2)} d\left( \sqrt{\frac{m}{2kT}} v_y \right) d\left( \sqrt{\frac{m}{2kT}} v_z \right)$   
 $= 2mn \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} \left( \frac{2kT}{m} \right)^{\frac{3}{2}} \cdot \frac{\sqrt{\pi}}{4} \cdot \frac{2kT}{m} \pi$   
 $= nkT.$

# Maxwell speed distribution

$$f(v_x, v_y, v_z) = n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)}$$

- Application 2: Pressure.

$$p = \frac{F}{A} = \frac{\Delta p_x}{\Delta t A} = \frac{\sum m v_x \cdot 2}{dA dt}$$

Number of particles  $f(v_x, v_y, v_z) v_x dv_x dv_y dv_z dA dt$ .

$$\begin{aligned} \bullet \therefore p &= \iiint f(v_x, v_y, v_z) v_x \cdot 2m v_x dv_x dv_y dv_z \\ &= n \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} \iiint 2m v_x^2 e^{-\frac{m}{2kT}(v_x^2 + v_y^2 + v_z^2)} dv_x dv_y dv_z \\ &= 2mn \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} \left( \frac{2kT}{m} \right)^{\frac{3}{2}} \int_0^\infty \left( \frac{m}{2kT} v_x^2 \right) e^{-\frac{m}{2kT} v_x^2} d\left( \sqrt{\frac{m}{2kT}} v_x \right) \cdot \\ &\quad \frac{2kT}{m} \iint_{-\infty}^\infty e^{-\frac{m}{2kT}(v_y^2 + v_z^2)} d\left( \sqrt{\frac{m}{2kT}} v_y \right) d\left( \sqrt{\frac{m}{2kT}} v_z \right) \\ &= 2mn \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} \left( \frac{2kT}{m} \right)^{\frac{3}{2}} \cdot \frac{\sqrt{\pi}}{4} \cdot \frac{2kT}{m} \pi \\ &= nkT. \end{aligned}$$

Same as eq (7.2.5).

# Table of contents

## 1 §7 Boltzmann Statistics

- 7.1 Thermal quantities in statistics
- 7.2 Equation of state of ideal gas
- 7.3 Maxwell speed distribution
- 7.4 Energy equipartition theorem
- 7.5 Internal energy and heat capacity of the ideal gas
- 7.6 Entropy of the ideal gas
- 7.7 Einstein's theory on heat capacity of solid
- 7.8 Paramagnetic solid
- 7.9 Negative temperature

## 7.4 Energy equipartition theorem

## 7.4 Energy equipartition theorem

- Theorem: any degree of freedom  $x$  which appears in the total energy  $\varepsilon$  only as a simple quadratic term, has an average energy of  $\frac{1}{2}kT$  in thermal equilibrium.

## 7.4 Energy equipartition theorem

- Theorem: any degree of freedom  $x$  which appears in the total energy  $\varepsilon$  only as a simple quadratic term, has an average energy of  $\frac{1}{2}kT$  in thermal equilibrium.
- Example: monatomic ideal gas.



## 7.4 Energy equipartition theorem

- Theorem: any degree of freedom  $x$  which appears in the total energy  $\varepsilon$  only as a simple quadratic term, has an average energy of  $\frac{1}{2}kT$  in thermal equilibrium.
- Example: monatomic ideal gas.  
$$\bar{\varepsilon} = \overline{\frac{1}{2}mv^2} = \overline{\frac{1}{2}m(v_x^2 + v_y^2 + v_z^2)}$$

## 7.4 Energy equipartition theorem

- Theorem: any degree of freedom  $x$  which appears in the total energy  $\varepsilon$  only as a simple quadratic term, has an average energy of  $\frac{1}{2}kT$  in thermal equilibrium.
- Example: monatomic ideal gas.

$$\begin{aligned}\bar{\varepsilon} &= \overline{\frac{1}{2}mv^2} = \overline{\frac{1}{2}m(v_x^2 + v_y^2 + v_z^2)} \\ &= \frac{1}{2}m\overline{v^2}\end{aligned}$$

## 7.4 Energy equipartition theorem

- Theorem: any degree of freedom  $x$  which appears in the total energy  $\varepsilon$  only as a simple quadratic term, has an average energy of  $\frac{1}{2}kT$  in thermal equilibrium.
- Example: monatomic ideal gas.

$$\begin{aligned}\bar{\varepsilon} &= \overline{\frac{1}{2}mv^2} = \overline{\frac{1}{2}m(v_x^2 + v_y^2 + v_z^2)} \\ &= \frac{1}{2}m\overline{v^2} = \frac{1}{2}m \cdot \frac{3kT}{m}\end{aligned}$$

## 7.4 Energy equipartition theorem

- Theorem: any degree of freedom  $x$  which appears in the total energy  $\varepsilon$  only as a simple quadratic term, has an average energy of  $\frac{1}{2}kT$  in thermal equilibrium.
- Example: monatomic ideal gas.

$$\begin{aligned}\bar{\varepsilon} &= \overline{\frac{1}{2}mv^2} = \overline{\frac{1}{2}m(v_x^2 + v_y^2 + v_z^2)} \\ &= \frac{1}{2}m\overline{v^2} = \frac{1}{2}m \cdot \frac{3kT}{m} = \frac{3}{2}kT.\end{aligned}$$

## 7.4 Energy equipartition theorem

- Theorem: any degree of freedom  $x$  which appears in the total energy  $\varepsilon$  only as a simple quadratic term, has an average energy of  $\frac{1}{2}kT$  in thermal equilibrium.
- Example: monatomic ideal gas.

$$\begin{aligned}\bar{\varepsilon} &= \overline{\frac{1}{2}mv^2} = \overline{\frac{1}{2}m(v_x^2 + v_y^2 + v_z^2)} \\ &= \frac{1}{2}m\overline{v^2} = \frac{1}{2}m \cdot \frac{3kT}{m} = \frac{3}{2}kT.\end{aligned}$$

- Proof:  $\varepsilon = \varepsilon_k + \varepsilon_p$ ,

## 7.4 Energy equipartition theorem

- Theorem: any degree of freedom  $x$  which appears in the total energy  $\varepsilon$  only as a simple quadratic term, has an average energy of  $\frac{1}{2}kT$  in thermal equilibrium.
- Example: monatomic ideal gas.

$$\begin{aligned}\bar{\varepsilon} &= \overline{\frac{1}{2}mv^2} = \overline{\frac{1}{2}m(v_x^2 + v_y^2 + v_z^2)} \\ &= \frac{1}{2}m\overline{v^2} = \frac{1}{2}m \cdot \frac{3kT}{m} = \frac{3}{2}kT.\end{aligned}$$

- Proof:  $\varepsilon = \varepsilon_k + \varepsilon_p$ , where  $\varepsilon_k = \frac{1}{2} \sum a_i p_i^2$ .

## 7.4 Energy equipartition theorem

- Theorem: any degree of freedom  $x$  which appears in the total energy  $\varepsilon$  only as a simple quadratic term, has an average energy of  $\frac{1}{2}kT$  in thermal equilibrium.
- Example: monatomic ideal gas.

$$\begin{aligned}\bar{\varepsilon} &= \overline{\frac{1}{2}mv^2} = \overline{\frac{1}{2}m(v_x^2 + v_y^2 + v_z^2)} \\ &= \frac{1}{2}m\overline{v^2} = \frac{1}{2}m \cdot \frac{3kT}{m} = \frac{3}{2}kT.\end{aligned}$$

- Proof:  $\varepsilon = \varepsilon_k + \varepsilon_p$ , where  $\varepsilon_k = \frac{1}{2} \sum a_i p_i^2$ .

Considering the 1st term,

$$\overline{\frac{1}{2}a_1 p_1^2}$$

## 7.4 Energy equipartition theorem

- Theorem: any degree of freedom  $x$  which appears in the total energy  $\varepsilon$  only as a simple quadratic term, has an average energy of  $\frac{1}{2}kT$  in thermal equilibrium.
- Example: monatomic ideal gas.

$$\begin{aligned}\bar{\varepsilon} &= \overline{\frac{1}{2}mv^2} = \overline{\frac{1}{2}m(v_x^2 + v_y^2 + v_z^2)} \\ &= \frac{1}{2}m\overline{v^2} = \frac{1}{2}m \cdot \frac{3kT}{m} = \frac{3}{2}kT.\end{aligned}$$

- Proof:  $\varepsilon = \varepsilon_k + \varepsilon_p$ , where  $\varepsilon_k = \frac{1}{2} \sum a_i p_i^2$ .

Considering the 1st term,

$$\overline{\frac{1}{2}a_1 p_1^2} = \frac{1}{N} \sum \frac{1}{2}a_1 p_1^2 \cdot a_l$$



## 7.4 Energy equipartition theorem

- Theorem: any degree of freedom  $x$  which appears in the total energy  $\varepsilon$  only as a simple quadratic term, has an average energy of  $\frac{1}{2}kT$  in thermal equilibrium.
- Example: monatomic ideal gas.

$$\begin{aligned}\bar{\varepsilon} &= \overline{\frac{1}{2}mv^2} = \overline{\frac{1}{2}m(v_x^2 + v_y^2 + v_z^2)} \\ &= \frac{1}{2}m\overline{v^2} = \frac{1}{2}m \cdot \frac{3kT}{m} = \frac{3}{2}kT.\end{aligned}$$

- Proof:  $\varepsilon = \varepsilon_k + \varepsilon_p$ , where  $\varepsilon_k = \frac{1}{2} \sum a_i p_i^2$ .

Considering the 1st term,

$$\begin{aligned}\overline{\frac{1}{2}a_1 p_1^2} &= \frac{1}{N} \sum \frac{1}{2}a_1 p_1^2 \cdot a_l \\ &= \frac{1}{N} \sum \frac{1}{2}a_1 p_1^2 \cdot \omega_l e^{-\alpha - \beta \varepsilon_l}\end{aligned}$$

## 7.4 Energy equipartition theorem

- Theorem: any degree of freedom  $x$  which appears in the total energy  $\varepsilon$  only as a simple quadratic term, has an average energy of  $\frac{1}{2}kT$  in thermal equilibrium.
- Example: monatomic ideal gas.

$$\begin{aligned}\bar{\varepsilon} &= \overline{\frac{1}{2}mv^2} = \overline{\frac{1}{2}m(v_x^2 + v_y^2 + v_z^2)} \\ &= \frac{1}{2}m\overline{v^2} = \frac{1}{2}m \cdot \frac{3kT}{m} = \frac{3}{2}kT.\end{aligned}$$

- Proof:  $\varepsilon = \varepsilon_k + \varepsilon_p$ , where  $\varepsilon_k = \frac{1}{2} \sum a_i p_i^2$ .

Considering the 1st term,

$$\begin{aligned}\overline{\frac{1}{2}a_1 p_1^2} &= \frac{1}{N} \sum \frac{1}{2}a_1 p_1^2 \cdot a_l \\ &= \frac{1}{N} \sum \frac{1}{2}a_1 p_1^2 \cdot \omega_l e^{-\alpha - \beta \varepsilon_l} \\ &= \frac{1}{N} \int \frac{1}{2}a_1 p_1^2 \cdot e^{-\alpha - \beta \varepsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r}\end{aligned}$$

## 7.4 Energy equipartition theorem

- Theorem: any degree of freedom  $x$  which appears in the total energy  $\varepsilon$  only as a simple quadratic term, has an average energy of  $\frac{1}{2}kT$  in thermal equilibrium.
- Example: monatomic ideal gas.

$$\begin{aligned}\bar{\varepsilon} &= \overline{\frac{1}{2}mv^2} = \overline{\frac{1}{2}m(v_x^2 + v_y^2 + v_z^2)} \\ &= \frac{1}{2}m\overline{v^2} = \frac{1}{2}m \cdot \frac{3kT}{m} = \frac{3}{2}kT.\end{aligned}$$

- Proof:  $\varepsilon = \varepsilon_k + \varepsilon_p$ , where  $\varepsilon_k = \frac{1}{2} \sum a_i p_i^2$ .

Considering the 1st term,

$$\begin{aligned}\overline{\frac{1}{2}a_1 p_1^2} &= \frac{1}{N} \sum \frac{1}{2}a_1 p_1^2 \cdot a_l \\ &= \frac{1}{N} \sum \frac{1}{2}a_1 p_1^2 \cdot \omega_l e^{-\alpha - \beta \varepsilon_l} \\ &= \frac{1}{N} \int \frac{1}{2}a_1 p_1^2 \cdot e^{-\alpha - \beta \varepsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r} \\ &= \frac{1}{Z} \int \frac{1}{2}a_1 p_1^2 \cdot e^{-\beta \varepsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h^r} \dots\end{aligned}$$

# Energy equipartition theorem

- $\overline{\frac{1}{2}a_1p_1^2} = \frac{1}{Z_1} \int \frac{1}{2}a_1p_1^2 \cdot e^{-\beta\epsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r}$

# Energy equipartition theorem

- $$\overline{\frac{1}{2}a_1 p_1^2} = \frac{1}{Z_1} \int \frac{1}{2}a_1 p_1^2 \cdot e^{-\beta \varepsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r}$$

$$= \int_{-\infty}^{\infty} \frac{1}{2}a_1 p_1^2 e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r}$$

# Energy equipartition theorem

$$\begin{aligned}
 \bullet \quad \overline{\frac{1}{2}a_1 p_1^2} &= \frac{1}{Z_1} \int \frac{1}{2}a_1 p_1^2 \cdot e^{-\beta \varepsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{2}a_1 p_1^2 e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{4}a_1 p_1 e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1^2 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r}
 \end{aligned}$$

# Energy equipartition theorem

$$\begin{aligned}
 \bullet \quad \overline{\frac{1}{2}a_1p_1^2} &= \frac{1}{Z_1} \int \frac{1}{2}a_1p_1^2 \cdot e^{-\beta\epsilon} \frac{dq_1\dots dq_r dp_1\dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{2}a_1p_1^2 e^{-\beta\frac{1}{2}a_1p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\epsilon - \frac{1}{2}a_1p_1^2)} \frac{dq_1\dots dq_r dp_2\dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{4}a_1p_1 e^{-\beta\frac{1}{2}a_1p_1^2} dp_1^2 \cdot \frac{1}{Z_1} \int e^{-\beta(\epsilon - \frac{1}{2}a_1p_1^2)} \frac{dq_1\dots dq_r dp_2\dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{4}a_1p_1 \cdot \left(\frac{-2}{\beta a_1}\right) de^{-\beta\frac{1}{2}a_1p_1^2} \cdot \\
 &\quad \frac{1}{Z_1} \int e^{-\beta(\epsilon - \frac{1}{2}a_1p_1^2)} \frac{dq_1\dots dq_r dp_2\dots dp_r}{h_0^r}
 \end{aligned}$$

# Energy equipartition theorem

$$\begin{aligned}
 \bullet \quad \overline{\frac{1}{2}a_1 p_1^2} &= \frac{1}{Z_1} \int \frac{1}{2}a_1 p_1^2 \cdot e^{-\beta \varepsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{2}a_1 p_1^2 e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{4}a_1 p_1 e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1^2 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{4}a_1 p_1 \cdot \left(\frac{-2}{\beta a_1}\right) de^{-\beta \frac{1}{2}a_1 p_1^2} \cdot \\
 &\quad \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \frac{1}{2\beta} \int_{-\infty}^{\infty} e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r}
 \end{aligned}$$



# Energy equipartition theorem

$$\begin{aligned}
 \bullet \quad \overline{\frac{1}{2}a_1 p_1^2} &= \frac{1}{Z_1} \int \frac{1}{2}a_1 p_1^2 \cdot e^{-\beta \varepsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{2}a_1 p_1^2 e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{4}a_1 p_1 e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1^2 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{4}a_1 p_1 \cdot \left(\frac{-2}{\beta a_1}\right) de^{-\beta \frac{1}{2}a_1 p_1^2} \cdot \\
 &\quad \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \frac{1}{2\beta} \int_{-\infty}^{\infty} e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \frac{1}{2\beta} \frac{1}{Z_1} \int e^{-\beta \varepsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r}
 \end{aligned}$$

# Energy equipartition theorem

$$\begin{aligned}
 \bullet \quad \overline{\frac{1}{2}a_1 p_1^2} &= \frac{1}{Z_1} \int \frac{1}{2}a_1 p_1^2 \cdot e^{-\beta \varepsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{2}a_1 p_1^2 e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{4}a_1 p_1 e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1^2 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{4}a_1 p_1 \cdot \left(\frac{-2}{\beta a_1}\right) de^{-\beta \frac{1}{2}a_1 p_1^2} \cdot \\
 &\quad \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \frac{1}{2\beta} \int_{-\infty}^{\infty} e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \frac{1}{2\beta} \frac{1}{Z_1} \int e^{-\beta \varepsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r} \\
 &= \frac{1}{2\beta}
 \end{aligned}$$

# Energy equipartition theorem

$$\begin{aligned}
 \bullet \quad \overline{\frac{1}{2}a_1 p_1^2} &= \frac{1}{Z_1} \int \frac{1}{2}a_1 p_1^2 \cdot e^{-\beta \varepsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{2}a_1 p_1^2 e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{4}a_1 p_1 e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1^2 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{4}a_1 p_1 \cdot \left(\frac{-2}{\beta a_1}\right) de^{-\beta \frac{1}{2}a_1 p_1^2} \cdot \\
 &\quad \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \frac{1}{2\beta} \int_{-\infty}^{\infty} e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\
 &= \frac{1}{2\beta} \frac{1}{Z_1} \int e^{-\beta \varepsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r} \\
 &= \frac{1}{2\beta} = \frac{1}{2}kT.
 \end{aligned}$$

# Energy equipartition theorem

- $$\begin{aligned}
 \overline{\frac{1}{2}a_1p_1^2} &= \frac{1}{Z_1} \int \frac{1}{2}a_1p_1^2 \cdot e^{-\beta\epsilon} \frac{dq_1\dots dq_r dp_1\dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{2}a_1p_1^2 e^{-\beta\frac{1}{2}a_1p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\epsilon - \frac{1}{2}a_1p_1^2)} \frac{dq_1\dots dq_r dp_2\dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{4}a_1p_1 e^{-\beta\frac{1}{2}a_1p_1^2} dp_1^2 \cdot \frac{1}{Z_1} \int e^{-\beta(\epsilon - \frac{1}{2}a_1p_1^2)} \frac{dq_1\dots dq_r dp_2\dots dp_r}{h_0^r} \\
 &= \int_{-\infty}^{\infty} \frac{1}{4}a_1p_1 \cdot \left(\frac{-2}{\beta a_1}\right) de^{-\beta\frac{1}{2}a_1p_1^2} \cdot \\
 &\quad \frac{1}{Z_1} \int e^{-\beta(\epsilon - \frac{1}{2}a_1p_1^2)} \frac{dq_1\dots dq_r dp_2\dots dp_r}{h_0^r} \\
 &= \frac{1}{2\beta} \int_{-\infty}^{\infty} e^{-\beta\frac{1}{2}a_1p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\epsilon - \frac{1}{2}a_1p_1^2)} \frac{dq_1\dots dq_r dp_2\dots dp_r}{h_0^r} \\
 &= \frac{1}{2\beta} \frac{1}{Z_1} \int e^{-\beta\epsilon} \frac{dq_1\dots dq_r dp_1\dots dp_r}{h_0^r} \\
 &= \frac{1}{2\beta} = \frac{1}{2}kT.
 \end{aligned}$$
- Similar for the other quadratic term.

# Energy equipartition theorem

- $$\begin{aligned} \overline{\frac{1}{2}a_1 p_1^2} &= \frac{1}{Z_1} \int \frac{1}{2}a_1 p_1^2 \cdot e^{-\beta \varepsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r} \\ &= \int_{-\infty}^{\infty} \frac{1}{2}a_1 p_1^2 e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\ &= \int_{-\infty}^{\infty} \frac{1}{4}a_1 p_1 e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1^2 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\ &= \int_{-\infty}^{\infty} \frac{1}{4}a_1 p_1 \cdot \left(\frac{-2}{\beta a_1}\right) de^{-\beta \frac{1}{2}a_1 p_1^2} \cdot \\ &\quad \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\ &= \frac{1}{2\beta} \int_{-\infty}^{\infty} e^{-\beta \frac{1}{2}a_1 p_1^2} dp_1 \cdot \frac{1}{Z_1} \int e^{-\beta(\varepsilon - \frac{1}{2}a_1 p_1^2)} \frac{dq_1 \dots dq_r dp_2 \dots dp_r}{h_0^r} \\ &= \frac{1}{2\beta} \frac{1}{Z_1} \int e^{-\beta \varepsilon} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r} \\ &= \frac{1}{2\beta} = \frac{1}{2}kT. \end{aligned}$$
- Similar for the other quadratic term.
- Application: internal energy  $U$ , thermal capacity  $C_V$ .

# Application 1. Monatomic molecules

# Application 1. Monatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$

# Application 1. Monatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$   
 $\Rightarrow \bar{\varepsilon} = \frac{3}{2}kT$  (3 quadratic terms).



# Application 1. Monatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$   
 $\Rightarrow \bar{\varepsilon} = \frac{3}{2}kT$  (3 quadratic terms).
- $U = N\bar{\varepsilon} = \frac{3}{2}NkT$ .

# Application 1. Monatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$   
 $\Rightarrow \bar{\varepsilon} = \frac{3}{2}kT$  (3 quadratic terms).
- $U = N\bar{\varepsilon} = \frac{3}{2}NkT.$
- $C_V = \frac{dU}{dT} = \frac{3}{2}Nk.$

# Application 1. Monatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$   
 $\Rightarrow \bar{\varepsilon} = \frac{3}{2}kT$  (3 quadratic terms).
- $U = N\bar{\varepsilon} = \frac{3}{2}NkT$ .
- $C_V = \frac{dU}{dT} = \frac{3}{2}Nk$ .
- As  $pV = nRT = NkT$ ,  $C_p - C_V = nR = Nk$ ,

# Application 1. Monatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$   
 $\Rightarrow \bar{\varepsilon} = \frac{3}{2}kT$  (3 quadratic terms).
- $U = N\bar{\varepsilon} = \frac{3}{2}NkT$ .
- $C_V = \frac{dU}{dT} = \frac{3}{2}Nk$ .
- As  $pV = nRT = NkT$ ,  $C_p - C_V = nR = Nk$ ,  
 $\therefore C_p = \frac{5}{2}Nk$ , and  $\gamma = \frac{C_p}{C_V} = \frac{5}{3}$ .

# Application 1. Monatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$   
 $\Rightarrow \bar{\varepsilon} = \frac{3}{2}kT$  (3 quadratic terms).
- $U = N\bar{\varepsilon} = \frac{3}{2}NkT$ .
- $C_V = \frac{dU}{dT} = \frac{3}{2}Nk$ .
- As  $pV = nRT = NkT$ ,  $C_p - C_V = nR = Nk$ ,  
 $\therefore C_p = \frac{5}{2}Nk$ , and  $\gamma = \frac{C_p}{C_V} = \frac{5}{3}$ .
- Table 7.2.

# Application 1. Monatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$   
 $\Rightarrow \bar{\varepsilon} = \frac{3}{2}kT$  (3 quadratic terms).
- $U = N\bar{\varepsilon} = \frac{3}{2}NkT$ .
- $C_V = \frac{dU}{dT} = \frac{3}{2}Nk$ .
- As  $pV = nRT = NkT$ ,  $C_p - C_V = nR = Nk$ ,  
 $\therefore C_p = \frac{5}{2}Nk$ , and  $\gamma = \frac{C_p}{C_V} = \frac{5}{3}$ .
- Table 7.2.

Did not consider the contribution of internal structure (electrons and protons), but consistent with the experiments. Why?

# Application 1. Monatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2)$   
 $\Rightarrow \bar{\varepsilon} = \frac{3}{2}kT$  (3 quadratic terms).
- $U = N\bar{\varepsilon} = \frac{3}{2}NkT$ .
- $C_V = \frac{dU}{dT} = \frac{3}{2}Nk$ .
- As  $pV = nRT = NkT$ ,  $C_p - C_V = nR = Nk$ ,  
 $\therefore C_p = \frac{5}{2}Nk$ , and  $\gamma = \frac{C_p}{C_V} = \frac{5}{3}$ .
- Table 7.2.

Did not consider the contribution of internal structure (electrons and protons), but consistent with the experiments. Why?

- Quantum effect. Energy level are discrete.

# Application 1. Monatomic molecules

- Such as potassium (K), the ionization energy  $418.8 \text{ kJ mol}^{-1} = \frac{418.8 \times 10^3 \text{ J}}{6.02 \times 10^{23}} \simeq 7 \times 10^{-19} \text{ J}$ .



# Application 1. Monatomic molecules

- Such as potassium (K), the ionization energy  $418.8 \text{ kJ mol}^{-1} = \frac{418.8 \times 10^3 \text{ J}}{6.02 \times 10^{23}} \simeq 7 \times 10^{-19} \text{ J}$ .

The transition energy (ground state to 1st excited state) is  $\frac{3}{4} \times 7 \times 10^{-19} \text{ J} \simeq 5 \times 10^{-19} \text{ J}$ .

# Application 1. Monatomic molecules

- Such as potassium (K), the ionization energy  $418.8 \text{ kJ mol}^{-1} = \frac{418.8 \times 10^3 \text{ J}}{6.02 \times 10^{23}} \simeq 7 \times 10^{-19} \text{ J}$ .  
The transition energy (ground state to 1st excited state) is  $\frac{3}{4} \times 7 \times 10^{-19} \text{ J} \simeq 5 \times 10^{-19} \text{ J}$ .
- Typical energy for  $T = 1000 \text{ K}$ ,  
 $kT = 1.38 \times 10^{-23} \text{ J K}^{-1} \times 10^3 \text{ K} \simeq 1.38 \times 10^{-20} \text{ J}$ .

# Application 1. Monatomic molecules

- Such as potassium (K), the ionization energy  $418.8 \text{ kJ mol}^{-1} = \frac{418.8 \times 10^3 \text{ J}}{6.02 \times 10^{23}} \simeq 7 \times 10^{-19} \text{ J}$ .  
The transition energy (ground state to 1st excited state) is  $\frac{3}{4} \times 7 \times 10^{-19} \text{ J} \simeq 5 \times 10^{-19} \text{ J}$ .
- Typical energy for  $T = 1000 \text{ K}$ ,  
 $kT = 1.38 \times 10^{-23} \text{ J K}^{-1} \times 10^3 \text{ K} \simeq 1.38 \times 10^{-20} \text{ J}$ .  
Smaller than the energy required by the transition energy.

# Application 1. Monatomic molecules

- Such as potassium (K), the ionization energy  $418.8 \text{ kJ mol}^{-1} = \frac{418.8 \times 10^3 \text{ J}}{6.02 \times 10^{23}} \simeq 7 \times 10^{-19} \text{ J}$ .  
The transition energy (ground state to 1st excited state) is  $\frac{3}{4} \times 7 \times 10^{-19} \text{ J} \simeq 5 \times 10^{-19} \text{ J}$ .
- Typical energy for  $T = 1000 \text{ K}$ ,  
 $kT = 1.38 \times 10^{-23} \text{ J K}^{-1} \times 10^3 \text{ K} \simeq 1.38 \times 10^{-20} \text{ J}$ .  
Smaller than the energy required by the transition energy.  
Therefore, the contribution of electrons should not be considered.

## Application 2. diatomic molecules

## Application 2. diatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2\theta}p_\varphi^2) + \frac{1}{2m_\mu}p_r^2 + u(r)$   
where  $m_\mu = \frac{m_1 m_2}{m_1 + m_2}$  is the reduced mass,  $I$  is the moment of inertia.  $\frac{1}{2m_\mu}p_r^2$  is the energy because of relative motion.

## Application 2. diatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2\theta}p_\varphi^2) + \frac{1}{2m_\mu}p_r^2 + u(r)$   
where  $m_\mu = \frac{m_1 m_2}{m_1 + m_2}$  is the reduced mass,  $I$  is the moment of inertia.  $\frac{1}{2m_\mu}p_r^2$  is the energy because of relative motion.
- Not considering  $p_r^2$ , there are 5 quadratic terms.

## Application 2. diatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2\theta}p_\varphi^2) + \frac{1}{2m_\mu}p_r^2 + u(r)$   
where  $m_\mu = \frac{m_1 m_2}{m_1 + m_2}$  is the reduced mass,  $I$  is the moment of inertia.  $\frac{1}{2m_\mu}p_r^2$  is the energy because of relative motion.
- Not considering  $p_r^2$ , there are 5 quadratic terms.  
 $\bar{\varepsilon} = \frac{5}{2}kT$ .



## Application 2. diatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2\theta}p_\varphi^2) + \frac{1}{2m_\mu}p_r^2 + u(r)$   
where  $m_\mu = \frac{m_1 m_2}{m_1 + m_2}$  is the reduced mass,  $I$  is the moment of inertia.  $\frac{1}{2m_\mu}p_r^2$  is the energy because of relative motion.
- Not considering  $p_r^2$ , there are 5 quadratic terms.  
 $\bar{\varepsilon} = \frac{5}{2}kT.$
- $U = \frac{5}{2}NkT,$

## Application 2. diatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2\theta}p_\varphi^2) + \frac{1}{2m_\mu}p_r^2 + u(r)$   
where  $m_\mu = \frac{m_1 m_2}{m_1 + m_2}$  is the reduced mass,  $I$  is the moment of inertia.  $\frac{1}{2m_\mu}p_r^2$  is the energy because of relative motion.
- Not considering  $p_r^2$ , there are 5 quadratic terms.  
 $\bar{\varepsilon} = \frac{5}{2}kT.$
- $U = \frac{5}{2}NkT, C_V = \frac{5}{2}Nk,$

## Application 2. diatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2\theta}p_\varphi^2) + \frac{1}{2m_\mu}p_r^2 + u(r)$   
where  $m_\mu = \frac{m_1 m_2}{m_1 + m_2}$  is the reduced mass,  $I$  is the moment of inertia.  $\frac{1}{2m_\mu}p_r^2$  is the energy because of relative motion.
- Not considering  $p_r^2$ , there are 5 quadratic terms.  
 $\bar{\varepsilon} = \frac{5}{2}kT$ .
- $U = \frac{5}{2}NkT$ ,  $C_V = \frac{5}{2}Nk$ ,  $C_p = \frac{7}{2}Nk$ ,

## Application 2. diatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2\theta}p_\varphi^2) + \frac{1}{2m_\mu}p_r^2 + u(r)$   
 where  $m_\mu = \frac{m_1 m_2}{m_1 + m_2}$  is the reduced mass,  $I$  is the moment of inertia.  $\frac{1}{2m_\mu}p_r^2$  is the energy because of relative motion.
- Not considering  $p_r^2$ , there are 5 quadratic terms.  
 $\bar{\varepsilon} = \frac{5}{2}kT$ .
- $U = \frac{5}{2}NkT$ ,  $C_V = \frac{5}{2}Nk$ ,  $C_p = \frac{7}{2}Nk$ ,  
 $\gamma = 1.4$ , see table 7.3.

## Application 2. diatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2\theta}p_\varphi^2) + \frac{1}{2m_\mu}p_r^2 + u(r)$   
 where  $m_\mu = \frac{m_1 m_2}{m_1 + m_2}$  is the reduced mass,  $I$  is the moment of inertia.  $\frac{1}{2m_\mu}p_r^2$  is the energy because of relative motion.
- Not considering  $p_r^2$ , there are 5 quadratic terms.  
 $\bar{\varepsilon} = \frac{5}{2}kT$ .
- $U = \frac{5}{2}NkT$ ,  $C_V = \frac{5}{2}Nk$ ,  $C_p = \frac{7}{2}Nk$ ,  
 $\gamma = 1.4$ , see table 7.3.
- i. Why no  $p_r^2$  ?

## Application 2. diatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2\theta}p_\varphi^2) + \frac{1}{2m_\mu}p_r^2 + u(r)$   
 where  $m_\mu = \frac{m_1 m_2}{m_1 + m_2}$  is the reduced mass,  $I$  is the moment of inertia.  $\frac{1}{2m_\mu}p_r^2$  is the energy because of relative motion.
- Not considering  $p_r^2$ , there are 5 quadratic terms.  
 $\bar{\varepsilon} = \frac{5}{2}kT$ .
- $U = \frac{5}{2}NkT$ ,  $C_V = \frac{5}{2}Nk$ ,  $C_p = \frac{7}{2}Nk$ ,  
 $\gamma = 1.4$ , see table 7.3.
- i. Why no  $p_r^2$  ? Discrete energy level.

## Application 2. diatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2\theta}p_\varphi^2) + \frac{1}{2m_\mu}p_r^2 + u(r)$   
where  $m_\mu = \frac{m_1 m_2}{m_1 + m_2}$  is the reduced mass,  $I$  is the moment of inertia.  $\frac{1}{2m_\mu}p_r^2$  is the energy because of relative motion.
- Not considering  $p_r^2$ , there are 5 quadratic terms.  
 $\bar{\varepsilon} = \frac{5}{2}kT$ .
- $U = \frac{5}{2}NkT$ ,  $C_V = \frac{5}{2}Nk$ ,  $C_p = \frac{7}{2}Nk$ ,  
 $\gamma = 1.4$ , see table 7.3.
- i. Why no  $p_r^2$  ? Discrete energy level.  
ii. Why  $H_2$  not good in low temperature?

## Application 2. diatomic molecules

- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2\theta}p_\varphi^2) + \frac{1}{2m_\mu}p_r^2 + u(r)$   
 where  $m_\mu = \frac{m_1 m_2}{m_1 + m_2}$  is the reduced mass,  $I$  is the moment of inertia.  $\frac{1}{2m_\mu}p_r^2$  is the energy because of relative motion.
- Not considering  $p_r^2$ , there are 5 quadratic terms.  
 $\bar{\varepsilon} = \frac{5}{2}kT$ .
- $U = \frac{5}{2}NkT$ ,  $C_V = \frac{5}{2}Nk$ ,  $C_p = \frac{7}{2}Nk$ ,  
 $\gamma = 1.4$ , see table 7.3.
- i. Why no  $p_r^2$  ? Discrete energy level.  
 ii. Why  $H_2$  not good in low temperature?  
 Eq. (7.2.6),  $\frac{V}{N}(\frac{2\pi mkT}{h^2})^{3/2}$ , classical limit.



## Application 3. Atoms in solid

## Application 3. Atoms in solid

- 1. Atoms are trapped.

## Application 3. Atoms in solid

- 1. Atoms are trapped.
- 2. Atom can vibrate near its equilibrium position.

## Application 3. Atoms in solid

- 1. Atoms are trapped.
- 2. Atom can vibrate near its equilibrium position.
- 3. Vibrations of each atom are independent.

## Application 3. Atoms in solid

- 1. Atoms are trapped.
- 2. Atom can vibrate near its equilibrium position.
- 3. Vibrations of each atom are independent.
- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2}m\omega^2(x^2 + y^2 + z^2).$

## Application 3. Atoms in solid

- 1. Atoms are trapped.
- 2. Atom can vibrate near its equilibrium position.
- 3. Vibrations of each atom are independent.
- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2}m\omega^2(x^2 + y^2 + z^2)$ .
- 6 quadratic terms,

## Application 3. Atoms in solid

- 1. Atoms are trapped.
- 2. Atom can vibrate near its equilibrium position.
- 3. Vibrations of each atom are independent.
- $\varepsilon = \frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2}m\omega^2(x^2 + y^2 + z^2)$ .
- 6 quadratic terms,  
 $\bar{\varepsilon} = 3kT, U = 3NkT, C_V = 3Nk$ .  
Consistent with the experiment at high temperature.

# Application 4. Equilibrium radiation



## Application 4. Equilibrium radiation

- What is the microscopic unit of thermal radiation?

## Application 4. Equilibrium radiation

- What is the microscopic unit of thermal radiation?
- View 1: consist of particles – bosons – photons (§8.4).

## Application 4. Equilibrium radiation

- What is the microscopic unit of thermal radiation?
- View 1: consist of particles – bosons – photons (§8.4).
- View 2: consist of electron-magnetic waves with different frequencies (discrete).

## Application 4. Equilibrium radiation

- What is the microscopic unit of thermal radiation?
- View 1: consist of particles – bosons – photons (§8.4).
- View 2: consist of electron-magnetic waves with different frequencies (discrete).

Each monochromatic wave with **a certain frequency** acts like a classical particle.

## Application 4. Equilibrium radiation

- What is the microscopic unit of thermal radiation?
- View 1: consist of particles – bosons – photons (§8.4).
- View 2: consist of electron-magnetic waves with different frequencies (discrete).

Each monochromatic wave with **a certain frequency** acts like a classical particle.

Each wave is equivalent to a harmonic oscillation.

## Application 4. Equilibrium radiation

- What is the microscopic unit of thermal radiation?
- View 1: consist of particles – bosons – photons (§8.4).
- View 2: consist of electron-magnetic waves with different frequencies (discrete).

Each monochromatic wave with **a certain frequency** acts like a classical particle.

Each wave is equivalent to a harmonic oscillation.

$$\varepsilon = \frac{p^2}{2m} + \frac{1}{2}m\omega^2 q^2.$$

## Application 4. Equilibrium radiation

- What is the microscopic unit of thermal radiation?
- View 1: consist of particles – bosons – photons (§8.4).
- View 2: consist of electron-magnetic waves with different frequencies (discrete).

Each monochromatic wave with **a certain frequency** acts like a classical particle.

Each wave is equivalent to a harmonic oscillation.

$$\varepsilon = \frac{p^2}{2m} + \frac{1}{2}m\omega^2 q^2.$$

- Two quadratic terms,  $\bar{\varepsilon} = kT$ .

## Application 4. Equilibrium radiation

- What is the microscopic unit of thermal radiation?
- View 1: consist of particles – bosons – photons (§8.4).
- View 2: consist of electron-magnetic waves with different frequencies (discrete).

Each monochromatic wave with **a certain frequency** acts like a classical particle.

Each wave is equivalent to a harmonic oscillation.

$$\varepsilon = \frac{p^2}{2m} + \frac{1}{2}m\omega^2 q^2.$$

- Two quadratic terms,  $\bar{\varepsilon} = kT$ .
- Using the periodic boundary condition:



## Application 4. Equilibrium radiation

- $k_x = \frac{2\pi}{L}n_x, n_x = 0, \pm 1, \pm 2, \dots$   
 $k_y = \frac{2\pi}{L}n_y, n_y = 0, \pm 1, \pm 2, \dots$   
 $k_z = \frac{2\pi}{L}n_z, n_z = 0, \pm 1, \pm 2, \dots$

## Application 4. Equilibrium radiation

- $k_x = \frac{2\pi}{L}n_x, n_x = 0, \pm 1, \pm 2, \dots$   
 $k_y = \frac{2\pi}{L}n_y, n_y = 0, \pm 1, \pm 2, \dots$   
 $k_z = \frac{2\pi}{L}n_z, n_z = 0, \pm 1, \pm 2, \dots$
- $dk_x dk_y dk_z = \left(\frac{2\pi}{L}\right)^3 \Delta n_x \Delta n_y \Delta n_z,$

## Application 4. Equilibrium radiation

- $k_x = \frac{2\pi}{L}n_x, n_x = 0, \pm 1, \pm 2, \dots$   
 $k_y = \frac{2\pi}{L}n_y, n_y = 0, \pm 1, \pm 2, \dots$   
 $k_z = \frac{2\pi}{L}n_z, n_z = 0, \pm 1, \pm 2, \dots$
- $dk_x dk_y dk_z = \left(\frac{2\pi}{L}\right)^3 \Delta n_x \Delta n_y \Delta n_z,$   
i.e., number of oscillations:  
$$\Delta n_x \Delta n_y \Delta n_z = \frac{V}{(2\pi)^3} dk_x dk_y dk_z.$$

## Application 4. Equilibrium radiation

- $k_x = \frac{2\pi}{L}n_x$ ,  $n_x = 0, \pm 1, \pm 2, \dots$   
 $k_y = \frac{2\pi}{L}n_y$ ,  $n_y = 0, \pm 1, \pm 2, \dots$   
 $k_z = \frac{2\pi}{L}n_z$ ,  $n_z = 0, \pm 1, \pm 2, \dots$
- $dk_x dk_y dk_z = \left(\frac{2\pi}{L}\right)^3 \Delta n_x \Delta n_y \Delta n_z$ ,  
i.e., number of oscillations:  
 $\Delta n_x \Delta n_y \Delta n_z = \frac{V}{(2\pi)^3} dk_x dk_y dk_z$ .
- In spheric polar coordinate, isotropic radiation,  
together with the polarization:

## Application 4. Equilibrium radiation

- $k_x = \frac{2\pi}{L}n_x$ ,  $n_x = 0, \pm 1, \pm 2, \dots$   
 $k_y = \frac{2\pi}{L}n_y$ ,  $n_y = 0, \pm 1, \pm 2, \dots$   
 $k_z = \frac{2\pi}{L}n_z$ ,  $n_z = 0, \pm 1, \pm 2, \dots$
- $dk_x dk_y dk_z = \left(\frac{2\pi}{L}\right)^3 \Delta n_x \Delta n_y \Delta n_z$ ,  
 i.e., number of oscillations:  
 $\Delta n_x \Delta n_y \Delta n_z = \frac{V}{(2\pi)^3} dk_x dk_y dk_z$ .
- In spheric polar coordinate, isotropic radiation,  
 together with the polarization:  
 $\Delta n_x \Delta n_y \Delta n_z = 2 \frac{V}{(2\pi)^3} k^2 dk \cdot 4\pi$

## Application 4. Equilibrium radiation

- $k_x = \frac{2\pi}{L}n_x$ ,  $n_x = 0, \pm 1, \pm 2, \dots$   
 $k_y = \frac{2\pi}{L}n_y$ ,  $n_y = 0, \pm 1, \pm 2, \dots$   
 $k_z = \frac{2\pi}{L}n_z$ ,  $n_z = 0, \pm 1, \pm 2, \dots$
- $dk_x dk_y dk_z = \left(\frac{2\pi}{L}\right)^3 \Delta n_x \Delta n_y \Delta n_z$ ,  
 i.e., number of oscillations:  
 $\Delta n_x \Delta n_y \Delta n_z = \frac{V}{(2\pi)^3} dk_x dk_y dk_z$ .
- In spheric polar coordinate, isotropic radiation,  
 together with the polarization:  
 $\Delta n_x \Delta n_y \Delta n_z = 2 \frac{V}{(2\pi)^3} k^2 dk \cdot 4\pi = 2 \frac{V}{(2\pi)^3} \left(\frac{\omega}{c}\right)^2 d\frac{\omega}{c} \cdot 4\pi$

## Application 4. Equilibrium radiation

- $k_x = \frac{2\pi}{L}n_x$ ,  $n_x = 0, \pm 1, \pm 2, \dots$   
 $k_y = \frac{2\pi}{L}n_y$ ,  $n_y = 0, \pm 1, \pm 2, \dots$   
 $k_z = \frac{2\pi}{L}n_z$ ,  $n_z = 0, \pm 1, \pm 2, \dots$
- $dk_x dk_y dk_z = \left(\frac{2\pi}{L}\right)^3 \Delta n_x \Delta n_y \Delta n_z$ ,  
 i.e., number of oscillations:  
 $\Delta n_x \Delta n_y \Delta n_z = \frac{V}{(2\pi)^3} dk_x dk_y dk_z$ .
- In spheric polar coordinate, isotropic radiation,  
 together with the polarization:  
 $\Delta n_x \Delta n_y \Delta n_z = 2 \frac{V}{(2\pi)^3} k^2 dk \cdot 4\pi = 2 \frac{V}{(2\pi)^3} \left(\frac{\omega}{c}\right)^2 d\frac{\omega}{c} \cdot 4\pi$   
 $= \frac{V}{\pi^2 c^3} \omega^2 d\omega \equiv D(\omega) d\omega$ .

# Application 4. Equilibrium radiation

- $k_x = \frac{2\pi}{L}n_x$ ,  $n_x = 0, \pm 1, \pm 2, \dots$   
 $k_y = \frac{2\pi}{L}n_y$ ,  $n_y = 0, \pm 1, \pm 2, \dots$   
 $k_z = \frac{2\pi}{L}n_z$ ,  $n_z = 0, \pm 1, \pm 2, \dots$
- $dk_x dk_y dk_z = \left(\frac{2\pi}{L}\right)^3 \Delta n_x \Delta n_y \Delta n_z$ ,  
 i.e., number of oscillations:  

$$\Delta n_x \Delta n_y \Delta n_z = \frac{V}{(2\pi)^3} dk_x dk_y dk_z.$$
- In spheric polar coordinate, isotropic radiation,  
 together with the polarization:  

$$\Delta n_x \Delta n_y \Delta n_z = 2 \frac{V}{(2\pi)^3} k^2 dk \cdot 4\pi = 2 \frac{V}{(2\pi)^3} \left(\frac{\omega}{c}\right)^2 d\frac{\omega}{c} \cdot 4\pi$$

$$= \frac{V}{\pi^2 c^3} \omega^2 d\omega \equiv D(\omega) d\omega.$$
- As for each vibration  $\bar{\epsilon} = kT$ ,



# Application 4. Equilibrium radiation

- $k_x = \frac{2\pi}{L}n_x$ ,  $n_x = 0, \pm 1, \pm 2, \dots$   
 $k_y = \frac{2\pi}{L}n_y$ ,  $n_y = 0, \pm 1, \pm 2, \dots$   
 $k_z = \frac{2\pi}{L}n_z$ ,  $n_z = 0, \pm 1, \pm 2, \dots$
- $dk_x dk_y dk_z = \left(\frac{2\pi}{L}\right)^3 \Delta n_x \Delta n_y \Delta n_z$ ,  
 i.e., number of oscillations:  

$$\Delta n_x \Delta n_y \Delta n_z = \frac{V}{(2\pi)^3} dk_x dk_y dk_z.$$
- In spheric polar coordinate, isotropic radiation,  
 together with the polarization:  

$$\Delta n_x \Delta n_y \Delta n_z = 2 \frac{V}{(2\pi)^3} k^2 dk \cdot 4\pi = 2 \frac{V}{(2\pi)^3} \left(\frac{\omega}{c}\right)^2 d\frac{\omega}{c} \cdot 4\pi$$

$$= \frac{V}{\pi^2 c^3} \omega^2 d\omega \equiv D(\omega) d\omega.$$
- As for each vibration  $\bar{\varepsilon} = kT$ , the internal energy  

$$U_\omega d\omega = \frac{V}{\pi^2 c^3} kT \omega^2 d\omega.$$

## Application 4. Equilibrium radiation

- $k_x = \frac{2\pi}{L}n_x$ ,  $n_x = 0, \pm 1, \pm 2, \dots$   
 $k_y = \frac{2\pi}{L}n_y$ ,  $n_y = 0, \pm 1, \pm 2, \dots$   
 $k_z = \frac{2\pi}{L}n_z$ ,  $n_z = 0, \pm 1, \pm 2, \dots$
- $dk_x dk_y dk_z = \left(\frac{2\pi}{L}\right)^3 \Delta n_x \Delta n_y \Delta n_z$ ,  
 i.e., number of oscillations:  

$$\Delta n_x \Delta n_y \Delta n_z = \frac{V}{(2\pi)^3} dk_x dk_y dk_z.$$
- In spheric polar coordinate, isotropic radiation, together with the polarization:  

$$\Delta n_x \Delta n_y \Delta n_z = 2 \frac{V}{(2\pi)^3} k^2 dk \cdot 4\pi = 2 \frac{V}{(2\pi)^3} \left(\frac{\omega}{c}\right)^2 d\frac{\omega}{c} \cdot 4\pi$$

$$= \frac{V}{\pi^2 c^3} \omega^2 d\omega \equiv D(\omega) d\omega.$$
- As for each vibration  $\bar{\varepsilon} = kT$ , the internal energy  

$$U_\omega d\omega = \frac{V}{\pi^2 c^3} kT \omega^2 d\omega.$$
 Only suitable for low frequency.

## Application 4. Equilibrium radiation

- Theory:  $U_\omega d\omega = \frac{V}{\pi^2 c^3} kT \omega^2 d\omega.$

## Application 4. Equilibrium radiation

- Theory:  $U_{\omega}d\omega = \frac{V}{\pi^2 c^3} kT \omega^2 d\omega$ .  
Experiment:  $U_{\omega}d\omega = \frac{V}{2\pi^3 c^3} \frac{h\omega^3 d\omega}{e^{\frac{h\omega}{2\pi kT}} - 1}$ .

## Application 4. Equilibrium radiation

- Theory:  $U_\omega d\omega = \frac{V}{\pi^2 c^3} kT \omega^2 d\omega$ .  
Experiment:  $U_\omega d\omega = \frac{V}{2\pi^3 c^3} \frac{h\omega^3 d\omega}{e^{\frac{h\omega}{2\pi kT}} - 1}$ .
- Planck's idea:  $D(\omega) d\omega = \frac{V}{\pi^2 c^3} \omega^2 d\omega$  still obeys.

## Application 4. Equilibrium radiation

- Theory:  $U_\omega d\omega = \frac{V}{\pi^2 c^3} kT \omega^2 d\omega$ .  
Experiment:  $U_\omega d\omega = \frac{V}{2\pi^3 c^3} \frac{h\omega^3 d\omega}{e^{\frac{h\omega}{2\pi kT}} - 1}$ .
- Planck's idea:  $D(\omega)d\omega = \frac{V}{\pi^2 c^3} \omega^2 d\omega$  still obeys. But the energy of the oscillator is NOT continuous, supposed to be multiples of  $\hbar\omega$ , i.e.,  $\varepsilon_n(\omega) = n\hbar\omega$ .

## Application 4. Equilibrium radiation

- Theory:  $U_\omega d\omega = \frac{V}{\pi^2 c^3} kT \omega^2 d\omega$ .  
Experiment:  $U_\omega d\omega = \frac{V}{2\pi^3 c^3} \frac{h\omega^3 d\omega}{e^{\frac{h\omega}{2\pi kT}} - 1}$ .
- Planck's idea:  $D(\omega)d\omega = \frac{V}{\pi^2 c^3} \omega^2 d\omega$  still obeys. But the energy of the oscillator is NOT continuous, supposed to be multiples of  $\hbar\omega$ , i.e.,  $\varepsilon_n(\omega) = n\hbar\omega$ .
- The most probable Boltzmann distribution:  
 $a_n(\omega) = e^{-\alpha - \beta\varepsilon_n},$

## Application 4. Equilibrium radiation

- Theory:  $U_\omega d\omega = \frac{V}{\pi^2 c^3} kT \omega^2 d\omega$ .  
Experiment:  $U_\omega d\omega = \frac{V}{2\pi^3 c^3} \frac{h\omega^3 d\omega}{e^{\frac{h\omega}{2\pi kT}} - 1}$ .
- Planck's idea:  $D(\omega)d\omega = \frac{V}{\pi^2 c^3} \omega^2 d\omega$  still obeys. But the energy of the oscillator is NOT continuous, supposed to be multiples of  $\hbar\omega$ , i.e.,  $\varepsilon_n(\omega) = n\hbar\omega$ .
- The most probable Boltzmann distribution:  
 $a_n(\omega) = e^{-\alpha - \beta\varepsilon_n},$   
degeneracy  $\omega_n(\omega) = 2$  (polarization),



## Application 4. Equilibrium radiation

- Theory:  $U_\omega d\omega = \frac{V}{\pi^2 c^3} kT \omega^2 d\omega$ .  
Experiment:  $U_\omega d\omega = \frac{V}{2\pi^3 c^3} \frac{h\omega^3 d\omega}{e^{\frac{h\omega}{2\pi kT}} - 1}$ .
- Planck's idea:  $D(\omega)d\omega = \frac{V}{\pi^2 c^3} \omega^2 d\omega$  still obeys. But the energy of the oscillator is NOT continuous, supposed to be multiples of  $\hbar\omega$ , i.e.,  $\varepsilon_n(\omega) = n\hbar\omega$ .
- The most probable Boltzmann distribution:  
 $a_n(\omega) = e^{-\alpha - \beta\varepsilon_n},$   
 degeneracy  $\omega_n(\omega) = 2$  (polarization),  
 partition function:  $Z_1 = \sum_{n=0}^{\infty} \omega_n e^{-\beta\varepsilon_n}$

# Application 4. Equilibrium radiation

- Theory:  $U_\omega d\omega = \frac{V}{\pi^2 c^3} kT \omega^2 d\omega$ .  
Experiment:  $U_\omega d\omega = \frac{V}{2\pi^3 c^3} \frac{h\omega^3 d\omega}{e^{\frac{h\omega}{2\pi kT}} - 1}$ .
- Planck's idea:  $D(\omega)d\omega = \frac{V}{\pi^2 c^3} \omega^2 d\omega$  still obeys. But the energy of the oscillator is NOT continuous, supposed to be multiples of  $\hbar\omega$ , i.e.,  $\varepsilon_n(\omega) = n\hbar\omega$ .
- The most probable Boltzmann distribution:  
 $a_n(\omega) = e^{-\alpha - \beta\varepsilon_n},$   
 degeneracy  $\omega_n(\omega) = 2$  (polarization),  
 partition function:  $Z_1 = \sum_{n=0}^{\infty} \omega_n e^{-\beta\varepsilon_n} = \sum_{n=0}^{\infty} 2e^{-\beta n\hbar\omega}$

# Application 4. Equilibrium radiation

- Theory:  $U_\omega d\omega = \frac{V}{\pi^2 c^3} kT \omega^2 d\omega$ .  
Experiment:  $U_\omega d\omega = \frac{V}{2\pi^3 c^3} \frac{h\omega^3 d\omega}{e^{\frac{h\omega}{2\pi kT}} - 1}$ .
- Planck's idea:  $D(\omega)d\omega = \frac{V}{\pi^2 c^3} \omega^2 d\omega$  still obeys. But the energy of the oscillator is NOT continuous, supposed to be multiples of  $\hbar\omega$ , i.e.,  $\varepsilon_n(\omega) = n\hbar\omega$ .
- The most probable Boltzmann distribution:  
 $a_n(\omega) = e^{-\alpha - \beta\varepsilon_n},$   
 degeneracy  $\omega_n(\omega) = 2$  (polarization),  
 partition function:  $Z_1 = \sum_{n=0}^{\infty} \omega_n e^{-\beta\varepsilon_n} = \sum_{n=0}^{\infty} 2e^{-\beta n\hbar\omega}$   
 $= \frac{2}{1 - e^{-\beta\hbar\omega}}.$

## Application 4. Equilibrium radiation

$$Z_1 = \sum_{n=0}^{\infty} 2e^{-\beta\varepsilon_n}, Z_1 = \frac{2}{1-e^{-\beta\hbar\omega}}$$

- The average energy of an oscillator with frequency  $\omega$ :

## Application 4. Equilibrium radiation

$$Z_1 = \sum_{n=0}^{\infty} 2e^{-\beta\varepsilon_n}, Z_1 = \frac{2}{1-e^{-\beta\hbar\omega}}$$

- The average energy of an oscillator with frequency  $\omega$ :

$$\bar{\varepsilon} = \frac{\sum a_n \varepsilon_n}{\sum a_n}$$

## Application 4. Equilibrium radiation

$$Z_1 = \sum_{n=0}^{\infty} 2e^{-\beta\varepsilon_n}, Z_1 = \frac{2}{1-e^{-\beta\hbar\omega}}$$

- The average energy of an oscillator with frequency  $\omega$ :

$$\bar{\varepsilon} = \frac{\sum a_n \varepsilon_n}{\sum a_n} = \frac{\sum e^{-\alpha-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\alpha-\beta\varepsilon_n}}$$

## Application 4. Equilibrium radiation

$$Z_1 = \sum_{n=0}^{\infty} 2e^{-\beta\varepsilon_n}, Z_1 = \frac{2}{1-e^{-\beta\hbar\omega}}$$

- The average energy of an oscillator with frequency  $\omega$ :

$$\bar{\varepsilon} = \frac{\sum a_n \varepsilon_n}{\sum a_n} = \frac{\sum e^{-\alpha-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\alpha-\beta\varepsilon_n}} = \frac{\sum e^{-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\beta\varepsilon_n}}$$

## Application 4. Equilibrium radiation

$$Z_1 = \sum_{n=0}^{\infty} 2e^{-\beta\varepsilon_n}, Z_1 = \frac{2}{1-e^{-\beta\hbar\omega}}$$

- The average energy of an oscillator with frequency  $\omega$ :

$$\bar{\varepsilon} = \frac{\sum a_n \varepsilon_n}{\sum a_n} = \frac{\sum e^{-\alpha-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\alpha-\beta\varepsilon_n}} = \frac{\sum e^{-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\beta\varepsilon_n}} = -\frac{\partial}{\partial\beta} \ln Z_1$$



## Application 4. Equilibrium radiation

$$Z_1 = \sum_{n=0}^{\infty} 2e^{-\beta\varepsilon_n}, Z_1 = \frac{2}{1-e^{-\beta\hbar\omega}}$$

- The average energy of an oscillator with frequency  $\omega$ :

$$\begin{aligned}\bar{\varepsilon} &= \frac{\sum a_n \varepsilon_n}{\sum a_n} = \frac{\sum e^{-\alpha-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\alpha-\beta\varepsilon_n}} = \frac{\sum e^{-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\beta\varepsilon_n}} = -\frac{\partial}{\partial\beta} \ln Z_1 \\ &= -\frac{\partial}{\partial\beta} \ln \frac{2}{1-e^{-\beta\hbar\omega}}\end{aligned}$$

## Application 4. Equilibrium radiation

$$Z_1 = \sum_{n=0}^{\infty} 2e^{-\beta\varepsilon_n}, Z_1 = \frac{2}{1-e^{-\beta\hbar\omega}}$$

- The average energy of an oscillator with frequency  $\omega$ :

$$\begin{aligned}\bar{\varepsilon} &= \frac{\sum a_n \varepsilon_n}{\sum a_n} = \frac{\sum e^{-\alpha-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\alpha-\beta\varepsilon_n}} = \frac{\sum e^{-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\beta\varepsilon_n}} = -\frac{\partial}{\partial\beta} \ln Z_1 \\ &= -\frac{\partial}{\partial\beta} \ln \frac{2}{1-e^{-\beta\hbar\omega}} = \frac{\partial}{\partial\beta} \ln(1 - e^{-\beta\hbar\omega})\end{aligned}$$

## Application 4. Equilibrium radiation

$$Z_1 = \sum_{n=0}^{\infty} 2e^{-\beta\varepsilon_n}, Z_1 = \frac{2}{1-e^{-\beta\hbar\omega}}$$

- The average energy of an oscillator with frequency  $\omega$ :

$$\begin{aligned}\bar{\varepsilon} &= \frac{\sum a_n \varepsilon_n}{\sum a_n} = \frac{\sum e^{-\alpha-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\alpha-\beta\varepsilon_n}} = \frac{\sum e^{-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\beta\varepsilon_n}} = -\frac{\partial}{\partial\beta} \ln Z_1 \\ &= -\frac{\partial}{\partial\beta} \ln \frac{2}{1-e^{-\beta\hbar\omega}} = \frac{\partial}{\partial\beta} \ln(1 - e^{-\beta\hbar\omega}) = \frac{\hbar\omega e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}\end{aligned}$$

# Application 4. Equilibrium radiation

$$Z_1 = \sum_{n=0}^{\infty} 2e^{-\beta\varepsilon_n}, Z_1 = \frac{2}{1-e^{-\beta\hbar\omega}}$$

- The average energy of an oscillator with frequency  $\omega$ :

$$\begin{aligned}\bar{\varepsilon} &= \frac{\sum a_n \varepsilon_n}{\sum a_n} = \frac{\sum e^{-\alpha-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\alpha-\beta\varepsilon_n}} = \frac{\sum e^{-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\beta\varepsilon_n}} = -\frac{\partial}{\partial\beta} \ln Z_1 \\ &= -\frac{\partial}{\partial\beta} \ln \frac{2}{1-e^{-\beta\hbar\omega}} = \frac{\partial}{\partial\beta} \ln(1 - e^{-\beta\hbar\omega}) = \frac{\hbar\omega e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \\ &= \frac{\hbar\omega}{e^{\beta\hbar\omega}-1}.\end{aligned}$$

# Application 4. Equilibrium radiation

$$Z_1 = \sum_{n=0}^{\infty} 2e^{-\beta\varepsilon_n}, Z_1 = \frac{2}{1-e^{-\beta\hbar\omega}}$$

- The average energy of an oscillator with frequency  $\omega$ :

$$\begin{aligned}\bar{\varepsilon} &= \frac{\sum a_n \varepsilon_n}{\sum a_n} = \frac{\sum e^{-\alpha-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\alpha-\beta\varepsilon_n}} = \frac{\sum e^{-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\beta\varepsilon_n}} = -\frac{\partial}{\partial\beta} \ln Z_1 \\ &= -\frac{\partial}{\partial\beta} \ln \frac{2}{1-e^{-\beta\hbar\omega}} = \frac{\partial}{\partial\beta} \ln(1 - e^{-\beta\hbar\omega}) = \frac{\hbar\omega e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \\ &= \frac{\hbar\omega}{e^{\beta\hbar\omega}-1}.\end{aligned}$$

- The internal energy:  $U_\omega d\omega = \bar{\varepsilon} D(\omega) d\omega$

# Application 4. Equilibrium radiation

$$Z_1 = \sum_{n=0}^{\infty} 2e^{-\beta\varepsilon_n}, Z_1 = \frac{2}{1-e^{-\beta\hbar\omega}}$$

- The average energy of an oscillator with frequency  $\omega$ :

$$\begin{aligned}\bar{\varepsilon} &= \frac{\sum a_n \varepsilon_n}{\sum a_n} = \frac{\sum e^{-\alpha-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\alpha-\beta\varepsilon_n}} = \frac{\sum e^{-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\beta\varepsilon_n}} = -\frac{\partial}{\partial\beta} \ln Z_1 \\ &= -\frac{\partial}{\partial\beta} \ln \frac{2}{1-e^{-\beta\hbar\omega}} = \frac{\partial}{\partial\beta} \ln(1 - e^{-\beta\hbar\omega}) = \frac{\hbar\omega e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \\ &= \frac{\hbar\omega}{e^{\beta\hbar\omega}-1}.\end{aligned}$$

- The internal energy:  $U_\omega d\omega = \bar{\varepsilon} D(\omega) d\omega$   

$$= \frac{\hbar\omega}{e^{\beta\hbar\omega}-1} \frac{V}{\pi^2 c^3} \omega^2 d\omega$$

# Application 4. Equilibrium radiation

$$Z_1 = \sum_{n=0}^{\infty} 2e^{-\beta\varepsilon_n}, Z_1 = \frac{2}{1-e^{-\beta\hbar\omega}}$$

- The average energy of an oscillator with frequency  $\omega$ :

$$\begin{aligned}\bar{\varepsilon} &= \frac{\sum a_n \varepsilon_n}{\sum a_n} = \frac{\sum e^{-\alpha-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\alpha-\beta\varepsilon_n}} = \frac{\sum e^{-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\beta\varepsilon_n}} = -\frac{\partial}{\partial\beta} \ln Z_1 \\ &= -\frac{\partial}{\partial\beta} \ln \frac{2}{1-e^{-\beta\hbar\omega}} = \frac{\partial}{\partial\beta} \ln(1-e^{-\beta\hbar\omega}) = \frac{\hbar\omega e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \\ &= \frac{\hbar\omega}{e^{\beta\hbar\omega}-1}.\end{aligned}$$

- The internal energy:  $U_\omega d\omega = \bar{\varepsilon} D(\omega) d\omega$   

$$= \frac{\hbar\omega}{e^{\beta\hbar\omega}-1} \frac{V}{\pi^2 c^3} \omega^2 d\omega = \frac{V}{2\pi^3 c^3} \frac{\hbar\omega^3}{e^{\frac{\hbar\omega}{2\pi kT}}-1} d\omega.$$

# Application 4. Equilibrium radiation

$$Z_1 = \sum_{n=0}^{\infty} 2e^{-\beta\varepsilon_n}, Z_1 = \frac{2}{1-e^{-\beta\hbar\omega}}$$

- The average energy of an oscillator with frequency  $\omega$ :

$$\begin{aligned}\bar{\varepsilon} &= \frac{\sum a_n \varepsilon_n}{\sum a_n} = \frac{\sum e^{-\alpha-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\alpha-\beta\varepsilon_n}} = \frac{\sum e^{-\beta\varepsilon_n} \varepsilon_n}{\sum e^{-\beta\varepsilon_n}} = -\frac{\partial}{\partial\beta} \ln Z_1 \\ &= -\frac{\partial}{\partial\beta} \ln \frac{2}{1-e^{-\beta\hbar\omega}} = \frac{\partial}{\partial\beta} \ln(1 - e^{-\beta\hbar\omega}) = \frac{\hbar\omega e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \\ &= \frac{\hbar\omega}{e^{\beta\hbar\omega}-1}.\end{aligned}$$

- The internal energy:  $U_\omega d\omega = \bar{\varepsilon} D(\omega) d\omega$   

$$= \frac{\hbar\omega}{e^{\beta\hbar\omega}-1} \frac{V}{\pi^2 c^3} \omega^2 d\omega = \frac{V}{2\pi^3 c^3} \frac{\hbar\omega^3}{e^{\frac{\hbar\omega}{2\pi kT}}-1} d\omega.$$

Exactly the black body radiation formula!



# Table of contents

## 1 §7 Boltzmann Statistics

- 7.1 Thermal quantities in statistics
- 7.2 Equation of state of ideal gas
- 7.3 Maxwell speed distribution
- 7.4 Energy equipartition theorem
- 7.5 Internal energy and heat capacity of the ideal gas
- 7.6 Entropy of the ideal gas
- 7.7 Einstein's theory on heat capacity of solid
- 7.8 Paramagnetic solid
- 7.9 Negative temperature

## 7.5 Internal energy and heat capacity of the ideal gas

## 7.5 Internal energy and heat capacity of the ideal gas

- Object: diatomic molecules.

## 7.5 Internal energy and heat capacity of the ideal gas

- Object: diatomic molecules.
- Method: directly use the partition function.

## 7.5 Internal energy and heat capacity of the ideal gas

- Object: diatomic molecules.
- Method: directly use the partition function.
- Reason: equipartition theorem only suitable for classical limits. However, in normal pressure and temperature, relative motion is not in classical limits.

## 7.5 Internal energy and heat capacity of the ideal gas

- Object: diatomic molecules.
- Method: directly use the partition function.
- Reason: equipartition theorem only suitable for classical limits. However, in normal pressure and temperature, relative motion is not in classical limits.
- Energy:  $\varepsilon = \varepsilon^t + \varepsilon^v + \varepsilon^r$ , where  $t$  for translational motion,  $v$  for vibration, and  $r$  for rotation.

## 7.5 Internal energy and heat capacity of the ideal gas

- Object: diatomic molecules.
- Method: directly use the partition function.
- Reason: equipartition theorem only suitable for classical limits. However, in normal pressure and temperature, relative motion is not in classical limits.
- Energy:  $\varepsilon = \varepsilon^t + \varepsilon^v + \varepsilon^r$ , where  $t$  for translational motion,  $v$  for vibration, and  $r$  for rotation.
- Partition function:  $Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$

## 7.5 Internal energy and heat capacity of the ideal gas

- Object: diatomic molecules.
- Method: directly use the partition function.
- Reason: equipartition theorem only suitable for classical limits. However, in normal pressure and temperature, relative motion is not in classical limits.
- Energy:  $\varepsilon = \varepsilon^t + \varepsilon^v + \varepsilon^r$ , where  $t$  for translational motion,  $v$  for vibration, and  $r$  for rotation.
- Partition function: 
$$Z_1 = \sum \omega_l e^{-\beta \varepsilon_l}$$
$$= \sum \omega^t \omega^v \omega^r e^{-\beta(\varepsilon^t + \varepsilon^v + \varepsilon^r)}$$



## 7.5 Internal energy and heat capacity of the ideal gas

- Object: diatomic molecules.
- Method: directly use the partition function.
- Reason: equipartition theorem only suitable for classical limits. However, in normal pressure and temperature, relative motion is not in classical limits.
- Energy:  $\varepsilon = \varepsilon^t + \varepsilon^v + \varepsilon^r$ , where  $t$  for translational motion,  $v$  for vibration, and  $r$  for rotation.
- Partition function: 
$$\begin{aligned} Z_1 &= \sum \omega_l e^{-\beta \varepsilon_l} \\ &= \sum \omega^t \omega^v \omega^r e^{-\beta(\varepsilon^t + \varepsilon^v + \varepsilon^r)} \\ &= \sum \omega^t e^{-\beta \varepsilon^t} \sum \omega^v e^{-\beta \varepsilon^v} \sum \omega^r e^{-\beta \varepsilon^r} \end{aligned}$$

## 7.5 Internal energy and heat capacity of the ideal gas

- Object: diatomic molecules.
- Method: directly use the partition function.
- Reason: equipartition theorem only suitable for classical limits. However, in normal pressure and temperature, relative motion is not in classical limits.
- Energy:  $\varepsilon = \varepsilon^t + \varepsilon^v + \varepsilon^r$ , where  $t$  for translational motion,  $v$  for vibration, and  $r$  for rotation.
- Partition function: 
$$\begin{aligned} Z_1 &= \sum \omega_l e^{-\beta \varepsilon_l} \\ &= \sum \omega^t \omega^v \omega^r e^{-\beta(\varepsilon^t + \varepsilon^v + \varepsilon^r)} \\ &= \sum \omega^t e^{-\beta \varepsilon^t} \sum \omega^v e^{-\beta \varepsilon^v} \sum \omega^r e^{-\beta \varepsilon^r} = Z_1^t \cdot Z_1^v \cdot Z_1^r. \end{aligned}$$

# Internal energy and heat capacity of the ideal gas

- Internal energy:  $U = -N \frac{\partial}{\partial \beta} \ln Z_1$

# Internal energy and heat capacity of the ideal gas

- Internal energy: 
$$U = -N \frac{\partial}{\partial \beta} \ln Z_1$$
$$= -N \frac{\partial}{\partial \beta} (\ln Z_1^t + \ln Z_1^v + \ln Z_1^r)$$

# Internal energy and heat capacity of the ideal gas

- Internal energy: 
$$U = -N \frac{\partial}{\partial \beta} \ln Z_1$$
$$= -N \frac{\partial}{\partial \beta} (\ln Z_1^t + \ln Z_1^v + \ln Z_1^r) = U^t + U^v + U^r.$$

# Internal energy and heat capacity of the ideal gas

- Internal energy:  $U = -N \frac{\partial}{\partial \beta} \ln Z_1$   
 $= -N \frac{\partial}{\partial \beta} (\ln Z_1^t + \ln Z_1^v + \ln Z_1^r) = U^t + U^v + U^r.$
- Heat capacity at constant volume:  
 $C_V = C_V^t + C_V^v + C_V^r.$

# Internal energy and heat capacity of the ideal gas

- Internal energy:  $U = -N \frac{\partial}{\partial \beta} \ln Z_1$   
 $= -N \frac{\partial}{\partial \beta} (\ln Z_1^t + \ln Z_1^v + \ln Z_1^r) = U^t + U^v + U^r.$
- Heat capacity at constant volume:  
 $C_V = C_V^t + C_V^v + C_V^r.$
- i. Translational motion part:

# Internal energy and heat capacity of the ideal gas

- Internal energy:  $U = -N \frac{\partial}{\partial \beta} \ln Z_1$   
 $= -N \frac{\partial}{\partial \beta} (\ln Z_1^t + \ln Z_1^v + \ln Z_1^r) = U^t + U^v + U^r.$
- Heat capacity at constant volume:  
 $C_V = C_V^t + C_V^v + C_V^r.$
- i. Translational motion part: (7.2.4)  $Z_1^t = V \left( \frac{2\pi m}{h^2 \beta} \right)^{3/2}.$



# Internal energy and heat capacity of the ideal gas

- Internal energy:  $U = -N \frac{\partial}{\partial \beta} \ln Z_1$   
 $= -N \frac{\partial}{\partial \beta} (\ln Z_1^t + \ln Z_1^v + \ln Z_1^r) = U^t + U^v + U^r.$
- Heat capacity at constant volume:  
 $C_V = C_V^t + C_V^v + C_V^r.$
- i. Translational motion part: (7.2.4)  $Z_1^t = V \left( \frac{2\pi m}{h^2 \beta} \right)^{3/2}.$   
 $U^t = \frac{3}{2} N k T,$

# Internal energy and heat capacity of the ideal gas

- Internal energy:  $U = -N \frac{\partial}{\partial \beta} \ln Z_1$   
 $= -N \frac{\partial}{\partial \beta} (\ln Z_1^t + \ln Z_1^v + \ln Z_1^r) = U^t + U^v + U^r.$
- Heat capacity at constant volume:  
 $C_V = C_V^t + C_V^v + C_V^r.$
- i. Translational motion part: (7.2.4)  $Z_1^t = V \left( \frac{2\pi m}{h^2 \beta} \right)^{3/2}.$   
 $U^t = \frac{3}{2} N k T, C_V^t = \frac{3}{2} N k.$

# Internal energy and heat capacity of the ideal gas

- Internal energy:  $U = -N \frac{\partial}{\partial \beta} \ln Z_1$   
 $= -N \frac{\partial}{\partial \beta} (\ln Z_1^t + \ln Z_1^v + \ln Z_1^r) = U^t + U^v + U^r.$
- Heat capacity at constant volume:  
 $C_V = C_V^t + C_V^v + C_V^r.$
- i. Translational motion part: (7.2.4)  $Z_1^t = V \left( \frac{2\pi m}{h^2 \beta} \right)^{3/2}.$   
 $U^t = \frac{3}{2} N k T, C_V^t = \frac{3}{2} N k.$
- ii. Vibration part:

# Internal energy and heat capacity of the ideal gas

- Internal energy:  $U = -N \frac{\partial}{\partial \beta} \ln Z_1$   
 $= -N \frac{\partial}{\partial \beta} (\ln Z_1^t + \ln Z_1^v + \ln Z_1^r) = U^t + U^v + U^r.$
- Heat capacity at constant volume:  
 $C_V = C_V^t + C_V^v + C_V^r.$
- i. Translational motion part: (7.2.4)  $Z_1^t = V \left( \frac{2\pi m}{h^2 \beta} \right)^{3/2}.$   
 $U^t = \frac{3}{2} N k T, C_V^t = \frac{3}{2} N k.$
- ii. Vibration part: taken as linear harmonic vibration,  
 $\varepsilon_n = (n + \frac{1}{2}) \hbar \omega.$

# Internal energy and heat capacity of the ideal gas

- Internal energy:  $U = -N \frac{\partial}{\partial \beta} \ln Z_1$   
 $= -N \frac{\partial}{\partial \beta} (\ln Z_1^t + \ln Z_1^v + \ln Z_1^r) = U^t + U^v + U^r.$
- Heat capacity at constant volume:  
 $C_V = C_V^t + C_V^v + C_V^r.$
- i. Translational motion part: (7.2.4)  $Z_1^t = V \left( \frac{2\pi m}{h^2 \beta} \right)^{3/2}.$   
 $U^t = \frac{3}{2} N k T, C_V^t = \frac{3}{2} N k.$
- ii. Vibration part: taken as linear harmonic vibration,  
 $\varepsilon_n = (n + \frac{1}{2}) \hbar \omega.$

Partition function:  $Z_1^v = \sum_{n=0}^{\infty} e^{-\beta \hbar \omega (n + \frac{1}{2})}$

# Internal energy and heat capacity of the ideal gas

- Internal energy:  $U = -N \frac{\partial}{\partial \beta} \ln Z_1$   
 $= -N \frac{\partial}{\partial \beta} (\ln Z_1^t + \ln Z_1^v + \ln Z_1^r) = U^t + U^v + U^r.$
- Heat capacity at constant volume:  
 $C_V = C_V^t + C_V^v + C_V^r.$
- i. Translational motion part: (7.2.4)  $Z_1^t = V \left( \frac{2\pi m}{h^2 \beta} \right)^{3/2}.$   
 $U^t = \frac{3}{2} N k T, C_V^t = \frac{3}{2} N k.$
- ii. Vibration part: taken as linear harmonic vibration,  
 $\varepsilon_n = (n + \frac{1}{2}) \hbar \omega.$

Partition function:  $Z_1^v = \sum_{n=0}^{\infty} e^{-\beta \hbar \omega (n + \frac{1}{2})}$

$$= e^{-\frac{1}{2} \beta \hbar \omega} \sum_{n=0}^{\infty} (e^{-\beta \hbar \omega})^n$$

# Internal energy and heat capacity of the ideal gas

- Internal energy:  $U = -N \frac{\partial}{\partial \beta} \ln Z_1$   
 $= -N \frac{\partial}{\partial \beta} (\ln Z_1^t + \ln Z_1^v + \ln Z_1^r) = U^t + U^v + U^r.$
- Heat capacity at constant volume:  
 $C_V = C_V^t + C_V^v + C_V^r.$
- i. Translational motion part: (7.2.4)  $Z_1^t = V \left( \frac{2\pi m}{h^2 \beta} \right)^{3/2}.$   
 $U^t = \frac{3}{2} N k T, C_V^t = \frac{3}{2} N k.$
- ii. Vibration part: taken as linear harmonic vibration,  
 $\varepsilon_n = (n + \frac{1}{2}) \hbar \omega.$

Partition function:  $Z_1^v = \sum_{n=0}^{\infty} e^{-\beta \hbar \omega (n + \frac{1}{2})}$

$$= e^{-\frac{1}{2} \beta \hbar \omega} \sum_{n=0}^{\infty} (e^{-\beta \hbar \omega})^n = \frac{e^{-\frac{1}{2} \beta \hbar \omega}}{1 - e^{-\beta \hbar \omega}}.$$

# Internal energy and heat capacity of the ideal gas

- $Z_1^v = \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}.$



# Internal energy and heat capacity of the ideal gas

- $Z_1^v = \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}.$
- $U^v = -N \frac{\partial}{\partial \beta} \ln Z_1^v$

# Internal energy and heat capacity of the ideal gas

- $Z_1^v = \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}.$
- $U^v = -N \frac{\partial}{\partial \beta} \ln Z_1^v = -N \frac{\partial}{\partial \beta} \ln\left(\frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}\right)$

# Internal energy and heat capacity of the ideal gas

- $Z_1^v = \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}.$
- $$U^v = -N \frac{\partial}{\partial \beta} \ln Z_1^v = -N \frac{\partial}{\partial \beta} \ln\left(\frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}\right)$$
$$= -N \frac{\partial}{\partial \beta} \left[ -\frac{1}{2}\beta\hbar\omega - \ln(1 - e^{-\beta\hbar\omega}) \right]$$

# Internal energy and heat capacity of the ideal gas

- $Z_1^v = \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}.$
- $$\begin{aligned} U^v &= -N \frac{\partial}{\partial \beta} \ln Z_1^v = -N \frac{\partial}{\partial \beta} \ln \left( \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \right) \\ &= -N \frac{\partial}{\partial \beta} \left[ -\frac{1}{2}\beta\hbar\omega - \ln(1 - e^{-\beta\hbar\omega}) \right] \\ &= \frac{1}{2}N\hbar\omega + N \frac{\hbar\omega e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \end{aligned}$$

# Internal energy and heat capacity of the ideal gas

- $Z_1^v = \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}.$
- $$\begin{aligned} U^v &= -N \frac{\partial}{\partial \beta} \ln Z_1^v = -N \frac{\partial}{\partial \beta} \ln \left( \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \right) \\ &= -N \frac{\partial}{\partial \beta} \left[ -\frac{1}{2}\beta\hbar\omega - \ln(1 - e^{-\beta\hbar\omega}) \right] \\ &= \frac{1}{2}N\hbar\omega + N \frac{\hbar\omega e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \\ &= \frac{1}{2}N\hbar\omega + \frac{N\hbar\omega}{e^{\beta\hbar\omega}-1}. \end{aligned}$$

# Internal energy and heat capacity of the ideal gas

- $Z_1^v = \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}.$
- $$\begin{aligned} U^v &= -N \frac{\partial}{\partial \beta} \ln Z_1^v = -N \frac{\partial}{\partial \beta} \ln \left( \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \right) \\ &= -N \frac{\partial}{\partial \beta} \left[ -\frac{1}{2}\beta\hbar\omega - \ln(1 - e^{-\beta\hbar\omega}) \right] \\ &= \frac{1}{2}N\hbar\omega + N \frac{\hbar\omega e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \\ &= \frac{1}{2}N\hbar\omega + \frac{N\hbar\omega}{e^{\beta\hbar\omega}-1}. \end{aligned}$$
- $\frac{1}{2}N\hbar\omega$  zero-point energy;

# Internal energy and heat capacity of the ideal gas

- $Z_1^v = \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}.$
- $$\begin{aligned} U^v &= -N \frac{\partial}{\partial \beta} \ln Z_1^v = -N \frac{\partial}{\partial \beta} \ln \left( \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \right) \\ &= -N \frac{\partial}{\partial \beta} \left[ -\frac{1}{2}\beta\hbar\omega - \ln(1 - e^{-\beta\hbar\omega}) \right] \\ &= \frac{1}{2}N\hbar\omega + N \frac{\hbar\omega e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \\ &= \frac{1}{2}N\hbar\omega + \frac{N\hbar\omega}{e^{\beta\hbar\omega}-1}. \end{aligned}$$
- $\frac{1}{2}N\hbar\omega$  zero-point energy;  
 $\frac{N\hbar\omega}{e^{\beta\hbar\omega}-1}$  energy excited by heat.

# Internal energy and heat capacity of the ideal gas

- Define a typical temperature:  $\theta_v$ :  $k\theta_v = \hbar\omega$ .



# Internal energy and heat capacity of the ideal gas

- Define a typical temperature:  $\theta_v$ :  $k\theta_v = \hbar\omega$ . Then
$$U^v = \frac{1}{2}Nk\theta_v + \frac{Nk\theta_v}{e^{\theta_v/T} - 1},$$

# Internal energy and heat capacity of the ideal gas

- Define a typical temperature:  $\theta_v$ :  $k\theta_v = \hbar\omega$ . Then

$$U^v = \frac{1}{2}Nk\theta_v + \frac{Nk\theta_v}{e^{\theta_v/T} - 1},$$

$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 \frac{e^{\theta_v/T}}{(e^{\theta_v/T} - 1)^2}.$$

# Internal energy and heat capacity of the ideal gas

- Define a typical temperature:  $\theta_v$ :  $k\theta_v = \hbar\omega$ . Then
$$U^v = \frac{1}{2}Nk\theta_v + \frac{Nk\theta_v}{e^{\theta_v/T}-1},$$
$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 \frac{e^{\theta_v/T}}{(e^{\theta_v/T}-1)^2}.$$
- $\theta_v$  is determined by the property of the material.

# Internal energy and heat capacity of the ideal gas

- Define a typical temperature:  $\theta_v$ :  $k\theta_v = \hbar\omega$ . Then
$$U^v = \frac{1}{2}Nk\theta_v + \frac{Nk\theta_v}{e^{\theta_v/T}-1},$$
$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 \frac{e^{\theta_v/T}}{(e^{\theta_v/T}-1)^2}.$$
- $\theta_v$  is determined by the property of the material.
- If  $T \ll \theta_v$ ,

# Internal energy and heat capacity of the ideal gas

- Define a typical temperature:  $\theta_v$ :  $k\theta_v = \hbar\omega$ . Then
$$U^v = \frac{1}{2}Nk\theta_v + \frac{Nk\theta_v}{e^{\theta_v/T}-1},$$
$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 \frac{e^{\theta_v/T}}{(e^{\theta_v/T}-1)^2}.$$
- $\theta_v$  is determined by the property of the material.
- If  $T \ll \theta_v$ ,  $U^v = \frac{1}{2}Nk\theta_v + Nk\theta_v e^{-\theta_v/T}$

# Internal energy and heat capacity of the ideal gas

- Define a typical temperature:  $\theta_v$ :  $k\theta_v = \hbar\omega$ . Then
$$U^v = \frac{1}{2}Nk\theta_v + \frac{Nk\theta_v}{e^{\theta_v/T} - 1},$$
$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 \frac{e^{\theta_v/T}}{(e^{\theta_v/T} - 1)^2}.$$
- $\theta_v$  is determined by the property of the material.
- If  $T \ll \theta_v$ ,  $U^v = \frac{1}{2}Nk\theta_v + Nk\theta_v e^{-\theta_v/T}$ 
$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 e^{-\theta_v/T}$$

# Internal energy and heat capacity of the ideal gas

- Define a typical temperature:  $\theta_v$ :  $k\theta_v = \hbar\omega$ . Then
$$U^v = \frac{1}{2}Nk\theta_v + \frac{Nk\theta_v}{e^{\theta_v/T}-1},$$
$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 \frac{e^{\theta_v/T}}{(e^{\theta_v/T}-1)^2}.$$
- $\theta_v$  is determined by the property of the material.
- If  $T \ll \theta_v$ ,  $U^v = \frac{1}{2}Nk\theta_v + Nk\theta_v e^{-\theta_v/T}$ 
$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 e^{-\theta_v/T} \rightarrow 0.$$

# Internal energy and heat capacity of the ideal gas

- Define a typical temperature:  $\theta_v$ :  $k\theta_v = \hbar\omega$ . Then

$$U^v = \frac{1}{2}Nk\theta_v + \frac{Nk\theta_v}{e^{\theta_v/T} - 1},$$

$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 \frac{e^{\theta_v/T}}{(e^{\theta_v/T} - 1)^2}.$$

- $\theta_v$  is determined by the property of the material.

- If  $T \ll \theta_v$ ,  $U^v = \frac{1}{2}Nk\theta_v + Nk\theta_v e^{-\theta_v/T}$

$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 e^{-\theta_v/T} \rightarrow 0.$$

Reason:  $\hbar\omega \gg kT$ .



# Internal energy and heat capacity of the ideal gas

- Define a typical temperature:  $\theta_v$ :  $k\theta_v = \hbar\omega$ . Then
$$U^v = \frac{1}{2}Nk\theta_v + \frac{Nk\theta_v}{e^{\theta_v/T} - 1},$$
$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 \frac{e^{\theta_v/T}}{(e^{\theta_v/T} - 1)^2}.$$
- $\theta_v$  is determined by the property of the material.
- If  $T \ll \theta_v$ ,  $U^v = \frac{1}{2}Nk\theta_v + Nk\theta_v e^{-\theta_v/T}$ 
$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 e^{-\theta_v/T} \rightarrow 0.$$
Reason:  $\hbar\omega \gg kT$ .
- iii. Rotational part.

# Internal energy and heat capacity of the ideal gas

- Define a typical temperature:  $\theta_v$ :  $k\theta_v = \hbar\omega$ . Then
 
$$U^v = \frac{1}{2}Nk\theta_v + \frac{Nk\theta_v}{e^{\theta_v/T} - 1},$$

$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 \frac{e^{\theta_v/T}}{(e^{\theta_v/T} - 1)^2}.$$
- $\theta_v$  is determined by the property of the material.
- If  $T \ll \theta_v$ ,  $U^v = \frac{1}{2}Nk\theta_v + Nk\theta_v e^{-\theta_v/T}$ 

$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 e^{-\theta_v/T} \rightarrow 0.$$
 Reason:  $\hbar\omega \gg kT$ .
- iii. Rotational part.
- iii.1. Diatomic molecule with different nuclei.

# Internal energy and heat capacity of the ideal gas

- Define a typical temperature:  $\theta_v$ :  $k\theta_v = \hbar\omega$ . Then

$$U^v = \frac{1}{2}Nk\theta_v + \frac{Nk\theta_v}{e^{\theta_v/T}-1},$$

$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 \frac{e^{\theta_v/T}}{(e^{\theta_v/T}-1)^2}.$$

- $\theta_v$  is determined by the property of the material.

- If  $T \ll \theta_v$ ,  $U^v = \frac{1}{2}Nk\theta_v + Nk\theta_v e^{-\theta_v/T}$

$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 e^{-\theta_v/T} \rightarrow 0.$$

Reason:  $\hbar\omega \gg kT$ .

- iii. Rotational part.
- iii.1. Diatomic molecule with different nuclei.

$$\varepsilon^r = \frac{l(l+1)}{2I} \hbar^2, \quad l = 0, 1, 2, \dots$$

# Internal energy and heat capacity of the ideal gas

- Define a typical temperature:  $\theta_v$ :  $k\theta_v = \hbar\omega$ . Then

$$U^v = \frac{1}{2}Nk\theta_v + \frac{Nk\theta_v}{e^{\theta_v/T} - 1},$$

$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 \frac{e^{\theta_v/T}}{(e^{\theta_v/T} - 1)^2}.$$

- $\theta_v$  is determined by the property of the material.

- If  $T \ll \theta_v$ ,  $U^v = \frac{1}{2}Nk\theta_v + Nk\theta_v e^{-\theta_v/T}$

$$C_V^v = Nk\left(\frac{\theta_v}{T}\right)^2 e^{-\theta_v/T} \rightarrow 0.$$

Reason:  $\hbar\omega \gg kT$ .

- iii. Rotational part.
- iii.1. Diatomic molecule with different nuclei.

$$\varepsilon^r = \frac{l(l+1)}{2I} \hbar^2, \quad l = 0, 1, 2, \dots$$

Degeneracy:  $\omega_l = 2l + 1$ .

# Internal energy and heat capacity of the ideal gas

- Partition function:  $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}.$

# Internal energy and heat capacity of the ideal gas

- Partition function:  $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$ .
- Define typical temperature  $\theta_r$ :  $k\theta_r = \frac{\hbar^2}{2I}$ .

# Internal energy and heat capacity of the ideal gas

- Partition function:  $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$ .
- Define typical temperature  $\theta_r$ :  $k\theta_r = \frac{\hbar^2}{2I}$ .
- $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$ .

# Internal energy and heat capacity of the ideal gas

- Partition function:  $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$ .
- Define typical temperature  $\theta_r$ :  $k\theta_r = \frac{\hbar^2}{2I}$ .
- $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$ .
- If  $\theta_r/T \ll 1$ ,  $\varepsilon^r$  is approximately continuous.



# Internal energy and heat capacity of the ideal gas

- Partition function:  $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$ .
- Define typical temperature  $\theta_r$ :  $k\theta_r = \frac{\hbar^2}{2I}$ .
- $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$ .
- If  $\theta_r/T \ll 1$ ,  $\varepsilon^r$  is approximately continuous. Set  $x = l(l+1)\frac{\theta_r}{T}$ , then  $dx = (2l+1)\frac{\theta_r}{T}\Delta l$ .

# Internal energy and heat capacity of the ideal gas

- Partition function:  $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$ .
- Define typical temperature  $\theta_r$ :  $k\theta_r = \frac{\hbar^2}{2I}$ .
- $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$ .
- If  $\theta_r/T \ll 1$ ,  $\varepsilon^r$  is approximately continuous. Set  $x = l(l+1)\frac{\theta_r}{T}$ , then  $dx = (2l+1)\frac{\theta_r}{T}\Delta l$ .  

$$Z_1^r = \int_0^{\infty} (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)}$$

# Internal energy and heat capacity of the ideal gas

- Partition function:  $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$ .
- Define typical temperature  $\theta_r$ :  $k\theta_r = \frac{\hbar^2}{2I}$ .
- $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$ .
- If  $\theta_r/T \ll 1$ ,  $\varepsilon^r$  is approximately continuous. Set  $x = l(l+1)\frac{\theta_r}{T}$ , then  $dx = (2l+1)\frac{\theta_r}{T}\Delta l$ .  

$$Z_1^r = \int_0^{\infty} (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{T}{\theta_r} \int_0^{\infty} e^{-x} dx$$

# Internal energy and heat capacity of the ideal gas

- Partition function:  $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$ .
- Define typical temperature  $\theta_r$ :  $k\theta_r = \frac{\hbar^2}{2I}$ .
- $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$ .
- If  $\theta_r/T \ll 1$ ,  $\varepsilon^r$  is approximately continuous. Set  $x = l(l+1)\frac{\theta_r}{T}$ , then  $dx = (2l+1)\frac{\theta_r}{T}\Delta l$ .  

$$Z_1^r = \int_0^{\infty} (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{T}{\theta_r} \int_0^{\infty} e^{-x} dx = \frac{T}{\theta_r}$$

# Internal energy and heat capacity of the ideal gas

- Partition function:  $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$ .
- Define typical temperature  $\theta_r$ :  $k\theta_r = \frac{\hbar^2}{2I}$ .
- $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$ .
- If  $\theta_r/T \ll 1$ ,  $\varepsilon^r$  is approximately continuous. Set  $x = l(l+1)\frac{\theta_r}{T}$ , then  $dx = (2l+1)\frac{\theta_r}{T}\Delta l$ .  

$$Z_1^r = \int_0^{\infty} (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{T}{\theta_r} \int_0^{\infty} e^{-x} dx = \frac{T}{\theta_r}$$

$$= T \frac{2Ik}{\hbar^2}$$

# Internal energy and heat capacity of the ideal gas

- Partition function:  $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$ .
- Define typical temperature  $\theta_r$ :  $k\theta_r = \frac{\hbar^2}{2I}$ .
- $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$ .
- If  $\theta_r/T \ll 1$ ,  $\varepsilon^r$  is approximately continuous. Set  $x = l(l+1)\frac{\theta_r}{T}$ , then  $dx = (2l+1)\frac{\theta_r}{T}\Delta l$ .  

$$Z_1^r = \int_0^{\infty} (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{T}{\theta_r} \int_0^{\infty} e^{-x} dx = \frac{T}{\theta_r}$$

$$= T \frac{2Ik}{\hbar^2} = \frac{2I}{\beta \hbar^2}.$$

# Internal energy and heat capacity of the ideal gas

- Partition function:  $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$ .
- Define typical temperature  $\theta_r$ :  $k\theta_r = \frac{\hbar^2}{2I}$ .
- $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$ .
- If  $\theta_r/T \ll 1$ ,  $\varepsilon^r$  is approximately continuous. Set  $x = l(l+1)\frac{\theta_r}{T}$ , then  $dx = (2l+1)\frac{\theta_r}{T}\Delta l$ .  

$$Z_1^r = \int_0^{\infty} (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{T}{\theta_r} \int_0^{\infty} e^{-x} dx = \frac{T}{\theta_r}$$

$$= T \frac{2Ik}{\hbar^2} = \frac{2I}{\beta \hbar^2}.$$
- $U^r = -N \frac{\partial}{\partial \beta} \ln Z_1^r$

# Internal energy and heat capacity of the ideal gas

- Partition function:  $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$ .
- Define typical temperature  $\theta_r$ :  $k\theta_r = \frac{\hbar^2}{2I}$ .
- $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$ .
- If  $\theta_r/T \ll 1$ ,  $\varepsilon^r$  is approximately continuous. Set  $x = l(l+1)\frac{\theta_r}{T}$ , then  $dx = (2l+1)\frac{\theta_r}{T}\Delta l$ .  

$$Z_1^r = \int_0^{\infty} (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{T}{\theta_r} \int_0^{\infty} e^{-x} dx = \frac{T}{\theta_r}$$

$$= T \frac{2Ik}{\hbar^2} = \frac{2I}{\beta\hbar^2}.$$
- $U^r = -N \frac{\partial}{\partial \beta} \ln Z_1^r = N \left( \frac{2I}{\beta\hbar^2} \right)^{-1} \cdot \frac{2I}{\beta^2\hbar^2}$



# Internal energy and heat capacity of the ideal gas

- Partition function:  $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$ .
- Define typical temperature  $\theta_r$ :  $k\theta_r = \frac{\hbar^2}{2I}$ .
- $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$ .
- If  $\theta_r/T \ll 1$ ,  $\varepsilon^r$  is approximately continuous. Set  $x = l(l+1)\frac{\theta_r}{T}$ , then  $dx = (2l+1)\frac{\theta_r}{T}\Delta l$ .  

$$Z_1^r = \int_0^{\infty} (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{T}{\theta_r} \int_0^{\infty} e^{-x} dx = \frac{T}{\theta_r}$$

$$= T \frac{2Ik}{\hbar^2} = \frac{2I}{\beta \hbar^2}.$$
- $U^r = -N \frac{\partial}{\partial \beta} \ln Z_1^r = N \left( \frac{2I}{\beta \hbar^2} \right)^{-1} \cdot \frac{2I}{\beta^2 \hbar^2} = \frac{N}{\beta}$

# Internal energy and heat capacity of the ideal gas

- Partition function:  $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$ .
- Define typical temperature  $\theta_r$ :  $k\theta_r = \frac{\hbar^2}{2I}$ .
- $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$ .
- If  $\theta_r/T \ll 1$ ,  $\varepsilon^r$  is approximately continuous. Set  $x = l(l+1)\frac{\theta_r}{T}$ , then  $dx = (2l+1)\frac{\theta_r}{T}\Delta l$ .  

$$Z_1^r = \int_0^{\infty} (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{T}{\theta_r} \int_0^{\infty} e^{-x} dx = \frac{T}{\theta_r}$$

$$= T \frac{2Ik}{\hbar^2} = \frac{2I}{\beta \hbar^2}.$$
- $U^r = -N \frac{\partial}{\partial \beta} \ln Z_1^r = N \left( \frac{2I}{\beta \hbar^2} \right)^{-1} \cdot \frac{2I}{\beta^2 \hbar^2} = \frac{N}{\beta} = NkT.$

# Internal energy and heat capacity of the ideal gas

- Partition function:  $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$ .
- Define typical temperature  $\theta_r$ :  $k\theta_r = \frac{\hbar^2}{2I}$ .
- $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$ .
- If  $\theta_r/T \ll 1$ ,  $\varepsilon^r$  is approximately continuous. Set  $x = l(l+1)\frac{\theta_r}{T}$ , then  $dx = (2l+1)\frac{\theta_r}{T}\Delta l$ .  

$$Z_1^r = \int_0^{\infty} (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{T}{\theta_r} \int_0^{\infty} e^{-x} dx = \frac{T}{\theta_r}$$

$$= T \frac{2Ik}{\hbar^2} = \frac{2I}{\beta \hbar^2}.$$
- $U^r = -N \frac{\partial}{\partial \beta} \ln Z_1^r = N \left( \frac{2I}{\beta \hbar^2} \right)^{-1} \cdot \frac{2I}{\beta^2 \hbar^2} = \frac{N}{\beta} = NkT.$
- $C_V^r = Nk.$

# Internal energy and heat capacity of the ideal gas

- Partition function:  $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\hbar^2}{2IkT}}$ .
- Define typical temperature  $\theta_r$ :  $k\theta_r = \frac{\hbar^2}{2I}$ .
- $Z_1^r = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$ .
- If  $\theta_r/T \ll 1$ ,  $\varepsilon^r$  is approximately continuous. Set  $x = l(l+1)\frac{\theta_r}{T}$ , then  $dx = (2l+1)\frac{\theta_r}{T}\Delta l$ .  

$$Z_1^r = \int_0^{\infty} (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{T}{\theta_r} \int_0^{\infty} e^{-x} dx = \frac{T}{\theta_r}$$

$$= T \frac{2Ik}{\hbar^2} = \frac{2I}{\beta \hbar^2}.$$
- $U^r = -N \frac{\partial}{\partial \beta} \ln Z_1^r = N \left( \frac{2I}{\beta \hbar^2} \right)^{-1} \cdot \frac{2I}{\beta^2 \hbar^2} = \frac{N}{\beta} = NkT$ .
- $C_V^r = Nk$ .
- At normal temperature (Table 7.5),  $\frac{\theta_r}{T} \ll 1$  holds.

# Internal energy and heat capacity of the ideal gas

- iii.2. Diatomic molecule with the same nuclei.  $H_2$  as an example.

# Internal energy and heat capacity of the ideal gas

- iii.2. Diatomic molecule with the same nuclei.  $H_2$  as an example.
- At normal temperature, there are  $3/4$  orthohydrogen ( $l$  odd) and  $1/4$  parahydrogen ( $l$  even).

# Internal energy and heat capacity of the ideal gas

- iii.2. Diatomic molecule with the same nuclei.  $H_2$  as an example.
- At normal temperature, there are 3/4 orthohydrogen ( $l$  odd) and 1/4 parahydrogen ( $l$  even).
- Orthohydrogen:  $Z_{1o}^r = \sum_{l=1,3,\dots} (2l+1)e^{-\frac{l(l+1)\theta_r}{T}};$

# Internal energy and heat capacity of the ideal gas

- iii.2. Diatomic molecule with the same nuclei.  $H_2$  as an example.
- At normal temperature, there are 3/4 orthohydrogen ( $l$  odd) and 1/4 parahydrogen ( $l$  even).
- Orthohydrogen:  $Z_{1o}^r = \sum_{l=1,3,\dots} (2l+1)e^{-\frac{l(l+1)\theta_r}{T}};$

$$\text{Parahydrogen: } Z_{1p}^r = \sum_{l=0,2,\dots} (2l+1)e^{-\frac{l(l+1)\theta_r}{T}}.$$



# Internal energy and heat capacity of the ideal gas

- iii.2. Diatomic molecule with the same nuclei.  $H_2$  as an example.
- At normal temperature, there are 3/4 orthohydrogen ( $l$  odd) and 1/4 parahydrogen ( $l$  even).
- Orthohydrogen:  $Z_{1o}^r = \sum_{l=1,3,\dots} (2l+1)e^{-\frac{l(l+1)\theta_r}{T}}$ ;  
Parahydrogen:  $Z_{1p}^r = \sum_{l=0,2,\dots} (2l+1)e^{-\frac{l(l+1)\theta_r}{T}}$ .
- $U^r = U_o^r + U_p^r$

# Internal energy and heat capacity of the ideal gas

- iii.2. Diatomic molecule with the same nuclei.  $H_2$  as an example.
- At normal temperature, there are 3/4 orthohydrogen ( $l$  odd) and 1/4 parahydrogen ( $l$  even).

- Orthohydrogen:  $Z_{1o}^r = \sum_{l=1,3,\dots} (2l+1)e^{-\frac{l(l+1)\theta_r}{T}};$

Parahydrogen:  $Z_{1p}^r = \sum_{l=0,2,\dots} (2l+1)e^{-\frac{l(l+1)\theta_r}{T}}.$

- $U^r = U_o^r + U_p^r = -\frac{3}{4}N \frac{\partial}{\partial \beta} \ln Z_{1o}^r - \frac{1}{4}N \frac{\partial}{\partial \beta} \ln Z_{1p}^r.$

# Internal energy and heat capacity of the ideal gas

- iii.2. Diatomic molecule with the same nuclei.  $H_2$  as an example.
- At normal temperature, there are 3/4 orthohydrogen ( $l$  odd) and 1/4 parahydrogen ( $l$  even).

- Orthohydrogen:  $Z_{1o}^r = \sum_{l=1,3,\dots} (2l+1)e^{-\frac{l(l+1)\theta_r}{T}};$

Parahydrogen:  $Z_{1p}^r = \sum_{l=0,2,\dots} (2l+1)e^{-\frac{l(l+1)\theta_r}{T}}.$

- $U^r = U_o^r + U_p^r = -\frac{3}{4}N \frac{\partial}{\partial \beta} \ln Z_{1o}^r - \frac{1}{4}N \frac{\partial}{\partial \beta} \ln Z_{1p}^r.$
- If  $\theta_r \ll T$ , the hydrogen molecule is in high  $l$  state,  
 $l \sim l+1,$

# Internal energy and heat capacity of the ideal gas

- iii.2. Diatomic molecule with the same nuclei.  $H_2$  as an example.
- At normal temperature, there are 3/4 orthohydrogen ( $l$  odd) and 1/4 parahydrogen ( $l$  even).
- Orthohydrogen:  $Z_{1o}^r = \sum_{l=1,3,\dots} (2l+1)e^{-\frac{l(l+1)\theta_r}{T}}$ ;

$$\text{Parahydrogen: } Z_{1p}^r = \sum_{l=0,2,\dots} (2l+1)e^{-\frac{l(l+1)\theta_r}{T}}.$$

- $U^r = U_o^r + U_p^r = -\frac{3}{4}N \frac{\partial}{\partial \beta} \ln Z_{1o}^r - \frac{1}{4}N \frac{\partial}{\partial \beta} \ln Z_{1p}^r.$
- If  $\theta_r \ll T$ , the hydrogen molecule is in high  $l$  state,  $l \sim l+1$ , then  $\sum_{l=0,2,\dots}^{\infty} \dots \simeq \sum_{l=1,3,\dots}^{\infty} \dots \simeq \frac{1}{2} \sum_{l=0,1,\dots}^{\infty} \dots$

# Internal energy and heat capacity of the ideal gas

- $Z_{1o}^r \simeq Z_{1p}^r \simeq \frac{1}{2} \sum_{l=0,1,\dots} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$

# Internal energy and heat capacity of the ideal gas

- $$\begin{aligned} Z_{1o}^r &\simeq Z_{1p}^r \simeq \frac{1}{2} \sum_{l=0,1,\dots} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}} \\ &\simeq \frac{1}{2} \int_0^\infty (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} \end{aligned}$$

# Internal energy and heat capacity of the ideal gas

- $$\begin{aligned} Z_{1o}^r &\simeq Z_{1p}^r \simeq \frac{1}{2} \sum_{l=0,1,\dots} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}} \\ &\simeq \frac{1}{2} \int_0^\infty (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{1}{2} \frac{T}{\theta_r} \end{aligned}$$

# Internal energy and heat capacity of the ideal gas

- $$\begin{aligned} Z_{1o}^r &\simeq Z_{1p}^r \simeq \frac{1}{2} \sum_{l=0,1,\dots} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}} \\ &\simeq \frac{1}{2} \int_0^\infty (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{1}{2} \frac{T}{\theta_r} = \frac{I}{\beta \hbar^2}. \end{aligned}$$



# Internal energy and heat capacity of the ideal gas

- $Z_{1o}^r \simeq Z_{1p}^r \simeq \frac{1}{2} \sum_{l=0,1,\dots} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$   
 $\simeq \frac{1}{2} \int_0^\infty (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{1}{2} \frac{T}{\theta_r} = \frac{I}{\beta \hbar^2}.$
- $U^r = -\frac{3}{4} N \frac{\partial}{\partial \beta} \ln Z_{1o}^r - \frac{1}{4} N \frac{\partial}{\partial \beta} \ln Z_{1p}^r$

# Internal energy and heat capacity of the ideal gas

- $Z_{1o}^r \simeq Z_{1p}^r \simeq \frac{1}{2} \sum_{l=0,1,\dots} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$   
 $\simeq \frac{1}{2} \int_0^\infty (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{1}{2} \frac{T}{\theta_r} = \frac{I}{\beta \hbar^2}.$
- $U^r = -\frac{3}{4} N \frac{\partial}{\partial \beta} \ln Z_{1o}^r - \frac{1}{4} N \frac{\partial}{\partial \beta} \ln Z_{1p}^r \simeq -N \frac{\partial}{\partial \beta} \ln Z_{1o}^r$

# Internal energy and heat capacity of the ideal gas

- $Z_{1o}^r \simeq Z_{1p}^r \simeq \frac{1}{2} \sum_{l=0,1,\dots} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$   
 $\simeq \frac{1}{2} \int_0^\infty (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{1}{2} \frac{T}{\theta_r} = \frac{I}{\beta \hbar^2}.$
- $U^r = -\frac{3}{4} N \frac{\partial}{\partial \beta} \ln Z_{1o}^r - \frac{1}{4} N \frac{\partial}{\partial \beta} \ln Z_{1p}^r \simeq -N \frac{\partial}{\partial \beta} \ln Z_{1o}^r$   
 $= -N \frac{\partial}{\partial \beta} \ln \frac{I}{\beta \hbar^2}$

# Internal energy and heat capacity of the ideal gas

- $$Z_{1o}^r \simeq Z_{1p}^r \simeq \frac{1}{2} \sum_{l=0,1,\dots} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$$

$$\simeq \frac{1}{2} \int_0^\infty (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{1}{2} \frac{T}{\theta_r} = \frac{I}{\beta \hbar^2}.$$
- $$U^r = -\frac{3}{4} N \frac{\partial}{\partial \beta} \ln Z_{1o}^r - \frac{1}{4} N \frac{\partial}{\partial \beta} \ln Z_{1p}^r \simeq -N \frac{\partial}{\partial \beta} \ln Z_{1o}^r$$

$$= -N \frac{\partial}{\partial \beta} \ln \frac{I}{\beta \hbar^2} = \frac{N}{\beta}$$

# Internal energy and heat capacity of the ideal gas

- $$Z_{1o}^r \simeq Z_{1p}^r \simeq \frac{1}{2} \sum_{l=0,1,\dots} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$$

$$\simeq \frac{1}{2} \int_0^\infty (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{1}{2} \frac{T}{\theta_r} = \frac{I}{\beta \hbar^2}.$$
- $$U^r = -\frac{3}{4} N \frac{\partial}{\partial \beta} \ln Z_{1o}^r - \frac{1}{4} N \frac{\partial}{\partial \beta} \ln Z_{1p}^r \simeq -N \frac{\partial}{\partial \beta} \ln Z_{1o}^r$$

$$= -N \frac{\partial}{\partial \beta} \ln \frac{I}{\beta \hbar^2} = \frac{N}{\beta} = NkT.$$

# Internal energy and heat capacity of the ideal gas

- $$Z_{1o}^r \simeq Z_{1p}^r \simeq \frac{1}{2} \sum_{l=0,1,\dots} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$$

$$\simeq \frac{1}{2} \int_0^\infty (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{1}{2} \frac{T}{\theta_r} = \frac{I}{\beta \hbar^2}.$$
- $$U^r = -\frac{3}{4} N \frac{\partial}{\partial \beta} \ln Z_{1o}^r - \frac{1}{4} N \frac{\partial}{\partial \beta} \ln Z_{1p}^r \simeq -N \frac{\partial}{\partial \beta} \ln Z_{1o}^r$$

$$= -N \frac{\partial}{\partial \beta} \ln \frac{I}{\beta \hbar^2} = \frac{N}{\beta} = NkT.$$
- $$C_V^r = Nk.$$

# Internal energy and heat capacity of the ideal gas

- $$Z_{1o}^r \simeq Z_{1p}^r \simeq \frac{1}{2} \sum_{l=0,1,\dots} (2l+1) e^{-\frac{l(l+1)\theta_r}{T}}$$

$$\simeq \frac{1}{2} \int_0^\infty (2l+1) e^{-x} \frac{dx}{(2l+1)(\theta_r/T)} = \frac{1}{2} \frac{T}{\theta_r} = \frac{I}{\beta \hbar^2}.$$
- $$U^r = -\frac{3}{4} N \frac{\partial}{\partial \beta} \ln Z_{1o}^r - \frac{1}{4} N \frac{\partial}{\partial \beta} \ln Z_{1p}^r \simeq -N \frac{\partial}{\partial \beta} \ln Z_{1o}^r$$

$$= -N \frac{\partial}{\partial \beta} \ln \frac{I}{\beta \hbar^2} = \frac{N}{\beta} = NkT.$$
- $$C_V^r = Nk.$$
- If  $\theta_r \ll T$  does not hold, the summation is necessary.

# General processes for **classical** statistics



# General processes for **classical** statistics

- Object as an example: diatomic molecules.

# General processes for **classical** statistics

- Object as an example: diatomic molecules.

- $\varepsilon =$

$$\frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2) + \frac{1}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2).$$

# General processes for **classical** statistics

- Object as an example: diatomic molecules.

- $\varepsilon =$

$$\frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2) + \frac{1}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2).$$

- $Z_1 = \int \dots \int e^{-\beta \varepsilon(q,p)} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r},$

# General processes for **classical** statistics

- Object as an example: diatomic molecules.

- $\varepsilon =$

$$\frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2) + \frac{1}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2).$$

- $Z_1 = \int \dots \int e^{-\beta \varepsilon(q,p)} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r},$   
 $Z_1^t = \int e^{-\frac{\beta}{2m}(p_x^2 + p_y^2 + p_z^2)} \frac{dx dy dz dp_x dp_y dp_z}{h_0^3},$

# General processes for **classical** statistics

- Object as an example: diatomic molecules.

- $\varepsilon =$

$$\frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2) + \frac{1}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2).$$

- $Z_1 = \int \dots \int e^{-\beta \varepsilon(q,p)} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r},$

$$Z_1^t = \int e^{-\frac{\beta}{2m}(p_x^2 + p_y^2 + p_z^2)} \frac{dx dy dz dp_x dp_y dp_z}{h_0^3},$$

$$Z_1^r = \int e^{-\frac{\beta}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2)} \frac{dp_\theta dp_\varphi d\theta d\varphi}{h_0^2},$$

# General processes for **classical** statistics

- Object as an example: diatomic molecules.

- $\varepsilon =$

$$\frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2) + \frac{1}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2).$$

- $Z_1 = \int \dots \int e^{-\beta \varepsilon(q,p)} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r},$

$$Z_1^t = \int e^{-\frac{\beta}{2m}(p_x^2 + p_y^2 + p_z^2)} \frac{dx dy dz dp_x dp_y dp_z}{h_0^3},$$

$$Z_1^r = \int e^{-\frac{\beta}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2)} \frac{dp_\theta dp_\varphi d\theta d\varphi}{h_0^2},$$

$$Z_1^v = \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0}.$$

# General processes for **classical** statistics

- Object as an example: diatomic molecules.

- $\varepsilon =$

$$\frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2) + \frac{1}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2).$$

- $Z_1 = \int \dots \int e^{-\beta \varepsilon(q,p)} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r},$

$$Z_1^t = \int e^{-\frac{\beta}{2m}(p_x^2 + p_y^2 + p_z^2)} \frac{dx dy dz dp_x dp_y dp_z}{h_0^3},$$

$$Z_1^r = \int e^{-\frac{\beta}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2)} \frac{dp_\theta dp_\varphi d\theta d\varphi}{h_0^2},$$

$$Z_1^v = \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0}.$$

- Integration:

# General processes for **classical** statistics

- Object as an example: diatomic molecules.

- $\varepsilon =$

$$\frac{1}{2m}(p_x^2 + p_y^2 + p_z^2) + \frac{1}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2) + \frac{1}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2).$$

- $Z_1 = \int \dots \int e^{-\beta \varepsilon(q,p)} \frac{dq_1 \dots dq_r dp_1 \dots dp_r}{h_0^r},$

$$Z_1^t = \int e^{-\frac{\beta}{2m}(p_x^2 + p_y^2 + p_z^2)} \frac{dx dy dz dp_x dp_y dp_z}{h_0^3},$$

$$Z_1^r = \int e^{-\frac{\beta}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2)} \frac{dp_\theta dp_\varphi d\theta d\varphi}{h_0^2},$$

$$Z_1^v = \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0}.$$

- Integration:

$$Z_1^t = V \left( \frac{2\pi m}{h_0^2 \beta} \right)^{3/2}.$$



# General processes for classical statistics

- $$Z_1^r = \int e^{-\frac{\beta}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2)} \frac{dp_\theta dp_\varphi d\theta d\varphi}{h_0^2}$$

# General processes for classical statistics

- $$Z_1^r = \int e^{-\frac{\beta}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2)} \frac{dp_\theta dp_\varphi d\theta d\varphi}{h_0^2}$$
$$= \frac{1}{h_0^2} \int_0^{2\pi} d\varphi \int_0^\pi d\theta \int_{-\infty}^\infty e^{-\frac{\beta}{2I} p_\theta^2} dp_\theta \int_{-\infty}^\infty e^{-\frac{\beta}{2I \sin^2 \theta} p_\varphi^2} dp_\varphi$$

# General processes for classical statistics

$$\begin{aligned}
 \bullet \quad Z_1^r &= \int e^{-\frac{\beta}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2)} \frac{dp_\theta dp_\varphi d\theta d\varphi}{h_0^2} \\
 &= \frac{1}{h_0^2} \int_0^{2\pi} d\varphi \int_0^\pi d\theta \int_{-\infty}^{\infty} e^{-\frac{\beta}{2I} p_\theta^2} dp_\theta \int_{-\infty}^{\infty} e^{-\frac{\beta}{2I \sin^2 \theta} p_\varphi^2} dp_\varphi \\
 &= \\
 &\frac{2\pi}{h_0^2} \int_0^\pi d\theta \sqrt{\frac{2I}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2I \sin^2 \theta}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2I \sin^2 \theta} p_\varphi^2} d\sqrt{\frac{\beta}{2I \sin^2 \theta} p_\varphi}
 \end{aligned}$$

# General processes for classical statistics

$$\begin{aligned}
 \bullet \quad Z_1^r &= \int e^{-\frac{\beta}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2)} \frac{dp_\theta dp_\varphi d\theta d\varphi}{h_0^2} \\
 &= \frac{1}{h_0^2} \int_0^{2\pi} d\varphi \int_0^\pi d\theta \int_{-\infty}^{\infty} e^{-\frac{\beta}{2I} p_\theta^2} dp_\theta \int_{-\infty}^{\infty} e^{-\frac{\beta}{2I \sin^2 \theta} p_\varphi^2} dp_\varphi \\
 &= \\
 &\frac{2\pi}{h_0^2} \int_0^\pi d\theta \sqrt{\frac{2I}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2I \sin^2 \theta}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2I \sin^2 \theta} p_\varphi^2} d\sqrt{\frac{\beta}{2I \sin^2 \theta} p_\varphi} \\
 &= \frac{2\pi}{h_0^2} \int_0^\pi d\theta \sqrt{\frac{2I}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2I \sin^2 \theta}{\beta}} \sqrt{\pi}
 \end{aligned}$$

# General processes for classical statistics

$$\begin{aligned}
 \bullet \quad Z_1^r &= \int e^{-\frac{\beta}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2)} \frac{dp_\theta dp_\varphi d\theta d\varphi}{h_0^2} \\
 &= \frac{1}{h_0^2} \int_0^{2\pi} d\varphi \int_0^\pi d\theta \int_{-\infty}^{\infty} e^{-\frac{\beta}{2I} p_\theta^2} dp_\theta \int_{-\infty}^{\infty} e^{-\frac{\beta}{2I \sin^2 \theta} p_\varphi^2} dp_\varphi \\
 &= \\
 &\frac{2\pi}{h_0^2} \int_0^\pi d\theta \sqrt{\frac{2I}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2I \sin^2 \theta}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2I \sin^2 \theta} p_\varphi^2} d\sqrt{\frac{\beta}{2I \sin^2 \theta} p_\varphi} \\
 &= \frac{2\pi}{h_0^2} \int_0^\pi d\theta \sqrt{\frac{2I}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2I \sin^2 \theta}{\beta}} \sqrt{\pi} \\
 &= \frac{2\pi^2}{h_0^2} \frac{2I}{\beta} \int_0^\pi \sin \theta d\theta
 \end{aligned}$$

# General processes for classical statistics

$$\begin{aligned}
 \bullet \quad Z_1^r &= \int e^{-\frac{\beta}{2I}(p_\theta^2 + \frac{1}{\sin^2 \theta} p_\varphi^2)} \frac{dp_\theta dp_\varphi d\theta d\varphi}{h_0^2} \\
 &= \frac{1}{h_0^2} \int_0^{2\pi} d\varphi \int_0^\pi d\theta \int_{-\infty}^{\infty} e^{-\frac{\beta}{2I} p_\theta^2} dp_\theta \int_{-\infty}^{\infty} e^{-\frac{\beta}{2I \sin^2 \theta} p_\varphi^2} dp_\varphi \\
 &= \\
 &\frac{2\pi}{h_0^2} \int_0^\pi d\theta \sqrt{\frac{2I}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2I \sin^2 \theta}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2I \sin^2 \theta} p_\varphi^2} d\sqrt{\frac{\beta}{2I \sin^2 \theta} p_\varphi} \\
 &= \frac{2\pi}{h_0^2} \int_0^\pi d\theta \sqrt{\frac{2I}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2I \sin^2 \theta}{\beta}} \sqrt{\pi} \\
 &= \frac{2\pi^2}{h_0^2} \frac{2I}{\beta} \int_0^\pi \sin \theta d\theta \\
 &= \frac{8\pi^2 I}{h_0^2 \beta}.
 \end{aligned}$$

# General processes for classical statistics

- $$Z_1^v = \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0}$$

# General processes for classical statistics

- $$Z_1^v = \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0}$$
$$= \frac{1}{h_0} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} dp_r \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} dr$$



# General processes for classical statistics

$$\begin{aligned}
 \bullet \quad Z_1^v &= \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0} \\
 &= \frac{1}{h_0} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} dp_r \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} dr \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} d\sqrt{\frac{\beta}{2m_\mu}} p_r \cdot \\
 &\quad \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} d\sqrt{\frac{m_\mu \omega^2 \beta}{2}} r
 \end{aligned}$$

# General processes for classical statistics

$$\begin{aligned}
 \bullet \quad Z_1^v &= \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0} \\
 &= \frac{1}{h_0} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} dp_r \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} dr \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} d\sqrt{\frac{\beta}{2m_\mu}} p_r \cdot \\
 &\quad \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} d\sqrt{\frac{m_\mu \omega^2 \beta}{2}} r \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \sqrt{\pi}
 \end{aligned}$$

# General processes for classical statistics

$$\begin{aligned}
 \bullet \quad Z_1^v &= \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0} \\
 &= \frac{1}{h_0} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} dp_r \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} dr \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} d\sqrt{\frac{\beta}{2m_\mu}} p_r \cdot \\
 &\quad \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} d\sqrt{\frac{m_\mu \omega^2 \beta}{2}} r \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \sqrt{\pi} \\
 &= \frac{2\pi}{h_0 \beta \omega}.
 \end{aligned}$$

# General processes for classical statistics

- $$\begin{aligned}
 Z_1^v &= \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0} \\
 &= \frac{1}{h_0} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} dp_r \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} dr \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} d\sqrt{\frac{\beta}{2m_\mu}} p_r \cdot \\
 &\quad \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} d\sqrt{\frac{m_\mu \omega^2 \beta}{2}} r \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \sqrt{\pi} \\
 &= \frac{2\pi}{h_0 \beta \omega}.
 \end{aligned}$$
- $$U^t = -N \frac{\partial}{\partial \beta} \ln Z_1^t$$

# General processes for classical statistics

- $$\begin{aligned}
 Z_1^v &= \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0} \\
 &= \frac{1}{h_0} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} dp_r \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} dr \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} d\sqrt{\frac{\beta}{2m_\mu}} p_r \cdot \\
 &\quad \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} d\sqrt{\frac{m_\mu \omega^2 \beta}{2}} r \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \sqrt{\pi} \\
 &= \frac{2\pi}{h_0 \beta \omega}.
 \end{aligned}$$
- $$U^t = -N \frac{\partial}{\partial \beta} \ln Z_1^t = -N \frac{\partial}{\partial \beta} \ln [V (\frac{2\pi m}{h_0^2 \beta})^{3/2}]$$

# General processes for classical statistics

- $$\begin{aligned}
 Z_1^v &= \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0} \\
 &= \frac{1}{h_0} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} dp_r \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} dr \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} d\sqrt{\frac{\beta}{2m_\mu}} p_r \cdot \\
 &\quad \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} d\sqrt{\frac{m_\mu \omega^2 \beta}{2}} r \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \sqrt{\pi} \\
 &= \frac{2\pi}{h_0 \beta \omega}.
 \end{aligned}$$
- $$U^t = -N \frac{\partial}{\partial \beta} \ln Z_1^t = -N \frac{\partial}{\partial \beta} \ln [V (\frac{2\pi m}{h_0^2 \beta})^{3/2}] = \frac{3}{2} N k T.$$

# General processes for classical statistics

- $$\begin{aligned}
 Z_1^v &= \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0} \\
 &= \frac{1}{h_0} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} dp_r \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} dr \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} d\sqrt{\frac{\beta}{2m_\mu}} p_r \cdot \\
 &\quad \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} d\sqrt{\frac{m_\mu \omega^2 \beta}{2}} r \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \sqrt{\pi} \\
 &= \frac{2\pi}{h_0 \beta \omega}.
 \end{aligned}$$
- $$\begin{aligned}
 U^t &= -N \frac{\partial}{\partial \beta} \ln Z_1^t = -N \frac{\partial}{\partial \beta} \ln [V (\frac{2\pi m}{h_0^2 \beta})^{3/2}] = \frac{3}{2} N k T. \\
 U^r &= -N \frac{\partial}{\partial \beta} \ln Z_1^r
 \end{aligned}$$

# General processes for classical statistics

- $$\begin{aligned}
 Z_1^v &= \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0} \\
 &= \frac{1}{h_0} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} dp_r \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} dr \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} d\sqrt{\frac{\beta}{2m_\mu}} p_r \cdot \\
 &\quad \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} d\sqrt{\frac{m_\mu \omega^2 \beta}{2}} r \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \sqrt{\pi} \\
 &= \frac{2\pi}{h_0 \beta \omega}.
 \end{aligned}$$
- $$\begin{aligned}
 U^t &= -N \frac{\partial}{\partial \beta} \ln Z_1^t = -N \frac{\partial}{\partial \beta} \ln [V (\frac{2\pi m}{h_0^2 \beta})^{3/2}] = \frac{3}{2} N k T. \\
 U^r &= -N \frac{\partial}{\partial \beta} \ln Z_1^r = -N \frac{\partial}{\partial \beta} \ln \frac{8\pi^2 I}{h_0^2 \beta}
 \end{aligned}$$



# General processes for classical statistics

- $$\begin{aligned}
 Z_1^v &= \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0} \\
 &= \frac{1}{h_0} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} dp_r \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} dr \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} d\sqrt{\frac{\beta}{2m_\mu}} p_r \cdot \\
 &\quad \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} d\sqrt{\frac{m_\mu \omega^2 \beta}{2}} r \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \sqrt{\pi} \\
 &= \frac{2\pi}{h_0 \beta \omega}.
 \end{aligned}$$
- $$\begin{aligned}
 U^t &= -N \frac{\partial}{\partial \beta} \ln Z_1^t = -N \frac{\partial}{\partial \beta} \ln [V (\frac{2\pi m}{h_0^2 \beta})^{3/2}] = \frac{3}{2} N k T. \\
 U^r &= -N \frac{\partial}{\partial \beta} \ln Z_1^r = -N \frac{\partial}{\partial \beta} \ln \frac{8\pi^2 I}{h_0^2 \beta} = N k T.
 \end{aligned}$$

# General processes for classical statistics

- $$\begin{aligned}
 Z_1^v &= \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0} \\
 &= \frac{1}{h_0} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} dp_r \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} dr \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} d\sqrt{\frac{\beta}{2m_\mu}} p_r \cdot \\
 &\quad \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} d\sqrt{\frac{m_\mu \omega^2 \beta}{2}} r \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \sqrt{\pi} \\
 &= \frac{2\pi}{h_0 \beta \omega}.
 \end{aligned}$$
- $$\begin{aligned}
 U^t &= -N \frac{\partial}{\partial \beta} \ln Z_1^t = -N \frac{\partial}{\partial \beta} \ln [V (\frac{2\pi m}{h_0^2 \beta})^{3/2}] = \frac{3}{2} N k T. \\
 U^r &= -N \frac{\partial}{\partial \beta} \ln Z_1^r = -N \frac{\partial}{\partial \beta} \ln \frac{8\pi^2 I}{h_0^2 \beta} = N k T. \\
 U^v &= -N \frac{\partial}{\partial \beta} \ln Z_1^v
 \end{aligned}$$

# General processes for classical statistics

- $$\begin{aligned}
 Z_1^v &= \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0} \\
 &= \frac{1}{h_0} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} dp_r \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} dr \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} d\sqrt{\frac{\beta}{2m_\mu}} p_r \cdot \\
 &\quad \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} d\sqrt{\frac{m_\mu \omega^2 \beta}{2}} r \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \sqrt{\pi} \\
 &= \frac{2\pi}{h_0 \beta \omega}.
 \end{aligned}$$
- $$\begin{aligned}
 U^t &= -N \frac{\partial}{\partial \beta} \ln Z_1^t = -N \frac{\partial}{\partial \beta} \ln [V (\frac{2\pi m}{h_0^2 \beta})^{3/2}] = \frac{3}{2} N k T. \\
 U^r &= -N \frac{\partial}{\partial \beta} \ln Z_1^r = -N \frac{\partial}{\partial \beta} \ln \frac{8\pi^2 I}{h_0^2 \beta} = N k T. \\
 U^v &= -N \frac{\partial}{\partial \beta} \ln Z_1^v = -N \frac{\partial}{\partial \beta} \ln \frac{2\pi}{h_0 \beta \omega}
 \end{aligned}$$

# General processes for classical statistics

- $$\begin{aligned}
 Z_1^v &= \int e^{-\frac{\beta}{2m_\mu}(p_r^2 + m_\mu^2 \omega^2 r^2)} \frac{dp_r dr}{h_0} \\
 &= \frac{1}{h_0} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} dp_r \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} dr \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} p_r^2} d\sqrt{\frac{\beta}{2m_\mu}} p_r \cdot \\
 &\quad \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \int_{-\infty}^{\infty} e^{-\frac{\beta}{2m_\mu} m_\mu^2 \omega^2 r^2} d\sqrt{\frac{m_\mu \omega^2 \beta}{2}} r \\
 &= \frac{1}{h_0} \sqrt{\frac{2m_\mu}{\beta}} \sqrt{\pi} \cdot \sqrt{\frac{2}{m_\mu \omega^2 \beta}} \sqrt{\pi} \\
 &= \frac{2\pi}{h_0 \beta \omega}.
 \end{aligned}$$
- $$\begin{aligned}
 U^t &= -N \frac{\partial}{\partial \beta} \ln Z_1^t = -N \frac{\partial}{\partial \beta} \ln [V (\frac{2\pi m}{h_0^2 \beta})^{3/2}] = \frac{3}{2} N k T. \\
 U^r &= -N \frac{\partial}{\partial \beta} \ln Z_1^r = -N \frac{\partial}{\partial \beta} \ln \frac{8\pi^2 I}{h_0^2 \beta} = N k T. \\
 U^v &= -N \frac{\partial}{\partial \beta} \ln Z_1^v = -N \frac{\partial}{\partial \beta} \ln \frac{2\pi}{h_0 \beta \omega} = N k T.
 \end{aligned}$$

# Table of contents

## 1 §7 Boltzmann Statistics

- 7.1 Thermal quantities in statistics
- 7.2 Equation of state of ideal gas
- 7.3 Maxwell speed distribution
- 7.4 Energy equipartition theorem
- 7.5 Internal energy and heat capacity of the ideal gas
- **7.6 Entropy of the ideal gas**
- 7.7 Einstein's theory on heat capacity of solid
- 7.8 Paramagnetic solid
- 7.9 Negative temperature

## 7.6 Entropy of the ideal gas

## 7.6 Entropy of the ideal gas

- Object: Monatomic molecules (translational motion).

## 7.6 Entropy of the ideal gas

- Object: Monatomic molecules (translational motion).
- (7.1.13):  $S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1)$ .



## 7.6 Entropy of the ideal gas

- Object: Monatomic molecules (translational motion).
- (7.1.13):  $S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1)$ .
- (7.5.21):  $Z_1^t = V(\frac{2\pi m}{h_0^2 \beta})^{3/2}$ .

## 7.6 Entropy of the ideal gas

- Object: Monatomic molecules (translational motion).
- (7.1.13):  $S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1)$ .
- (7.5.21):  $Z_1^t = V(\frac{2\pi m}{h_0^2 \beta})^{3/2}$ .
- $S = Nk\{\ln[V(\frac{2\pi m}{h_0^2 \beta})^{3/2}] - \beta \frac{\partial}{\partial \beta} \ln[V(\frac{2\pi m}{h_0^2 \beta})^{3/2}]\}$

## 7.6 Entropy of the ideal gas

- Object: Monatomic molecules (translational motion).
- (7.1.13):  $S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1)$ .
- (7.5.21):  $Z_1^t = V(\frac{2\pi m}{h_0^2 \beta})^{3/2}$ .
- $$S = Nk\{\ln[V(\frac{2\pi m}{h_0^2 \beta})^{3/2}] - \beta \frac{\partial}{\partial \beta} \ln[V(\frac{2\pi m}{h_0^2 \beta})^{3/2}]\}$$
$$= Nk\{\ln V + \frac{3}{2} \ln(\frac{2\pi m}{h_0^2 \beta}) - \beta \frac{\partial}{\partial \beta} [\ln V + \frac{3}{2} \ln(\frac{2\pi m}{h_0^2 \beta})]\}$$

## 7.6 Entropy of the ideal gas

- Object: Monatomic molecules (translational motion).
- (7.1.13):  $S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1)$ .
- (7.5.21):  $Z_1^t = V(\frac{2\pi m}{h_0^2 \beta})^{3/2}$ .
- $$\begin{aligned} S &= Nk \left\{ \ln \left[ V \left( \frac{2\pi m}{h_0^2 \beta} \right)^{3/2} \right] - \beta \frac{\partial}{\partial \beta} \ln \left[ V \left( \frac{2\pi m}{h_0^2 \beta} \right)^{3/2} \right] \right\} \\ &= Nk \left\{ \ln V + \frac{3}{2} \ln \left( \frac{2\pi m}{h_0^2 \beta} \right) - \beta \frac{\partial}{\partial \beta} \left[ \ln V + \frac{3}{2} \ln \left( \frac{2\pi m}{h_0^2 \beta} \right) \right] \right\} \\ &= Nk \left[ \ln V + \frac{3}{2} \ln \left( \frac{2\pi m}{h_0^2 \beta} \right) - 0 - \frac{3}{2} \beta \cdot \left( -\frac{1}{\beta} \right) \right] \end{aligned}$$

## 7.6 Entropy of the ideal gas

- Object: Monatomic molecules (translational motion).
- (7.1.13):  $S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1)$ .
- (7.5.21):  $Z_1^t = V(\frac{2\pi m}{h_0^2 \beta})^{3/2}$ .
- $$\begin{aligned} S &= Nk \left\{ \ln \left[ V \left( \frac{2\pi m}{h_0^2 \beta} \right)^{3/2} \right] - \beta \frac{\partial}{\partial \beta} \ln \left[ V \left( \frac{2\pi m}{h_0^2 \beta} \right)^{3/2} \right] \right\} \\ &= Nk \left\{ \ln V + \frac{3}{2} \ln \left( \frac{2\pi m}{h_0^2 \beta} \right) - \beta \frac{\partial}{\partial \beta} \left[ \ln V + \frac{3}{2} \ln \left( \frac{2\pi m}{h_0^2 \beta} \right) \right] \right\} \\ &= Nk \left[ \ln V + \frac{3}{2} \ln \left( \frac{2\pi m}{h_0^2 \beta} \right) - 0 - \frac{3}{2} \beta \cdot \left( -\frac{1}{\beta} \right) \right] \\ &= Nk \left[ \ln V + \frac{3}{2} \ln \left( \frac{2\pi m}{h_0^2 \beta} \right) + \frac{3}{2} \right] \end{aligned}$$

## 7.6 Entropy of the ideal gas

- Object: Monatomic molecules (translational motion).
- (7.1.13):  $S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1)$ .
- (7.5.21):  $Z_1^t = V(\frac{2\pi m}{h_0^2 \beta})^{3/2}$ .
- $$\begin{aligned} S &= Nk \left\{ \ln \left[ V \left( \frac{2\pi m}{h_0^2 \beta} \right)^{3/2} \right] - \beta \frac{\partial}{\partial \beta} \ln \left[ V \left( \frac{2\pi m}{h_0^2 \beta} \right)^{3/2} \right] \right\} \\ &= Nk \left\{ \ln V + \frac{3}{2} \ln \left( \frac{2\pi m}{h_0^2 \beta} \right) - \beta \frac{\partial}{\partial \beta} \left[ \ln V + \frac{3}{2} \ln \left( \frac{2\pi m}{h_0^2 \beta} \right) \right] \right\} \\ &= Nk \left[ \ln V + \frac{3}{2} \ln \left( \frac{2\pi m}{h_0^2 \beta} \right) - 0 - \frac{3}{2} \beta \cdot \left( -\frac{1}{\beta} \right) \right] \\ &= Nk \left[ \ln V + \frac{3}{2} \ln \left( \frac{2\pi m}{h_0^2 \beta} \right) + \frac{3}{2} \right] \\ &= \frac{3}{2} Nk \ln T + Nk \ln V + \frac{3}{2} Nk \left[ 1 + \ln \frac{2\pi m k}{h_0^2} \right]. \end{aligned}$$

## 7.6 Entropy of the ideal gas

- Object: Monatomic molecules (translational motion).
- (7.1.13):  $S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1)$ .
- (7.5.21):  $Z_1^t = V(\frac{2\pi m}{h_0^2 \beta})^{3/2}$ .
- $$\begin{aligned} S &= Nk \left\{ \ln \left[ V \left( \frac{2\pi m}{h_0^2 \beta} \right)^{3/2} \right] - \beta \frac{\partial}{\partial \beta} \ln \left[ V \left( \frac{2\pi m}{h_0^2 \beta} \right)^{3/2} \right] \right\} \\ &= Nk \left\{ \ln V + \frac{3}{2} \ln \left( \frac{2\pi m}{h_0^2 \beta} \right) - \beta \frac{\partial}{\partial \beta} \left[ \ln V + \frac{3}{2} \ln \left( \frac{2\pi m}{h_0^2 \beta} \right) \right] \right\} \\ &= Nk \left[ \ln V + \frac{3}{2} \ln \left( \frac{2\pi m}{h_0^2 \beta} \right) - 0 - \frac{3}{2} \beta \cdot \left( -\frac{1}{\beta} \right) \right] \\ &= Nk \left[ \ln V + \frac{3}{2} \ln \left( \frac{2\pi m}{h_0^2 \beta} \right) + \frac{3}{2} \right] \\ &= \frac{3}{2} Nk \ln T + Nk \ln V + \frac{3}{2} Nk \left[ 1 + \ln \frac{2\pi m k}{h_0^2} \right]. \end{aligned}$$
- Problem 1.  $S$  depends on  $h_0$ ;

## 7.6 Entropy of the ideal gas

- Object: Monatomic molecules (translational motion).
- (7.1.13):  $S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1)$ .
- (7.5.21):  $Z_1^t = V(\frac{2\pi m}{h_0^2 \beta})^{3/2}$ .
- $$\begin{aligned} S &= Nk \left\{ \ln \left[ V \left( \frac{2\pi m}{h_0^2 \beta} \right)^{3/2} \right] - \beta \frac{\partial}{\partial \beta} \ln \left[ V \left( \frac{2\pi m}{h_0^2 \beta} \right)^{3/2} \right] \right\} \\ &= Nk \left\{ \ln V + \frac{3}{2} \ln \left( \frac{2\pi m}{h_0^2 \beta} \right) - \beta \frac{\partial}{\partial \beta} \left[ \ln V + \frac{3}{2} \ln \left( \frac{2\pi m}{h_0^2 \beta} \right) \right] \right\} \\ &= Nk \left[ \ln V + \frac{3}{2} \ln \left( \frac{2\pi m}{h_0^2 \beta} \right) - 0 - \frac{3}{2} \beta \cdot \left( -\frac{1}{\beta} \right) \right] \\ &= Nk \left[ \ln V + \frac{3}{2} \ln \left( \frac{2\pi m}{h_0^2 \beta} \right) + \frac{3}{2} \right] \\ &= \frac{3}{2} Nk \ln T + Nk \ln V + \frac{3}{2} Nk \left[ 1 + \ln \frac{2\pi m k}{h_0^2} \right]. \end{aligned}$$
- Problem 1.  $S$  depends on  $h_0$ ;
- Problem 2.  $S$  is not extensive.



# Entropy of the ideal gas

- Using (7.1.13') instead:

$$S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) - k \ln N!$$

# Entropy of the ideal gas

- Using (7.1.13') instead:

$$S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) - k \ln N!$$

Reason: Monatomic molecules are identical particles.

# Entropy of the ideal gas

- Using (7.1.13') instead:

$$S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) - k \ln N!$$

Reason: Monatomic molecules are identical particles.

$$S = k \ln \Omega_{\text{M.B.}} \rightarrow S = k \ln \frac{\Omega_{\text{M.B.}}}{N!} = k \ln \Omega_{\text{M.B.}} - k \ln N!.$$

# Entropy of the ideal gas

- Using (7.1.13') instead:

$$S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) - k \ln N!$$

Reason: Monatomic molecules are identical particles.

$$S = k \ln \Omega_{\text{M.B.}} \rightarrow S = k \ln \frac{\Omega_{\text{M.B.}}}{N!} = k \ln \Omega_{\text{M.B.}} - k \ln N!.$$

- $$S = \frac{3}{2}Nk \ln T + Nk \ln V + \frac{3}{2}Nk \left[ 1 + \ln \frac{2\pi mk}{h^2} \right] - k \ln N!$$

# Entropy of the ideal gas

- Using (7.1.13') instead:

$$S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) - k \ln N!$$

Reason: Monatomic molecules are identical particles.

$$S = k \ln \Omega_{\text{M.B.}} \rightarrow S = k \ln \frac{\Omega_{\text{M.B.}}}{N!} = k \ln \Omega_{\text{M.B.}} - k \ln N!.$$

$$\begin{aligned} \bullet S &= \frac{3}{2}Nk \ln T + Nk \ln V + \frac{3}{2}Nk \left[ 1 + \ln \frac{2\pi mk}{h^2} \right] - k \ln N! \\ &= \frac{3}{2}Nk \ln T + Nk \ln V + \frac{3}{2}Nk \left[ 1 + \ln \frac{2\pi mk}{h^2} \right] - kN(\ln N - 1) \end{aligned}$$

# Entropy of the ideal gas

- Using (7.1.13') instead:

$$S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) - k \ln N!$$

Reason: Monatomic molecules are identical particles.

$$S = k \ln \Omega_{\text{M.B.}} \rightarrow S = k \ln \frac{\Omega_{\text{M.B.}}}{N!} = k \ln \Omega_{\text{M.B.}} - k \ln N!.$$

$$\begin{aligned} \bullet S &= \frac{3}{2}Nk \ln T + Nk \ln V + \frac{3}{2}Nk \left[ 1 + \ln \frac{2\pi mk}{h^2} \right] - k \ln N! \\ &= \frac{3}{2}Nk \ln T + Nk \ln V + \frac{3}{2}Nk \left[ 1 + \ln \frac{2\pi mk}{h^2} \right] - kN(\ln N - 1) \\ &= \frac{3}{2}Nk \ln T + Nk \ln \frac{V}{N} + \frac{3}{2}Nk \left[ \frac{5}{2} + \ln \frac{2\pi mk}{h^2} \right]. \end{aligned}$$

# Entropy of the ideal gas

- Using (7.1.13') instead:

$$S = Nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) - k \ln N!$$

Reason: Monatomic molecules are identical particles.

$$S = k \ln \Omega_{\text{M.B.}} \rightarrow S = k \ln \frac{\Omega_{\text{M.B.}}}{N!} = k \ln \Omega_{\text{M.B.}} - k \ln N!.$$

$$\begin{aligned} \bullet S &= \frac{3}{2}Nk \ln T + Nk \ln V + \frac{3}{2}Nk \left[ 1 + \ln \frac{2\pi mk}{h^2} \right] - k \ln N! \\ &= \frac{3}{2}Nk \ln T + Nk \ln V + \frac{3}{2}Nk \left[ 1 + \ln \frac{2\pi mk}{h^2} \right] - kN(\ln N - 1) \\ &= \frac{3}{2}Nk \ln T + Nk \ln \frac{V}{N} + \frac{3}{2}Nk \left[ \frac{5}{2} + \ln \frac{2\pi mk}{h^2} \right]. \end{aligned}$$

- Not depend on  $h_0$ , extensive.

# Entropy of the ideal gas

- Compare with the experiment.



# Entropy of the ideal gas

- Compare with the experiment.
- Object: gas and condensed phase in equilibrium.

# Entropy of the ideal gas

- Compare with the experiment.
- Object: gas and condensed phase in equilibrium.
- For gas,  $pV = NkT$ ,

# Entropy of the ideal gas

- Compare with the experiment.
- Object: gas and condensed phase in equilibrium.
- For gas,  $pV = NkT$ ,  
$$\frac{S_{\text{gas}}}{Nk} = \frac{3}{2} \ln T + \ln \frac{V}{N} + \frac{3}{2} \left[ \frac{5}{2} + \ln \frac{2\pi mk}{h^2} \right]$$

# Entropy of the ideal gas

- Compare with the experiment.
- Object: gas and condensed phase in equilibrium.

- For gas,  $pV = NkT$ ,

$$\begin{aligned}\frac{S_{\text{gas}}}{Nk} &= \frac{3}{2} \ln T + \ln \frac{V}{N} + \frac{3}{2} \left[ \frac{5}{2} + \ln \frac{2\pi mk}{h^2} \right] \\ &= \frac{5}{2} \ln T + \ln \frac{V}{NkT} + \ln k + \frac{3}{2} \left[ \frac{5}{2} + \ln \frac{2\pi mk}{h^2} \right]\end{aligned}$$

# Entropy of the ideal gas

- Compare with the experiment.
- Object: gas and condensed phase in equilibrium.
- For gas,  $pV = NkT$ ,

$$\begin{aligned}\frac{S_{\text{gas}}}{Nk} &= \frac{3}{2} \ln T + \ln \frac{V}{N} + \frac{3}{2} \left[ \frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] \\ &= \frac{5}{2} \ln T + \ln \frac{V}{NkT} + \ln k + \frac{3}{2} \left[ \frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] \\ &= \frac{5}{2} \ln T + \ln \frac{1}{p} + \ln k + \frac{3}{2} \left[ \frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right]\end{aligned}$$

# Entropy of the ideal gas

- Compare with the experiment.
- Object: gas and condensed phase in equilibrium.

- For gas,  $pV = NkT$ ,

$$\begin{aligned}\frac{S_{\text{gas}}}{Nk} &= \frac{3}{2} \ln T + \ln \frac{V}{N} + \frac{3}{2} \left[ \frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] \\ &= \frac{5}{2} \ln T + \ln \frac{V}{NkT} + \ln k + \frac{3}{2} \left[ \frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] \\ &= \frac{5}{2} \ln T + \ln \frac{1}{p} + \ln k + \frac{3}{2} \left[ \frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right]\end{aligned}$$

- i.e.,  $\ln p = \frac{5}{2} \ln T + \ln k + \frac{3}{2} \left[ \frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] - \frac{S_{\text{gas}}}{Nk}$

# Entropy of the ideal gas

- Compare with the experiment.
- Object: gas and condensed phase in equilibrium.

- For gas,  $pV = NkT$ ,

$$\begin{aligned}\frac{S_{\text{gas}}}{Nk} &= \frac{3}{2} \ln T + \ln \frac{V}{N} + \frac{3}{2} \left[ \frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] \\ &= \frac{5}{2} \ln T + \ln \frac{V}{NkT} + \ln k + \frac{3}{2} \left[ \frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] \\ &= \frac{5}{2} \ln T + \ln \frac{1}{p} + \ln k + \frac{3}{2} \left[ \frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right]\end{aligned}$$

- i.e.,  $\ln p = \frac{5}{2} \ln T + \ln k + \frac{3}{2} \left[ \frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] - \frac{S_{\text{gas}}}{Nk}$

- $dS = \frac{\delta Q}{T}$ ,  $\rightarrow$  Latent  $L = T(S_{\text{gas}} - S_{\text{con}})$ .

# Entropy of the ideal gas

- Compare with the experiment.
- Object: gas and condensed phase in equilibrium.

- For gas,  $pV = NkT$ ,

$$\begin{aligned}\frac{S_{\text{gas}}}{Nk} &= \frac{3}{2} \ln T + \ln \frac{V}{N} + \frac{3}{2} \left[ \frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] \\ &= \frac{5}{2} \ln T + \ln \frac{V}{NkT} + \ln k + \frac{3}{2} \left[ \frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] \\ &= \frac{5}{2} \ln T + \ln \frac{1}{p} + \ln k + \frac{3}{2} \left[ \frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right]\end{aligned}$$

- i.e.,  $\ln p = \frac{5}{2} \ln T + \ln k + \frac{3}{2} \left[ \frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] - \frac{S_{\text{gas}}}{Nk}$

- $dS = \frac{\delta Q}{T}$ ,  $\rightarrow$  Latent  $L = T(S_{\text{gas}} - S_{\text{con}})$ .

- If  $T$  is low enough,  $S_{\text{gas}} \gg S_{\text{con}}$ ,



# Entropy of the ideal gas

- Compare with the experiment.
- Object: gas and condensed phase in equilibrium.

- For gas,  $pV = NkT$ ,

$$\begin{aligned}\frac{S_{\text{gas}}}{Nk} &= \frac{3}{2} \ln T + \ln \frac{V}{N} + \frac{3}{2} \left[ \frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] \\ &= \frac{5}{2} \ln T + \ln \frac{V}{NkT} + \ln k + \frac{3}{2} \left[ \frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] \\ &= \frac{5}{2} \ln T + \ln \frac{1}{p} + \ln k + \frac{3}{2} \left[ \frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right]\end{aligned}$$

- i.e.,  $\ln p = \frac{5}{2} \ln T + \ln k + \frac{3}{2} \left[ \frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] - \frac{S_{\text{gas}}}{Nk}$

- $dS = \frac{\delta Q}{T}$ ,  $\rightarrow$  Latent  $L = T(S_{\text{gas}} - S_{\text{con}})$ .

- If  $T$  is low enough,  $S_{\text{gas}} \gg S_{\text{con}}$ ,  $L = TS_{\text{gas}}$ .

# Entropy of the ideal gas

- Compare with the experiment.
- Object: gas and condensed phase in equilibrium.

- For gas,  $pV = NkT$ ,

$$\begin{aligned}\frac{S_{\text{gas}}}{Nk} &= \frac{3}{2} \ln T + \ln \frac{V}{N} + \frac{3}{2} \left[ \frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] \\ &= \frac{5}{2} \ln T + \ln \frac{V}{NkT} + \ln k + \frac{3}{2} \left[ \frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] \\ &= \frac{5}{2} \ln T + \ln \frac{1}{p} + \ln k + \frac{3}{2} \left[ \frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right]\end{aligned}$$

- I.e.,  $\ln p = \frac{5}{2} \ln T + \ln k + \frac{3}{2} \left[ \frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] - \frac{S_{\text{gas}}}{Nk}$
- $dS = \frac{\delta Q}{T}$ ,  $\rightarrow$  Latent  $L = T(S_{\text{gas}} - S_{\text{con}})$ .
- If  $T$  is low enough,  $S_{\text{gas}} \gg S_{\text{con}}$ ,  $L = TS_{\text{gas}}$ . Then  $\ln p = \frac{5}{2} \ln T + \ln k + \frac{3}{2} \left[ \frac{5}{3} + \ln \frac{2\pi mk}{h^2} \right] - \frac{L}{RT}$ , comparable.

# Chemical potential of the monatomic ideal gas

# Chemical potential of the monatomic ideal gas

- $dF = -SdT - pdV + \mu dN, \quad (3.2.9)$

# Chemical potential of the monatomic ideal gas

- $dF = -SdT - pdV + \mu dN, \quad (3.2.9)$   
 $\mu = \left(\frac{\partial F}{\partial N}\right)_{T,V}.$

# Chemical potential of the monatomic ideal gas

- $dF = -SdT - pdV + \mu dN$ , (3.2.9)  
 $\mu = \left(\frac{\partial F}{\partial N}\right)_{T,V}.$
- (7.1.16'),  
 $F = -NkT \ln Z_1 + kT \ln N!$

# Chemical potential of the monatomic ideal gas

- $dF = -SdT - pdV + \mu dN, \quad (3.2.9)$

$$\mu = \left(\frac{\partial F}{\partial N}\right)_{T,V}.$$

- (7.1.16'),

$$F = -NkT \ln Z_1 + kT \ln N!$$

$$= -NkT \ln Z_1 + kTN(\ln N - 1)$$

# Chemical potential of the monatomic ideal gas

- $dF = -SdT - pdV + \mu dN, \quad (3.2.9)$

$$\mu = \left(\frac{\partial F}{\partial N}\right)_{T,V}.$$

- (7.1.16'),

$$F = -NkT \ln Z_1 + kT \ln N!$$

$$= -NkT \ln Z_1 + kTN(\ln N - 1)$$

$$= NkT \ln \frac{N}{Z_1} - NkT.$$



# Chemical potential of the monatomic ideal gas

- $dF = -SdT - pdV + \mu dN, \quad (3.2.9)$

$$\mu = \left(\frac{\partial F}{\partial N}\right)_{T,V}.$$

- (7.1.16'),

$$F = -NkT \ln Z_1 + kT \ln N!$$

$$= -NkT \ln Z_1 + kTN(\ln N - 1)$$

$$= NkT \ln \frac{N}{Z_1} - NkT.$$

$$(Z_1^t = V(\frac{2\pi m}{h^2\beta})^{3/2} \quad (7.2.4))$$

# Chemical potential of the monatomic ideal gas

- $dF = -SdT - pdV + \mu dN, \quad (3.2.9)$

$$\mu = \left(\frac{\partial F}{\partial N}\right)_{T,V}.$$

- (7.1.16'),

$$F = -NkT \ln Z_1 + kT \ln N!$$

$$= -NkT \ln Z_1 + kTN(\ln N - 1)$$

$$= NkT \ln \frac{N}{Z_1} - NkT.$$

$$(Z_1^t = V(\frac{2\pi m}{h^2\beta})^{3/2} \quad (7.2.4))$$

- $\mu = \left(\frac{\partial F}{\partial N}\right)_{T,V}$

# Chemical potential of the monatomic ideal gas

- $dF = -SdT - pdV + \mu dN, \quad (3.2.9)$

$$\mu = \left(\frac{\partial F}{\partial N}\right)_{T,V}.$$

- (7.1.16'),

$$F = -NkT \ln Z_1 + kT \ln N!$$

$$= -NkT \ln Z_1 + kTN(\ln N - 1)$$

$$= NkT \ln \frac{N}{Z_1} - NkT.$$

$$(Z_1^t = V(\frac{2\pi m}{h^2\beta})^{3/2} \quad (7.2.4))$$

- $\mu = \left(\frac{\partial F}{\partial N}\right)_{T,V} = kT \ln \frac{N}{Z_1} + NkT \frac{Z_1}{N} \frac{1}{Z_1} - kT$

# Chemical potential of the monatomic ideal gas

- $dF = -SdT - pdV + \mu dN, \quad (3.2.9)$

$$\mu = \left(\frac{\partial F}{\partial N}\right)_{T,V}.$$

- (7.1.16'),

$$F = -NkT \ln Z_1 + kT \ln N!$$

$$= -NkT \ln Z_1 + kTN(\ln N - 1)$$

$$= NkT \ln \frac{N}{Z_1} - NkT.$$

$$(Z_1^t = V(\frac{2\pi m}{h^2\beta})^{3/2} \quad (7.2.4))$$

- $\mu = \left(\frac{\partial F}{\partial N}\right)_{T,V} = kT \ln \frac{N}{Z_1} + NkT \frac{Z_1}{N} \frac{1}{Z_1} - kT$   
 $= kT \ln \frac{N}{Z_1}$

# Chemical potential of the monatomic ideal gas

- $dF = -SdT - pdV + \mu dN, \quad (3.2.9)$

$$\mu = \left(\frac{\partial F}{\partial N}\right)_{T,V}.$$

- (7.1.16'),

$$F = -NkT \ln Z_1 + kT \ln N!$$

$$= -NkT \ln Z_1 + kTN(\ln N - 1)$$

$$= NkT \ln \frac{N}{Z_1} - NkT.$$

$$(Z_1^t = V(\frac{2\pi m}{h^2\beta})^{3/2} \quad (7.2.4))$$

- $\mu = \left(\frac{\partial F}{\partial N}\right)_{T,V} = kT \ln \frac{N}{Z_1} + NkT \frac{Z_1}{N} \frac{1}{Z_1} - kT$

$$= kT \ln \frac{N}{Z_1}$$

$$= kT \ln \left[ \frac{N}{V} \left( \frac{h^2}{2\pi m kT} \right)^{3/2} \right].$$

# Table of contents

## 1 §7 Boltzmann Statistics

- 7.1 Thermal quantities in statistics
- 7.2 Equation of state of ideal gas
- 7.3 Maxwell speed distribution
- 7.4 Energy equipartition theorem
- 7.5 Internal energy and heat capacity of the ideal gas
- 7.6 Entropy of the ideal gas
- 7.7 Einstein's theory on heat capacity of solid
- 7.8 Paramagnetic solid
- 7.9 Negative temperature

## 7.7 Einstein's theory on heat capacity of solid

## 7.7 Einstein's theory on heat capacity of solid

- Energy equipartition theorem:  $U = 3NkT$ ,  
 $C_V = 3Nk$ .



## 7.7 Einstein's theory on heat capacity of solid

- Energy equipartition theorem:  $U = 3NkT$ ,  
 $C_V = 3Nk$ .  
Only suitable for high temperature.

## 7.7 Einstein's theory on heat capacity of solid

- Energy equipartition theorem:  $U = 3NkT$ ,  
 $C_V = 3Nk$ .  
Only suitable for high temperature.
- General: taken the thermal motion as  $3N$  harmonic oscillators.

## 7.7 Einstein's theory on heat capacity of solid

- Energy equipartition theorem:  $U = 3NkT$ ,  
 $C_V = 3Nk$ .  
Only suitable for high temperature.
- General: taken the thermal motion as  $3N$  harmonic oscillators.
- Energy level:  $\varepsilon_n = (n + \frac{1}{2})\hbar\omega$ ,  $n = 0, 1, 2, \dots$

## 7.7 Einstein's theory on heat capacity of solid

- Energy equipartition theorem:  $U = 3NkT$ ,  
 $C_V = 3Nk$ .  
Only suitable for high temperature.
- General: taken the thermal motion as  $3N$  harmonic oscillators.
- Energy level:  $\varepsilon_n = (n + \frac{1}{2})\hbar\omega$ ,  $n = 0, 1, 2\ldots$
- Local, distinguishable, semi-independent  $\rightarrow$  Boltzmann distribution:

## 7.7 Einstein's theory on heat capacity of solid

- Energy equipartition theorem:  $U = 3NkT$ ,  
 $C_V = 3Nk$ .  
Only suitable for high temperature.
- General: taken the thermal motion as  $3N$  harmonic oscillators.
- Energy level:  $\varepsilon_n = (n + \frac{1}{2})\hbar\omega$ ,  $n = 0, 1, 2, \dots$
- Local, distinguishable, semi-independent  $\rightarrow$  Boltzmann distribution:

$$Z_1 = \sum_{n=0}^{\infty} \omega_l e^{-\beta\hbar\omega(n+\frac{1}{2})}$$

## 7.7 Einstein's theory on heat capacity of solid

- Energy equipartition theorem:  $U = 3NkT$ ,  
 $C_V = 3Nk$ .  
Only suitable for high temperature.
- General: taken the thermal motion as  $3N$  harmonic oscillators.
- Energy level:  $\varepsilon_n = (n + \frac{1}{2})\hbar\omega$ ,  $n = 0, 1, 2, \dots$
- Local, distinguishable, semi-independent  $\rightarrow$  Boltzmann distribution:

$$Z_1 = \sum_{n=0}^{\infty} \omega_l e^{-\beta\hbar\omega(n+\frac{1}{2})} = \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}.$$

# Einstein's theory on heat capacity of solid

- Internal energy:  $U = -3N \frac{\partial}{\partial \beta} \ln Z_1$

# Einstein's theory on heat capacity of solid

- Internal energy: 
$$U = -3N \frac{\partial}{\partial \beta} \ln Z_1$$
$$= -3N \frac{\partial}{\partial \beta} \ln \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}$$



# Einstein's theory on heat capacity of solid

- Internal energy: 
$$\begin{aligned} U &= -3N \frac{\partial}{\partial \beta} \ln Z_1 \\ &= -3N \frac{\partial}{\partial \beta} \ln \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1 - e^{-\beta\hbar\omega}} \\ &= -3N \frac{\partial}{\partial \beta} [\ln e^{-\frac{1}{2}\beta\hbar\omega} - \ln(1 - e^{-\beta\hbar\omega})] \end{aligned}$$

# Einstein's theory on heat capacity of solid

- Internal energy: 
$$\begin{aligned} U &= -3N \frac{\partial}{\partial \beta} \ln Z_1 \\ &= -3N \frac{\partial}{\partial \beta} \ln \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \\ &= -3N \frac{\partial}{\partial \beta} [\ln e^{-\frac{1}{2}\beta\hbar\omega} - \ln(1 - e^{-\beta\hbar\omega})] \\ &= 3N \left( \frac{1}{2}\hbar\omega + \frac{e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \hbar\omega \right) \end{aligned}$$

# Einstein's theory on heat capacity of solid

- Internal energy: 
$$\begin{aligned} U &= -3N \frac{\partial}{\partial \beta} \ln Z_1 \\ &= -3N \frac{\partial}{\partial \beta} \ln \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \\ &= -3N \frac{\partial}{\partial \beta} [\ln e^{-\frac{1}{2}\beta\hbar\omega} - \ln(1 - e^{-\beta\hbar\omega})] \\ &= 3N \left( \frac{1}{2}\hbar\omega + \frac{e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \hbar\omega \right) = \frac{3}{2}N\hbar\omega + \frac{3N\hbar\omega}{e^{\beta\hbar\omega} - 1}. \end{aligned}$$

# Einstein's theory on heat capacity of solid

- Internal energy:  $U = -3N \frac{\partial}{\partial \beta} \ln Z_1$ 
$$= -3N \frac{\partial}{\partial \beta} \ln \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1 - e^{-\beta\hbar\omega}}$$
$$= -3N \frac{\partial}{\partial \beta} [\ln e^{-\frac{1}{2}\beta\hbar\omega} - \ln(1 - e^{-\beta\hbar\omega})]$$
$$= 3N \left( \frac{1}{2}\hbar\omega + \frac{e^{-\beta\hbar\omega}}{1 - e^{-\beta\hbar\omega}} \hbar\omega \right) = \frac{3}{2}N\hbar\omega + \frac{3N\hbar\omega}{e^{\beta\hbar\omega} - 1}.$$
- Introduce typical temperature  $\theta_E$ :  $k\theta_E = \hbar\omega$ .

# Einstein's theory on heat capacity of solid

- Internal energy:  $U = -3N \frac{\partial}{\partial \beta} \ln Z_1$ 
$$\begin{aligned} &= -3N \frac{\partial}{\partial \beta} \ln \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \\ &= -3N \frac{\partial}{\partial \beta} [\ln e^{-\frac{1}{2}\beta\hbar\omega} - \ln(1 - e^{-\beta\hbar\omega})] \\ &= 3N \left( \frac{1}{2}\hbar\omega + \frac{e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \hbar\omega \right) = \frac{3}{2}N\hbar\omega + \frac{3N\hbar\omega}{e^{\beta\hbar\omega}-1}. \end{aligned}$$
- Introduce typical temperature  $\theta_E$ :  $k\theta_E = \hbar\omega$ .
$$U = \frac{3}{2}Nk\theta_E + \frac{3Nk\theta_E}{e^{\frac{\theta_E}{T}}-1}.$$

# Einstein's theory on heat capacity of solid

- Internal energy:  $U = -3N \frac{\partial}{\partial \beta} \ln Z_1$ 
$$= -3N \frac{\partial}{\partial \beta} \ln \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1 - e^{-\beta\hbar\omega}}$$
$$= -3N \frac{\partial}{\partial \beta} [\ln e^{-\frac{1}{2}\beta\hbar\omega} - \ln(1 - e^{-\beta\hbar\omega})]$$
$$= 3N \left( \frac{1}{2}\hbar\omega + \frac{e^{-\beta\hbar\omega}}{1 - e^{-\beta\hbar\omega}} \hbar\omega \right) = \frac{3}{2}N\hbar\omega + \frac{3N\hbar\omega}{e^{\beta\hbar\omega} - 1}.$$
- Introduce typical temperature  $\theta_E$ :  $k\theta_E = \hbar\omega$ .
$$U = \frac{3}{2}Nk\theta_E + \frac{3Nk\theta_E}{e^{\frac{\theta_E}{T}} - 1}.$$
$$C_V = \left( \frac{\partial U}{\partial T} \right)_V$$

# Einstein's theory on heat capacity of solid

- Internal energy:  $U = -3N \frac{\partial}{\partial \beta} \ln Z_1$ 

$$= -3N \frac{\partial}{\partial \beta} \ln \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}$$

$$= -3N \frac{\partial}{\partial \beta} [\ln e^{-\frac{1}{2}\beta\hbar\omega} - \ln(1 - e^{-\beta\hbar\omega})]$$

$$= 3N \left( \frac{1}{2}\hbar\omega + \frac{e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \hbar\omega \right) = \frac{3}{2}N\hbar\omega + \frac{3N\hbar\omega}{e^{\beta\hbar\omega}-1}.$$
- Introduce typical temperature  $\theta_E$ :  $k\theta_E = \hbar\omega$ .
 
$$U = \frac{3}{2}Nk\theta_E + \frac{3Nk\theta_E}{e^{\frac{\theta_E}{T}}-1}.$$

$$C_V = \left(\frac{\partial U}{\partial T}\right)_V = 0 - \frac{3Nk\theta_E}{(e^{\frac{\theta_E}{T}}-1)^2} \cdot e^{\frac{\theta_E}{T}} \cdot \left(-\frac{\theta_E}{T^2}\right)$$

# Einstein's theory on heat capacity of solid

- Internal energy:  $U = -3N \frac{\partial}{\partial \beta} \ln Z_1$ 

$$= -3N \frac{\partial}{\partial \beta} \ln \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}}$$

$$= -3N \frac{\partial}{\partial \beta} [\ln e^{-\frac{1}{2}\beta\hbar\omega} - \ln(1 - e^{-\beta\hbar\omega})]$$

$$= 3N \left( \frac{1}{2}\hbar\omega + \frac{e^{-\beta\hbar\omega}}{1-e^{-\beta\hbar\omega}} \hbar\omega \right) = \frac{3}{2}N\hbar\omega + \frac{3N\hbar\omega}{e^{\beta\hbar\omega}-1}.$$
- Introduce typical temperature  $\theta_E$ :  $k\theta_E = \hbar\omega$ .
 
$$U = \frac{3}{2}Nk\theta_E + \frac{3Nk\theta_E}{e^{\frac{\theta_E}{T}}-1}.$$

$$C_V = \left(\frac{\partial U}{\partial T}\right)_V = 0 - \frac{3Nk\theta_E}{(e^{\frac{\theta_E}{T}}-1)^2} \cdot e^{\frac{\theta_E}{T}} \cdot \left(-\frac{\theta_E}{T^2}\right)$$

$$= 3Nk\left(\frac{\theta_E}{T}\right)^2 \frac{e^{\frac{\theta_E}{T}}}{(e^{\frac{\theta_E}{T}}-1)^2}.$$



# Einstein's theory on heat capacity of solid

- $$C_V = 3Nk\left(\frac{\theta_E}{T}\right)^2 \frac{e^{\frac{\theta_E}{T}}}{(e^{\frac{\theta_E}{T}} - 1)^2},$$

# Einstein's theory on heat capacity of solid

- $$C_V = 3Nk \left( \frac{\theta_E}{T} \right)^2 \frac{e^{\frac{\theta_E}{T}}}{(e^{\frac{\theta_E}{T}} - 1)^2},$$
$$\theta_E \sim 1320\text{K for diamond.}$$

# Einstein's theory on heat capacity of solid

- $C_V = 3Nk \left(\frac{\theta_E}{T}\right)^2 \frac{e^{\frac{\theta_E}{T}}}{(e^{\frac{\theta_E}{T}} - 1)^2},$   
 $\theta_E \sim 1320\text{K}$  for diamond.
- High temperature:  $T \gg \theta_E$ ,  $e^{\frac{\theta_E}{T}} \simeq 1 + \frac{\theta_E}{T}.$

# Einstein's theory on heat capacity of solid

- $C_V = 3Nk \left(\frac{\theta_E}{T}\right)^2 \frac{e^{\frac{\theta_E}{T}}}{(e^{\frac{\theta_E}{T}} - 1)^2},$   
 $\theta_E \sim 1320\text{K}$  for diamond.
- High temperature:  $T \gg \theta_E$ ,  $e^{\frac{\theta_E}{T}} \simeq 1 + \frac{\theta_E}{T}.$   
 $C_V \simeq 3Nk$ , equipartition theorem.

# Einstein's theory on heat capacity of solid

- $C_V = 3Nk \left( \frac{\theta_E}{T} \right)^2 \frac{e^{\frac{\theta_E}{T}}}{(e^{\frac{\theta_E}{T}} - 1)^2},$   
 $\theta_E \sim 1320\text{K}$  for diamond.
- High temperature:  $T \gg \theta_E$ ,  $e^{\frac{\theta_E}{T}} \simeq 1 + \frac{\theta_E}{T}$ .  
 $C_V \simeq 3Nk$ , equipartition theorem.
- Low temperature:  $T \ll \theta_E$ ,  $e^{\frac{\theta_E}{T}} - 1 \simeq e^{\frac{\theta_E}{T}}$ .

# Einstein's theory on heat capacity of solid

- $C_V = 3Nk \left(\frac{\theta_E}{T}\right)^2 \frac{e^{\frac{\theta_E}{T}}}{(e^{\frac{\theta_E}{T}} - 1)^2},$   
 $\theta_E \sim 1320\text{K}$  for diamond.
- High temperature:  $T \gg \theta_E$ ,  $e^{\frac{\theta_E}{T}} \simeq 1 + \frac{\theta_E}{T}$ .  
 $C_V \simeq 3Nk$ , equipartition theorem.
- Low temperature:  $T \ll \theta_E$ ,  $e^{\frac{\theta_E}{T}} - 1 \simeq e^{\frac{\theta_E}{T}}$ .  
 $C_V \simeq 3Nk \left(\frac{\theta_E}{T}\right)^2 e^{-\frac{\theta_E}{T}} \rightarrow 0.$

# Table of contents

## 1 §7 Boltzmann Statistics

- 7.1 Thermal quantities in statistics
- 7.2 Equation of state of ideal gas
- 7.3 Maxwell speed distribution
- 7.4 Energy equipartition theorem
- 7.5 Internal energy and heat capacity of the ideal gas
- 7.6 Entropy of the ideal gas
- 7.7 Einstein's theory on heat capacity of solid
- **7.8 Paramagnetic solid**
- 7.9 Negative temperature

## 7.8 Paramagnetic solid



## 7.8 Paramagnetic solid

- Object: paramagnetic particles fixed on the crystal lattice.

## 7.8 Paramagnetic solid

- Object: paramagnetic particles fixed on the crystal lattice.
- Simplification: Magnetic quantum number  $\frac{1}{2}$ .

## 7.8 Paramagnetic solid

- Object: paramagnetic particles fixed on the crystal lattice.
- Simplification: Magnetic quantum number  $\frac{1}{2}$ .
- In the external magnetic field  $B$ , only two choices of quantum number  $\pm\frac{1}{2}$  (parallel or anti-parallel to  $B$ ).

## 7.8 Paramagnetic solid

- Object: paramagnetic particles fixed on the crystal lattice.
- Simplification: Magnetic quantum number  $\frac{1}{2}$ .
- In the external magnetic field  $B$ , only two choices of quantum number  $\pm\frac{1}{2}$  (parallel or anti-parallel to  $B$ ).
- Energy level:  $\pm\mu B$ . ( $\mu$  is the magnetic momentum.)

## 7.8 Paramagnetic solid

- Object: paramagnetic particles fixed on the crystal lattice.
- Simplification: Magnetic quantum number  $\frac{1}{2}$ .
- In the external magnetic field  $B$ , only two choices of quantum number  $\pm\frac{1}{2}$  (parallel or anti-parallel to  $B$ ).
- Energy level:  $\pm\mu B$ . ( $\mu$  is the magnetic momentum.)
- Partition function:  $Z_1 = \sum \omega_l e^{-\beta\epsilon_l}$

## 7.8 Paramagnetic solid

- Object: paramagnetic particles fixed on the crystal lattice.
- Simplification: Magnetic quantum number  $\frac{1}{2}$ .
- In the external magnetic field  $B$ , only two choices of quantum number  $\pm\frac{1}{2}$  (parallel or anti-parallel to  $B$ ).
- Energy level:  $\pm\mu B$ . ( $\mu$  is the magnetic momentum.)
- Partition function:  $Z_1 = \sum \omega_l e^{-\beta\epsilon_l} = e^{\beta\mu B} + e^{-\beta\mu B}$ .

## 7.8 Paramagnetic solid

- Object: paramagnetic particles fixed on the crystal lattice.
- Simplification: Magnetic quantum number  $\frac{1}{2}$ .
- In the external magnetic field  $B$ , only two choices of quantum number  $\pm\frac{1}{2}$  (parallel or anti-parallel to  $B$ ).
- Energy level:  $\pm\mu B$ . ( $\mu$  is the magnetic momentum.)
- Partition function:  $Z_1 = \sum \omega_l e^{-\beta \epsilon_l} = e^{\beta \mu B} + e^{-\beta \mu B}$ .
- Remind eq.(2.7.19),  $\text{d}W = -\mu_0 m \text{d}H$ .

## 7.8 Paramagnetic solid

- Object: paramagnetic particles fixed on the crystal lattice.
- Simplification: Magnetic quantum number  $\frac{1}{2}$ .
- In the external magnetic field  $B$ , only two choices of quantum number  $\pm\frac{1}{2}$  (parallel or anti-parallel to  $B$ ).
- Energy level:  $\pm\mu B$ . ( $\mu$  is the magnetic momentum.)
- Partition function:  $Z_1 = \sum \omega_l e^{-\beta\epsilon_l} = e^{\beta\mu B} + e^{-\beta\mu B}$ .
- Remind eq.(2.7.19),  $\text{d}W = -\mu_0 m \text{d}H$ . ( $-\mu_0 m$  generalized force,  $H$  generalized displacement.)



## 7.8 Paramagnetic solid

- Object: paramagnetic particles fixed on the crystal lattice.
- Simplification: Magnetic quantum number  $\frac{1}{2}$ .
- In the external magnetic field  $B$ , only two choices of quantum number  $\pm\frac{1}{2}$  (parallel or anti-parallel to  $B$ ).
- Energy level:  $\pm\mu B$ . ( $\mu$  is the magnetic momentum.)
- Partition function:  $Z_1 = \sum \omega_l e^{-\beta \epsilon_l} = e^{\beta \mu B} + e^{-\beta \mu B}$ .
- Remind eq.(2.7.19),  $\text{d}W = -\mu_0 m \text{d}H$ . ( $-\mu_0 m$  generalized force,  $H$  generalized displacement.)  
eq(7.1.6),  $Y = -\frac{N}{\beta} \frac{\partial}{\partial y} \ln Z_1$ .

## 7.8 Paramagnetic solid

- Object: paramagnetic particles fixed on the crystal lattice.
- Simplification: Magnetic quantum number  $\frac{1}{2}$ .
- In the external magnetic field  $B$ , only two choices of quantum number  $\pm\frac{1}{2}$  (parallel or anti-parallel to  $B$ ).
- Energy level:  $\pm\mu B$ . ( $\mu$  is the magnetic momentum.)
- Partition function:  $Z_1 = \sum \omega_l e^{-\beta \epsilon_l} = e^{\beta \mu B} + e^{-\beta \mu B}$ .
- Remind eq.(2.7.19),  $\text{d}W = -\mu_0 m \text{d}H$ . ( $-\mu_0 m$  generalized force,  $H$  generalized displacement.)  
eq(7.1.6),  $Y = -\frac{N}{\beta} \frac{\partial}{\partial y} \ln Z_1$ .  
 $\therefore -\mu_0 m = -\frac{N}{\beta} \frac{\partial}{\partial H} \ln Z_1$ .

# Paramagnetic solid

- $-\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1,$

# Paramagnetic solid

- $$-\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1,$$
$$\Rightarrow M = \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1$$

# Paramagnetic solid

- $-\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1,$   
 $\Rightarrow M = \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B})$

# Paramagnetic solid

- $$\begin{aligned} -\mu_0 \frac{m}{V} &= -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1, \\ \Rightarrow M &= \frac{n}{\beta} \frac{\partial}{\partial(\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B}) \\ &= \frac{n}{\beta} \frac{\beta\mu e^{\beta\mu B} - \beta\mu e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} \end{aligned}$$

# Paramagnetic solid

- $$\begin{aligned} -\mu_0 \frac{m}{V} &= -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1, \\ \Rightarrow M &= \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B}) \\ &= \frac{n}{\beta} \frac{\beta\mu e^{\beta\mu B} - \beta\mu e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = n\mu \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} \end{aligned}$$

# Paramagnetic solid

$$\begin{aligned}
 \bullet \quad & -\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1, \\
 \Rightarrow \quad & M = \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B}) \\
 = \quad & \frac{n}{\beta} \frac{\beta\mu e^{\beta\mu B} - \beta\mu e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = n\mu \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} = n\mu \frac{e^{\frac{2\mu B}{kT}} - 1}{e^{\frac{2\mu B}{kT}} + 1}.
 \end{aligned}$$



# Paramagnetic solid

- $$-\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1,$$

$$\Rightarrow M = \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B})$$

$$= \frac{n}{\beta} \frac{\beta\mu e^{\beta\mu B} - \beta\mu e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = n\mu \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} = n\mu \frac{e^{\frac{2\mu B}{kT}} - 1}{e^{\frac{2\mu B}{kT}} + 1}.$$
- $$\frac{\mu B}{kT} \ll 1,$$

# Paramagnetic solid

- $$-\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1,$$

$$\Rightarrow M = \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B})$$

$$= \frac{n}{\beta} \frac{\beta\mu e^{\beta\mu B} - \beta\mu e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = n\mu \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} = n\mu \frac{e^{\frac{2\mu B}{kT}} - 1}{e^{\frac{2\mu B}{kT}} + 1}.$$
- $$\frac{\mu B}{kT} \ll 1, \quad M \simeq n\mu \frac{\frac{2\mu B}{kT}}{2}$$

# Paramagnetic solid

- $$-\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1,$$

$$\Rightarrow M = \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B})$$

$$= \frac{n}{\beta} \frac{\beta\mu e^{\beta\mu B} - \beta\mu e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = n\mu \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} = n\mu \frac{e^{\frac{2\mu B}{kT}} - 1}{e^{\frac{2\mu B}{kT}} + 1}.$$
- $$\frac{\mu B}{kT} \ll 1, \quad M \simeq n\mu \frac{\frac{2\mu B}{kT}}{2} = \frac{n\mu^2}{kT} B, \text{ Curie's law.}$$

# Paramagnetic solid

- $$-\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1,$$

$$\Rightarrow M = \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B})$$

$$= \frac{n}{\beta} \frac{\beta\mu e^{\beta\mu B} - \beta\mu e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = n\mu \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} = n\mu \frac{e^{\frac{2\mu B}{kT}} - 1}{e^{\frac{2\mu B}{kT}} + 1}.$$
- $$\frac{\mu B}{kT} \ll 1, \quad M \simeq n\mu \frac{\frac{2\mu B}{kT}}{2} = \frac{n\mu^2}{kT} B, \text{ Curie's law.}$$
- $$\frac{\mu B}{kT} \gg 1,$$

# Paramagnetic solid

- $$-\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1,$$

$$\Rightarrow M = \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B})$$

$$= \frac{n}{\beta} \frac{\beta\mu e^{\beta\mu B} - \beta\mu e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = n\mu \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} = n\mu \frac{e^{\frac{2\mu B}{kT}} - 1}{e^{\frac{2\mu B}{kT}} + 1}.$$
- $$\frac{\mu B}{kT} \ll 1, M \simeq n\mu \frac{\frac{2\mu B}{kT}}{2} = \frac{n\mu^2}{kT} B, \text{ Curie's law.}$$
- $$\frac{\mu B}{kT} \gg 1, M \simeq n\mu, \text{ saturation.}$$

# Paramagnetic solid

- $$-\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1,$$

$$\Rightarrow M = \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B})$$

$$= \frac{n}{\beta} \frac{\beta\mu e^{\beta\mu B} - \beta\mu e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = n\mu \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} = n\mu \frac{e^{\frac{2\mu B}{kT}} - 1}{e^{\frac{2\mu B}{kT}} + 1}.$$
- $$\frac{\mu B}{kT} \ll 1, M \simeq n\mu \frac{\frac{2\mu B}{kT}}{2} = \frac{n\mu^2}{kT} B, \text{ Curie's law.}$$
- $$\frac{\mu B}{kT} \gg 1, M \simeq n\mu, \text{ saturation.}$$
- Internal energy density:  $u = -n \frac{\partial}{\partial \beta} \ln Z_1$

# Paramagnetic solid

- $$-\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1,$$

$$\Rightarrow M = \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B})$$

$$= \frac{n}{\beta} \frac{\beta\mu e^{\beta\mu B} - \beta\mu e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = n\mu \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} = n\mu \frac{e^{\frac{2\mu B}{kT}} - 1}{e^{\frac{2\mu B}{kT}} + 1}.$$
- $$\frac{\mu B}{kT} \ll 1, M \simeq n\mu \frac{\frac{2\mu B}{kT}}{2} = \frac{n\mu^2}{kT} B, \text{ Curie's law.}$$
- $$\frac{\mu B}{kT} \gg 1, M \simeq n\mu, \text{ saturation.}$$
- Internal energy density:  $u = -n \frac{\partial}{\partial \beta} \ln Z_1$

$$= -n \frac{\partial}{\partial \beta} \ln(e^{\beta\mu B} + e^{-\beta\mu B})$$

# Paramagnetic solid

- $$-\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1,$$

$$\Rightarrow M = \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B})$$

$$= \frac{n}{\beta} \frac{\beta\mu e^{\beta\mu B} - \beta\mu e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = n\mu \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} = n\mu \frac{e^{\frac{2\mu B}{kT}} - 1}{e^{\frac{2\mu B}{kT}} + 1}.$$
- $$\frac{\mu B}{kT} \ll 1, M \simeq n\mu \frac{\frac{2\mu B}{kT}}{2} = \frac{n\mu^2}{kT} B, \text{ Curie's law.}$$
- $$\frac{\mu B}{kT} \gg 1, M \simeq n\mu, \text{ saturation.}$$
- Internal energy density:  $u = -n \frac{\partial}{\partial \beta} \ln Z_1$

$$= -n \frac{\partial}{\partial \beta} \ln(e^{\beta\mu B} + e^{-\beta\mu B}) = -n \frac{\mu B e^{\beta\mu B} - \mu B e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}}$$



# Paramagnetic solid

- $$-\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1,$$

$$\Rightarrow M = \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B})$$

$$= \frac{n}{\beta} \frac{\beta\mu e^{\beta\mu B} - \beta\mu e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = n\mu \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} = n\mu \frac{e^{\frac{2\mu B}{kT}} - 1}{e^{\frac{2\mu B}{kT}} + 1}.$$
- $$\frac{\mu B}{kT} \ll 1, M \simeq n\mu \frac{\frac{2\mu B}{kT}}{2} = \frac{n\mu^2}{kT} B, \text{ Curie's law.}$$
- $$\frac{\mu B}{kT} \gg 1, M \simeq n\mu, \text{ saturation.}$$
- Internal energy density:  $u = -n \frac{\partial}{\partial \beta} \ln Z_1$

$$= -n \frac{\partial}{\partial \beta} \ln(e^{\beta\mu B} + e^{-\beta\mu B}) = -n \frac{\mu B e^{\beta\mu B} - \mu B e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}}$$

$$= -n\mu B \frac{e^{\beta\mu B} - e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}}$$

# Paramagnetic solid

- $$-\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1,$$

$$\Rightarrow M = \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B})$$

$$= \frac{n}{\beta} \frac{\beta\mu e^{\beta\mu B} - \beta\mu e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = n\mu \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} = n\mu \frac{e^{\frac{2\mu B}{kT}} - 1}{e^{\frac{2\mu B}{kT}} + 1}.$$
- $$\frac{\mu B}{kT} \ll 1, M \simeq n\mu \frac{\frac{2\mu B}{kT}}{2} = \frac{n\mu^2}{kT} B, \text{ Curie's law.}$$
- $$\frac{\mu B}{kT} \gg 1, M \simeq n\mu, \text{ saturation.}$$
- Internal energy density:  $u = -n \frac{\partial}{\partial \beta} \ln Z_1$

$$= -n \frac{\partial}{\partial \beta} \ln(e^{\beta\mu B} + e^{-\beta\mu B}) = -n \frac{\mu B e^{\beta\mu B} - \mu B e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}}$$

$$= -n\mu B \frac{e^{\beta\mu B} - e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = -MB,$$

# Paramagnetic solid

- $$-\mu_0 \frac{m}{V} = -\frac{N}{V\beta} \frac{\partial}{\partial H} \ln Z_1,$$

$$\Rightarrow M = \frac{n}{\beta} \frac{\partial}{\partial (\mu_0 H)} \ln Z_1 = \frac{n}{\beta} \frac{\partial}{\partial B} \ln(e^{\beta\mu B} + e^{-\beta\mu B})$$

$$= \frac{n}{\beta} \frac{\beta\mu e^{\beta\mu B} - \beta\mu e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = n\mu \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} = n\mu \frac{e^{\frac{2\mu B}{kT}} - 1}{e^{\frac{2\mu B}{kT}} + 1}.$$
- $$\frac{\mu B}{kT} \ll 1, M \simeq n\mu \frac{\frac{2\mu B}{kT}}{2} = \frac{n\mu^2}{kT} B, \text{ Curie's law.}$$
- $$\frac{\mu B}{kT} \gg 1, M \simeq n\mu, \text{ saturation.}$$
- Internal energy density:  $u = -n \frac{\partial}{\partial \beta} \ln Z_1$

$$= -n \frac{\partial}{\partial \beta} \ln(e^{\beta\mu B} + e^{-\beta\mu B}) = -n \frac{\mu B e^{\beta\mu B} - \mu B e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}}$$

$$= -n\mu B \frac{e^{\beta\mu B} - e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}} = -MB,$$

potential energy in the external field.

# Paramagnetic solid

- Entropy density eq.(7.1.13):

$$s = nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1)$$

# Paramagnetic solid

- Entropy density eq.(7.1.13):

$$\begin{aligned} s &= nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) \\ &= nk[\ln(e^{\beta\mu B} + e^{-\beta\mu B}) - \beta\mu B \frac{e^{\beta\mu B} - e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}}] \end{aligned}$$

# Paramagnetic solid

- Entropy density eq.(7.1.13):

$$\begin{aligned} s &= nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) \\ &= nk[\ln(e^{\beta\mu B} + e^{-\beta\mu B}) - \beta\mu B \frac{e^{\beta\mu B} - e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}}] \\ &= nk \left[ \ln(e^{2\beta\mu B} + 1) - \beta\mu B - \beta\mu B \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} \right] \end{aligned}$$

# Paramagnetic solid

- Entropy density eq.(7.1.13):

$$\begin{aligned} s &= nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) \\ &= nk[\ln(e^{\beta\mu B} + e^{-\beta\mu B}) - \beta\mu B \frac{e^{\beta\mu B} - e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}}] \\ &= nk \left[ \ln(e^{2\beta\mu B} + 1) - \beta\mu B - \beta\mu B \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} \right] \\ &= nk \left[ \ln(e^{2\beta\mu B} + 1) - \beta\mu B \frac{2e^{2\beta\mu B}}{e^{2\beta\mu B} + 1} \right] \end{aligned}$$

# Paramagnetic solid

- Entropy density eq.(7.1.13):

$$\begin{aligned} s &= nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) \\ &= nk[\ln(e^{\beta\mu B} + e^{-\beta\mu B}) - \beta\mu B \frac{e^{\beta\mu B} - e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}}] \\ &= nk \left[ \ln(e^{2\beta\mu B} + 1) - \beta\mu B - \beta\mu B \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} \right] \\ &= nk \left[ \ln(e^{2\beta\mu B} + 1) - \beta\mu B \frac{2e^{2\beta\mu B}}{e^{2\beta\mu B} + 1} \right] \\ &= nk \left[ \ln(e^{\frac{2\mu B}{kT}} + 1) - \frac{\mu B}{kT} \frac{2e^{\frac{2\mu B}{kT}}}{e^{\frac{2\mu B}{kT}} + 1} \right]. \end{aligned}$$



# Paramagnetic solid

- Entropy density eq.(7.1.13):

$$\begin{aligned}
 s &= nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1) \\
 &= nk[\ln(e^{\beta\mu B} + e^{-\beta\mu B}) - \beta\mu B \frac{e^{\beta\mu B} - e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}}] \\
 &= nk \left[ \ln(e^{2\beta\mu B} + 1) - \beta\mu B - \beta\mu B \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} \right] \\
 &= nk \left[ \ln(e^{2\beta\mu B} + 1) - \beta\mu B \frac{2e^{2\beta\mu B}}{e^{2\beta\mu B} + 1} \right] \\
 &= nk \left[ \ln(e^{\frac{2\mu B}{kT}} + 1) - \frac{\mu B}{kT} \frac{2e^{\frac{2\mu B}{kT}}}{e^{\frac{2\mu B}{kT}} + 1} \right].
 \end{aligned}$$

- $\frac{\mu B}{kT} \ll 1$ ,  $s = nk \ln 2 = k \ln 2^n$ ,

# Paramagnetic solid

- Entropy density eq.(7.1.13):

$$s = nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1)$$

$$= nk[\ln(e^{\beta\mu B} + e^{-\beta\mu B}) - \beta\mu B \frac{e^{\beta\mu B} - e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}}]$$

$$= nk \left[ \ln(e^{2\beta\mu B} + 1) - \beta\mu B - \beta\mu B \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} \right]$$

$$= nk \left[ \ln(e^{2\beta\mu B} + 1) - \beta\mu B \frac{2e^{2\beta\mu B}}{e^{2\beta\mu B} + 1} \right]$$

$$= nk \left[ \ln\left(e^{\frac{2\mu B}{kT}} + 1\right) - \frac{\mu B}{kT} \frac{2e^{\frac{2\mu B}{kT}}}{e^{\frac{2\mu B}{kT}} + 1} \right].$$

- $\frac{\mu B}{kT} \ll 1$ ,  $s = nk \ln 2 = k \ln 2^n$ , high temperature, two states are both possible, total number of states:  $2^n$ .

# Paramagnetic solid

- Entropy density eq.(7.1.13):

$$s = nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1)$$

$$= nk[\ln(e^{\beta\mu B} + e^{-\beta\mu B}) - \beta\mu B \frac{e^{\beta\mu B} - e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}}]$$

$$= nk \left[ \ln(e^{2\beta\mu B} + 1) - \beta\mu B - \beta\mu B \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} \right]$$

$$= nk \left[ \ln(e^{2\beta\mu B} + 1) - \beta\mu B \frac{2e^{2\beta\mu B}}{e^{2\beta\mu B} + 1} \right]$$

$$= nk \left[ \ln\left(e^{\frac{2\mu B}{kT}} + 1\right) - \frac{\mu B}{kT} \frac{2e^{\frac{2\mu B}{kT}}}{e^{\frac{2\mu B}{kT}} + 1} \right].$$

- $\frac{\mu B}{kT} \ll 1$ ,  $s = nk \ln 2 = k \ln 2^n$ , high temperature, two states are both possible, total number of states:  $2^n$ .
- $\frac{\mu B}{kT} \gg 1$ ,  $s \simeq 0$ ,

# Paramagnetic solid

- Entropy density eq.(7.1.13):

$$s = nk(\ln Z_1 - \beta \frac{\partial}{\partial \beta} \ln Z_1)$$

$$= nk[\ln(e^{\beta\mu B} + e^{-\beta\mu B}) - \beta\mu B \frac{e^{\beta\mu B} - e^{-\beta\mu B}}{e^{\beta\mu B} + e^{-\beta\mu B}}]$$

$$= nk \left[ \ln(e^{2\beta\mu B} + 1) - \beta\mu B - \beta\mu B \frac{e^{2\beta\mu B} - 1}{e^{2\beta\mu B} + 1} \right]$$

$$= nk \left[ \ln(e^{2\beta\mu B} + 1) - \beta\mu B \frac{2e^{2\beta\mu B}}{e^{2\beta\mu B} + 1} \right]$$

$$= nk \left[ \ln\left(e^{\frac{2\mu B}{kT}} + 1\right) - \frac{\mu B}{kT} \frac{2e^{\frac{2\mu B}{kT}}}{e^{\frac{2\mu B}{kT}} + 1} \right].$$

- $\frac{\mu B}{kT} \ll 1$ ,  $s = nk \ln 2 = k \ln 2^n$ , high temperature, two states are both possible, total number of states:  $2^n$ .
- $\frac{\mu B}{kT} \gg 1$ ,  $s \simeq 0$ , only one micro-state: all magnetic moments are aligned along the external field.

# Table of contents

## 1 §7 Boltzmann Statistics

- 7.1 Thermal quantities in statistics
- 7.2 Equation of state of ideal gas
- 7.3 Maxwell speed distribution
- 7.4 Energy equipartition theorem
- 7.5 Internal energy and heat capacity of the ideal gas
- 7.6 Entropy of the ideal gas
- 7.7 Einstein's theory on heat capacity of solid
- 7.8 Paramagnetic solid
- 7.9 Negative temperature

## 7.9 Negative temperature

## 7.9 Negative temperature

- Basic function of thermodynamics:  $dU = TdS - pdV$ .

## 7.9 Negative temperature

- Basic function of thermodynamics:  $dU = TdS - pdV$ .  
More general:  $dU = TdS + Ydy$ ,



## 7.9 Negative temperature

- Basic function of thermodynamics:  $dU = TdS - pdV$ .  
More general:  $dU = TdS + Ydy$ ,  
 $\Rightarrow \left(\frac{\partial S}{\partial U}\right)_y = \frac{1}{T}$ .

## 7.9 Negative temperature

- Basic function of thermodynamics:  $dU = TdS - pdV$ .

More general:  $dU = TdS + Ydy$ ,

$$\Rightarrow \left(\frac{\partial S}{\partial U}\right)_y = \frac{1}{T}.$$

If  $U$  increases but  $S$  decreases (micro-states),  $T$  is negative.

## 7.9 Negative temperature

- Basic function of thermodynamics:  $dU = TdS - pdV$ .  
More general:  $dU = TdS + Ydy$ ,  
 $\Rightarrow \left(\frac{\partial S}{\partial U}\right)_y = \frac{1}{T}$ .  
If  $U$  increases but  $S$  decreases (micro-states),  $T$  is negative.
- Example: isolated nuclear spin system.

## 7.9 Negative temperature

- Basic function of thermodynamics:  $dU = TdS - pdV$ .  
More general:  $dU = TdS + Ydy$ ,  
 $\Rightarrow \left(\frac{\partial S}{\partial U}\right)_y = \frac{1}{T}$ .  
If  $U$  increases but  $S$  decreases (micro-states),  $T$  is negative.
- Example: isolated nuclear spin system.
- In external magnetic field  $B$ , the energy  
 $\pm\varepsilon = \pm\mu B = \pm\frac{e\hbar}{2m}B$ .

## 7.9 Negative temperature

- Basic function of thermodynamics:  $dU = TdS - pdV$ .  
More general:  $dU = TdS + Ydy$ ,  
 $\Rightarrow \left(\frac{\partial S}{\partial U}\right)_y = \frac{1}{T}$ .  
If  $U$  increases but  $S$  decreases (micro-states),  $T$  is negative.
- Example: isolated nuclear spin system.
- In external magnetic field  $B$ , the energy  
 $\pm\varepsilon = \pm\mu B = \pm\frac{e\hbar}{2m}B$ .
- Number of spins:  $N$ , where  $N_+$  with  $\varepsilon$ ,  $N_-$  with  $-\varepsilon$ .

## 7.9 Negative temperature

- Basic function of thermodynamics:  $dU = TdS - pdV$ .  
More general:  $dU = TdS + Ydy$ ,  
 $\Rightarrow \left(\frac{\partial S}{\partial U}\right)_y = \frac{1}{T}$ .  
If  $U$  increases but  $S$  decreases (micro-states),  $T$  is negative.
- Example: isolated nuclear spin system.
- In external magnetic field  $B$ , the energy  
 $\pm\varepsilon = \pm\mu B = \pm\frac{e\hbar}{2m}B$ .
- Number of spins:  $N$ , where  $N_+$  with  $\varepsilon$ ,  $N_-$  with  $-\varepsilon$ .
- $N = N_+ + N_-$ ,  $E = N_+\varepsilon + N_-(-\varepsilon)$

## 7.9 Negative temperature

- Basic function of thermodynamics:  $dU = TdS - pdV$ .  
More general:  $dU = TdS + Ydy$ ,  
 $\Rightarrow (\frac{\partial S}{\partial U})_y = \frac{1}{T}$ .  
If  $U$  increases but  $S$  decreases (micro-states),  $T$  is negative.
- Example: isolated nuclear spin system.
- In external magnetic field  $B$ , the energy  
 $\pm\varepsilon = \pm\mu B = \pm\frac{e\hbar}{2m}B$ .
- Number of spins:  $N$ , where  $N_+$  with  $\varepsilon$ ,  $N_-$  with  $-\varepsilon$ .
- $N = N_+ + N_-$ ,  $E = N_+\varepsilon + N_-(-\varepsilon)$   
 $\Rightarrow N_+ = \frac{N}{2}(1 + \frac{E}{N\varepsilon})$ ,  $N_- = \frac{N}{2}(1 - \frac{E}{N\varepsilon})$ .

Negative temperature  $N_{\pm} = \frac{N}{2} \left( 1 \pm \frac{E}{N\epsilon} \right)$

- Entropy  $S = k \ln \Omega$



Negative temperature  $N_{\pm} = \frac{N}{2} \left( 1 \pm \frac{E}{N\varepsilon} \right)$

- Entropy  $S = k \ln \Omega = k \ln \frac{N!}{N_+! N_-!}$

Negative temperature  $N_{\pm} = \frac{N}{2} \left( 1 \pm \frac{E}{N\varepsilon} \right)$

- Entropy  $S = k \ln \Omega = k \ln \frac{N!}{N_+! N_-!}$   
 $\simeq k[N(\ln N - 1) - N_+(\ln N_+ - 1) - N_-(\ln N_- - 1)]$

# Negative temperature

$$N_{\pm} = \frac{N}{2} \left( 1 \pm \frac{E}{N\varepsilon} \right)$$

- Entropy  $S = k \ln \Omega = k \ln \frac{N!}{N_+! N_-!}$   
 $\simeq k[N(\ln N - 1) - N_+(\ln N_+ - 1) - N_-(\ln N_- - 1)]$   
 $= k(N \ln N - N_+ \ln N_+ - N_- \ln N_-)$

## Negative temperature

$$N_{\pm} = \frac{N}{2} \left( 1 \pm \frac{E}{N\varepsilon} \right)$$

- Entropy  $S = k \ln \Omega = k \ln \frac{N!}{N_+! N_-!}$ 

$$\begin{aligned} &\simeq k[N(\ln N - 1) - N_+(\ln N_+ - 1) - N_-(\ln N_- - 1)] \\ &= k(N \ln N - N_+ \ln N_+ - N_- \ln N_-) \\ &= k\left\{ N \ln N - \frac{N}{2} \left( 1 + \frac{E}{N\varepsilon} \right) \ln \left[ \frac{N}{2} \left( 1 + \frac{E}{N\varepsilon} \right) \right] \right. \\ &\quad \left. - \frac{N}{2} \left( 1 - \frac{E}{N\varepsilon} \right) \ln \left[ \frac{N}{2} \left( 1 - \frac{E}{N\varepsilon} \right) \right] \right\} \end{aligned}$$

## Negative temperature

$$N_{\pm} = \frac{N}{2} \left( 1 \pm \frac{E}{N\varepsilon} \right)$$

- Entropy  $S = k \ln \Omega = k \ln \frac{N!}{N_+! N_-!}$ 

$$\begin{aligned} &\simeq k[N(\ln N - 1) - N_+(\ln N_+ - 1) - N_-(\ln N_- - 1)] \\ &= k(N \ln N - N_+ \ln N_+ - N_- \ln N_-) \\ &= k\left\{ N \ln N - \frac{N}{2} \left( 1 + \frac{E}{N\varepsilon} \right) \ln \left[ \frac{N}{2} \left( 1 + \frac{E}{N\varepsilon} \right) \right] \right. \\ &\quad \left. - \frac{N}{2} \left( 1 - \frac{E}{N\varepsilon} \right) \ln \left[ \frac{N}{2} \left( 1 - \frac{E}{N\varepsilon} \right) \right] \right\} \\ &= k\left\{ N \ln N - \frac{N}{2} \left( 1 + \frac{E}{N\varepsilon} \right) \left[ \ln \frac{N}{2} + \ln \left( 1 + \frac{E}{N\varepsilon} \right) \right] \right. \\ &\quad \left. - \frac{N}{2} \left( 1 - \frac{E}{N\varepsilon} \right) \left[ \ln \frac{N}{2} + \ln \left( 1 - \frac{E}{N\varepsilon} \right) \right] \right\} \end{aligned}$$

## Negative temperature

$$N_{\pm} = \frac{N}{2} \left( 1 \pm \frac{E}{N\varepsilon} \right)$$

- Entropy  $S = k \ln \Omega = k \ln \frac{N!}{N_+! N_-!}$ 

$$\begin{aligned} &\simeq k[N(\ln N - 1) - N_+(\ln N_+ - 1) - N_-(\ln N_- - 1)] \\ &= k(N \ln N - N_+ \ln N_+ - N_- \ln N_-) \\ &= k\left\{ N \ln N - \frac{N}{2} \left( 1 + \frac{E}{N\varepsilon} \right) \ln \left[ \frac{N}{2} \left( 1 + \frac{E}{N\varepsilon} \right) \right] \right. \\ &\quad \left. - \frac{N}{2} \left( 1 - \frac{E}{N\varepsilon} \right) \ln \left[ \frac{N}{2} \left( 1 - \frac{E}{N\varepsilon} \right) \right] \right\} \\ &= k\left\{ N \ln N - \frac{N}{2} \left( 1 + \frac{E}{N\varepsilon} \right) \left[ \ln \frac{N}{2} + \ln \left( 1 + \frac{E}{N\varepsilon} \right) \right] \right. \\ &\quad \left. - \frac{N}{2} \left( 1 - \frac{E}{N\varepsilon} \right) \left[ \ln \frac{N}{2} + \ln \left( 1 - \frac{E}{N\varepsilon} \right) \right] \right\} \\ &= k\left\{ N \ln N - \frac{N}{2} \ln \frac{N}{2} \left[ \left( 1 + \frac{E}{N\varepsilon} \right) + \left( 1 - \frac{E}{N\varepsilon} \right) \right] \right. \\ &\quad \left. - \frac{N}{2} \left( 1 + \frac{E}{N\varepsilon} \right) \ln \left( 1 + \frac{E}{N\varepsilon} \right) - \frac{N}{2} \left( 1 - \frac{E}{N\varepsilon} \right) \ln \left( 1 - \frac{E}{N\varepsilon} \right) \right\} \end{aligned}$$

## Negative temperature

$$N_{\pm} = \frac{N}{2} \left( 1 \pm \frac{E}{N\varepsilon} \right)$$

- Entropy  $S = k \ln \Omega = k \ln \frac{N!}{N_+! N_-!}$ 

$$\begin{aligned} &\simeq k[N(\ln N - 1) - N_+(\ln N_+ - 1) - N_-(\ln N_- - 1)] \\ &= k(N \ln N - N_+ \ln N_+ - N_- \ln N_-) \\ &= k\left\{ N \ln N - \frac{N}{2} \left( 1 + \frac{E}{N\varepsilon} \right) \ln \left[ \frac{N}{2} \left( 1 + \frac{E}{N\varepsilon} \right) \right] \right. \\ &\quad \left. - \frac{N}{2} \left( 1 - \frac{E}{N\varepsilon} \right) \ln \left[ \frac{N}{2} \left( 1 - \frac{E}{N\varepsilon} \right) \right] \right\} \\ &= k\left\{ N \ln N - \frac{N}{2} \left( 1 + \frac{E}{N\varepsilon} \right) \left[ \ln \frac{N}{2} + \ln \left( 1 + \frac{E}{N\varepsilon} \right) \right] \right. \\ &\quad \left. - \frac{N}{2} \left( 1 - \frac{E}{N\varepsilon} \right) \left[ \ln \frac{N}{2} + \ln \left( 1 - \frac{E}{N\varepsilon} \right) \right] \right\} \\ &= k\left\{ N \ln N - \frac{N}{2} \ln \frac{N}{2} \left[ \left( 1 + \frac{E}{N\varepsilon} \right) + \left( 1 - \frac{E}{N\varepsilon} \right) \right] \right. \\ &\quad \left. - \frac{N}{2} \left( 1 + \frac{E}{N\varepsilon} \right) \ln \left( 1 + \frac{E}{N\varepsilon} \right) - \frac{N}{2} \left( 1 - \frac{E}{N\varepsilon} \right) \ln \left( 1 - \frac{E}{N\varepsilon} \right) \right\} \\ &= k\left\{ N \ln N - N \ln \frac{N}{2} \right. \\ &\quad \left. - \frac{N}{2} \left( 1 + \frac{E}{N\varepsilon} \right) \ln \left( 1 + \frac{E}{N\varepsilon} \right) - \frac{N}{2} \left( 1 - \frac{E}{N\varepsilon} \right) \ln \left( 1 - \frac{E}{N\varepsilon} \right) \right\} \end{aligned}$$

## Negative temperature

$$N_{\pm} = \frac{N}{2} \left(1 \pm \frac{E}{N\varepsilon}\right)$$

- Entropy  $S = k \ln \Omega = k \ln \frac{N!}{N_+! N_-!}$ 

$$\begin{aligned} &\simeq k[N(\ln N - 1) - N_+(\ln N_+ - 1) - N_-(\ln N_- - 1)] \\ &= k(N \ln N - N_+ \ln N_+ - N_- \ln N_-) \\ &= k\left\{N \ln N - \frac{N}{2}\left(1 + \frac{E}{N\varepsilon}\right) \ln\left[\frac{N}{2}\left(1 + \frac{E}{N\varepsilon}\right)\right] \right. \\ &\quad \left. - \frac{N}{2}\left(1 - \frac{E}{N\varepsilon}\right) \ln\left[\frac{N}{2}\left(1 - \frac{E}{N\varepsilon}\right)\right]\right\} \\ &= k\left\{N \ln N - \frac{N}{2}\left(1 + \frac{E}{N\varepsilon}\right) \left[\ln \frac{N}{2} + \ln\left(1 + \frac{E}{N\varepsilon}\right)\right] \right. \\ &\quad \left. - \frac{N}{2}\left(1 - \frac{E}{N\varepsilon}\right) \left[\ln \frac{N}{2} + \ln\left(1 - \frac{E}{N\varepsilon}\right)\right]\right\} \\ &= k\left\{N \ln N - \frac{N}{2} \ln \frac{N}{2} \left[\left(1 + \frac{E}{N\varepsilon}\right) + \left(1 - \frac{E}{N\varepsilon}\right)\right] \right. \\ &\quad \left. - \frac{N}{2}\left(1 + \frac{E}{N\varepsilon}\right) \ln\left(1 + \frac{E}{N\varepsilon}\right) - \frac{N}{2}\left(1 - \frac{E}{N\varepsilon}\right) \ln\left(1 - \frac{E}{N\varepsilon}\right)\right\} \\ &= k\left\{N \ln N - N \ln \frac{N}{2} \right. \\ &\quad \left. - \frac{N}{2}\left(1 + \frac{E}{N\varepsilon}\right) \ln\left(1 + \frac{E}{N\varepsilon}\right) - \frac{N}{2}\left(1 - \frac{E}{N\varepsilon}\right) \ln\left(1 - \frac{E}{N\varepsilon}\right)\right\} \\ &= Nk\left[\ln 2 - \frac{1}{2}\left(1 + \frac{E}{N\varepsilon}\right) \ln\left(1 + \frac{E}{N\varepsilon}\right) - \frac{1}{2}\left(1 - \frac{E}{N\varepsilon}\right) \ln\left(1 - \frac{E}{N\varepsilon}\right)\right]. \end{aligned}$$



# Negative temperature

Eq.(1.4.7)  $\delta W = V H dB$ ,  $B$  generalized displacement.

# Negative temperature

Eq.(1.4.7)  $\delta W = V H dB$ ,  $B$  generalized displacement.

$$\frac{1}{T} = \left( \frac{\partial S}{\partial E} \right)_B$$

# Negative temperature

Eq.(1.4.7)  $\delta W = V H dB$ ,  $B$  generalized displacement.

$$\begin{aligned} \frac{1}{T} &= \left( \frac{\partial S}{\partial E} \right)_B \\ &= \frac{\partial}{\partial E} \left\{ Nk \left[ \ln 2 - \frac{1}{2} \left( 1 + \frac{E}{N\epsilon} \right) \ln \left( 1 + \frac{E}{N\epsilon} \right) - \frac{1}{2} \left( 1 - \frac{E}{N\epsilon} \right) \ln \left( 1 - \frac{E}{N\epsilon} \right) \right] \right\} \end{aligned}$$

# Negative temperature

Eq.(1.4.7)  $\delta W = V H dB$ ,  $B$  generalized displacement.

$$\begin{aligned}
 \frac{1}{T} &= \left( \frac{\partial S}{\partial E} \right)_B \\
 &= \frac{\partial}{\partial E} \left\{ Nk \left[ \ln 2 - \frac{1}{2} \left( 1 + \frac{E}{N\varepsilon} \right) \ln \left( 1 + \frac{E}{N\varepsilon} \right) - \frac{1}{2} \left( 1 - \frac{E}{N\varepsilon} \right) \ln \left( 1 - \frac{E}{N\varepsilon} \right) \right] \right\} \\
 &= -\frac{Nk}{2} \left[ \left( 1 + \frac{E}{N\varepsilon} \right) \cdot \frac{\frac{1}{N\varepsilon}}{1 + \frac{E}{N\varepsilon}} + \frac{1}{N\varepsilon} \ln \left( 1 + \frac{E}{N\varepsilon} \right) + \left( 1 - \frac{E}{N\varepsilon} \right) \cdot \right. \\
 &\quad \left. \frac{-\frac{1}{N\varepsilon}}{1 - \frac{E}{N\varepsilon}} - \frac{1}{N\varepsilon} \ln \left( 1 - \frac{E}{N\varepsilon} \right) \right]
 \end{aligned}$$

# Negative temperature

Eq.(1.4.7)  $\delta W = V H dB$ ,  $B$  generalized displacement.

$$\begin{aligned}
 \frac{1}{T} &= \left( \frac{\partial S}{\partial E} \right)_B \\
 &= \frac{\partial}{\partial E} \left\{ Nk \left[ \ln 2 - \frac{1}{2} \left( 1 + \frac{E}{N\varepsilon} \right) \ln \left( 1 + \frac{E}{N\varepsilon} \right) - \frac{1}{2} \left( 1 - \frac{E}{N\varepsilon} \right) \ln \left( 1 - \frac{E}{N\varepsilon} \right) \right] \right\} \\
 &= -\frac{Nk}{2} \left[ \left( 1 + \frac{E}{N\varepsilon} \right) \cdot \frac{\frac{1}{N\varepsilon}}{1 + \frac{E}{N\varepsilon}} + \frac{1}{N\varepsilon} \ln \left( 1 + \frac{E}{N\varepsilon} \right) + \left( 1 - \frac{E}{N\varepsilon} \right) \cdot \right. \\
 &\quad \left. -\frac{\frac{1}{N\varepsilon}}{1 - \frac{E}{N\varepsilon}} - \frac{1}{N\varepsilon} \ln \left( 1 - \frac{E}{N\varepsilon} \right) \right] \\
 &= -\frac{Nk}{2} \left[ \frac{1}{N\varepsilon} + \frac{1}{N\varepsilon} \ln \left( 1 + \frac{E}{N\varepsilon} \right) - \frac{1}{N\varepsilon} - \frac{1}{N\varepsilon} \ln \left( 1 - \frac{E}{N\varepsilon} \right) \right]
 \end{aligned}$$

# Negative temperature

Eq.(1.4.7)  $\delta W = V H dB$ ,  $B$  generalized displacement.

$$\begin{aligned}
 \frac{1}{T} &= \left( \frac{\partial S}{\partial E} \right)_B \\
 &= \frac{\partial}{\partial E} \left\{ Nk \left[ \ln 2 - \frac{1}{2} \left( 1 + \frac{E}{N\varepsilon} \right) \ln \left( 1 + \frac{E}{N\varepsilon} \right) - \frac{1}{2} \left( 1 - \frac{E}{N\varepsilon} \right) \ln \left( 1 - \frac{E}{N\varepsilon} \right) \right] \right\} \\
 &= -\frac{Nk}{2} \left[ \left( 1 + \frac{E}{N\varepsilon} \right) \cdot \frac{\frac{1}{N\varepsilon}}{1 + \frac{E}{N\varepsilon}} + \frac{1}{N\varepsilon} \ln \left( 1 + \frac{E}{N\varepsilon} \right) + \left( 1 - \frac{E}{N\varepsilon} \right) \cdot \right. \\
 &\quad \left. -\frac{\frac{1}{N\varepsilon}}{1 - \frac{E}{N\varepsilon}} - \frac{1}{N\varepsilon} \ln \left( 1 - \frac{E}{N\varepsilon} \right) \right] \\
 &= -\frac{Nk}{2} \left[ \frac{1}{N\varepsilon} + \frac{1}{N\varepsilon} \ln \left( 1 + \frac{E}{N\varepsilon} \right) - \frac{1}{N\varepsilon} - \frac{1}{N\varepsilon} \ln \left( 1 - \frac{E}{N\varepsilon} \right) \right] \\
 &= -\frac{k}{2\varepsilon} \left[ \ln \left( 1 + \frac{E}{N\varepsilon} \right) - \ln \left( 1 - \frac{E}{N\varepsilon} \right) \right]
 \end{aligned}$$

# Negative temperature



Eq.(1.4.7)  $\delta W = V H dB$ ,  $B$  generalized displacement.

$$\begin{aligned}
 \frac{1}{T} &= \left( \frac{\partial S}{\partial E} \right)_B \\
 &= \frac{\partial}{\partial E} \left\{ Nk \left[ \ln 2 - \frac{1}{2} \left( 1 + \frac{E}{N\varepsilon} \right) \ln \left( 1 + \frac{E}{N\varepsilon} \right) - \frac{1}{2} \left( 1 - \frac{E}{N\varepsilon} \right) \ln \left( 1 - \frac{E}{N\varepsilon} \right) \right] \right\} \\
 &= -\frac{Nk}{2} \left[ \left( 1 + \frac{E}{N\varepsilon} \right) \cdot \frac{\frac{1}{N\varepsilon}}{1 + \frac{E}{N\varepsilon}} + \frac{1}{N\varepsilon} \ln \left( 1 + \frac{E}{N\varepsilon} \right) + \left( 1 - \frac{E}{N\varepsilon} \right) \cdot \right. \\
 &\quad \left. -\frac{\frac{1}{N\varepsilon}}{1 - \frac{E}{N\varepsilon}} - \frac{1}{N\varepsilon} \ln \left( 1 - \frac{E}{N\varepsilon} \right) \right] \\
 &= -\frac{Nk}{2} \left[ \frac{1}{N\varepsilon} + \frac{1}{N\varepsilon} \ln \left( 1 + \frac{E}{N\varepsilon} \right) - \frac{1}{N\varepsilon} - \frac{1}{N\varepsilon} \ln \left( 1 - \frac{E}{N\varepsilon} \right) \right] \\
 &= -\frac{k}{2\varepsilon} \left[ \ln \left( 1 + \frac{E}{N\varepsilon} \right) - \ln \left( 1 - \frac{E}{N\varepsilon} \right) \right] = \frac{k}{2\varepsilon} \ln \frac{N\varepsilon - E}{N\varepsilon + E}.
 \end{aligned}$$

# Negative temperature

Eq.(1.4.7)  $\text{d}W = V H \text{d}B$ ,  $B$  generalized displacement.

$$\begin{aligned}
 \frac{1}{T} &= \left( \frac{\partial S}{\partial E} \right)_B \\
 &= \frac{\partial}{\partial E} \left\{ Nk \left[ \ln 2 - \frac{1}{2} \left( 1 + \frac{E}{N\varepsilon} \right) \ln \left( 1 + \frac{E}{N\varepsilon} \right) - \frac{1}{2} \left( 1 - \frac{E}{N\varepsilon} \right) \ln \left( 1 - \frac{E}{N\varepsilon} \right) \right] \right\} \\
 &= -\frac{Nk}{2} \left[ \left( 1 + \frac{E}{N\varepsilon} \right) \cdot \frac{\frac{1}{N\varepsilon}}{1 + \frac{E}{N\varepsilon}} + \frac{1}{N\varepsilon} \ln \left( 1 + \frac{E}{N\varepsilon} \right) + \left( 1 - \frac{E}{N\varepsilon} \right) \cdot \right. \\
 &\quad \left. -\frac{\frac{1}{N\varepsilon}}{1 - \frac{E}{N\varepsilon}} - \frac{1}{N\varepsilon} \ln \left( 1 - \frac{E}{N\varepsilon} \right) \right] \\
 &= -\frac{Nk}{2} \left[ \frac{1}{N\varepsilon} + \frac{1}{N\varepsilon} \ln \left( 1 + \frac{E}{N\varepsilon} \right) - \frac{1}{N\varepsilon} - \frac{1}{N\varepsilon} \ln \left( 1 - \frac{E}{N\varepsilon} \right) \right] \\
 &= -\frac{k}{2\varepsilon} \left[ \ln \left( 1 + \frac{E}{N\varepsilon} \right) - \ln \left( 1 - \frac{E}{N\varepsilon} \right) \right] = \frac{k}{2\varepsilon} \ln \frac{N\varepsilon - E}{N\varepsilon + E}.
 \end{aligned}$$





# Table of contents

## 1 §7 Boltzmann Statistics

- 7.1 Thermal quantities in statistics
- 7.2 Equation of state of ideal gas
- 7.3 Maxwell speed distribution
- 7.4 Energy equipartition theorem
- 7.5 Internal energy and heat capacity of the ideal gas
- 7.6 Entropy of the ideal gas
- 7.7 Einstein's theory on heat capacity of solid
- 7.8 Paramagnetic solid
- 7.9 Negative temperature