

some bundle stuff

ho boon suan

November 25, 2020

Proposition 10.26. *Suppose E and E' are smooth vector bundles over a smooth manifold M with or without boundary, and $F: E \rightarrow E'$ is a bijective smooth bundle homomorphism over M . Then F is a smooth bundle isomorphism.*

Proof. We first show that F is a bundle isomorphism. Since $\pi' \circ F = \pi$, we have $\pi' = \pi \circ F^{-1}$. It then suffices to show that $F|_{E_p}: E_p \rightarrow E'_p$ is an isomorphism of vector spaces. Since F is a bijection, its restriction $F|_{E_p}$ is a bijection onto its image $\text{im}(F|_{E_p}) \subset E'_p$. Since $x \notin E_p$ implies that $F(x) \notin E'_p$, it follows that $E'_p \subset \text{im}(F|_{E_p})$. In particular, E and E' are vector bundles of the same rank k .

We now show that F is a local diffeomorphism. Since bijective local diffeomorphisms are diffeomorphisms, this will complete the proof. Let $p \in U \subset M$ for an appropriate neighborhood U of p , and choose local trivializations Φ and Ψ of E and E' as in the diagram

$$\begin{array}{ccccc} U \times \mathbf{R}^k & \xleftarrow{\Phi} & \pi^{-1}(U) & \xrightarrow{F} & \pi'^{-1}(U) & \xrightarrow{\Psi} & U \times \mathbf{R}^k \\ & \searrow \pi_U & \searrow \pi & & \swarrow \pi' & \swarrow \pi_U & \\ & & U & & & & \end{array} .$$

We then have

$$(\Psi \circ F \circ \Phi^{-1})(p, v) = (p, \tau(p)v)$$

for some smooth $\tau: U \rightarrow \text{GL}(n, \mathbf{R})$. It follows that

$$(\Phi \circ F^{-1} \circ \Psi^{-1})(p, v) = (p, \tau(p)^{-1}v),$$

which implies that F^{-1} is smooth on $\pi'^{-1}(U)$, since inversion is a diffeomorphism in $\text{GL}(n, \mathbf{R})$. This completes the proof. \square