






EGFR · AS intronic lncRNA
EGFR driver locus
log2FC (TKI_resistant / TKI_sensitive)
0
-2
-4
-6
EGFR
BambuTx285_i
EGFR-223
EGFR-204
EGFR-201
EGFR-220
EGFR-222
EGFR-207
DE (FDR < 0.05)
No
Yes
Feature type
gene
novel lncRNA
protein coding
protein coding CDS not defined



NAPSA · lineage marker loss
LUAD lineage identity
log2FC (TKI_resistant / TKI_sensitive)
0
-2
-4
-6
-8
NAPSA
NAPSA-215
NAPSA-220
NAPSA-216
NAPSA-206
NAPSA-227
NAPSA-201
NAPSA-207
DE (FDR < 0.05)
No
Yes
Feature type
gene
protein coding
retained intron



CEACAM5 · lineage marker loss
LUAD lineage identity
log2FC (TKI_resistant / TKI_sensitive)
0.0
-2.5
-5.0
-7.5
CEACAM5
CEACAM5-216
CEACAM5-212
CEACAM5-246
DE (FDR < 0.05)
No
Yes
Feature type
gene
protein coding



KRT7 · lineage marker
LUAD lineage identity · 10 of 23 isoforms shown
log2FC (TKI_resistant / TKI_sensitive)
0
-2
-4
-6
-8
KRT7
KRT7-233
KRT7-231
KRT7-228
KRT7-215
KRT7-236
KRT7-201
KRT7-238
KRT7-214
KRT7-226
KRT7-206
DE (FDR < 0.05)
No
Yes
Feature type
gene
nonsense mediated decay
protein coding
protein coding CDS not defined



NDUFA12 · isoform switch
OXPHOS isoform switching · 10 of 11 isoforms shown
log2FC (TKI_resistant / TKI_sensitive)
10
5
0
-5
-10
NDUFA12
NDUFA12-208
NDUFA12-216
NDUFA12-227
NDUFA12-211
NDUFA12-206
NDUFA12-204
NDUFA12-224
NDUFA12-222
NDUFA12-201
NDUFA12-212
DE (FDR < 0.05)
No
Yes
Feature type
gene
nonsense mediated decay
protein coding
retained intron



COX6B1 · isoform switch
OXPHOS isoform switching
log2FC (TKI_resistant / TKI_sensitive)
5
0
-5
-10
COX6B1
COX6B1-221
COX6B1-215
COX6B1-202
COX6B1-222
COX6B1-219
COX6B1-220
COX6B1-216
COX6B1-208
COX6B1-204
DE (FDR < 0.05)
No
Yes
Feature type
gene
nonsense mediated decay
protein coding



NDUFA6 · isoform switch
OXPHOS isoform switching
log2FC (TKI_resistant / TKI_sensitive)
2.5
0.0
-2.5
-5.0
-7.5
NDUFA6
NDUFA6-210
NDUFA6-206
NDUFA6-207
NDUFA6-211
NDUFA6-203
NDUFA6-205
NDUFA6-202
DE (FDR < 0.05)
No
Yes
Feature type
gene
nonsense mediated decay
protein coding



ELOC · switch + AS intronic lncRNA
Hypoxia (VHL-HIF axis) · 10 of 34 isoforms shown
log2FC (TKI_resistant / TKI_sensitive)
5
0
-5
-10
ELOC
BambuTx326_i
ELOC-206
ELOC-230
ELOC-301
ELOC-304
ELOC-261
ELOC-290
ELOC-287
ELOC-210
ELOC-303
DE (FDR < 0.05)
No
Yes
Feature type
gene
novel lncRNA
protein coding



MIR210HG · hypoxia lncRNA
Hypoxia (VHL-HIF axis)
log2FC (TKI_resistant / TKI_sensitive)
2
1
0
MIR210HG
MIR210HG-205
MIR210HG-201
DE (FDR < 0.05)
No
Yes
Feature type
gene
lncRNA



UBE2I / UBE2I-AS1 · AS lncRNA
lncRNA regulation · 10 of 18 isoforms shown
log2FC (TKI_resistant / TKI_sensitive)
8
4
0
-4
UBE2I
UBE2I-AS1-209
UBE2I-AS1-215
UBE2I-AS1-214
UBE2I-AS1-205
UBE2I-AS1-204
UBE2I-AS1-216
UBE2I-AS1-217
UBE2I-214
UBE2I-246
UBE2I-206
DE (FDR < 0.05)
No
Yes
Feature type
gene
lncRNA
protein coding


SNHG17 · lncRNA isoforms
lncRNA regulation · 10 of 27 isoforms shown
log2FC (TKI_resistant / TKI_sensitive)
6
4
2
0
SNHG17
SNHG17-251
SNHG17-319
SNHG17-248
SNHG17-226
SNHG17-245
SNHG17-297
SNHG17-337
SNHG17-232
SNHG17-321
SNHG17-237
DE (FDR < 0.05)
No
Yes
Feature type
gene
lncRNA


TPT1 · switch + NMD isoform loss
Translation / survival · 10 of 43 isoforms shown
log2FC (TKI_resistant / TKI_sensitive)
5
0
-5
-10
TPT1
TPT1-228
TPT1-250
TPT1-242
TPT1-201
TPT1-203
TPT1-231
TPT1-266
TPT1-218
TPT1-268
TPT1-227
DE (FDR < 0.05)
No
Yes
Feature type
gene
nonsense mediated decay
protein coding


POU2F3 · tuft-cell marker
Lineage plasticity
log2FC (TKI_resistant / TKI_sensitive)
0
-1
-2
POU2F3
POU2F3-208
POU2F3-203
DE (FDR < 0.05)
No
Yes
Feature type
gene
protein coding
retained intron


HMGA2 · EMT marker
Lineage plasticity
log2FC (TKI_resistant / TKI_sensitive)
1.5
1.0
0.5
0.0
-0.5
-1.0
HMGA2
HMGA2-202
HMGA2-214
HMGA2-204
HMGA2-201
DE (FDR < 0.05)
No
Yes
Feature type
gene
protein coding


SEC61G · EGFR-amplicon neighbour
EGFR driver locus · 10 of 18 isoforms shown
log2FC (TKI_resistant / TKI_sensitive)
5
0
-5
-10
SEC61G
SEC61G-213
SEC61G-231
SEC61G-201
SEC61G-228
SEC61G-225
SEC61G-241
SEC61G-218
SEC61G-210
SEC61G-202
SEC61G-209
DE (FDR < 0.05)
No
Yes
Feature type
gene
protein coding