

a




Filters reads, aligns them in splice-aware mode, and estimate mapping coverage per sample; unstranded ONT cDNA is first reoriented.




Displays novel transcripts in their genomic context and summarizes the transcriptome in an integrated report.


integrativebioinformatics/

pulposeq





Transcriptome assembly, novel isoform disovery and quantifiction. Novel isoforms are classified according to coding potential prediction scores.




Novel transcripts ≥ 500 nt are assigned a biotype (novel lncRNA, novel non-coding or novel protein-coding) based on its structural relationship with reference annotations defined by GffCompare's classification system.

reference *novel*


b Length bias

Equimolar SIRVs; 0 = expected abundance; within 2× = E0 isoforms with median abundance within 2-fold of expected; F1 = SIRV identification after validation




c ERCC dose response

Detected / in mix; R^2; LOD (amol/µl)




d Sequin ground truth

Mix B (SCLC-A) vs Mix A; edgeR glmTreat (H0 |log2FC| <= 0.5), FDR < 0.05; shaded: not truly DE




e Read-level support across biotypes




f Novel transcripts' length

Medians




g UIC sharing

UIC = Unique Intron Chain




h UIC sharing by length

Facet = source protocol; point size = n; bins with n < 10 omitted



i Transcript ends within an identical UIC

Share of protocol pairs whose ends lie within a distance; grey lines = 50 and 500 bp



97.5% shared UICs agree on biotype, 98.7% on class code and 98.4% host gene