

# Recent results and ongoing research

## 1 Sublinear Explicit Incremental Planar Voronoi Diagrams

Let  $S := \{s_1, s_2, \dots, s_N\}$  be a set of  $N$  distinct points in the Euclidean plane. *Voronoi diagram* of  $S$  is the subdivision of the plane into  $N$  *cells*, one for each site in  $S$ , with the property that a point  $q$  lies in the cell corresponding to a site  $s_i$  if and only if  $\text{dist}(q, s_i) < \text{dist}(q, s_j)$  for each  $s_j \in S$  with  $j \neq i$ .

Voronoi diagrams have been of a great interest in TCS (computational geometry and other domains) since their first thorough study [8] and the research on the subject is still going on [5]. The aim of this work is to maintain the graph of a Voronoi diagram under insertion of new sites in time sublinear in  $N$ .

It was observed previously [2] that while there could be a linear number of changes to the embedded Voronoi diagram with each site insertion, this is not equivalent to the number of combinatorial changes to the graph structure of the Voronoi diagram. It was then proved by showing that the amortized number of combinatorial changes is only  $\Theta(N^{\frac{1}{2}})$  per site insertion.

This observation opened the possibility to maintain the Voronoi diagram graph in faster than linear time per insertion, which was consequently done for the restricted case where the sites are in convex position [2].

We create a data structure that explicitly maintains the graph of a Voronoi diagram. The diagram is stored in adjacency list format on which primitive operations, including links and cuts, are performed. Our data structure allows insertions of new sites in  $\tilde{O}(N^{\frac{3}{4}})$  expected amortized time, where  $\tilde{O}$  suppresses polylogarithmic terms. Previously, no sublinear method was known for this problem. The result has been presented at JCDCGGG<sup>3</sup> conference [3] and is now awaiting publication in the special issue of Journal of Information Processing.

## 2 Ongoing research: $\tilde{O}$ ptimal-Time Incremental Combinatorial Voronoi Diagrams

The time per insertion of a new site in our previous work is sublinear yet not optimal. We are now aiming at maintaining the Voronoi diagram while having optimal,  $\tilde{O}(N^{\frac{1}{2}})$  time per insertion.

The idea is to use divide-and-conquer strategy. There is a known divide-and-conquer algorithm for the static case of the Voronoi diagram: on each level the sites are divided into two roughly equal subsets by a vertical line, for each of them the diagram is calculated recursively, the two Voronoi diagrams are merged afterwards. The algorithm does not fit directly for semi-dynamic case. I am now working on figuring this out and hope to get the final result soon.

## 3 Previous research: All convex polyhedra that can be glued from regular pentagons

A *net* is a set of polygons equipped with a number of rules describing the way edges of these polygons must be glued to each other. Alexandrov's Uniqueness Theorem [1] says that if a net is homeomorphic to a sphere and the sum of angles at each of its vertices is at most  $360^\circ$  then there is a single convex polyhedron  $P$  that can be glued from this net.

However, the proof of this theorem is highly non-constructive, and the algorithmic question of *constructing* the polyhedron  $P$ , coordinates of its vertices or even its 1-dimensional skeleton in polynomial time remains open for around 60 years.

To get closer to the solution of this problem one can solve some of its restricted cases. One way is to consider a single polygon of a particular type, i.e. a regular polygon, and *edge-to-edge* gluings (an edge of a polygon  $T_i$  needs to be glued to an entire other edge of another polygon  $T_j$ , possibly  $j = i$ ) of its several copies [6].

Using a method of enumerating all graph structures [7], we are able to solve the problem of establishing the graph structure of convex polyhedra that are glued from regular pentagons edge-to-edge and list all such polyhedra.

We derive an upper bound for discrepancy in vertex coordinates between the unique convex polyhedron corresponding to a given polyhedral metric and a given approximate polyhedron, this implies a sufficient condition for the polyhedron to have a certain edge. The program is developed that checks this condition. We also apply geometric methods to establish the graph structures that are non-simplicial.

The main outcome of this work is the full list of the polyhedra that are obtained by edge-to-edge gluings of regular pentagons. However, the methods for obtaining it are of independent interest and may be applied to other problems of the same flavour which gives this work a lot of implications related to considering different polygons and gluings composed of them. I presented this result at EGC 2019 [4]. It is now awaiting publication in the special issue of Journal of Information Processing.

## References

- [1] A. Alexandrov. *Convex Polyhedra*. Springer-Verlag, Berlin, 2005. Translation from Russian.
- [2] S. R. Allen, L. Barba, J. Iacono, and S. Langerman. Incremental voronoi diagrams. *Discrete & Computational Geometry*, 58(4):822–848, 2017.
- [3] E. Arseneva, J. Iacono, G. Koumoutsos, S. Langerman, and B. Zolotov. Sublinear explicit incremental planar voronoi diagrams. In *Japan Conference on Discrete and Computational Geometry, Graphs, and Games*, page 35–36, Tokyo University of Science, September 2019.
- [4] E. Arseneva, S. Langerman, and B. Zolotov. A complete list of all convex shapes made by gluing regular pentagons. In *XVIII Spanish Meeting on Computational Geometry*, page 26–29, Girona, Spain, July 2019.
- [5] F. Aurenhammer, R. Klein, and D.-T. Lee. *Voronoi Diagrams and Delaunay Triangulations*. World Scientific Publishing Company, August 2013.
- [6] E. Demaine and J. O’Rourke. *Geometric folding algorithms*. Cambridge University Press, 2007.
- [7] E. Khramtcova and S. Langerman. Which convex polyhedra can be made by gluing regular hexagons? In *JCDCG<sup>3</sup>*, 2017.
- [8] G. Voronoi. Nouvelles applications des paramètres continus à la théorie de formes quadratiques. *Journal für die reine und angewandte Mathematik*, 134:198–287, 1908.