

Rest
Stim 8 h
Stim 48 h
A Out-degree
P(degree ≥ k)
CD3E (8 h)
ZAP70 (8 h)
LCP2 (8 h)
Genes changed per knockdown, k
B In-degree
TXNIP (Rest)
SESN1 (48 h)
FDXR (48 h)
Knockdowns changing the gene, k
C Genome coverage
IMD
Non-IMD control
Reference SNPs (%)
Anchors
Regulators
Targets
R 8h 48h R 8h 48h
D Expanded set sizes
Anchor set size
All knockdowns with an effect
Regulators
Targets
Autoimmune thyroid disease
Crohn's disease
Childhood-onset asthma
Type 1 diabetes
Rheumatoid arthritis (EUR)
Inflammatory bowel disease
Adult-onset asthma
Atopic dermatitis
Rheumatoid arthritis (EAS)
Asthma (EAS)
Ulcerative colitis
Systemic lupus erythematosus
Type 2 diabetes
Myocardial infarction
Schizophrenia
Osteoarthritis
Alzheimer's disease (proxy)
Ischaemic stroke
Genes in set
E Share of set from largest anchor
RASGRP1/CD247
UBE2L3
TRAF3/CD247
TUFM
RASGRP1
TUFM
SIK2/CD247
TRAF3
RASGRP1
UBE2L3
FBXW7
REST/FDX1
KMT2E
NDUFS2
Share