

| Equation of Motion Time Discretization

| Newton's 2nd Law as a
first-order system

$$\begin{cases} \dot{\mathbf{x}} = \mathbf{v} \\ \mathbf{M}\dot{\mathbf{v}} = \mathbf{f}(\mathbf{x}, \mathbf{v}) \end{cases}$$

BDF1

$$\dot{\mathbf{y}}_t \approx \frac{\mathbf{y}_{t+1} - \mathbf{y}_t}{h}$$

| Explicit Euler

$$\begin{cases} \mathbf{x}^+ = \mathbf{x}^- + h \mathbf{v}^- \\ \mathbf{v}^+ = \mathbf{v}^- + h \mathbf{M}^{-1} \mathbf{f}(\mathbf{x}^-, \mathbf{v}^-) \end{cases}$$

| Implicit Euler

$$\begin{cases} \mathbf{x}^+ = \mathbf{x}^- + h \mathbf{v}^+ \\ \mathbf{M}(\mathbf{v}^+ - \mathbf{v}^-) = h \mathbf{f}(\mathbf{x}^+, \mathbf{v}^+) \end{cases}$$

| Implicit Euler

$$\begin{cases} \mathbf{x}^+ = \mathbf{x}^- + h \mathbf{v}^+ \\ \mathbf{M}(\mathbf{v}^+ - \mathbf{v}^-) = h \mathbf{f}(\mathbf{x}^+, \mathbf{v}^+) \end{cases}$$



$$\mathbf{r}(\mathbf{x}^+) = \frac{1}{h^2} \mathbf{M}(\mathbf{x}^+ - \tilde{\mathbf{x}}) - \mathbf{f}(\mathbf{x}^+) = \mathbf{0}$$



**Non-Linear
System of
Equations**

$$\frac{1}{h^2} \begin{bmatrix} m_1 & & & \\ & m_2 & & \\ & & \ddots & \\ & & & m_n \end{bmatrix} \begin{bmatrix} \mathbf{x}_1^+ - \tilde{\mathbf{x}}_1 \\ \mathbf{x}_2^+ - \tilde{\mathbf{x}}_2 \\ \vdots \\ \mathbf{x}_n^+ - \tilde{\mathbf{x}}_n \end{bmatrix} - \begin{bmatrix} \mathbf{f}_1(\mathbf{x}^+) \\ \mathbf{f}_2(\mathbf{x}^+) \\ \vdots \\ \mathbf{f}_n(\mathbf{x}^+) \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

| Implicit Euler

$$\mathbf{r}(\mathbf{x}^+) = \frac{1}{h^2} \mathbf{M} (\mathbf{x}^+ - \tilde{\mathbf{x}}) - \mathbf{f}(\mathbf{x}^+) = \mathbf{0}$$

Unknown

Timestep Length

$\mathbf{x}^- + h \mathbf{v}^-$

Force: Non-Linear Function

Diagonal Mass Matrix: $\begin{bmatrix} m_1 & & \\ & \ddots & \\ & & m_n \end{bmatrix}$

The diagram illustrates the Implicit Euler method equation: $\mathbf{r}(\mathbf{x}^+) = \frac{1}{h^2} \mathbf{M} (\mathbf{x}^+ - \tilde{\mathbf{x}}) - \mathbf{f}(\mathbf{x}^+) = \mathbf{0}$. Annotations include: 'Unknown' pointing to $\mathbf{x}^+$; 'Timestep Length' pointing to h ; $\mathbf{x}^- + h \mathbf{v}^-$ pointing to $\tilde{\mathbf{x}}$; 'Force: Non-Linear Function' pointing to $\mathbf{f}(\mathbf{x}^+)$; and 'Diagonal Mass Matrix' pointing to $\mathbf{M}$, which is represented by a diagonal matrix with elements $m_1, \dots, m_n$.

| Newton-Raphson Method

We have:

$$\mathbf{g} : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

We want:


$$\mathbf{g}(\mathbf{z}^*) = \mathbf{0}$$

Linearize around a guess:

$$\mathbf{g}(\mathbf{z}^{(k)} + \Delta \mathbf{z}) \approx \mathbf{g}(\mathbf{z}^{(k)}) + \frac{\partial \mathbf{g}}{\partial \mathbf{z}}(\mathbf{z}^{(k)}) \Delta \mathbf{z} = \mathbf{0}$$

Linear System of Equations

$$\frac{\partial \mathbf{g}}{\partial \mathbf{z}} \Delta \mathbf{z} = -\mathbf{g}(\mathbf{z}^{(k)})$$
$$\mathbf{z}^{(k+1)} = \mathbf{z}^{(k)} + \Delta \mathbf{z}$$



| Newton-Raphson Method for Implicit Dynamic System

We have:


$$\mathbf{r} : \mathbb{R}^{3n} \rightarrow \mathbb{R}^{3n} \quad \text{such that:} \quad \mathbf{r}(\mathbf{x}) = \frac{1}{h^2} \mathbf{M}(\mathbf{x} - \tilde{\mathbf{x}}) - \mathbf{f}(\mathbf{x})$$

We want:

$$\mathbf{r}(\mathbf{x}^+) = \mathbf{0}$$

Linearize around a guess:

$$\frac{\partial \mathbf{r}}{\partial \mathbf{x}} = \underbrace{\frac{1}{h^2} \mathbf{M} - \frac{\partial \mathbf{f}}{\partial \mathbf{x}}}_{\mathbf{A} \in \mathbb{R}^{3n \times 3n}}$$



| Newton-Raphson Method Single Iteration

Expanded:

$$\underbrace{\left[\mathbf{M} - h^2 \frac{\partial \mathbf{f}}{\partial \mathbf{x}} \right]}_{\mathbf{A}(\mathbf{x}^-)} \Delta \mathbf{x} = \underbrace{h^2 \mathbf{f}(\mathbf{x}^-) + h \mathbf{M} \mathbf{v}^-}_{\mathbf{b}(\mathbf{x}^-)}$$


Algorithm:

- 1 **build:** $\mathbf{A}, \mathbf{b}$
- 2 **solve:** $\mathbf{A} \Delta \mathbf{x} = \mathbf{b}$
- 3 $\mathbf{x}^+ = \mathbf{x}^- + \Delta \mathbf{x}$
- 4 $\mathbf{v}^+ = \frac{1}{h} (\mathbf{x}^+ - \mathbf{x}^-)$

| Recasting Root-Finding as Minimization

Does exist a: $g(\mathbf{x})$ such that: $\nabla g(\mathbf{x}) = \mathbf{r}(\mathbf{x})$

$$g(\mathbf{x}) = \frac{1}{2h^2} (\mathbf{x} - \tilde{\mathbf{x}})^\top \mathbf{M} (\mathbf{x} - \tilde{\mathbf{x}}) + U(\mathbf{x})$$



$$\mathbf{f}(\mathbf{x}) = -\frac{\partial U}{\partial \mathbf{x}}(\mathbf{x}) \quad \text{Conservative Forces}$$

$$\nabla g(\mathbf{x}) = \frac{1}{h^2} \mathbf{M} (\mathbf{x} - \tilde{\mathbf{x}}) - \mathbf{f}(\mathbf{x}) = \mathbf{r}(\mathbf{x})$$

| Recasting Root-Finding as Minimization

Does exists a: $g(\mathbf{x})$ such that: $\nabla g(\mathbf{x}) = \mathbf{r}(\mathbf{x})$

$$g(\mathbf{x}) = \frac{1}{2h^2} (\mathbf{x} - \tilde{\mathbf{x}})^\top \mathbf{M} (\mathbf{x} - \tilde{\mathbf{x}}) + U(\mathbf{x})$$


$$\mathbf{f}(\mathbf{x}) = -\frac{\partial U}{\partial \mathbf{x}}(\mathbf{x})$$


Conservative Forces

$$\mathbf{r}(\mathbf{x}) = \mathbf{0} \quad \Longleftrightarrow \quad \nabla g(\mathbf{x}) = \mathbf{0} \quad \begin{array}{c} \text{if Convex} \\ \Longleftrightarrow \end{array} \quad \min_{\mathbf{x}} g(\mathbf{x})$$


| Root-Finding and Minimization Equivalence

Equivalent Problems:

$$\mathbf{x}^+ = \arg \min_{\mathbf{x}} g(\mathbf{x})$$



$$\mathbf{r}(\mathbf{x}^+) = 0$$



| Descent Methods How to Solve the Minimization Problem

```
1 Initialize  $\mathbf{x}^{(0)}$ 
2 for  $k = 0, 1, \dots, N - 1$  do
3    $\Delta \mathbf{x} = \text{DescentDirection}()$ 
4    $\mathbf{x}^{(k+1)} = \mathbf{x}^{(k)} + \alpha^{(k)} \cdot \Delta \mathbf{x}$ 
5  $\mathbf{x}^+ = \mathbf{x}^{(N)}$ 
6  $\mathbf{v}^+ = \frac{1}{h}(\mathbf{x}^+ - \mathbf{x}^-)$ 
```

| Jacobian:

$$\mathbf{J} = \nabla g(\mathbf{x}) = \mathbf{r}(\mathbf{x}) \in \mathbb{R}^{3n}$$

| Hessian:

$$\mathbf{H} = \nabla^2 g(\mathbf{x}) = \mathbf{A}(\mathbf{x}) \in \mathbb{R}^{3n \times 3n}$$

| Gradient Descent:

$$\Delta \mathbf{x}^{(k)} = -\mathbf{J}^{(k)}$$

| Newton:

$$\Delta \mathbf{x}^{(k)} = -[\mathbf{H}^{(k)}]^{-1} \mathbf{J}^{(k)}$$

| Quasi-Newton:

$$\Delta \mathbf{x}^{(k)} = -\mathbf{P}^{(k)} \mathbf{J}^{(k)}, \quad \mathbf{P}^{(k)} \approx [\mathbf{H}^{(k)}]^{-1}$$

| Projective Dynamics:

$$\Delta \mathbf{x}^{(k)} = -\mathbf{L}_{\text{PD}}^{-1} \mathbf{J}_{\text{PD}}^{(k)}$$

$$\mathbf{J}_{\text{PD}} = \nabla g_{\text{PD}}(\mathbf{x}), \quad \mathbf{L}_{\text{PD}} \approx \mathbf{H} \text{ (fixed)}$$

| Baraff-Witkin:

$$\Delta \mathbf{x} = -\mathbf{H}^{-1} \mathbf{J}, \quad k = 1$$

| Newton's Minimization Method = Newton-Raphson Root Finding's Method



function NEWTONMETHOD($\mathbf{x}^-$, $\mathbf{v}^-$, N)

```
1 Initialize  $\mathbf{x}^{(0)}$ 
2 for  $k = 0, 1, \dots, N - 1$  do
3     BUILD  $\mathbf{J}^{(k)}$ ,  $\mathbf{H}^{(k)}$ 
4     SOLVE  $\mathbf{H}^{(k)} \Delta \mathbf{x} = -\mathbf{J}^{(k)}$ 
5      $\mathbf{x}^{(k+1)} \leftarrow \mathbf{x}^{(k)} + \Delta \mathbf{x}$ 
6  $\mathbf{x}^+ \leftarrow \mathbf{x}^{(N)}$ 
7  $\mathbf{v}^+ \leftarrow \frac{1}{h}(\mathbf{x}^+ - \mathbf{x}^-)$ 
```

function NEWTONRAPHSON($\mathbf{x}^-$, $\mathbf{v}^-$, N)



```
1 Initialize  $\mathbf{x}^{(0)}$ 
2 for  $k = 0, 1, \dots, N - 1$  do
3     BUILD  $\mathbf{A}^{(k)}$ ,  $\mathbf{b}^{(k)}$ 
4     SOLVE  $\mathbf{A}^{(k)} \Delta \mathbf{x} = \mathbf{b}^{(k)}$ 
5      $\mathbf{x}^{(k+1)} \leftarrow \mathbf{x}^{(k)} + \Delta \mathbf{x}$ 
6  $\mathbf{x}^+ \leftarrow \mathbf{x}^{(N)}$ 
7  $\mathbf{v}^+ \leftarrow \frac{1}{h}(\mathbf{x}^+ - \mathbf{x}^-)$ 
```

| Convexity and Newton's Method When the Energy **IS** Convex





| Convexity and Newton's Method When the Energy **IS** Convex

$$g(\mathbf{x}^{(k+1)}) < g(\mathbf{x}^{(k)})$$





| Convexity and Newton's Method When the Energy **IS NOT** Convex

- There might be multiple stationary points
- There might be areas where the curvature is negative





| Convexity and Newton's Method: Multiple Stationary Points



| Convexity and Newton's Method: Negative Curvature

If the curvature is negative, by taking the Newton's step, there might be an **increase** in Energy

$$g(\mathbf{x}^{(k+1)}) > g(\mathbf{x}^{(k)})$$



| Convexity and Newton's Method: Solve the Negative Curvature Problem


A matrix is SPD (**S**ymmetric **P**ositive **D**efinite), written $\mathbf{A} \succ 0$ if it is symmetric and all its eigenvalues are positive:

$$\mathbf{A} = \mathbf{A}^\top \quad \text{and} \quad \mathbf{A} = \mathbf{Q} \mathbf{\Lambda} \mathbf{Q}^\top, \quad \lambda_i > 0 \quad \forall i.$$

- $\mathbf{A} \succ 0, \mathbf{B} \succ 0 \Rightarrow \mathbf{A} + \mathbf{B} \succ 0$
- $\mathbf{A} \succ 0, \mathbf{B} \not\succ 0 \Rightarrow \mathbf{A} + \mathbf{B}$ no guarantee

$\mathbf{H} \not\succ 0$: $\Delta \mathbf{x} = -\mathbf{H}^{-1} \mathbf{J}$ then $\Delta \mathbf{x}$ may not be a descent direction

$\bar{\mathbf{H}} \succ 0$: $\Delta \mathbf{x} = -\bar{\mathbf{H}}^{-1} \mathbf{J}$ then $\alpha \cdot \Delta \mathbf{x}$ is guaranteed to be a descent direction for a sufficiently small $\alpha > 0$

 is the SPD-Projection of $\mathbf{H}$

Convexity and Newton's Method: Solve the Negative Curvature Problem

How to fix the Energy's Hessian

Always Convex

$$g(\mathbf{x}) = \frac{1}{2h^2} (\mathbf{x} - \tilde{\mathbf{x}})^\top \mathbf{M} (\mathbf{x} - \tilde{\mathbf{x}}) + U(\mathbf{x})$$

$\mathbf{H} = \frac{1}{h^2} \mathbf{M} + \nabla^2 U(\mathbf{x})$

Always SPD Might not be SPD:
we must **SPD-Project**

$\nabla^2 U(\mathbf{x}) \in \mathbb{R}^{3n \times 3n}$: it is very expensive to SPD-Project such a big matrix

Convexity and Newton's Method: Solve the Negative Curvature Problem

How is the Potential Energy Factor Constructed

$$U(\mathbf{x}) = \sum_e U_e(\mathbf{x}_e)$$

The total energy is the sum of single element energies. Each element (edge, triangle, tetrahedron) involves only a few particles (2, 3 or 4).

$$\nabla^2 U(\mathbf{x}) = \sum_e \nabla^2 U_e(\mathbf{x}_e)$$

$$\nabla^2 U_e(\mathbf{x}_e) \in \mathbb{R}^{6 \times 6 | 9 \times 9 | 12 \times 12}$$

So the Hessian is also a sum of small element Hessians. Each one can be easily SPD-projected independently and in parallel, then assembled into the full Hessian.

Convexity and Newton's Method: Solve the Negative Curvature Problem

How an SPD-Projection is computed

How do we SPD-Project the single element energies:

$$\nabla^2 U_e(\mathbf{x}_e) \in \mathbb{R}^{6 \times 6 | 9 \times 9 | 12 \times 12}$$

Two ways:

- **Numerical:**

$$\nabla^2 U_e = \mathbf{Q} \mathbf{\Lambda} \mathbf{Q}^\top \longrightarrow \bar{\lambda}_i = \max(\lambda_i, 0) \longrightarrow \overline{\nabla^2 U_e} = \mathbf{Q} \bar{\mathbf{\Lambda}} \mathbf{Q}^\top$$

- **Analytical:**

for many frequently used energies the eigenvalues and eigenvectors are known in closed form. This route is usually faster and more robust.

| Analytical SPD Projection Quadratic Spring Potential


Spring:

$$\mathbf{x}_e = \begin{bmatrix} \mathbf{x}_1 \\ \mathbf{x}_2 \end{bmatrix} \in \mathbb{R}^6 \quad \mathbf{e} = \mathbf{x}_1 - \mathbf{x}_2$$

$$\hat{\mathbf{e}} = \frac{\mathbf{e}}{\|\mathbf{e}\|}$$

Energy:

$$U_e(\mathbf{x}_e) = \frac{k}{2} (\|\mathbf{e}\| - l_{\text{rest}})^2$$



Hessian:

$$\mathbf{H}_e = \begin{bmatrix} \mathbf{K} & -\mathbf{K} \\ -\mathbf{K} & \mathbf{K} \end{bmatrix} \in \mathbb{R}^{6 \times 6}$$

$$\mathbf{K} = k \left[\hat{\mathbf{e}} \hat{\mathbf{e}}^\top + \left(1 - \frac{l}{\|\mathbf{e}\|} \right) (\mathbf{I} - \hat{\mathbf{e}} \hat{\mathbf{e}}^\top) \right]$$

Eigenvalues:

$$\lambda_1 \quad \lambda_2 \quad \lambda_3 \quad \lambda_4 \quad \lambda_5 = \lambda_6$$

$$0, \quad 0, \quad 0, \quad 2k, \quad 2k \left(1 - \frac{l}{\|\mathbf{e}\|} \right)$$

Can be negative

SPD-Hessian:

$$\bar{\mathbf{H}}_e = \begin{bmatrix} \bar{\mathbf{K}} & -\bar{\mathbf{K}} \\ -\bar{\mathbf{K}} & \bar{\mathbf{K}} \end{bmatrix} \in \mathbb{R}^{6 \times 6} \quad \bar{\mathbf{H}}_e \succeq 0$$

$$\bar{\mathbf{K}} = k \left[\hat{\mathbf{e}} \hat{\mathbf{e}}^\top + \max \left(1 - \frac{l}{\|\mathbf{e}\|}, 0 \right) (\mathbf{I} - \hat{\mathbf{e}} \hat{\mathbf{e}}^\top) \right]$$

| SPD-Aware Newton's Method Algorithm

```
function SPDAWARENEWTONMETHOD( $\mathbf{x}^-$ ,  $\mathbf{v}^-$ )  
  2: Initialize  $\mathbf{x}^{(0)}$   
  3: for  $k = 0, 1, \dots, N - 1$  do  
    4:  $\mathbf{J} \leftarrow \nabla g(\mathbf{x}^{(k)})$   
    5:  $\bar{\mathbf{H}} \leftarrow \text{SPDPROJECTION}(\nabla^2 g(\mathbf{x}^{(k)}))$   
    6: SOLVE  $\bar{\mathbf{H}} \Delta \mathbf{x} = -\mathbf{J}$   
    7:  $\alpha \leftarrow \text{LINESEARCH}(\mathbf{x}^{(k)}, \Delta \mathbf{x}, \mathbf{J})$   
    8:  $\mathbf{x}^{(k+1)} \leftarrow \mathbf{x}^{(k)} + \alpha \Delta \mathbf{x}$   
  
  9:  $\mathbf{x}^+ \leftarrow \mathbf{x}^{(N)}$   
 10:  $\mathbf{v}^+ \leftarrow \frac{1}{h}(\mathbf{x}^+ - \mathbf{x}^-)$   
end function
```

| SPD-Aware Newton's Method → Projective Dynamics (PD)

At each iteration k we have to:

- $\bar{\mathbf{H}}^{(k)} = \text{SPDProjection}(\nabla^2 g(\mathbf{x}^{(k)}))$
- $\mathbf{C}^{(k)} = \text{Factorize}(\bar{\mathbf{H}}^{(k)})$
- $\Delta \mathbf{x} = \text{SolveLinearSystem}(\mathbf{C}^{(k)}, -\mathbf{J}^{(k)})$

What is Factorization:

Factorization rewrites the matrix as a triangular matrix times its transpose:

$$\bar{\mathbf{H}} = \mathbf{C} \mathbf{C}^\top, \quad \mathbf{C} \text{ lower triangular}$$

The system is then solved with two cheap triangular substitutions:

$$\mathbf{C} \mathbf{y} = -\mathbf{J} \longrightarrow \mathbf{C}^\top \Delta \mathbf{x} = \mathbf{y}$$

The **factorization** is the expensive part.

These are very expensive operations (no Real-Time)

Idea of PD: restrict the energies to a special form, so that the system matrix is **constant** and **SPD**, then **factorized once** and reused forever.

| Projective Dynamics (PD)

Each element energy is the squared distance between a linear function of the positions and a set of admissible configurations:

$$U_e(\mathbf{x}) = \min_{\mathbf{p}_e \in \mathcal{M}_e} \frac{k_e}{2} \left\| \mathbf{G}_e \mathbf{x} - \mathbf{p}_e \right\|^2$$

- $\mathbf{G}_e$: **constant** linear operator (selects and combines the element's particles)
- $\mathcal{M}_e$: **constraint manifold**, the set of undeformed (zero-energy) configurations
- $\mathbf{p}_e$: auxiliary variable, the **projection** of $\mathbf{G}_e \mathbf{x}$ onto $\mathcal{M}_e$

| Projective Dynamics (PD)


Augmented Energy:

$$g(\mathbf{x}, \mathbf{p}) = \frac{1}{2h^2} (\mathbf{x} - \tilde{\mathbf{x}})^\top \mathbf{M} (\mathbf{x} - \tilde{\mathbf{x}}) + \sum_e \frac{k_e}{2} \|\mathbf{G}_e \mathbf{x} - \mathbf{p}_e\|^2$$

Minimization Problem:

$$\min_{\mathbf{x}, \mathbf{p}_e \in \mathcal{M}_e} g(\mathbf{x}, \mathbf{p})$$

We now minimize over two sets of variables, positions and projections. Instead of solving for both together, we alternate: minimize over one set while keeping the other fixed. This is alternating minimization, it turns the problem into a two-step iterative solver in which each step never increases the energy



| Projective Dynamics (PD)

$$\min_{\mathbf{x}, \mathbf{p}_e \in \mathcal{M}_e} g(\mathbf{x}, \mathbf{p}) = \frac{1}{2h^2} (\mathbf{x} - \tilde{\mathbf{x}})^\top \mathbf{M} (\mathbf{x} - \tilde{\mathbf{x}}) + \sum_e \frac{k_e}{2} \|\mathbf{G}_e \mathbf{x} - \mathbf{p}_e\|^2$$

Local step: fix $\mathbf{x}$, minimize over each $\mathbf{p}_e$. Each is a small **independent projection**, done in **parallel**:

$$\min_{\mathbf{p}_e \in \mathcal{M}_e} g(\mathbf{x}, \mathbf{p}) \longrightarrow \mathbf{p}_e = \text{Proj}_{\mathcal{M}_e} (\mathbf{G}_e \mathbf{x}^{(k)})$$

Global step: fix $\mathbf{p}_e$, minimize over $\mathbf{x}$. The objective is quadratic, so we set its gradient to zero, which gives the linear system:

$$\underbrace{\left[\frac{1}{h^2} \mathbf{M} + \sum_e k_e \mathbf{G}_e^\top \mathbf{G}_e \right]}_{\mathbf{L}_{\text{PD}} : \text{is constant and SPD}} \mathbf{x}^{(k+1)} = \underbrace{\frac{1}{h^2} \mathbf{M} \tilde{\mathbf{x}} + \sum_e k_e \mathbf{G}_e^\top \mathbf{p}_e}_{\mathbf{b}}$$

| Projective Dynamics (PD)


for $k \in [0, 1, \dots, N - 1]$ **do**:

for each element e **do**:

Compute $\mathbf{G}_e \mathbf{x}^{(k)}$

$\mathbf{p}_e = \text{Proj}_{\mathcal{M}_e} (\mathbf{G}_e \mathbf{x}^{(k)})$

Solve $\mathbf{L}_{\text{PD}} \mathbf{x}^{(k+1)} = \mathbf{b}$



| Projective Dynamics (PD)


for $k \in [0, 1, \dots, N - 1]$ **do**:

for each element e **do**:

Compute $\mathbf{G}_e \mathbf{x}^{(k)}$

$\mathbf{p}_e = \text{Proj}_{\mathcal{M}_e} (\mathbf{G}_e \mathbf{x}^{(k)})$

Solve $\mathbf{L}_{\text{PD}} \mathbf{x}^{(k+1)} = \mathbf{b}$



| Projective Dynamics (PD)


for $k \in [0, 1, \dots, N - 1]$ **do**:

for each element e **do**:

Compute $\mathbf{G}_e \mathbf{x}^{(k)}$

$$\mathbf{p}_e = \text{Proj}_{\mathcal{M}_e} (\mathbf{G}_e \mathbf{x}^{(k)})$$

Solve $\mathbf{L}_{\text{PD}} \mathbf{x}^{(k+1)} = \mathbf{b}$



| Projective Dynamics (PD)


for $k \in [0, 1, \dots, N - 1]$ **do**:

for each element e **do**:

Compute $\mathbf{G}_e \mathbf{x}^{(k)}$

$$\mathbf{p}_e = \text{Proj}_{\mathcal{M}_e} (\mathbf{G}_e \mathbf{x}^{(k)})$$

Solve $\mathbf{L}_{\text{PD}} \mathbf{x}^{(k+1)} = \mathbf{b}$



| Projective Dynamics (PD)


for $k \in [0, 1, \dots, N - 1]$ **do**:

for each element e **do**:

Compute $\mathbf{G}_e \mathbf{x}^{(k)}$

$$\mathbf{p}_e = \text{Proj}_{\mathcal{M}_e} (\mathbf{G}_e \mathbf{x}^{(k)})$$

Solve $\mathbf{L}_{\text{PD}} \mathbf{x}^{(k+1)} = \mathbf{b}$



| Projective Dynamics (PD)


for $k \in [0, 1, \dots, N - 1]$ **do**:

for each element e **do**:

Compute $\mathbf{G}_e \mathbf{x}^{(k)}$

$\mathbf{p}_e = \text{Proj}_{\mathcal{M}_e} (\mathbf{G}_e \mathbf{x}^{(k)})$

Solve $\mathbf{L}_{\text{PD}} \mathbf{x}^{(k+1)} = \mathbf{b}$



| Projective Dynamics (PD)


for $k \in [0, 1, \dots, N - 1]$ **do**:

for each element e **do**:

Compute $\mathbf{G}_e \mathbf{x}^{(k)}$

$$\mathbf{p}_e = \text{Proj}_{\mathcal{M}_e} (\mathbf{G}_e \mathbf{x}^{(k)})$$

$$\text{Solve } \mathbf{L}_{\text{PD}} \mathbf{x}^{(k+1)} = \mathbf{b}$$



| Projective Dynamics (PD)


for $k \in [0, 1, \dots, N - 1]$ **do:**

for each element e **do:**

Compute $\mathbf{G}_e \mathbf{x}^{(k)}$

$\mathbf{p}_e = \text{Proj}_{\mathcal{M}_e} (\mathbf{G}_e \mathbf{x}^{(k)})$

Solve $\mathbf{L}_{\text{PD}} \mathbf{x}^{(k+1)} = \mathbf{b}$



| Projective Dynamics (PD)

function PROJECTIVEDYNAMICS($\mathbf{x}^-, \mathbf{v}^-$)

1: $\tilde{\mathbf{x}} \leftarrow \mathbf{x}^- + h \mathbf{v}^-$

2: Initialize $\mathbf{x}^{(0)}$

3: **for** $k = 0, 1, \dots, N - 1$ **do**

4: **for** each element e **in parallel do**

5: $\mathbf{p}_e \leftarrow \text{PROJ}_{\mathcal{M}_e}(\mathbf{G}_e \mathbf{x}^{(k)})$

6: $\mathbf{b} \leftarrow \frac{1}{h^2} \mathbf{M} \tilde{\mathbf{x}} + \sum_e k_e \mathbf{G}_e^\top \mathbf{p}_e$

7: **SOLVE** $\mathbf{L}_{\text{PD}} \mathbf{x}^{(k+1)} = \mathbf{b}$

8: $\mathbf{x}^+ \leftarrow \mathbf{x}^{(N)}$

9: $\mathbf{v}^+ \leftarrow \frac{1}{h}(\mathbf{x}^+ - \mathbf{x}^-)$

end function

Local step

Each element is projected onto its constraint set independently and in parallel.


Global step

Here, the linear solve is fast because the constant system matrix $\mathbf{L}_{\text{PD}}$ has already been factorized once, at initialization.

| Projective Dynamics (PD) Spring Constraint

Spring:

i, j : indices of particles l_{rest} : rest length k : stiffness



Projective Dynamics (PD) Spring Constraint


Spring:

i, j : indices of particles l_{rest} : rest length k : stiffness

Linear Operator:

$$\mathbf{G}_e \in \mathbb{R}^{3 \times 3n}$$

$$\mathbf{G}_e \mathbf{x} = \begin{bmatrix} \cdots & \overset{i}{\mathbf{I}} & \cdots & \overset{j}{-\mathbf{I}} & \cdots \end{bmatrix}^\top \begin{bmatrix} \vdots \\ \mathbf{x}_i \\ \vdots \\ \mathbf{x}_j \\ \vdots \end{bmatrix} = \mathbf{x}_i - \mathbf{x}_j = \mathbf{e}$$



Projective Dynamics (PD) Spring Constraint

Spring:

i, j : indices of particles l_{rest} : rest length k : stiffness


Linear Operator:

$$\mathbf{G}_e = \begin{bmatrix} \cdots & \overset{i}{\mathbf{I}} & \cdots & \overset{j}{-\mathbf{I}} & \cdots \end{bmatrix} \in \mathbb{R}^{3 \times 3n}$$

$$\mathbf{G}_e \mathbf{x} = \mathbf{x}_i - \mathbf{x}_j = \mathbf{e}$$

Constraint Manifold:

$$\mathcal{M}_e = \{ \mathbf{p} \in \mathbb{R}^3 : \|\mathbf{p}\| = l_{\text{rest}} \}$$



Projective Dynamics (PD) Spring Constraint

Spring:

i, j : indices of particles l_{rest} : rest length k : stiffness

Linear Operator:

$$\mathbf{G}_e = \begin{bmatrix} \cdots & \overset{i}{\mathbf{I}} & \cdots & \overset{j}{-\mathbf{I}} & \cdots \end{bmatrix} \in \mathbb{R}^{3 \times 3n}$$

$$\mathbf{G}_e \mathbf{x} = \mathbf{x}_i - \mathbf{x}_j = \mathbf{e}$$

Constraint Manifold:

$$\mathcal{M}_e = \left\{ \mathbf{p} \in \mathbb{R}^3 : \|\mathbf{p}\| = l_{\text{rest}} \right\}$$


Local Step (Projection):

$$\mathbf{p}_e = l_{\text{rest}} \frac{\mathbf{G}_e \mathbf{x}}{\|\mathbf{G}_e \mathbf{x}\|}$$

Global Step:

$$\mathbf{L}_{\text{PD}} += k_e \mathbf{G}_e^\top \mathbf{G}_e$$

$$\mathbf{b} += k_e \mathbf{G}_e^\top \mathbf{p}_e$$



Projective Dynamics (PD) Spring Constraint

Spring:

i, j : indices of particles l_{rest} : rest length k : stiffness

Linear Operator:

$$\mathbf{G}_e = \begin{bmatrix} & i & & j & \\ \cdots & \mathbf{I} & \cdots & -\mathbf{I} & \cdots \end{bmatrix} \in \mathbb{R}^{3 \times 3n}$$

$$\mathbf{G}_e \mathbf{x} = \mathbf{x}_i - \mathbf{x}_j = \mathbf{e}$$

Constraint Manifold:

$$\mathcal{M}_e = \{ \mathbf{p} \in \mathbb{R}^3 : \|\mathbf{p}\| = l_{\text{rest}} \}$$


Local Step (Projection):

$$\mathbf{p}_e = l_{\text{rest}} \frac{\mathbf{G}_e \mathbf{x}}{\|\mathbf{G}_e \mathbf{x}\|}$$

Global Step:

$$\mathbf{L}_{\text{PD}} += k_e \mathbf{G}_e^\top \mathbf{G}_e$$

$$\mathbf{b} += k_e \mathbf{G}_e^\top \mathbf{p}_e$$



$$\begin{matrix} & i & & j & \\ & \vdots & & \vdots & \\ i & \cdots & \mathbf{I} & \cdots & -\mathbf{I} & \cdots \\ & \vdots & & \vdots & & \\ j & \cdots & -\mathbf{I} & \cdots & \mathbf{I} & \cdots \\ & \vdots & & \vdots & & \end{matrix} \in \mathbb{R}^{3n \times 3n}$$

$$\begin{bmatrix} \cdots & \mathbf{p}_e & \cdots & -\mathbf{p}_e & \cdots \end{bmatrix}^\top \in \mathbb{R}^{3n}$$

| Differentiating Through Variational Implicit Euler Solver

The loss is measured on the simulated trajectory. We know the gradient of the loss after a step and want it before the step:

$$\text{known: } \frac{\partial \phi}{\partial \mathbf{x}^+} \longrightarrow \text{wanted: } \frac{\partial \phi}{\partial \mathbf{x}^-}$$

Repeating this step by step, from the last to first frame, gives the gradient with respect to the initial state:

$$\frac{\partial \phi}{\partial \mathbf{x}_T} \xrightarrow{\text{step } T} \frac{\partial \phi}{\partial \mathbf{x}_{T-1}} \xrightarrow{\text{step } T-1} \frac{\partial \phi}{\partial \mathbf{x}_{T-2}} \longrightarrow \dots \longrightarrow \frac{\partial \phi}{\partial \mathbf{x}_0}$$

The building block we would like to compute then is:

$$\frac{\partial \mathbf{x}^+}{\partial \mathbf{x}^-} \xrightarrow{\text{which through chain rule gives us:}} \frac{\partial \phi}{\partial \mathbf{x}^-} = \frac{\partial \phi}{\partial \mathbf{x}^+} \frac{\partial \mathbf{x}^+}{\partial \mathbf{x}^-}$$

| Differentiating Through Variational Implicit Euler Solver

Energy and its Derivatives:


$$g(\mathbf{x}) = \frac{1}{2h^2} (\mathbf{x} - \tilde{\mathbf{x}})^\top \mathbf{M} (\mathbf{x} - \tilde{\mathbf{x}}) + U(\mathbf{x})$$

$$\mathbf{J} = \nabla g(\mathbf{x}) = \frac{1}{h^2} \mathbf{M} (\mathbf{x} - \tilde{\mathbf{x}}) + \nabla U(\mathbf{x})$$

$$\mathbf{H} = \nabla^2 g(\mathbf{x}) = \frac{1}{h^2} \mathbf{M} + \nabla^2 U(\mathbf{x})$$

At the end of the step the converged solution $\mathbf{x}^+$ satisfy the **optimality condition**:

$$\nabla g(\mathbf{x}^+) = \mathbf{0}$$



This holds at convergence whatever solver was used (Newton, PD, ...)

We differentiate this condition (**implicit differentiation**), not the solver's iterations.

| Differentiating Through Variational Implicit Euler Solver

Energy and its Derivatives:

$$g(\mathbf{x}) = \frac{1}{2h^2} (\mathbf{x} - \tilde{\mathbf{x}})^\top \mathbf{M} (\mathbf{x} - \tilde{\mathbf{x}}) + U(\mathbf{x})$$

$$\mathbf{J} = \nabla g(\mathbf{x}) = \frac{1}{h^2} \mathbf{M} (\mathbf{x} - \tilde{\mathbf{x}}) + \nabla U(\mathbf{x})$$

$$\mathbf{H} = \nabla^2 g(\mathbf{x}) = \frac{1}{h^2} \mathbf{M} + \nabla^2 U(\mathbf{x})$$

Differentiating the optimality condition with respect to $\mathbf{x}^-$ gives:

$$\underbrace{\left[\frac{1}{h^2} \mathbf{M} + \nabla^2 U(\mathbf{x}^+) \right]}_{\mathbf{H} = \nabla^2 g(\mathbf{x}^+)} \frac{\partial \mathbf{x}^+}{\partial \mathbf{x}^-} - \frac{1}{h^2} \mathbf{M} = \mathbf{0} \quad \implies \quad \frac{\partial \mathbf{x}^+}{\partial \mathbf{x}^-} = \frac{1}{h^2} \mathbf{H}^{-1} \mathbf{M}$$

Then the chain rule gives:

$$\frac{\partial \phi}{\partial \mathbf{x}^-} = \frac{\partial \phi}{\partial \mathbf{x}^+} \frac{\partial \mathbf{x}^+}{\partial \mathbf{x}^-} = \frac{1}{h^2} \frac{\partial \phi}{\partial \mathbf{x}^+} \mathbf{H}^{-1} \mathbf{M}$$

↓
computed once at convergence

| Differentiating Through Variational Implicit Euler Solver

$$\frac{\partial \phi}{\partial \mathbf{x}^-} = \frac{1}{h^2} \underbrace{\frac{\partial \phi}{\partial \mathbf{x}^+} \mathbf{H}^{-1}}_{\mathbf{z}^\top} \mathbf{M}$$

We never compute the dense matrix $\mathbf{H}^{-1}$. Instead, we group the first two factors into the **adjoint vector** $\mathbf{z}$, which comes from a single linear solve:

$$\mathbf{z}^\top = \frac{\partial \phi}{\partial \mathbf{x}^+} \mathbf{H}^{-1} \iff \mathbf{H} \mathbf{z} = \left(\frac{\partial \phi}{\partial \mathbf{x}^+} \right)^\top \longrightarrow \text{Single Linear System Solve}$$

Then both positions and velocities gradients are cheap matrix–vector products:

$$\frac{\partial \phi}{\partial \mathbf{x}^-} = \frac{1}{h^2} \mathbf{z}^\top \mathbf{M} \qquad \frac{\partial \phi}{\partial \mathbf{v}^-} = \frac{1}{h} \mathbf{z}^\top \mathbf{M}$$