

Tikhonov Regularisation for (Large) Inverse Problems

Melina Freitag

Department of Mathematical Sciences
University of Bath

17th ILAS Conference
Braunschweig, Germany
23rd August 2011

joint work with C.J. Budd (Bath) and N.K. Nichols (Reading)



Inverse Problems

Data Assimilation as a Large Inverse Problem

Regularisation Parameter estimation in 4DVar

Regularisation Parameter estimation

Example

Application of L_1 -norm regularisation in 4DVar

Motivation: Results from image processing

L_1 -norm regularisation in 4DVar

Examples

Ill-posed Problems

Given an operator A we wish to solve

$$Af = g.$$

It is **well-posed** if

- solution exists
- solution is unique
- is stable (A^{-1} continuous)

but ..

In finite dimensions existence and uniqueness can be imposed, but

- discrete problem of underlying ill-posed problem becomes **ill-conditioned**
- **singular values of A decay to zero**

An Illustrative Example

Fredholm first kind integral equation in 1D

$$g(x) = \int_0^1 k(x-x')f(x')dx' =: (Af)(x), \quad 0 < x < 1$$

- f light source intensity as a function of x
- g image intensity
- k kernel representing blurring effects, e.g. $k(x) = C \exp\left(-\frac{x^2}{2\gamma^2}\right)$, C, γ are positive parameters.

Discretisation

- use a piecewise smooth source \mathbf{f}
- determine \mathbf{A} using standard numerical quadrature;

$$(\mathbf{A})_{ij} = hC \exp\left(-\frac{((i-j)h)^2}{2\gamma^2}\right), \quad 1 \leq i, j \leq n, \quad h = \frac{1}{n}$$

$$\gamma = 0.05, \quad C = \frac{1}{\gamma\sqrt{2\pi}}.$$

Inverse Problem

Given the kernel k , and the blurred image g , determine the source f from $g = Af$, solve the discrete linear system

$$\mathbf{g} = \mathbf{A}\mathbf{f}.$$




Figure: True solution and blurred image \mathbf{g}

Inverse Problem

Given the kernel k , and the blurred image g , determine the source f from $g = Af$, solve the discrete linear system

$$\mathbf{g} = \mathbf{A}\mathbf{f}.$$




Figure: Naive Solution

Inverse Problem

Problem: data \mathbf{g} are observed and contain **noise** and \mathbf{A} is **ill-conditioned**:

$$\mathbf{g}_{\text{exact}} + \mathbf{e} = \mathbf{A}\mathbf{f},$$

\mathbf{e} is unknown white noise.




Figure: True solution and discrete noisy data

Inverse Problem

Problem: data \mathbf{g} are observed and contain **noise** and \mathbf{A} is **ill-conditioned**:

$$\mathbf{g}_{\text{exact}} + \mathbf{e} = \mathbf{A}\mathbf{f},$$

\mathbf{e} is unknown white noise.

Singular Value Decomposition

Let

$$\mathbf{A} = \mathbf{U}\mathbf{\Sigma}\mathbf{V}^T = \sum_{i=1}^r \sigma_i \mathbf{u}_i \mathbf{v}_i^T$$

where

- $\mathbf{\Sigma} = \text{diag}(\sigma_1, \dots, \sigma_r)$ and $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_r > 0$
- $\mathbf{U}^T \mathbf{U} = \mathbf{I}$ and $\mathbf{V}^T \mathbf{V} = \mathbf{I}$


Inverse Problem - Regularisation needed

Least squares solution (with and without noise)

$$\mathbf{f}_{\text{exact}} = \mathbf{A}^\dagger \mathbf{g}_{\text{exact}} = \sum_{i=1}^r \frac{\mathbf{u}_i^T \mathbf{g}_{\text{exact}}}{\sigma_i} \mathbf{v}_i$$

$$\mathbf{f} = \mathbf{A}^\dagger \mathbf{g} = \mathbf{A}^\dagger (\mathbf{g}_{\text{exact}} + \mathbf{e}) = \sum_{i=1}^r \frac{\mathbf{u}_i^T \mathbf{g}_{\text{exact}}}{\sigma_i} \mathbf{v}_i + \sum_{i=1}^r \frac{\mathbf{u}_i^T \mathbf{e}}{\sigma_i} \mathbf{v}_i$$

$$= \mathbf{f}_{\text{exact}} + \sum_{i=1}^r \frac{\mathbf{u}_i^T \mathbf{e}}{\sigma_i} \mathbf{v}_i$$



Tikhonov Regularisation

Regularised solution of the form

$$\mathbf{f}_\alpha = \sum_{i=1}^r \frac{\sigma_i^2}{\sigma_i^2 + \alpha^2} \frac{\mathbf{u}_i^T \mathbf{g}}{\sigma_i} \mathbf{v}_i$$

α regularisation parameter.

Solution \mathbf{f}_α to the minimisation problem

$$\min_{\mathbf{f}} \{ \|\mathbf{g} - \mathbf{A}\mathbf{f}\|_2^2 + \alpha^2 \|\mathbf{f}\|_2^2 \}.$$

Least squares solution \mathbf{f}_α to the linear system

$$\begin{bmatrix} \mathbf{A} \\ \alpha \mathbf{I} \end{bmatrix} \mathbf{f} = \begin{bmatrix} \mathbf{g} \\ \mathbf{0} \end{bmatrix}.$$

Normal equations

$$(\mathbf{A}^T \mathbf{A} + \alpha^2 \mathbf{I}) \mathbf{f}_\alpha = \mathbf{A}^T \mathbf{g}.$$

Tikhonov Regularisation

Regularisation parameter α

Regularised solution of the form

$$\mathbf{f}_\alpha = \sum_{i=1}^r \frac{\sigma_i^2}{\sigma_i^2 + \alpha^2} \frac{\mathbf{u}_i^T \mathbf{g}}{\sigma_i} \mathbf{v}_i$$

Filter factor $\frac{\sigma_i^2}{\sigma_i^2 + \alpha^2}$ as diagonal entries of the filter matrix Ψ

Regularisation and perturbation error

$$\begin{aligned} \mathbf{f}_\alpha &= \mathbf{V}\Psi\mathbf{U}^T \mathbf{g}, \quad \mathbf{g} = \mathbf{g}_{\text{exact}} + \mathbf{e} \\ &= \mathbf{V}\Psi\Sigma^{-1}\mathbf{U}^T \mathbf{g}_{\text{exact}} + \mathbf{V}\Psi\Sigma^{-1}\mathbf{U}^T \mathbf{e} \\ &= \mathbf{V}\Psi\Sigma^{-1}\mathbf{U}^T \mathbf{U}\Sigma\mathbf{V}^T \mathbf{f}_{\text{exact}} + \mathbf{V}\Psi\Sigma^{-1}\mathbf{U}^T \mathbf{e} \\ &= \mathbf{V}\Psi\mathbf{V}^T \mathbf{f}_{\text{exact}} + \mathbf{V}\Psi\Sigma^{-1}\mathbf{U}^T \mathbf{e} \\ \mathbf{f}_{\text{exact}} - \mathbf{f}_\alpha &= (\mathbf{I} - \mathbf{V}\Psi\mathbf{V}^T) \mathbf{f}_{\text{exact}} - \mathbf{V}\Psi\Sigma^{-1}\mathbf{U}^T \mathbf{e} \end{aligned}$$

Regularisation error

Perturbation error

Tikhonov Regularisation

Regularisation and perturbation error

$$\mathbf{f}_{\text{exact}} - \mathbf{f}_\alpha = (\mathbf{I} - \mathbf{V}\Psi\mathbf{V}^T) \mathbf{f}_{\text{exact}} - \mathbf{V}\Psi\Sigma^{-1}\mathbf{U}^T \mathbf{e}$$




Figure: Regularisation and perturbation error

Tikhonov Regularisation

Illustrative example




Figure: α too small

Illustrative example




Figure: α too large

Illustrative example




Figure: Good Value for α