

Introduction to Bayesian analysis for medical studies

Part III: Bayesian applications in biomedical sciences

Boris Hejblum

Université de Bordeaux, ISPED, Inserm BPH U1219/Inria SISTM, Bordeaux, France

boris.hejblum@u-bordeaux.fr

<https://borishejblum.science>

Graduate School of Health and Medical Sciences
at the University of Copenhagen

May 14th, 2019

Introduction

Examples of Bayesian applications

3 different real-world use cases in biomedical sciences:
illustrations where the Bayesian approach can be particularly useful

⚠ **Disclaimer:** this course is NOT

- ~~a meta-analysis short course~~
- ~~an adaptive design in clinical trials short course~~

Post-mortem re-analysis
of an under-powered randomized trial

Original analysis of EOLIA

EOLIA (Combes *et al.*, NEJM, 2018):

- randomized clinical trial
- evaluation of a new treatment for severe acute respiratory distress syndrome
- outcome: mortality rate after 60 days
- 249 patients:
 - 125 controls
⇒ mechanical ventilation (conventional treatment)
 - 124 treated
⇒ ECMO (extracorporeal membrane oxygenation – new(er) treatment)

Frequentist analysis:

⇒ Relative Risk of death at 60 days for ECMO compared to control: 0.76

$CI_{95\%} = [0.55, 1.04]$

p-value = 0.09.

Bayesian re-analysis of EOLIA data

Goligher *et al.* (*JAMA*, 2018)

	Group	
	ECMO	Control
group size n	124	125
number of deaths at 60 days	44	57

Observed data in the EOLIA trial

Your turn !



Practical: exercise 5

What is a meta-analysis

“An analysis of analyses”

⇒ a single quantitative summary of studies answering the *same research question*

Ex: medical therapies effects are often evaluated in multiple different studies.

What is a meta-analysis

“An analysis of analyses”

⇒ a single quantitative summary of studies answering the *same research question*

Ex: medical therapies effects are often evaluated in multiple different studies.

⇒ pool individual observations from multiple studies ?

What is a meta-analysis

“An analysis of analyses”

⇒ a single quantitative summary of studies answering the *same research question*

Ex: medical therapies effects are often evaluated in multiple different studies.

⇒ pool individual observations from multiple studies ?

- ⚠ potential differences in the pooled experiments
- ⚠ only aggregated summary statistics estimates (“effect sizes”) available
 - alongside uncertainty (e.g. standard errors)

Study Heterogeneity

⚠ variations of the observed effects. . .

Study Heterogeneity

⚠ variations of the observed effects. . .

- within-study uncertainty, or
- real heterogeneity in effect size between the different studies

Study Heterogeneity

⚠ variations of the observed effects. . .

- within-study uncertainty, or
- real heterogeneity in effect size between the different studies

Often, different studies used different populations

⇒ potential extra-variability

+ different sample sizes ⇒ also impact the estimate and its variability

Meta-analysis random effects model

Common approach for meta analysis:

$$y_i \sim \mathcal{N}(\theta_i, \sigma_i^2)$$

$$\theta_i \sim \mathcal{N}(\mu, \tau^2)$$

Meta-analysis random effects model

Common approach for meta analysis:

$$y_i \sim \mathcal{N}(\theta_i, \sigma_i^2)$$

$$\theta_i \sim \mathcal{N}(\mu, \tau^2)$$

Hierarchical generalization of the fixed effect model:

$$y_i \sim \mathcal{N}(\mu, \sigma_i^2)$$

Meta-analysis random effects model

Common approach for meta analysis:

$$y_i \sim \mathcal{N}(\theta_i, \sigma_i^2)$$

$$\theta_i \sim \mathcal{N}(\mu, \tau^2)$$

⇒ between study variability: $y_i \sim \mathcal{N}(\theta, \sigma_i^2 + \tau)$

Hierarchical generalization of the fixed effect model:

$$y_i \sim \mathcal{N}(\mu, \sigma_i^2)$$

⇒ assume same average effect for each study

Bayesian meta-analysis in practice

Meta-analysis: a perfect usecase for Bayesian analysis ?

Bayesian meta-analysis in practice

Meta-analysis: a perfect usecase for Bayesian analysis ?

- few observations
- informative *prior*
- sequential

Going further

Scientific literature search

⚠ FIRST (!) exhaustive search of the scientific literature

Going further

Scientific literature search

⚠ FIRST (!) exhaustive search of the scientific literature: hard !!!

⚠ effect size estimate () along with their standard errors) must often be **transformed before** the meta-analysis

Going further

Scientific literature search

⚠ FIRST (!) exhaustive search of the scientific literature: hard !!!

⚠ effect size estimate () along with their standard errors) must often be **transformed before** the meta-analysis

Evidence synthesis

Meta-analysis \in evidence synthesis

e.g. meta-regression, mechanistic modeling, ...

Going further

Scientific literature search

⚠ FIRST (!) exhaustive search of the scientific literature: hard !!!

⚠ effect size estimate () along with their standard errors) must often be **transformed before** the meta-analysis

Evidence synthesis

Meta-analysis \in evidence synthesis

e.g. meta-regression, mechanistic modeling, ...

Still active research domains:

- random effects model will down-weight studies with larger sample sizes
 - Serghiou & Goodman, *JAMA*, 2018

Going further

Scientific literature search

⚠ FIRST (!) exhaustive search of the scientific literature: hard !!!

⚠ effect size estimate () along with their standard errors) must often be **transformed before** the meta-analysis

Evidence synthesis

Meta-analysis \in evidence synthesis

e.g. meta-regression, mechanistic modeling, ...

Still active research domains:

- random effects model will down-weight studies with larger sample sizes
 - Serghiou & Goodman, *JAMA*, 2018
 - a bug or a feature ?*

Your turn !



Practical: exercise 6

Continuous Reassessment method

CRM [O'Quigley *at al.*, 1990]

Objective: identify the optimal dose

(i.e. Minimum Efficient Dose or Maximum Tolerated Dose)

⇒ select iteratively the dose for the next (batch of) recruited patient(s)
based accumulating observations from previously included patients

Continuous Reassessment method

CRM [O'Quigley *at al.*, 1990]

Objective: identify the optimal dose

(i.e. Minimum Efficient Dose or Maximum Tolerated Dose)

⇒ select iteratively the dose for the next (batch of) recruited patient(s)
based accumulating observations from previously included patients

- 😊 treat each patient ethically (dose best supported by the current evidence)
- 😊 *prior* knowledge
- 😊 sequential Bayesian: online update of the posterior

increasingly used (but still minority. . .)

Your turn !



Practical: exercise 7