

Connection of Vacuum Energy with Inflation: The Elastic Potential, Initial Conditions, and the Origin of Primordial Fluctuations

Boris Kriger^{1,2}

¹ Information Physics Institute, Gosport, Hampshire, United Kingdom
boris.kriger@informationphysicsinstitute.net

² Institute of Integrative and Interdisciplinary Research, Toronto, Canada
boriskriger@interdisciplinary-institute.org

ORCID: [0009-0001-0034-2903](https://orcid.org/0009-0001-0034-2903)

Abstract

The elastic vacuum framework—developed in a companion paper as a dynamical sequestering mechanism for vacuum energy—implies that spacetime possesses intrinsic elasticity established by the Einstein–Hilbert action, confirmed by LIGO, and derivable from Sakharov’s induced gravity programme. This paper develops three consequences for inflationary cosmology. First, we argue on quantum-mechanical, thermodynamic, and structural grounds that the initial state of the universe cannot be exactly homogeneous: differentiation is a precondition for any describable physical state, and primordial inhomogeneities are ontologically prior to the homogeneous background, not perturbations upon it. Second, we derive the scalar field potential $V(\psi)$ from the elastic properties of spacetime: the Einstein–Hilbert action supplemented by the leading quantum correction (R^2) yields, upon conformal transformation, the Starobinsky potential $V(\psi) = V_0(1 - e^{-\sqrt{2/3}\psi/M_{\text{Pl}}})^2$. This potential is not an arbitrary choice but the leading-order elastic potential of the vacuum within the R^2 truncation of the effective action—quadratic near equilibrium, saturating at large deformation. We show that it reproduces the Planck-favored values $n_s \approx 0.964$ and $r \approx 0.004$ and reinterpret inflation as the elastic recoil of spacetime from extreme initial deformation. Third, we derive the flatness feedback parameter $\beta > 0$ from the convexity of $V(\psi)$, completing the proof that spatial flatness is a late-time attractor of the elastic vacuum. The framework uses only general relativity with standard quantum corrections, introducing no new matter particles beyond the screening scalar field and requiring no fine-tuning. A three-dimensional elastic analogy— injection of a substance into a transparent elastic medium—replaces and extends the membrane model of the companion paper, resolving its principal limitations.

Keywords: vacuum energy, cosmic inflation, elastic vacuum, Starobinsky potential, primordial fluctuations, flatness problem, initial conditions, cosmological constant problem, R^2 gravity

Contents

1	Introduction	4
2	Logical Structure of the Argument	5
3	The Inhomogeneous Origin: Why Initial Conditions Cannot Be Homogeneous	6
3.1	The Quantum-Mechanical Argument	6
3.2	The Thermodynamic Argument	7
3.3	The Structural Argument: A Deeper Perspective	7
4	The Three-Dimensional Elastic Analogy	8
4.1	The Model	9
4.2	What Happens After Injection	9
4.3	Advantages Over the Membrane Model	9
4.4	Formal Correspondence	10
5	The Elastic Potential: Deriving $V(\psi)$ from Spacetime Elasticity	11
5.1	Linear Elasticity: The Einstein–Hilbert Action	11
5.2	Nonlinear Elasticity: Quantum Corrections	11
5.3	Conformal Transformation: The Starobinsky Potential	12
5.4	Physical Interpretation: The Elastic Potential	13
6	Modified Inflationary Dynamics in the Elastic Vacuum	15
6.1	The Slow-Roll Equations	15
6.2	Number of e-Folds	15
6.3	Observational Predictions	16
6.4	Physical Interpretation: Inflation as Elastic Recoil	16
6.5	The Mass Hierarchy: What Is Unified and What Is Not	17
7	Primordial Fluctuations from Vacuum Inhomogeneities	19
7.1	Standard Mechanism: Quantum Fluctuations of ψ	19
7.2	The New Element: Primordial Inhomogeneities	19
7.3	Quantitative Estimate	20
7.4	Observable Consequences	21
8	The Flatness Feedback: Deriving β from $V(\psi)$	22
8.1	The Scalar Field in Curved FLRW Spacetime	22
8.2	Quasi-Static Approximation	22
8.3	Scalar Field Energy Density	22
8.4	Derivation of β	23
8.5	Epoch Dependence	24

9	Observational Consistency and Predictions	25
9.1	Inherited Predictions: The n_s - r Plane	25
9.2	Inherited Predictions: The Equation of State	25
9.3	New Predictions: Large-Scale CMB Anomalies	26
9.4	New Predictions: Non-Gaussianity	26
9.5	Falsifiable Predictions	26
10	Discussion	27
11	Limitations	27
12	Future Directions	28
13	Position Within the Research Programme	28
14	Recommended Journal for Publication	29
	Conclusions	30
	Acknowledgments	30
	References	30
A	Computational Code	34
	Peer Reviews and Revision History	36

1 Introduction

The preceding paper in this programme [Kriger, 2025a] developed the elastic vacuum framework: a dynamical picture in which spacetime geometry is not a passive background but an elastic medium that resists deformation, relaxes toward flatness, and whose late-time elastic recovery drives the observed cosmic acceleration. Three results were established. First, a sequestering mechanism separates the cosmological constant Λ from the quantum field theory vacuum energy ρ_{vac} , preserving the equivalence principle locally while screening vacuum energy from cosmological expansion. Second, a scalar field ψ with a horizon-sensitive mass $m_\psi \sim H_0$ produces scale-dependent gravitational screening. Third, a positive feedback loop—decreasing curvature facilitates faster expansion, which further decreases curvature—drives the universe toward spatial flatness.

Three open questions were explicitly identified and deferred to the present paper:

- (i) What is the specific form of the scalar field potential $V(\psi)$?
- (ii) Can the feedback parameter $\beta > 0$ be derived from the action rather than postulated?
- (iii) How does the elastic vacuum framework connect to inflationary cosmology?

This paper answers all three. We show that $V(\psi)$ is not a free function to be chosen for phenomenological convenience: it is the elastic potential of spacetime, determined by the same physics—the Einstein–Hilbert action and its quantum corrections—that establishes the elasticity of the vacuum. The result is the Starobinsky potential [Starobinsky, 1980], currently the inflationary model most favored by Planck data [Planck Collaboration, 2020]. This identification is not a coincidence. Starobinsky’s R^2 inflation was derived from quantum corrections to gravity—precisely the higher-order elastic terms in the effective action. Our contribution is to recognize this derivation as the nonlinear elastic response of the vacuum: just as a rubber band has a yield stress beyond which further stretching costs less energy per unit strain, spacetime has an elastic limit beyond which further curvature costs less energy per unit curvature. The inflationary plateau is the yield plateau of spacetime elasticity.

We also address a deeper question: the origin of primordial fluctuations and the problem of initial conditions. Standard inflation assumes that the inflaton field begins in a nearly homogeneous state atop a flat potential, and that quantum fluctuations of this field generate the seeds of structure. But the homogeneous initial state is itself a fine-tuned assumption—one that inflation was supposed to render unnecessary. We argue, primarily on quantum-mechanical grounds and with support from thermodynamic and structural considerations, that exact homogeneity is not merely unlikely but physically unrealizable: the Heisenberg uncertainty principle forbids any finite-energy field configuration from having zero spatial variance. Primordial inhomogeneities are not perturbations imposed on a homogeneous background; they are constitutive of the initial state itself. Inflation—understood as elastic recoil—then smooths these primordial inhomogeneities to the observed level, producing the nearly scale-invariant spectrum without requiring the inflaton to start in a special state.

The paper is organized as follows. Section 2 maps the logical structure of the argument. Section 3 establishes the impossibility of exactly homogeneous initial conditions. Section 4

develops the three-dimensional elastic analogy. Section 5 derives $V(\psi)$ from spacetime elasticity. Section 6 analyzes the modified inflationary dynamics. Section 7 addresses primordial fluctuations. Section 8 derives $\beta > 0$ from $V(\psi)$. Section 9 identifies observational predictions. Sections 10–12 discuss implications, limitations, and future directions.

2 Logical Structure of the Argument

The reasoning proceeds through seven steps, each building on the previous.

Step 1 (§3): The initial state cannot be homogeneous. The Heisenberg uncertainty principle forbids any quantum field from occupying an exactly homogeneous state in a finite-energy configuration. This quantum-mechanical result is reinforced by a thermodynamic argument (zero-entropy initial states are statistically unrealizable) and a structural observation (exact homogeneity contains no internal distinctions and cannot serve as an initial condition for structure formation).

Step 2 (§4): The three-dimensional elastic analogy. We extend the two-dimensional membrane model of Paper 5 to a three-dimensional elastic medium into which energy–matter is injected. This resolves the principal limitations of the membrane analogy (no preferred direction, no embedding space) while preserving its key dynamical features: active geometry, mutual feedback, and local/global asymmetry.

Step 3 (§5): Derivation of $V(\psi)$ from elastic physics. The Einstein–Hilbert action, supplemented by the leading quantum correction (R^2), defines the elastic energy of spacetime. Conformal transformation to the Einstein frame yields the Starobinsky potential $V(\psi) = V_0(1 - e^{-\sqrt{2/3}\psi/M_{\text{Pl}}})^2$. This is the leading-order elastic potential within the R^2 truncation: quadratic near equilibrium, saturating at large deformation.

Step 4 (§6): Inflationary dynamics as elastic recoil. With $V(\psi)$ specified, the slow-roll equations are solved. The predictions— $n_s \approx 1 - 2/N_*$ and $r \approx 12/N_*^2$ —match Planck observations. Inflation is reinterpreted as the elastic recoil of spacetime from extreme initial deformation. The 10^{55} mass hierarchy between the inflationary and late-time sectors is identified as the programme’s deepest unsolved problem.

Step 5 (§7): Primordial fluctuations from elastic physics. The necessarily inhomogeneous initial state (Step 1) provides the seed perturbations. The elastic recoil (Step 4) stretches and smooths them. A quantitative estimate shows that for $N_{\text{tot}} \gtrsim 69$ e-folds, even Planck-amplitude initial perturbations are subdominant on CMB scales.

Step 6 (§8): Derivation of $\beta > 0$. In the quasi-static limit, the convexity of $V(\psi)$ at its minimum—guaranteed by the elastic potential structure—implies that the scalar field back-reaction on the Friedmann equations reduces the effective spatial curvature. The feedback parameter β is computed explicitly and shown to be positive, completing the conditional flatness attractor proof of Paper 5. Full dynamical confirmation is deferred to numerical work.

Step 7 (§9): Observational predictions. The framework makes specific predictions in the n_s – r plane, for the equation of state $w(a)$, for non-Gaussianity, and for the ISW effect. The Starobinsky model’s predictions are inherited and reinterpreted as consequences of spacetime elasticity.

Synthesis. The seven steps converge on a unified picture: inflation is not driven by a hypothetical scalar field with an ad hoc potential, but by the elastic recoil of spacetime—whose elastic properties are established by the Einstein–Hilbert action and its quantum corrections. The initial conditions need not be fine-tuned because exact homogeneity is impossible; what inflation produces is the observed *approximate* homogeneity, starting from any physically realizable initial state.

3 The Inhomogeneous Origin: Why Initial Conditions Cannot Be Homogeneous

Standard inflationary cosmology assumes that the inflaton field ϕ begins in a nearly homogeneous configuration on the plateau of its potential $V(\phi)$, with quantum fluctuations providing the small seed perturbations that grow into the observed large-scale structure [Mukhanov & Chibisov, 1981, Hawking, 1982, Guth, 1981, Linde, 1982]. The success of this picture is remarkable: it predicts a nearly scale-invariant, nearly Gaussian, nearly adiabatic spectrum of curvature perturbations, in excellent agreement with CMB observations [Planck Collaboration, 2020].

However, the assumption of initial homogeneity is itself a fine-tuning problem. Why should the inflaton start in a homogeneous state over a region much larger than the Hubble radius? This is precisely the kind of special initial condition that inflation was designed to explain away [Penrose, 1989, Brandenberger, 2017]. We now present three independent arguments that exact homogeneity is not merely unlikely but physically impossible.

3.1 The Quantum-Mechanical Argument

The Heisenberg uncertainty principle forbids any quantum field from occupying a state with exactly zero variance in both field amplitude and conjugate momentum. For a scalar field ϕ in a finite region of volume V_R , the spatially averaged field $\bar{\phi} = V_R^{-1} \int_{V_R} \phi(\mathbf{x}) d^3x$ and its conjugate $\bar{\pi}$ satisfy

$$\Delta\bar{\phi} \Delta\bar{\pi} \geq \frac{1}{2}. \quad (1)$$

A perfectly homogeneous field configuration corresponds to $\Delta\bar{\phi} = 0$ for all spatial averages—a delta-function state in field space, which has infinite momentum uncertainty and infinite energy. Such a state is not normalizable and does not belong to the physical Hilbert space.

More concretely, in any state with finite energy, the field exhibits vacuum fluctuations with variance [Birrell & Davies, 1982]

$$\langle\phi^2(\mathbf{x})\rangle = \int \frac{d^3k}{(2\pi)^3} \frac{1}{2\omega_k}, \quad (2)$$

where $\omega_k = \sqrt{k^2 + m^2}$. This integral diverges, requiring regularization, but the key point is structural: the variance is nonzero for any finite-energy state. The quantum vacuum is irreducibly inhomogeneous.

Counterarguments and Responses. *Objection: Coherent states can have arbitrarily small spatial fluctuations at the cost of large momentum fluctuations. Isn't a coherent state effectively homogeneous?*

Response: A coherent state $|\alpha\rangle$ of the zero mode is homogeneous in expectation, $\langle\phi(\mathbf{x})\rangle = \bar{\phi}_0$, but it still contains nonzero fluctuations $\Delta\phi \neq 0$ in every finite region. The fluctuations are suppressed by $V_R^{-1/2}$ for large regions but never vanish. Moreover, the higher (nonzero) modes are in their vacuum state, contributing the full spectrum of vacuum fluctuations. A coherent state is the *closest* a quantum field can come to homogeneity—and it is not homogeneous.

3.2 The Thermodynamic Argument

A perfectly homogeneous state is a state of zero configurational entropy. The Boltzmann entropy of a field configuration is $S = k_B \ln \Omega$, where Ω is the number of microstates consistent with the macroscopic description. A single, exactly specified homogeneous configuration has $\Omega = 1$, hence $S = 0$.

The second law of thermodynamics requires that the entropy of any closed system be non-decreasing. A universe that begins in a zero-entropy state and evolves to the high-entropy state we observe today is thermodynamically consistent but statistically extraordinary. Penrose has estimated the probability of such an initial state as $\exp(-10^{123})$ [Penrose, 1989]—a number so small that it renders the homogeneous initial condition effectively impossible within any statistical framework.

More importantly, a zero-entropy initial state contains no information. It specifies the macrostate completely but contains no internal structure—no gradients, no boundaries, no distinctions. As we argue in the next subsection, such a state cannot serve as the initial condition for a system that is to develop structure.

3.3 The Structural Argument: A Deeper Perspective

The quantum-mechanical and thermodynamic arguments above are sufficient to establish the physical impossibility of exact homogeneity. We note, however, that a deeper structural argument—prior to and independent of any specific physics—reaches the same conclusion.

Remark 3.1 (Structural necessity of differentiation). Any describable physical state must contain at least two distinguishable sub-states. A perfectly homogeneous state—one in which every spatial region is identical to every other—contains no internal distinctions. Such a state cannot serve as the initial condition for a system that develops structure, because structure presupposes distinctions, and a homogeneous state contains none.

This observation does not depend on quantum mechanics or thermodynamics. It is a precondition for description itself: if nothing can be distinguished from anything else, no proposition can be formulated about the system. The philosophical lineage extends through Spencer-Brown's *Laws of Form* [Spencer-Brown, 1969], where the act of drawing a distinction is the primitive operation from which logical structure arises, and through Bateson's definition of information as “a difference which makes a difference” [Bateson, 1972]. A systematic development of this argument, showing that differentiation is a necessary condition for any system capable of sustaining complex structure, is given in Kriger [2026].

The structural argument is not invoked as a premise of the physical derivations that follow—the quantum-mechanical argument (§3.1) provides the necessary physical grounding. It is recorded here because it illuminates the conceptual status of the “homogeneous initial state”: not as a physical possibility that happens to be improbable, but as a mathematical idealization that cannot correspond to any describable state of affairs.

Counterarguments and Responses. *Objection: The structural argument is philosophical, not physical. A physics paper should not rest on ontological claims.*

Response: We agree, and we do not rest the paper’s physical results on this argument. The quantum-mechanical argument of §3.1 is the operative premise: the Heisenberg uncertainty principle forbids exact homogeneity in any finite-energy state. The structural observation is recorded because it explains *why* the quantum-mechanical result should be expected on general grounds, not because the subsequent derivations depend on it.

Objection: The universe could begin homogeneous and develop inhomogeneity through dynamical evolution.

Response: This is precisely the standard inflationary picture, and it faces two problems. First, as argued in §3.1, quantum mechanics forbids exact homogeneity. Second, dynamical evolution requires equations of motion, which require a state to evolve *from*—and that state, being physical, must already contain distinctions at the quantum level. The “homogeneous” initial state is a mathematical idealization (the Bunch–Davies vacuum on exact de Sitter space) that approximates but does not coincide with any physical state.

Section Result and Implications

The quantum-mechanical argument establishes, as a consequence of the Heisenberg uncertainty principle, that the initial state of the universe cannot be exactly homogeneous: any finite-energy field configuration has nonzero spatial variance. The thermodynamic argument reinforces this conclusion from the perspective of statistical mechanics, and the structural observation illuminates the conceptual status of exact homogeneity as an idealization. Together, they transform the problem of initial conditions: inflation need not create homogeneity from a homogeneous state (a tautology) but must create *approximate* homogeneity from an *inherently inhomogeneous* initial state. The elastic recoil mechanism of the following sections provides precisely this.

4 The Three-Dimensional Elastic Analogy

Paper 5 developed the elastic membrane model—a two-dimensional rubber sheet with viscous cream—as a physical analogy for the elastic vacuum. The analogy captured four essential features: active geometry, mutual feedback between matter and curvature, local/global asymmetry of vacuum effects, and the tendency toward flatness. However, it suffered from well-documented limitations (Paper 5, §4.5): the membrane curves extrinsically into an ambient space that does not exist in general relativity; it requires an external “down” direction; and it is two-dimensional while spacetime is four-dimensional with Lorentzian signature.

We now extend the analogy to three dimensions, resolving the most severe of these limitations.

4.1 The Model

Consider a transparent, three-dimensional elastic medium—a block of silicone or ballistic gel—uniform, isotropic, and initially undeformed. Into the center of this block, a syringe injects a dense, viscous substance: a mixture representing the total energy content of the universe (matter, radiation, and vacuum energy).

This is our model. The elastic medium is spacetime geometry. The injected substance is the energy–matter content. The act of injection is the Big Bang.

4.2 What Happens After Injection

The injected substance expands outward from the injection point, driven by its own internal pressure. Simultaneously, the elastic medium resists the deformation and pushes back—the elastic restoring force. Two processes drive the expansion: the substance spreading under its own pressure, and the medium recovering from its deformation.

In the first moments, the medium near the injection point is *extremely* deformed—stressed far beyond the linear elastic regime. The elastic response is nonlinear: the medium’s resistance saturates at a maximum stress (the yield plateau). This means the substance expands almost freely, encountering less resistance per unit expansion than a linear elastic model would predict. This is inflationary expansion: a brief epoch of rapid, nearly exponential growth driven by the elastic medium’s nonlinear response at extreme deformation.

As the substance spreads and thins, the deformation decreases. The medium’s response transitions from nonlinear (plateau) to linear (Hooke’s law). The restoring force now increases proportionally to the remaining deformation, and expansion decelerates. This is the radiation and matter-dominated era.

Eventually, the substance has spread so thin that its weight can no longer resist the medium’s residual elasticity. The medium continues to flatten, driving late-time accelerated expansion. This is the current epoch.

4.3 Advantages Over the Membrane Model

The three-dimensional model resolves three limitations of the membrane analogy:

1. **No preferred direction.** The substance expands isotropically from the injection point. There is no “down,” no external gravitational field, no embedding space. The expansion is driven by internal pressure and elastic recovery, not by an external force.
2. **No extrinsic curvature.** The elastic medium deforms *internally*—its density and stress fields change—without bending into an ambient space. This is closer to the intrinsic curvature of general relativity.
3. **Inhomogeneity is natural.** The injected substance cannot be perfectly homogeneous (Section 3). The injection process itself introduces gradients in pressure, density, and

flow velocity. These gradients—smoothed but not eliminated by the elastic response—become the primordial perturbations.

4.4 Formal Correspondence

Table 1: Correspondence between the elastic medium and spacetime. Compare with the two-dimensional correspondence in Paper 5, §4.4.

Elastic medium	Spacetime	Evidence
Elastic modulus κ	$1/(16\pi G)$	Einstein–Hilbert action
Undeformed equilibrium	Minkowski space ($R = 0$)	Vacuum field equations
Nonlinear elastic limit	Inflationary plateau	Starobinsky R^2
Elastic waves	Gravitational waves	LIGO (GW150914)
Energy in deformation	Isaacson $t_{\mu\nu}^{\text{GW}}$	LIGO energy measurement
Yield plateau	Slow-roll inflation	Planck CMB data
Injection inhomogeneities	Primordial perturbations	CMB anisotropies
Microstructural elasticity	Induced gravity	Sakharov (1967)

Counterarguments and Responses. *Objection: The injection model introduces an “injector”—an external agent that has no counterpart in cosmology.*

Response: The injector is a visualization aid, not a physical claim. What it represents is the initial condition: the energy content exists at $t = 0$ in a highly concentrated configuration. The origin of this initial condition is not addressed by inflation either—inflation pushes the question back but does not answer it. The model’s physical content begins *after* the injection, when the elastic medium responds to the deformation.

Objection: The injection introduces a preferred spatial point—a center—whereas the FRW metric is homogeneous and isotropic everywhere, not merely isotropic from one point.

Response: This is a genuine limitation. The injection picture is isotropic from the injection point but not homogeneous: regions near the center differ from regions far away. The FRW metric, by contrast, looks the same from every point. The analogy captures the *temporal* narrative (initial concentration $\rightarrow$ expansion $\rightarrow$ thinning) but not the spatial homogeneity of the cosmological solution. In the formal framework, homogeneity is imposed by the FRW ansatz (Eq. 33); the injection picture should be understood as illustrating the dynamics at any single comoving point, not the global spatial geometry.

Objection: The three-dimensional elastic medium has a rest frame. Spacetime in general relativity does not.

Response: This limitation is real and shared by all elastic analogies, including acoustic analogies for black holes [Unruh, 1981]. The rest frame of the medium corresponds to the comoving frame in cosmology, which is physically distinguished (it is the frame in which the CMB dipole vanishes). The analogy does not capture Lorentz invariance, which must be supplied by the formal framework.

Section Result and Implications

The three-dimensional elastic analogy resolves the principal limitations of the membrane model while preserving its four key dynamical features. The injection picture provides a natural origin for inhomogeneity (the substance cannot be homogeneous), a mechanism for inflation (the nonlinear elastic response at extreme deformation), and a unified narrative from Big Bang to present.

5 The Elastic Potential: Deriving $V(\psi)$ from Spacetime Elasticity

This section contains the central technical result of the paper: the derivation of the scalar field potential $V(\psi)$ from the established elastic properties of spacetime. We show that $V(\psi)$ is not a free function but is determined, at leading order in the curvature expansion, by the same physics that establishes the elasticity of the vacuum.

5.1 Linear Elasticity: The Einstein–Hilbert Action

As established in Paper 5 (§4.1), the gravitational action

$$S_{\text{EH}} = \frac{1}{16\pi G} \int d^4x \sqrt{-g} R \quad (3)$$

is an elastic energy functional: it vanishes at zero curvature ($R = 0$), generates restoring dynamics that drive the geometry toward flatness, and describes a medium whose elastic modulus is $1/(16\pi G)$.

In the language of elasticity, the Einstein–Hilbert action describes the *linear* elastic response of spacetime. It is the gravitational analog of Hooke’s law: the energy cost of deformation is proportional to the first power of the curvature invariant. (We emphasize: the field equations derived from (3) are second-order and highly nonlinear. The “linearity” refers to the action’s dependence on R , not to the dynamics. This subtlety is discussed in the Counterarguments below.)

5.2 Nonlinear Elasticity: Quantum Corrections

Real elastic materials are not exactly linear. At small deformations, Hooke’s law holds; at large deformations, the material enters a nonlinear regime where the stress-strain relation deviates from proportionality. The elastic energy acquires higher-order terms:

$$E_{\text{elastic}} = \frac{1}{2}\kappa\epsilon^2 + \frac{1}{3}\kappa_3\epsilon^3 + \frac{1}{4}\kappa_4\epsilon^4 + \cdots \quad (4)$$

where ϵ is the strain and κ_n are higher-order elastic moduli.

For spacetime, the analogous higher-order terms arise from quantum corrections to the gravitational action. The one-loop effective action for gravity, computed by integrating out

matter fields on a curved background, takes the form [Stelle, 1977, Birrell & Davies, 1982, Barvinsky & Vilkovisky, 1985]:

$$S_{\text{eff}} = \int d^4x \sqrt{-g} \left[\frac{R}{16\pi G} + \frac{1}{6M^2} R^2 + c_1 R_{\mu\nu} R^{\mu\nu} + c_2 R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} + \dots \right] \quad (5)$$

where M is a mass scale and c_1, c_2 are dimensionless coefficients determined by the field content.

By the Gauss–Bonnet theorem in four dimensions, the $R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma}$ and $R_{\mu\nu} R^{\mu\nu}$ terms can be combined with the R^2 term (up to a topological invariant that does not affect the equations of motion). The leading correction is therefore R^2 :

$$S_{\text{eff}} = \int d^4x \sqrt{-g} \left[\frac{R}{16\pi G} + \frac{R^2}{6M^2} \right] \quad (6)$$

This is the effective action of gravity at scales below the Planck scale but above the characteristic scale M . It includes both the linear elastic response (R term) and the leading nonlinear correction (R^2 term).

5.3 Conformal Transformation: The Starobinsky Potential

The $R + R^2$ action (6) is mathematically equivalent to general relativity coupled to a scalar field [Whitt, 1984, Maeda, 1988, Barrow & Cotsakis, 1988]. We present the derivation in detail, since the claim is that $V(\psi)$ is derived rather than assumed.

Step 1: Auxiliary field. Introduce the auxiliary field Φ and consider:

$$S = \int d^4x \sqrt{-g} \left[\frac{1}{16\pi G} \Phi R - \frac{3M^2 M_{\text{Pl}}^2}{4} (\Phi - 1)^2 \right]. \quad (7)$$

The equation of motion for Φ gives $\Phi = 1 + R/(3M^2)$, and substituting back into (7) recovers (6) identically. The two actions are classically equivalent.

Step 2: Conformal transformation. Define $\tilde{g}_{\mu\nu} = \Phi g_{\mu\nu}$. Under this transformation, the Ricci scalar transforms as (see, e.g., Whitt [1984], Eq. (2.5)):

$$\sqrt{-g} \Phi R = \sqrt{-\tilde{g}} \left[\tilde{R} - \frac{3}{2} \frac{\tilde{g}^{\mu\nu} \partial_\mu \Phi \partial_\nu \Phi}{\Phi^2} \right]. \quad (8)$$

Step 3: Scalar field redefinition. Define ψ by $\Phi = e^{\sqrt{2/3} \psi / M_{\text{Pl}}}$, or equivalently:

$$\psi = \sqrt{\frac{3}{2}} M_{\text{Pl}} \ln \Phi. \quad (9)$$

The kinetic term becomes $\frac{3}{2} \frac{(\partial\Phi)^2}{\Phi^2} = (\partial\psi)^2$, and the potential becomes:

$$V(\psi) = \frac{3M^2 M_{\text{Pl}}^2}{4} \left(1 - e^{-\sqrt{2/3} \psi / M_{\text{Pl}}} \right)^2 = \frac{3M^2 M_{\text{Pl}}^2}{4} (1 - \Phi^{-1})^2. \quad (10)$$

Step 4: Einstein frame action. The final result is:

$$S = \int d^4x \sqrt{-\tilde{g}} \left[\frac{\tilde{R}}{16\pi G} - \frac{1}{2}(\tilde{\nabla}\psi)^2 - V(\psi) \right] \quad (11)$$

with the potential

$$V(\psi) = \frac{3M^2 M_{\text{Pl}}^2}{4} \left(1 - e^{-\sqrt{2/3}\psi/M_{\text{Pl}}} \right)^2 \quad (12)$$

This is the Starobinsky potential [Starobinsky, 1980]. Its structure is illustrated in Figure 1.

5.4 Physical Interpretation: The Elastic Potential

The Starobinsky potential (12) has precisely the structure of a nonlinear elastic energy:

- **Near equilibrium** ($\psi \ll M_{\text{Pl}}$): expanding the exponential gives $V(\psi) \approx \frac{1}{2}m_\psi^2 \psi^2$ with $m_\psi^2 = \frac{4}{3} \frac{M^2 M_{\text{Pl}}^2}{M_{\text{Pl}}^2} \cdot \frac{2}{3M_{\text{Pl}}^2} = \frac{2M^2}{3}$. This is Hooke's law—quadratic elastic energy with spring constant m_ψ^2 .
- **At large deformation** ($\psi \gg M_{\text{Pl}}$): the potential saturates at $V_0 = \frac{3}{4}M^2 M_{\text{Pl}}^2$. Further increasing ψ costs negligible additional energy. This is the *yield plateau*—the elastic limit of the vacuum.
- **The transition between regimes** occurs at $\psi \sim M_{\text{Pl}}$, where the exponential factor transitions from ≈ 1 (linear) to ≈ 0 (saturated).

The identification is summarized as follows:

Proposition 5.1 (Elastic potential of spacetime). *Within the R^2 truncation of the gravitational effective action—i.e., retaining the Einstein–Hilbert term and its first quantum correction—the conformal transformation to the Einstein frame yields a unique scalar field potential: the Starobinsky potential $V(\psi) = V_0(1 - e^{-\sqrt{2/3}\psi/M_{\text{Pl}}})^2$. This potential has the universal structure of a nonlinear elastic energy: quadratic near equilibrium, saturating at large deformation. Higher-order corrections (R^3 , R^4 , $\nabla^2 R$) modify the potential at curvatures approaching the Planck scale but are suppressed by $R/M_{\text{Pl}}^2 \sim 10^{-6}$ during inflation.*

Proof. The Einstein–Hilbert action is the leading term in the derivative expansion of the gravitational effective action (established in Paper 5, §4.1–4.3). The next-to-leading term, by dimensional analysis and the Gauss–Bonnet identity, is R^2/M^2 [Stelle, 1977]. The conformal transformation to the Einstein frame [Whitt, 1984] converts this $R + R^2/M^2$ action into GR coupled to a scalar field with the potential (12). The derivation uses only standard results of quantum field theory in curved spacetime and the mathematical equivalence of $f(R)$ gravity with scalar-tensor theory. The truncation at R^2 is justified by the effective field theory principle: at inflationary energies $H_{\text{inf}}^2 \sim M^2 \sim 10^{-6}M_{\text{Pl}}^2$, higher-order terms contribute corrections of order $(H_{\text{inf}}/M_{\text{Pl}})^2 \sim 10^{-6}$, which are negligible. $\square$




Figure 1: (a) Elastic energy of a nonlinear material: quadratic (Hooke’s law) at small strain, saturating at the yield stress V_0 at large strain. (b) The Starobinsky potential $V(\psi)$, derived from the $R+R^2$ action. The structural correspondence is exact: quadratic near the minimum (late-time screening regime), saturating at a plateau (inflationary regime). The potential is not chosen for phenomenological convenience—it is the elastic energy of spacetime geometry.

Counterarguments and Responses. *Objection: The Einstein–Hilbert action is linear in R , while elastic energy is quadratic in strain. The “elastic” interpretation is imprecise.*

Response: This objection was addressed in Paper 5, §4.5. We do not claim algebraic identity between the Einstein equations and Hooke’s law. The structural correspondence is: both functionals are minimized at zero deformation, both generate restoring dynamics, and both admit nonlinear extensions at large deformation. The R^2 term—which is quadratic in curvature—completes the elastic picture at the action level. At the level of the equations of motion, the Starobinsky model produces dynamics (slow-roll, oscillation, relaxation) that are precisely those of a nonlinear elastic system.

Objection: The mass scale M is a free parameter. The potential is not fully determined.

Response: The form of $V(\psi)$ is determined; the scale M is not. This is analogous to Hooke’s law: the form $F = -kx$ is universal for elastic materials, but the spring constant k depends on the specific material. For spacetime, M is fixed by observations: the amplitude of the CMB power spectrum requires $M \approx 1.3 \times 10^{13} \text{ GeV}$ [Planck Collaboration, 2020]. This is three orders of magnitude below the Planck scale, consistent with R^2 being a perturbative correction.

Objection: Higher-order terms (R^3 , R^4 , $\nabla^2 R$) are neglected. Why is the R^2 truncation justified?

Response: By the effective field theory principle, higher-order terms are suppressed by additional powers of R/M_{Pl}^2 or ∇^2/M_{Pl}^2 . During inflation, $R \sim H_{\text{inf}}^2 \sim M^2 \sim 10^{-6} M_{\text{Pl}}^2$, so $R^3/M_{\text{Pl}}^4 \sim 10^{-12}$ corrections are negligible. The R^2 truncation is the leading-order nonlinear elastic approximation, valid as long as curvatures remain below the Planck scale—a condition satisfied throughout inflation.

Section Result and Implications

The scalar field potential $V(\psi)$ is derived, not postulated. It is the Starobinsky potential, arising uniquely from the Einstein–Hilbert action plus its first quantum correction. The elastic interpretation—quadratic near equilibrium, saturating at large deformation—explains *why* this potential has the right structure for inflation: it describes the nonlinear elastic response of spacetime at extreme curvature.

6 Modified Inflationary Dynamics in the Elastic Vacuum

With the potential $V(\psi)$ established, we now analyze the inflationary dynamics.

6.1 The Slow-Roll Equations

In the Einstein frame, the Friedmann equation and the scalar field equation of motion are:

$$H^2 = \frac{1}{3M_{\text{Pl}}^2} \left[\frac{1}{2}\dot{\psi}^2 + V(\psi) \right] \quad (13)$$

$$\ddot{\psi} + 3H\dot{\psi} + V'(\psi) = 0 \quad (14)$$

where $V(\psi)$ is the Starobinsky potential (12) and primes denote derivatives with respect to ψ .

The slow-roll approximation ($\dot{\psi}^2 \ll V$ and $|\ddot{\psi}| \ll |3H\dot{\psi}|$) yields:

$$H^2 \approx \frac{V(\psi)}{3M_{\text{Pl}}^2}, \quad (15)$$

$$3H\dot{\psi} \approx -V'(\psi). \quad (16)$$

The slow-roll parameters are:

$$\epsilon = \frac{M_{\text{Pl}}^2}{2} \left(\frac{V'}{V} \right)^2 = \frac{4}{3} \frac{e^{-2\sqrt{2/3}\psi/M_{\text{Pl}}}}{(1 - e^{-\sqrt{2/3}\psi/M_{\text{Pl}}})^2}, \quad (17)$$

$$\eta = M_{\text{Pl}}^2 \frac{V''}{V} = \frac{4}{3} \frac{e^{-\sqrt{2/3}\psi/M_{\text{Pl}}}(2e^{-\sqrt{2/3}\psi/M_{\text{Pl}}} - 1)}{(1 - e^{-\sqrt{2/3}\psi/M_{\text{Pl}}})^2}. \quad (18)$$

Inflation ends when $\epsilon(\psi_{\text{end}}) = 1$, giving $\psi_{\text{end}} \approx 0.94 M_{\text{Pl}}$.

6.2 Number of e-Folds

The number of e-folds between the field value ψ_* (when the pivot scale exits the horizon) and the end of inflation is:

$$N_* = \int_{\psi_{\text{end}}}^{\psi_*} \frac{V}{V'} \frac{d\psi}{M_{\text{Pl}}^2} = \frac{3}{4} \left(e^{\sqrt{2/3}\psi_*/M_{\text{Pl}}} - \frac{\sqrt{2/3}\psi_*}{M_{\text{Pl}}} \right) - \text{const.} \quad (19)$$

For large ψ_*/M_{Pl} , the dominant term is $N_* \approx \frac{3}{4}e^{\sqrt{2/3}\psi_*/M_{\text{Pl}}}$, giving $\psi_* \approx \sqrt{3/2} M_{\text{Pl}} \ln(4N_*/3)$. For $N_* = 55$: $\psi_* \approx 5.29 M_{\text{Pl}}$; for $N_* = 60$: $\psi_* \approx 5.36 M_{\text{Pl}}$.

6.3 Observational Predictions

The spectral index and tensor-to-scalar ratio, evaluated at $\psi = \psi_*$, are [Starobinsky, 1980, Mukhanov & Chibisov, 1981b]:

$$n_s = 1 - 6\epsilon + 2\eta \approx 1 - \frac{2}{N_*}, \quad (20)$$

$$r = 16\epsilon \approx \frac{12}{N_*^2}. \quad (21)$$

Proposition 6.1 (Planck compatibility). *For $N_* = 55 \pm 5$ e -folds (the range consistent with standard reheating scenarios), the elastic vacuum predicts:*

$$n_s = 0.964 \pm 0.003, \quad r = (3.3\text{--}4.8) \times 10^{-3}. \quad (22)$$

These values are consistent with Planck 2018 results: $n_s = 0.9649 \pm 0.0042$ and $r < 0.056$ at 95% confidence [Planck Collaboration, 2020].

The predictions are illustrated in Figures 2 and 3.

6.4 Physical Interpretation: Inflation as Elastic Recoil

In the three-dimensional elastic analogy of Section 4, inflation corresponds to the first moments after injection: the elastic medium is extremely deformed (high curvature, large ψ), and its response saturates at the yield plateau (the flat region of $V(\psi)$). The substance expands nearly freely because the medium’s resistance has plateaued—further deformation costs negligible additional energy. This is exponential expansion.

As the deformation decreases (ψ rolls off the plateau), the medium’s response becomes linear again (quadratic V), and expansion decelerates. The transition from plateau to quadratic is the graceful exit from inflation—a natural consequence of the elastic potential structure.

Definition 6.2 (Elastic recoil inflation). Inflation is the elastic recoil of spacetime geometry from a state of extreme initial deformation. The inflationary plateau is the yield stress of spacetime elasticity. The slow roll is the transition from nonlinear to linear elastic response. Reheating corresponds to oscillations of the scalar field about the equilibrium—the elastic vibrations of the relaxing medium.

Counterarguments and Responses. *Objection: This is just Starobinsky inflation with a new name. What is gained by calling it “elastic recoil”?*

Response: Three things. First, *physical motivation*: Starobinsky derived his model from quantum corrections to gravity, but the reason these corrections produce a viable inflationary potential was unexplained. The elastic interpretation provides the reason: the potential has the universal structure of nonlinear elastic energy. Second, *connection to late-time physics*: the same elastic framework explains dark energy (Paper 5) and inflation (this paper) as different regimes of the same elastic potential—low deformation and high deformation, respectively. Third, *predictive unity*: the framework predicts that $V(\psi)$ has the Starobinsky form, not as one choice among many inflationary models, but as the leading-order elastic potential of spacetime within the R^2 truncation.




Figure 2: Inflationary dynamics in the elastic vacuum. (a) Evolution of the scalar field ψ in e-folds N . (b) Slow-roll parameters ϵ and $|\eta|$; inflation ends when $\epsilon = 1$. (c) Trajectory on the potential. (d) Spectral index n_s and tensor-to-scalar ratio r as functions of N_* , with the Planck constraint band for n_s .

6.5 The Mass Hierarchy: What Is Unified and What Is Not

We must address directly the most significant open tension in the elastic vacuum programme. The inflationary scalaron has mass $m_\psi \sim M \sim 1.3 \times 10^{13}$ GeV, fixed by the CMB power spectrum amplitude. The late-time screening field of Paper 5 requires $m_\psi \sim H_0 \sim 10^{-33}$ eV. The ratio is:

$$\frac{m_{\psi,\text{inf}}}{m_{\psi,\text{late}}} \sim \frac{10^{13} \text{ GeV}}{10^{-33} \text{ eV}} \sim 10^{55}. \quad (23)$$

This is not a quantitative discrepancy amenable to perturbative correction. It is a qualitative chasm.

What is unified: the interpretation. At both epochs, the dynamics are driven by the elastic response of spacetime geometry to deformation. The Einstein–Hilbert action provides the elastic modulus. The quantum corrections provide the nonlinear extension. The physical picture—recoil from deformation (inflation) and relaxation toward equilibrium (late-time acceleration)—is the same.

What is not unified: the dynamics. The inflationary and late-time sectors involve




Figure 3: Predictions in the n_s – r plane. The elastic vacuum / Starobinsky model (red line, with markers at $N_* = 50, 55, 60$) falls within the Planck-favored region. The quadratic potential $V \propto \phi^2$ (dashed green) is ruled out.

effective scalar degrees of freedom with masses differing by 55 orders of magnitude. Presenting these as “different excitations of the same elastic medium” is suggestive—in the same way that acoustic and seismic waves in a solid are different excitations of the same elastic medium. But in a solid, the frequency ratio between acoustic and seismic modes is at most a few orders of magnitude, not 10^{55} . The analogy is therefore aspirational, not demonstrated.

Possible resolution: renormalization group running. The mass parameter M in the gravitational effective action may run with the energy scale. Asymptotic safety calculations [Codello et al., 2009, Falls et al., 2014] show that gravitational couplings run between UV and IR, but the running factors are typically $\mathcal{O}(1)$ to $\mathcal{O}(10)$ —far short of the required 10^{55} . A complete bridge would require either (a) a mechanism that drives M to extreme IR values through cosmological evolution (not merely RG running), (b) a multi-field extension where the inflationary scalaron and the late-time screening field are different fields with different masses, related by the common elastic principle but not by a single effective potential, or (c) a qualitative change in the effective action between the inflationary and late-time regimes, perhaps driven by phase transitions in the vacuum structure.

Honest assessment. The elastic vacuum programme currently achieves *unification of interpretation*—the same physical picture applies at both epochs—but not *unification of dynamics*. The two sectors have different effective parameters by many orders of magnitude, and bridging this gap is the programme’s deepest unsolved problem. Until it is solved, the claim of a “unified elastic framework” should be understood as describing a common physical principle, not a single dynamical theory with one set of parameters.

Section Result and Implications

The elastic vacuum framework derives Starobinsky inflation from the elastic properties of spacetime, producing predictions ($n_s \approx 0.964$, $r \approx 0.004$) that match Planck observations. The interpretation of inflation as elastic recoil is physically motivated and connects early- and late-time cosmology through a common principle. However, the dynamical unification of the inflationary and late-time sectors remains open: the mass hierarchy of 10^{55} between the two regimes is the programme’s most significant unsolved problem, and this paper achieves interpretive rather than dynamical unification.

7 Primordial Fluctuations from Vacuum Inhomogeneities

In the standard inflationary picture, primordial curvature perturbations arise from quantum fluctuations of the inflaton field in the Bunch–Davies vacuum state on de Sitter space [Mukhanov & Chibisov, 1981, Hawking, 1982, Bardeen et al., 1983]. The power spectrum of these perturbations is:

$$\Delta_\zeta^2(k) = \frac{1}{24\pi^2 M_{\text{Pl}}^4} \left. \frac{V}{\epsilon} \right|_{k=aH}, \quad (24)$$

and it is nearly scale-invariant because V and ϵ vary slowly during inflation.

In the elastic vacuum framework, this mechanism is preserved but supplemented by a deeper physical picture.

7.1 Standard Mechanism: Quantum Fluctuations of ψ

The scalar field ψ undergoes quantum fluctuations in the inflationary de Sitter background, with the standard mode equation [Mukhanov & Sasaki, 1986]:

$$v_k'' + \left(k^2 - \frac{z''}{z} \right) v_k = 0, \quad (25)$$

where $v_k = z \zeta_k$ is the Mukhanov–Sasaki variable, $z = a\dot{\psi}/H$, and primes denote conformal time derivatives. This is identical to the standard calculation, and the result is (24) with the Starobinsky potential.

For $N_* = 55$: $\Delta_\zeta^2 \approx 2.1 \times 10^{-9}$, matching the Planck measurement [Planck Collaboration, 2020]. This fixes the mass scale: $M \approx 1.3 \times 10^{13}$ GeV.

7.2 The New Element: Primordial Inhomogeneities

The analysis of Section 3 establishes that the initial state is necessarily inhomogeneous. This means the field ψ does not start in the Bunch–Davies vacuum (which is the ground state of quantum fluctuations on an exact de Sitter background) but in a more general state that includes pre-existing inhomogeneities.

Let the initial field configuration be:

$$\psi(\mathbf{x}, t_i) = \bar{\psi}_i + \delta\psi_i(\mathbf{x}), \quad (26)$$

where $\bar{\psi}_i$ is the spatial average and $\delta\psi_i$ represents the irreducible inhomogeneities.

During inflation, these pre-existing perturbations are stretched to super-horizon scales and redshifted. Modes that were sub-Hubble before inflation are stretched beyond the horizon, where they freeze as classical curvature perturbations. The total curvature perturbation is:

$$\zeta_k = \zeta_k^{(\text{QM})} + \zeta_k^{(\text{init})}, \quad (27)$$

where $\zeta_k^{(\text{QM})}$ is the standard quantum contribution (24) and $\zeta_k^{(\text{init})}$ arises from the primordial inhomogeneities.

Proposition 7.1 (Primordial inhomogeneity suppression). *If inflation lasts N_* e-folds, the primordial inhomogeneity contribution to the curvature perturbation on scale k is suppressed relative to the quantum contribution by a factor:*

$$\frac{\zeta_k^{(\text{init})}}{\zeta_k^{(\text{QM})}} \sim \frac{\delta\psi_i(k)}{H/(2\pi)} \cdot e^{-\Delta N(k)}, \quad (28)$$

where $\Delta N(k)$ is the number of e-folds between the time the pre-existing mode was the same size as the Hubble radius and the time it re-enters the quantum regime.

Proof. A pre-existing perturbation $\delta\psi_i$ on scale λ is stretched during inflation to $\lambda \cdot e^N$. Its amplitude decays as $\delta\psi \propto e^{-N}$ (diluted by expansion) while quantum fluctuations are continuously generated with amplitude $H/(2\pi)$. After N e-folds, the ratio of initial to quantum perturbations on the same scale goes as $e^{-\Delta N}$. For $\Delta N \gg 1$, the initial perturbations are negligible on observable scales. $\square$

This result explains why inflation produces a nearly scale-invariant spectrum regardless of initial conditions: whatever the initial inhomogeneities, they are exponentially suppressed and replaced by the quantum fluctuations generated during the slow roll. The key insight from Section 3 is that *some* initial inhomogeneities must exist—but inflation renders them unobservable on CMB scales, leaving only the quantum-generated spectrum.

7.3 Quantitative Estimate

To move beyond the scaling argument, consider a concrete example. Suppose the pre-inflationary state has an initial perturbation spectrum with characteristic amplitude $\delta\psi_i/M_{\text{Pl}} \sim \mathcal{O}(1)$ (Planck-scale inhomogeneities, the most extreme physically reasonable case) on a co-moving scale λ_i comparable to the Hubble radius at the onset of inflation, $\lambda_i \sim H_{\text{inf}}^{-1}$.

During N_{tot} total e-folds of inflation, this mode is stretched to a physical scale $\lambda_{\text{final}} = \lambda_i \cdot e^{N_{\text{tot}}}$. The mode corresponding to the CMB pivot scale $k_* = 0.05 \text{ Mpc}^{-1}$ exits the horizon at $N_* \approx 55$ e-folds before the end of inflation. If inflation lasts N_{tot} total e-folds, the pre-inflationary mode was at the Hubble scale $\Delta N = N_{\text{tot}} - N_*$ e-folds before it would re-enter the quantum regime. The suppression factor is:

$$\frac{\zeta_*^{(\text{init})}}{\zeta_*^{(\text{QM})}} \sim \frac{\delta\psi_i}{H_{\text{inf}}/(2\pi)} e^{-\Delta N} \sim \frac{M_{\text{Pl}}}{H_{\text{inf}}/(2\pi)} e^{-(N_{\text{tot}}-55)}. \quad (29)$$

For $H_{\text{inf}} \sim 10^{13}$ GeV (Starobinsky inflation) and $M_{\text{Pl}} \sim 2.4 \times 10^{18}$ GeV:

$$\frac{\zeta_*^{(\text{init})}}{\zeta_*^{(\text{QM})}} \sim 10^6 e^{-(N_{\text{tot}}-55)}. \quad (30)$$

For this ratio to be $\lesssim 1$ (quantum fluctuations dominate), we need $N_{\text{tot}} - 55 \gtrsim 14$, i.e., $N_{\text{tot}} \gtrsim 69$. For $N_{\text{tot}} = 70$: the ratio is $\sim 10^6 e^{-15} \sim 3$. For $N_{\text{tot}} = 80$: the ratio is $\sim 10^6 e^{-25} \sim 10^{-5}$. For $N_{\text{tot}} = 100$: the ratio is $\sim 10^6 e^{-45} \sim 10^{-14}$.

The conclusion is robust: if inflation lasts more than ~ 70 e-folds, even Planck-amplitude pre-inflationary perturbations are subdominant on CMB scales. The marginal case ($N_{\text{tot}} \approx 65$ – 70) is where residual effects might appear at the largest angular scales ($\ell \lesssim 10$), potentially contributing to observed CMB anomalies.

7.4 Observable Consequences

The elastic vacuum framework introduces two potential deviations from the standard prediction:

(i) Large-scale anomalies. If inflation lasted only marginally more than the minimum ~ 60 e-folds, modes corresponding to the largest observable scales ($\ell \lesssim 10$ in the CMB) may retain a residual contribution from pre-inflationary inhomogeneities. This would appear as anomalous power on the largest angular scales—qualitatively consistent with the observed low- ℓ power deficit and hemispherical asymmetry in the CMB [Planck Collaboration, 2020b].

(ii) Non-Gaussianity. The Starobinsky model predicts $f_{\text{NL}} \sim \mathcal{O}(1/N_*) \sim 0.02$, well below current observational limits [Planck Collaboration, 2020c]. Pre-inflationary inhomogeneities could contribute additional non-Gaussianity, but only on the largest scales (by the same suppression argument).

Counterarguments and Responses. *Objection: If initial inhomogeneities are exponentially suppressed, they are unobservable. What is the physical content of this section?*

Response: The physical content is conceptual, not observational on most scales. The point is that inflation does not require a homogeneous initial state—it produces approximate homogeneity from *any* initial state with sufficient duration. This resolves the initial conditions problem that standard inflation merely relocates. The residual observational signature (large-scale anomalies) is a bonus, not the main result.

Section Result and Implications

The elastic vacuum framework preserves the standard mechanism for generating primordial fluctuations (quantum fluctuations of ψ in slow roll) while providing a deeper physical picture: the initial state is necessarily inhomogeneous, and inflation smooths these inhomogeneities exponentially, replacing them with the quantum spectrum. The initial conditions problem is resolved: any physically realizable initial state, processed through elastic recoil, produces the observed near-scale-invariant spectrum.

8 The Flatness Feedback: Deriving β from $V(\psi)$

Paper 5 (§8) introduced the flatness feedback mechanism: a positive feedback loop in which decreasing curvature facilitates faster expansion, which further decreases curvature. The mechanism was encoded in the modified Friedmann equation:

$$H^2 = \frac{8\pi G}{3}\rho + \frac{\Lambda_0}{3} - \frac{Kc^2}{a^2} + \beta \frac{|K|}{a^2} H^2, \quad (31)$$

where $\beta > 0$ is the feedback parameter. Paper 5 established the flatness attractor *conditional* on $\beta > 0$ but did not derive β from the action. We now complete this derivation.

8.1 The Scalar Field in Curved FLRW Spacetime

Consider the full action from Paper 5 (§6.1):

$$S = \int d^4x \sqrt{-g} \left[\frac{R - 2\Lambda_0}{16\pi G} + \mathcal{L}_m + \mathcal{L}_{\text{vac}} - \frac{1}{2}(\nabla\psi)^2 - V(\psi) - \xi \psi T^\mu_\mu \right] \quad (32)$$

in a Friedmann–Lemaître–Robertson–Walker spacetime with spatial curvature K :

$$ds^2 = -dt^2 + a(t)^2 \left[\frac{dr^2}{1 - Kr^2} + r^2 d\Omega^2 \right]. \quad (33)$$

The scalar field equation of motion is:

$$\ddot{\psi} + 3H\dot{\psi} + V'(\psi) = -\xi T^\mu_\mu. \quad (34)$$

For pressureless matter ($p_m = 0$) and radiation ($T^\mu_{\mu, \text{rad}} = 0$), the trace is $T^\mu_\mu = -\rho_m$, so:

$$\ddot{\psi} + 3H\dot{\psi} + V'(\psi) = \xi \rho_m. \quad (35)$$

8.2 Quasi-Static Approximation

At late times ($z \lesssim 0.7$), when the elastic vacuum dominates and ψ is near its minimum, we write $\psi = \psi_0 + \delta\psi$ with $\delta\psi$ small. In the quasi-static approximation ($\ddot{\psi}$ and $3H\dot{\psi}$ are small compared to $m_\psi^2 \delta\psi$):

$$m_\psi^2 \delta\psi \approx \xi \rho_m, \quad \Rightarrow \quad \delta\psi \approx \frac{\xi \rho_m}{m_\psi^2}, \quad (36)$$

where $m_\psi^2 = V''(\psi_0) > 0$ for any convex potential.

8.3 Scalar Field Energy Density

The energy density of ψ is:

$$\rho_\psi = \frac{1}{2}\dot{\psi}^2 + V(\psi) \approx V(\psi_0) + \frac{1}{2}m_\psi^2 (\delta\psi)^2 = V_0 + \frac{\xi^2 \rho_m^2}{2m_\psi^2}. \quad (37)$$

The Friedmann equation with curvature is:

$$H^2 + \frac{K}{a^2} = \frac{8\pi G}{3}(\rho_m + \rho_\psi) + \frac{\Lambda_0}{3} = \frac{8\pi G}{3} \left(\rho_m + V_0 + \frac{\xi^2 \rho_m^2}{2m_\psi^2} \right) + \frac{\Lambda_0}{3}. \quad (38)$$

8.4 Derivation of β

Now consider the dependence of ρ_ψ on spatial curvature. From the Friedmann constraint, the matter density ρ_m depends implicitly on K/a^2 : in a universe with $K \neq 0$, the same expansion rate H corresponds to a different matter density than in a flat universe.

Differentiating (38) with respect to K/a^2 at fixed H :

$$1 = \frac{8\pi G}{3} \left(\frac{\partial \rho_m}{\partial(K/a^2)} + \frac{\xi^2 \rho_m}{m_\psi^2} \frac{\partial \rho_m}{\partial(K/a^2)} \right). \quad (39)$$

Solving:

$$\frac{\partial \rho_m}{\partial(K/a^2)} = \frac{3/(8\pi G)}{1 + \xi^2 \rho_m / m_\psi^2}. \quad (40)$$

The effective Friedmann equation can be rewritten as:

$$H^2 = \frac{8\pi G}{3} \rho_{\text{eff}} + \frac{\Lambda_{\text{eff}}}{3} - \frac{K_{\text{eff}}}{a^2}, \quad (41)$$

where the effective curvature parameter is:

$$K_{\text{eff}} = K \left(1 - \frac{\xi^2 \rho_m}{m_\psi^2 + \xi^2 \rho_m} \right) = K \frac{m_\psi^2}{m_\psi^2 + \xi^2 \rho_m}. \quad (42)$$

Proposition 8.1 (Positive feedback parameter — quasi-static limit). *For the action (32) with a convex potential ($V''(\psi_0) > 0$), in the quasi-static regime ($m_\psi \gg H$, $|\dot{\delta\psi}| \ll m_\psi^2 |\delta\psi|$), the flatness feedback parameter is:*

$$\boxed{\beta = \frac{\xi^2 \rho_m}{m_\psi^2 + \xi^2 \rho_m} > 0} \quad (43)$$

for any $\xi \neq 0$ and $m_\psi^2 > 0$.

Proof. From (42), $K_{\text{eff}} = K(1 - \beta)$ with β given by (43). Since $\xi^2 \geq 0$, $\rho_m > 0$, and $m_\psi^2 > 0$ (by the convexity of V at its minimum), we have $0 < \beta < 1$. The effective curvature K_{eff} has the same sign as K but smaller magnitude: $|K_{\text{eff}}| < |K|$. The scalar field backreaction reduces the effective spatial curvature, driving the universe toward flatness. $\square$

Validity of the quasi-static approximation. The quasi-static limit requires $m_\psi \gg H$, i.e., that the scalar field responds to changes in ρ_m faster than the cosmological expansion timescale. Paper 5’s screening mechanism, however, requires $m_\psi \sim H_0$. In this marginal regime, the quasi-static approximation breaks down: the scalar field evolves on the same timescale as the expansion, and the full dynamical equations (35) must be solved. The sign of β is robust—it follows from the convexity of $V(\psi)$ at its minimum, which guarantees that the scalar field’s energy increases with $|\rho_m|$, irrespective of the approximation—but the magnitude and the detailed time dependence of $\beta(a)$ may differ substantially from (44) in the full dynamical regime. A numerical solution of the coupled (ψ, a, K) system with $m_\psi \sim H_0$ is required to confirm the quantitative flatness attractor, and is identified as a priority for Paper 7.




Figure 4: Evolution of $|\Omega_K|$ with scale factor. (a) Standard Λ CDM: curvature decreases at late times due to Λ domination but is not actively driven to zero. (b) Elastic vacuum with feedback parameter γ (related to β through the elastic response): curvature is exponentially suppressed after the deceleration-to-acceleration transition at $z \approx 0.7$. The green dashed line shows the Planck bound $|\Omega_K| < 0.001$.

8.5 Epoch Dependence

The parameter β is not constant; it depends on $\rho_m(a)$:

$$\beta(a) = \frac{\xi^2 \rho_{m,0} a^{-3}}{m_\psi^2 + \xi^2 \rho_{m,0} a^{-3}}. \quad (44)$$

At early times ($a \rightarrow 0$, high density): $\beta \rightarrow 1$. The curvature is almost completely screened. At late times ($a \rightarrow \infty$, low density): $\beta \rightarrow 0$. The screening diminishes.

This epoch dependence resolves the consistency concern raised in Paper 5 (§8.3): during matter domination, ρ_m is large, so β is significant, and curvature is actively suppressed. During the late-time accelerated epoch, ρ_m has diluted, but the curvature has already been driven to near-zero values. The flatness is maintained by its own stability (once $|K|$ is small, the feedback is no longer needed to keep it small).

Counterarguments and Responses. *Objection: The derivation assumes quasi-static ψ . What if ψ oscillates or evolves dynamically?*

Response: The quasi-static approximation is valid when $m_\psi \gg H$, which is the regime of Paper 5’s screening mechanism. If $m_\psi \sim H$ (as required for Hubble-scale screening), the approximation is marginal, and the full dynamical equations must be solved. A numerical analysis with specific $V(\psi)$ and ξ values is identified as future work. However, the key qualitative result— $\beta > 0$ when $V''(\psi_0) > 0$ —does not depend on the quasi-static assumption; it follows from the convexity of the potential, which guarantees that the scalar field’s energy increases with $|\rho_m|$, which in turn is enhanced by $|K|$.

Objection: β depends on ξ and m_ψ , which are free parameters. The derivation is not fully determined.

Response: The form of β is derived, not postulated. The sign ($\beta > 0$) is guaranteed for any convex $V(\psi)$ with $\xi \neq 0$ —this is a *structural* result, independent of specific parameter values. The magnitude of β depends on $\xi^2 \rho_m / m_\psi^2$, which can in principle be constrained by rotation curve data (Paper 5, §9.2) or ISW measurements.

Section Result and Implications

The flatness feedback parameter β is derived from the action and shown to be positive for any convex scalar field potential with nonzero matter coupling. This completes the proof deferred from Paper 5: spatial flatness is a stable late-time attractor of the elastic vacuum. The β parameter is epoch-dependent, being largest during matter domination (when curvature suppression is most needed) and diminishing at late times (when flatness has been established).

9 Observational Consistency and Predictions

The elastic vacuum framework inherits the observational predictions of Starobinsky inflation—the model from which $V(\psi)$ is derived—and supplements them with qualitative predictions arising from the elastic interpretation. We distinguish these two categories explicitly.

9.1 Inherited Predictions: The n_s – r Plane

The following predictions are identical to Starobinsky R^2 inflation and are inherited by construction:

The elastic vacuum predicts the Starobinsky model values:

$$n_s \approx 1 - \frac{2}{N_*}, \quad r \approx \frac{12}{N_*^2}, \quad (45)$$

placing the model squarely within the Planck-favored region (Figure 3). The prediction $r \sim 3\text{--}5 \times 10^{-3}$ is testable by next-generation CMB experiments including LiteBIRD [LiteBIRD Collaboration, 2023] and CMB-S4 [CMB-S4 Collaboration, 2019], which will achieve sensitivity $\sigma(r) \sim 10^{-3}$.

9.2 Inherited Predictions: The Equation of State

At late times, the elastic vacuum predicts $w \neq -1$ with $w_0 > -1$ and $w_a < 0$ in the CPL parameterization (Paper 5, §9.1). This prediction is inherited from the dynamical dark energy sector of the framework. Recent DESI BAO data [DESI Collaboration, 2024] provide mild ($2\text{--}3\sigma$) evidence for dynamical dark energy consistent with this prediction.

9.3 New Predictions: Large-Scale CMB Anomalies

If inflation lasted only marginally more than 60 e-folds, the elastic vacuum framework predicts residual pre-inflationary inhomogeneities on the largest angular scales, potentially explaining the observed low quadrupole, hemispherical asymmetry, and cold spot anomalies [Planck Collaboration, 2020b].

9.4 New Predictions: Non-Gaussianity

The Starobinsky model predicts $f_{\text{NL}}^{\text{local}} \sim 0.02$, well below current limits but potentially detectable by future large-scale structure surveys.

9.5 Falsifiable Predictions

Following Paper 5 (§9.5):

$$\mu(a, k) = \Sigma(a, k) = 1, \quad \eta(a, k) = 0, \quad c_T^2 = 1. \quad (46)$$

Any detection of anomalous gravitational slip, modified light propagation, or anomalous tensor speed would falsify the framework. Additionally:

- Phantom behavior ($w < -1$) is excluded.
- The tensor-to-scalar ratio must satisfy $r > 10^{-3}$ (for $N_* < 70$).
- The running of the spectral index is $dn_s/d \ln k \approx -2/N_*^2 \approx -6 \times 10^{-4}$.

Counterarguments and Responses. *Objection: These predictions are identical to Starobinsky inflation. Where is the new physics?*

Response: We agree, and the section above distinguishes these explicitly. The inherited predictions (Starobinsky n_s , r ; late-time $w \neq -1$) are observational *consistency checks*, not new physics. The genuinely new predictions are: (1) large-scale CMB anomalies from pre-inflationary inhomogeneities (quantified in §7); (2) the physical explanation of *why* the Starobinsky model is singled out among hundreds of inflationary models (elastic recoil); (3) the connection between inflationary and late-time physics through a common elastic principle. At nonlinear order, the pre-inflationary inhomogeneity contribution may produce distinctive signatures in the bispectrum, though this requires the full computation deferred to future work.

Section Result and Implications

The elastic vacuum makes specific, falsifiable predictions across multiple observational domains. The inflationary predictions coincide with the Starobinsky model (which is their physical origin) and are testable by LiteBIRD and CMB-S4. The late-time predictions are testable by DESI and Euclid.

10 Discussion

The elastic vacuum framework, extended to the inflationary epoch, produces a unified physical picture in which a single principle—the intrinsic elasticity of spacetime—governs both the earliest and the latest epochs of cosmic history.

The deepest result of this paper is Proposition 5.1: the Starobinsky potential is not one choice among hundreds of inflationary models but the leading-order elastic potential of the vacuum within the R^2 truncation of the effective action. This reduces the inflationary model-selection problem to a physical derivation—within the effective field theory framework where the truncation is justified. We note the distinction between what is derived (the Starobinsky potential from $R + R^2$, via standard conformal transformation) and what is interpreted (the elastic recoil picture, the connection to late-time physics). The derivation is standard and uncontroversial; the interpretation is the contribution of this paper. The Starobinsky model’s empirical success—its consistency with all Planck data, its economy of assumptions, its derivability from quantum corrections to gravity—is explained by the elastic interpretation: it describes the nonlinear elastic response of spacetime, and no other structure is available at leading order in the curvature expansion.

The connection between epochs is interpretive rather than dynamical, as discussed candidly in §6.5. The same elastic physics that produces inflationary recoil at high curvature produces late-time acceleration at low curvature (Paper 5). But the effective mass scales differ by 55 orders of magnitude, and this gap is not yet bridged. The elastic vacuum programme currently achieves unification of physical picture, not unification of dynamics.

The initial conditions problem is resolved at the conceptual level. Inflation does not require homogeneous initial conditions; it requires only that the elastic medium be sufficiently deformed (large R , large ψ). The initial inhomogeneities, which are physically unavoidable (Section 3), are exponentially smoothed by the elastic recoil, leaving only the quantum fluctuation spectrum on observable scales.

The historical irony noted in Paper 5 deepens. In 1967, Zel’dovich identified vacuum energy with the cosmological constant, producing the worst prediction in physics. In the same year, Sakharov showed that gravity is the elastic response of the vacuum. Thirteen years later, Starobinsky used quantum corrections to gravity—the nonlinear extension of Sakharov’s elasticity—to construct the first viable inflationary model. The elastic vacuum framework recognizes these as aspects of a single physical picture that has been available, in pieces, for over half a century.

11 Limitations

The mass hierarchy. The inflationary scalaron has $m_\psi \sim M \sim 10^{13}$ GeV; the late-time screening field requires $m_\psi \sim H_0 \sim 10^{-33}$ eV. This 55-order-of-magnitude gap is the programme’s most significant open problem, discussed in detail in §6.5.

The quasi-static approximation. The derivation of β in Section 8 assumes quasi-static evolution of ψ , valid only for $m_\psi \gg H$. The sign of β is robust (it follows from convexity of V), but the magnitude requires full dynamical confirmation. A numerical solution of the coupled (ψ, a, K) system is needed.

No numerical MCMC analysis. Compatibility with CMB data is argued analytically through the Starobinsky model equivalence but not verified through a full Boltzmann code (CLASS/CAMB) analysis.

The coupling parameter ξ . The matter coupling ξ remains undetermined from first principles. Its value is constrained by solar system tests ($\xi \lesssim 10^{-3}$) and by the requirement of galactic-scale effects.

Derivation versus interpretation. The derivation of the Starobinsky potential from $R + R^2$ gravity via conformal transformation is standard. The elastic recoil interpretation—while physically motivated and pedagogically powerful—is a layer of narrative placed atop this standard result. The paper is transparent about this distinction (see Discussion), but readers should note that the interpretive claims (“inflation is elastic recoil,” “the yield plateau of spacetime”) are not independently testable beyond the Starobinsky model’s predictions.

The scalar field ψ is a new degree of freedom. Although we describe ψ as a collective mode of the elastic vacuum (analogous to a phonon in condensed matter), it is operationally a new scalar field with its own equation of motion and coupling constant ξ . The framework does not contain “no new degrees of freedom”—it contains one new scalar degree of freedom whose physical interpretation differs from standard quintessence.

12 Future Directions

Renormalization group bridge. The most urgent priority is to demonstrate whether the Starobinsky mass M can evolve to the Hubble scale through renormalization group running in curved spacetime, connecting the inflationary and late-time regimes. Preliminary results in asymptotic safety [Codello et al., 2009, Falls et al., 2014] suggest that gravitational couplings run significantly between the inflationary scale and the IR, but a complete calculation is needed.

Pre-inflationary power spectrum. Computing the full power spectrum including pre-inflationary inhomogeneities, with realistic initial states, would produce quantitative predictions for large-scale CMB anomalies.

Full Boltzmann code analysis. Implementing the elastic vacuum modifications in CLASS or CAMB and running MCMC against Planck, DESI, and ACT data.

N-body simulations. Implementing the scale-dependent screening in cosmological simulation codes to study structure formation.

Connection to Volovik. The condensed-matter analog of the elastic vacuum—superfluid helium—provides a laboratory system where the elastic potential and screening mechanism may be studied experimentally [Volovik, 2003].

13 Position Within the Research Programme

This paper is the sixth in a programme of seventeen papers developing the hypothesis that quantum vacuum energy gravitates locally. Its specific role is to connect the elastic vacuum framework (Paper 5) to inflationary cosmology.

What this paper accomplishes:

- (i) Derives $V(\psi)$ from spacetime elasticity—the Starobinsky potential, within the R^2 truncation of the effective action.
- (ii) Derives $\beta > 0$ from the action in the quasi-static limit, completing the conditional flatness attractor proof of Paper 5.
- (iii) Establishes, via the Heisenberg uncertainty principle, that the initial state of the universe cannot be exactly homogeneous, resolving the initial conditions problem at the conceptual level.
- (iv) Provides the three-dimensional elastic analogy, resolving the principal limitations of Paper 5’s membrane model.
- (v) Identifies the mass hierarchy ($\sim 10^{55}$) between inflationary and late-time sectors as the programme’s most significant open problem, distinguishing interpretive from dynamical unification.

What is deferred:

- (i) Full dynamical confirmation of $\beta > 0$ beyond the quasi-static limit; numerical solution of the coupled (ψ, a, K) system (Paper 7).
- (ii) CMB MCMC analysis with Boltzmann code (Paper 7).
- (iii) Structure growth and N-body simulations (Papers 8–9).
- (iv) Galaxy rotation curves (Paper 12).
- (v) The mass hierarchy problem: bridging the inflationary and late-time sectors dynamically (future work, possibly beyond the current programme).

14 Recommended Journal for Publication

This paper is recommended for submission to the **Journal of Cosmology and Astroparticle Physics (JCAP)**.

The primary reasons are as follows. The paper derives a specific inflationary potential from physical principles, computes observational predictions (n_s, r, f_{NL}), and connects inflation to late-time dark energy within a unified framework. This combination of inflationary model building, observational cosmology, and theoretical foundations falls squarely within JCAP’s editorial scope.

The paper would also be appropriate for Physical Review D (given the mathematical development and the connection to $f(R)$ gravity) or Classical and Quantum Gravity (given the foundational aspects of the elastic vacuum interpretation).

Conclusions

The elastic vacuum framework provides a physical explanation for inflation.

The Starobinsky potential—the inflationary model most favored by Planck data—is not an arbitrary choice from a landscape of hundreds of scalar field models. Within the R^2 truncation of the gravitational effective action, it is the leading-order elastic potential of space-time geometry, arising from the Einstein–Hilbert action supplemented by its first quantum correction. The inflationary plateau is the yield stress of spacetime elasticity: beyond a critical deformation, further curvature costs negligible additional energy, producing exponential expansion.

Inflation, in this picture, is elastic recoil. The universe began in a state of extreme geometric deformation—necessarily inhomogeneous, as the Heisenberg uncertainty principle requires for any finite-energy field configuration. The elastic response of spacetime drove exponential expansion, smoothing the initial inhomogeneities and replacing them with the quantum fluctuation spectrum that we observe in the CMB.

The framework yields two concrete results deferred from Paper 5. First, $V(\psi) = V_0(1 - e^{-\sqrt{2/3}\psi/M_{\text{Pl}}})^2$: derived from the elastic properties of spacetime, not postulated. Second, $\beta > 0$ in the quasi-static limit: proved from the convexity of $V(\psi)$, completing the demonstration that spatial flatness is a late-time attractor—with full dynamical confirmation deferred to numerical work.

The screening scalar field ψ is a new degree of freedom—we do not deny this—but it is not a dark matter candidate; it is a mediator of the vacuum’s elastic response, with a role analogous to the phonon in condensed matter: not a new constituent of the universe but a collective mode of the medium that is already there. The framework requires no fine-tuning of initial conditions or vacuum energy. Only the consistent application of general relativity, quantum field theory, and the recognition that spacetime is elastic—a fact established by the Einstein–Hilbert action, confirmed by LIGO, derived from quantum field theory by Sakharov—and that the natural response of an elastic medium to extreme deformation is, first, to recoil (inflation), and then, to relax (the observed universe).

The physics to see this has been available since Starobinsky’s 1980 paper, which derived R^2 inflation from quantum corrections to gravity. What was missing was the recognition that these quantum corrections are the nonlinear elastic terms in the effective action of a medium whose linear elasticity is the Einstein–Hilbert action itself. The dynamical unification of the inflationary and late-time regimes—bridging the 55-order-of-magnitude mass gap—remains the programme’s deepest open problem. But the interpretive unification is already in place: the same elastic principle governs both epochs, and the Starobinsky potential is its leading-order expression.

Acknowledgments

The author thanks George F. R. Ellis for consultations on the trace-free Einstein equations. The author is grateful to Joan Solà Peracaula for discussions on the running vacuum model. The structural observation in Remark 3.1 draws on Kriger [2026], whose philosophical foundations were developed independently. Any errors are the author’s own.

References

- Kriger, B. (2025). The Elastic Vacuum: A Sequestering Mechanism for Vacuum Energy and the Complete Dynamical History of Cosmic Expansion. Manuscript submitted for publication.
- Kriger, B. (2026). Structural Genesis of Dynamical Architecture: Deriving the Necessary Form of Physical Law from Conditions on Sustainable Complexity. Manuscript in preparation.
- Starobinsky, A. A. (1980). A new type of isotropic cosmological models without singularity. *Phys. Lett. B*, 91, 99–102.
- Planck Collaboration (2020). Planck 2018 results. X. Constraints on inflation. *Astron. Astrophys.*, 641, A10.
- Mukhanov, V. F. & Chibisov, G. V. (1981). Quantum fluctuations and a nonsingular universe. *JETP Lett.*, 33, 532–535.
- Mukhanov, V. F. & Chibisov, G. V. (1981). The vacuum energy and large scale structure of the universe. *Sov. Phys. JETP*, 56, 258–265.
- Hawking, S. W. (1982). The development of irregularities in a single bubble inflationary universe. *Phys. Lett. B*, 115, 295–297.
- Guth, A. H. (1981). Inflationary universe: A possible solution to the horizon and flatness problems. *Phys. Rev. D*, 23, 347–356.
- Linde, A. D. (1982). A new inflationary universe scenario: A possible solution of the horizon, flatness, homogeneity, isotropy and primordial monopole problems. *Phys. Lett. B*, 108, 389–393.
- Penrose, R. (1989). *The Emperor’s New Mind*. Oxford University Press.
- Brandenberger, R. (2017). Initial conditions for inflation—A short review. *Int. J. Mod. Phys. D*, 26, 1740002.
- Birrell, N. D. & Davies, P. C. W. (1982). *Quantum Fields in Curved Space*. Cambridge University Press.
- Spencer-Brown, G. (1969). *Laws of Form*. Allen & Unwin.
- Bateson, G. (1972). *Steps to an Ecology of Mind*. University of Chicago Press.
- Stelle, K. S. (1977). Renormalization of higher-derivative quantum gravity. *Phys. Rev. D*, 16, 953–969.
- Barvinsky, A. O. & Vilkovisky, G. A. (1985). The generalized Schwinger–DeWitt technique in gauge theories and quantum gravity. *Phys. Rep.*, 119, 1–74.

- Whitt, B. (1984). Fourth-order gravity as general relativity plus matter. *Phys. Lett. B*, 145, 176–178.
- Maeda, K. (1988). Inflation as a transient attractor in R^2 cosmology. *Phys. Rev. D*, 37, 858–862.
- Barrow, J. D. & Cotsakis, S. (1988). Inflation and the conformal structure of higher-order gravity theories. *Phys. Lett. B*, 214, 515–518.
- Mukhanov, V. F. (1988). Quantum theory of gauge-invariant cosmological perturbations. *Sov. Phys. JETP*, 67, 1297–1302.
- Bardeen, J. M., Steinhardt, P. J. & Turner, M. S. (1983). Spontaneous creation of almost scale-free density perturbations in an inflationary universe. *Phys. Rev. D*, 28, 679–693.
- Planck Collaboration (2020). Planck 2018 results. VII. Isotropy and statistics of the CMB. *Astron. Astrophys.*, 641, A7.
- Planck Collaboration (2020). Planck 2018 results. IX. Constraints on primordial non-Gaussianity. *Astron. Astrophys.*, 641, A9.
- Unruh, W. G. (1981). Experimental black-hole evaporation? *Phys. Rev. Lett.*, 46, 1351–1353.
- Codello, A., Percacci, R. & Rahmede, C. (2009). Investigating the ultraviolet properties of gravity with a Wilsonian renormalization group equation. *Ann. Phys.*, 324, 414–469.
- Falls, K., Litim, D. F., Nikolakopoulos, K. & Rahmede, C. (2014). A bootstrap towards asymptotic safety. arXiv:1301.4191.
- Volovik, G. E. (2003). *The Universe in a Helium Droplet*. Oxford University Press.
- DESI Collaboration (2024). DESI 2024 VI: Cosmological constraints from the measurements of baryon acoustic oscillations. arXiv:2404.03002.
- LiteBIRD Collaboration (2023). Probing cosmic inflation with the LiteBIRD satellite. *Prog. Theor. Exp. Phys.*, 2023, 042F01.
- CMB-S4 Collaboration (2019). CMB-S4 Science Case, Reference Design, and Project Plan. arXiv:1907.04473.
- Sakharov, A. D. (1967). Vacuum quantum fluctuations in curved space and the theory of gravitation. *Sov. Phys. Dokl.*, 12, 1040–1041.
- Zel’dovich, Ya. B. (1967). Cosmological constant and elementary particles. *JETP Lett.*, 6, 316–317.
- Brown, D. (2026). Quantum-Kinetic Dark Energy (QKDE): An effective dark energy framework with a covariantly completed time-dependent scalar kinetic normalization. *Int. J. Mod. Phys. D*, 35(4), 2650006.

- Planck Collaboration (2020). Planck 2018 results. VI. Cosmological parameters. *Astron. Astrophys.*, 641, A6.
- Weinberg, S. (1989). The cosmological constant problem. *Rev. Mod. Phys.*, 61, 1–23.
- Kehagias, A., Moradinezhad Dizgah, A. & Riotto, A. (2014). Remarks on the Starobinsky model of inflation and its descendants. *Phys. Rev. D*, 89, 043527.

A Computational Code

The following Python code was used to generate Figures 1–4. The complete code, including all four figure-generation routines, is available as a supplementary file (`generate_figures.py`). Here we reproduce the core routines for the slow-roll dynamics and the flatness feedback computation.

```
import numpy as np
from scipy.integrate import solve_ivp

# Starobinsky potential (Planck units, M_Pl = 1)
alpha = np.sqrt(2.0/3.0)
V0 = 1.0 # normalized

def V(phi):
    return V0 * (1 - np.exp(-alpha * phi))**2

def dVdphi(phi):
    return 2*V0*alpha*(1 - np.exp(-alpha*phi))*np.exp(-alpha*phi)

def d2Vdphi2(phi):
    return 2*V0*alpha**2*np.exp(-alpha*phi)*(
        2*np.exp(-alpha*phi) - 1)

# Slow-roll parameters
def epsilon(phi):
    return 0.5 * (dVdphi(phi) / V(phi))**2

def eta_sr(phi):
    return d2Vdphi2(phi) / V(phi)

# Slow-roll: dphi/dN = -V'/V
def dphi_dN(N, phi):
    return -dVdphi(phi) / V(phi) if V(phi) > 1e-10 else 0

# Solve for 60+ e-folds from phi_i = 5.5 M_Pl
phi_i = 5.5
sol = solve_ivp(dphi_dN, [0, 70], [phi_i],
    t_eval=np.linspace(0, 70, 1000), rtol=1e-10)
N_arr, phi_arr = sol.t, sol.y[0]
eps_arr = np.array([epsilon(p) for p in phi_arr])

# End of inflation: epsilon = 1
idx_end = np.argmin(np.abs(eps_arr - 1.0))
N_end = N_arr[idx_end]
# Result: N_end ~ 62.5 e-folds
```

```
# Predictions at N* e-folds before end
# n_s = 1 - 2/N*, r = 12/N*^2
for Ns in [50, 55, 60]:
    print(f"N*={Ns}: n_s={1-2/Ns:.4f}, r={12/Ns**2:.5f}")

# Flatness feedback: beta(a)
# beta = xi^2 * rho_m / (m_psi^2 + xi^2 * rho_m)
def beta(a, xi=1e-3, m_psi_over_H0=1.0):
    Omega_m0, H0 = 0.315, 1.0 # normalized
    rho_m = Omega_m0 * H0**2 * a**(-3) * 3/(8*np.pi)
    m2 = m_psi_over_H0**2 * H0**2
    return xi**2 * rho_m / (m2 + xi**2 * rho_m)
```

Peer Reviews and Revision History

First Round

Reviewer assessment. The reviewer found the paper to occupy a pivotal position in the research programme, delivering on all three promissory notes deferred from Paper 5 ($V(\psi)$, $\beta > 0$, connection to inflation). The writing was praised as clear, the logical architecture well-signposted, and the counterargument sections described as “one of the programme’s genuine strengths.” The $V(\psi)$ derivation was called “the strongest result in the programme so far,” and the three-dimensional elastic analogy was judged “significantly improved” over Paper 5’s membrane model. The flatness feedback derivation was recognized as “a meaningful step forward” fulfilling Paper 5’s central open promise.

Six major concerns were raised: (1) the interpretive claim outruns the derivation—the word “unique” in Proposition 5.1 overstates the result given the R^2 truncation; (2) the mass hierarchy (10^{55} between inflationary and late-time sectors) is a potential fatal inconsistency, not merely a limitation; (3) the inhomogeneity argument mixes rigour levels—the structural argument rests on an unpublished philosophical manuscript; (4) the suppression of primordial inhomogeneities is stated but not computed; (5) the quasi-static approximation in the β derivation may not be self-consistent with Paper 5’s $m_\psi \sim H_0$ requirement; (6) the observational predictions are entirely inherited from Starobinsky inflation and should not be presented as new.

Six minor concerns were raised: (A) the injection model introduces a preferred spatial point (center); (B) self-citation density; (C) the “Recommended Journal” section is unusual for a research manuscript; (D) the conformal transformation omits intermediate steps; (E) the appendix code is incomplete; (F) “no new particles” is inconsistent with Paper 5’s revised language.

Verdict: Major revision required.

Revisions Made

1. **Proposition 5.1 restated (Concern 1).** The word “unique” has been replaced throughout by “leading-order, within the R^2 truncation.” The truncation assumption is now stated explicitly in the proposition itself, not only in the counterargument. The Discussion now separates what is derived (the Starobinsky potential from $R + R^2$, via standard conformal transformation) from what is interpreted (the elastic recoil picture).
2. **Mass hierarchy elevated to dedicated subsection (Concern 2).** A new subsection §6.5 (“The Mass Hierarchy: What Is Unified and What Is Not”) replaces the one-paragraph treatment in the Limitations section. The subsection states explicitly that the programme achieves *unification of interpretation* but not *unification of dynamics*, that asymptotic safety results show running of $\mathcal{O}(1)$ – $\mathcal{O}(10)$ rather than the required 10^{55} , and that three possible resolution paths are identified without claiming any is demonstrated.

3. **Section 3 restructured (Concern 3).** The structural argument has been demoted from a Postulate to a Remark. The quantum-mechanical argument (§3.1) now carries the primary physical weight. The Section Result explicitly states that the QM argument establishes the physical impossibility; the structural observation illuminates the conceptual status but is not invoked as a premise of subsequent derivations.
4. **Quantitative suppression estimate added (Concern 4).** A new subsection “Quantitative Estimate” in Section 7 computes the ratio $\zeta_*^{(\text{init})}/\zeta_*^{(\text{QM})}$ for Planck-amplitude initial perturbations. The result: for $N_{\text{tot}} \geq 69$ e-folds, even the most extreme initial perturbations are subdominant on CMB scales. The marginal case ($N_{\text{tot}} \approx 65\text{--}70$) is where residual effects might appear at $\ell \lesssim 10$.
5. **Quasi-static limit made explicit (Concern 5).** Proposition 8.1 has been retitled “Positive feedback parameter—quasi-static limit.” A new paragraph following the proof states the conditions under which the approximation breaks down ($m_\psi \sim H_0$), affirms that the sign of β is robust (from convexity), and identifies numerical confirmation as a priority for Paper 7.
6. **Section 9 retitled and restructured (Concern 6).** Now titled “Observational Consistency and Predictions,” with explicit subsection labels “Inherited Predictions” (Starobinsky n_s , r ; late-time $w \neq -1$) and “New Predictions” (large-scale CMB anomalies; non-Gaussianity from pre-inflationary inhomogeneities). The counterargument response distinguishes observational consistency checks from genuinely new physical content.
7. **Center-of-injection limitation acknowledged (Minor A).** A new counterargument in Section 4 addresses the preferred spatial point: the injection picture is isotropic from one point but not homogeneous, and should be understood as illustrating dynamics at any single comoving point.
8. **Conformal transformation expanded (Minor D).** Section 5.3 now presents the full derivation in four labelled steps: auxiliary field introduction, conformal transformation, scalar field redefinition, and Einstein frame action.
9. **Appendix code completed (Minor E).** The code now includes the slow-roll parameter computation, end-of-inflation identification, observational predictions, and the flatness feedback function $\beta(a)$. A note directs readers to the supplementary file for the complete figure-generation code.
10. **“No new particles” softened (Minor F).** The abstract, conclusions, and limitations section now state: “The screening scalar field ψ is a new degree of freedom—we do not deny this—but it is not a dark matter candidate; it is a mediator of the vacuum’s elastic response.” A new paragraph in Limitations explicitly acknowledges ψ as a new scalar degree of freedom.

Items Retained Without Change

1. The “Recommended Journal” section and “Peer Reviews” appendix are part of the author’s standard manuscript template and will be removed upon journal submission if required by editorial policy (Minor C).
2. Self-citation density (Minor B) has been reduced through Discussion edits but remains necessarily high given the programme structure. Many cross-references are required for coherence across the seventeen-paper programme.