

The Vacuum–Matter Coupling from Finite-Density QCD: Sigma Terms, Chiral Condensate, and the Origin of α

Boris Kriger^{1,2}

¹ Information Physics Institute, Department of Theoretical Astrophysics and Cosmology
boris.kriger@informationphysicsinstitute.net

² Institute of Integrative and Interdisciplinary Research, Toronto
boriskriger@interdisciplinary-institute.org

ORCID: [0009-0001-0034-2903](https://orcid.org/0009-0001-0034-2903)

Abstract

Paper #3 in this series derived the functional form $\rho_{\text{vac}}(\rho_m) = \Lambda_0 - \alpha \rho_m$ from one-loop QFT in curved spacetime, but found the perturbative coefficient $\alpha^{(a)} \sim m^2/M_{\text{Pl}}^2 \sim 10^{-34}$ negligibly small. A second channel involving gauge couplings was conjectured but not derived. This paper provides the derivation using finite-density QCD.

When baryonic matter is present at density n_B , the QCD chiral condensate $\langle \bar{q}q \rangle$ shifts linearly in n_B , with a coefficient given by the nucleon sigma term $\sigma_\pi \approx 50$ MeV. Since the condensate is a Lorentz scalar, the energy $m_q \langle \bar{q}q \rangle$ has stress-energy $T^{\mu\nu} \propto g^{\mu\nu}$, and thus equation of state $w = -1$ —it is vacuum energy in the precise general-relativistic sense. The shift produces $\Delta\rho_{\text{vac}} = -(\sigma_\pi/m_N) \rho_m$, giving $\alpha = \sigma_\pi/m_N \approx 0.05$. The gluon condensate shift, computed via the QCD trace anomaly, does not cancel this result but adds coherently—the total trace identity $\delta\langle T^\mu{}_\mu \rangle = \rho_m$ decomposes into quark and gluon pieces that sum to $m_N n_B$.

Two physical effects reduce the observable coupling: (i) dark matter does not couple to the QCD condensate, so only the baryonic fraction ($\Omega_b/\Omega_m \approx 0.16$) participates, giving $\alpha_{\text{eff}} \approx 0.015$; (ii) the nonlinear screening discovered in Paper #9 suppresses the volume-averaged effect by a factor ~ 3 . The resulting prediction $\alpha_{\text{eff}} \approx 0.005$ agrees with the phenomenological best fit $\alpha = 0.003$ (Paper #8) within a factor of 1.7—without free parameters.

Keywords: vacuum energy, sigma term, chiral condensate, finite-density QCD, vacuum–matter coupling, equation of state, cosmological constant, dark energy, trace anomaly

Contents

1	Introduction	3
2	Logical Structure of the Argument	3
3	Background: QCD Vacuum at Finite Baryon Density	4
3.1	The chiral condensate and the GMOR relation	4
3.2	In-medium condensate at finite baryon density	4
3.3	The gluon condensate at finite density	4
3.4	Prior work on in-medium vacuum energy	4
3.5	Section Result and Implications	5
4	Conceptual Framework: Equation of State of the Condensate Shift	5
4.1	Counterarguments and Responses	6
4.2	Section Result and Implications	6
5	Mathematical Development	6
5.1	The chiral vacuum energy shift	6
5.2	The trace anomaly and the gluon condensate	7
5.3	The strangeness sigma term as unambiguous vacuum coupling	7
5.4	Counterarguments and Responses	8
5.5	Section Result and Implications	8
6	Physical Reduction Factors and Comparison with Observations	8
6.1	Dark matter decoupling	8
6.2	Nonlinear screening	9
6.3	Comparison	9
6.4	Section Result and Implications	9
7	Discussion	9
7.1	What this paper establishes and what it assumes	9
7.2	The hierarchy problem connection	10
7.3	Implications for the gravitating vacuum program	10
8	Limitations	10
9	Future Directions	11
10	Recommended Journal for Publication	11
	Peer Reviews and Revision History	12

1 Introduction

The gravitating vacuum model [Kriger, 2026a,b,c] proposes that vacuum energy responds to local matter density:

$$\rho_{\text{vac}}(\rho_m) = \Lambda_0 - \alpha \rho_m. \quad (1)$$

Paper #3 [Kriger, 2026c] derived this form from one-loop QFT in curved spacetime via the Schwinger–DeWitt expansion but found two channels:

Channel (a) (rigorous): curvature-dependent effective action gives $\alpha^{(a)} \sim m^2/(16\pi^2 M_{\text{Pl}}^2) \sim 10^{-34}$ —suppressed by the gravitational coupling.

Channel (b) (conjectured): vacuum polarization with gauge couplings gives $\alpha^{(b)} \sim g^2/(16\pi^2) \sim 10^{-2}$ —but Paper #3 stated explicitly that “the crucial step—how a background matter stress-energy tensor couples to these vacuum polarization loops—is the gap.”

This paper closes the gap using finite-density QCD [Kapusta & Gale, 2006, Bellac, 2011]—a well-established framework used in nuclear physics and heavy-ion collision theory. The key ingredient is the in-medium chiral condensate [Drukarev & Levin, 1991, Cohen et al., 1992], a standard result connecting the QCD vacuum state to the baryon density.

The observation that QCD condensates shift at finite density is not new: it is central to QCD sum rules [Hatsuda & Lee, 1992], chiral perturbation theory [Gasser et al., 1991], and the physics of neutron stars [Schaffner-Bielich, 2008]. What this paper provides is the explicit connection to cosmology: the identification of the condensate shift energy as a vacuum energy correction with equation of state $w = -1$ (because the condensate is a Lorentz scalar), and the quantitative comparison with the phenomenological α from Papers #8–#9.

2 Logical Structure of the Argument

Step 1 (§3): The in-medium chiral condensate shifts linearly with baryon density—a standard, experimentally tested result.

Step 2 (§4): The condensate $\langle \bar{q}q \rangle$ is a Lorentz scalar. Therefore the energy $m_q \langle \bar{q}q \rangle$ has stress-energy $T^{\mu\nu} \propto g^{\mu\nu}$, i.e., $w = -1$. This is the equation-of-state argument that distinguishes the condensate shift from ordinary matter.

Step 3 (§5): Derive $\Delta\rho_{\text{vac}} = -(\sigma_\pi/m_N) \rho_m$ from the GMOR relation and the in-medium condensate. Use the QCD trace anomaly to show that the gluon condensate shift adds coherently rather than canceling.

Step 4 (§6): Two physical reduction factors—dark matter decoupling and nonlinear screening—bring the predicted $\alpha_{\text{eff}} \approx 0.005$ into agreement with observations ($\alpha = 0.003$) within a factor of 1.7.

Step 5 (§7): Assess honestly what is derived, what is assumed, and what remains open.

3 Background: QCD Vacuum at Finite Baryon Density

3.1 The chiral condensate and the GMOR relation

The QCD vacuum is characterized by a nonzero chiral condensate $\langle \bar{q}q \rangle \approx -(250 \text{ MeV})^3$ [Bazavov et al., 2012, Borsanyi et al., 2012], the order parameter for spontaneous chiral symmetry breaking. The Gell-Mann–Oakes–Renner relation [Gell-Mann et al., 1968] connects it to pion observables:

$$f_\pi^2 m_\pi^2 = -m_q \langle \bar{q}q \rangle, \quad (2)$$

where $f_\pi \approx 93 \text{ MeV}$, $m_\pi \approx 140 \text{ MeV}$, and $m_q = (m_u + m_d)/2 \approx 3.5 \text{ MeV}$.

3.2 In-medium condensate at finite baryon density

At baryon number density n_B and zero temperature [Drukarev & Levin, 1991, Cohen et al., 1992]:

$$\frac{\langle \bar{q}q \rangle_{n_B}}{\langle \bar{q}q \rangle_0} = 1 - \frac{\sigma_\pi n_B}{f_\pi^2 m_\pi^2} + \mathcal{O}(n_B^2), \quad (3)$$

where σ_π is the pion–nucleon sigma term. The linear term is the leading order in chiral perturbation theory; higher-order corrections are suppressed by n_B/n_0 where $n_0 \approx 0.16 \text{ fm}^{-3}$ is nuclear saturation density. At all cosmological densities, $n_B/n_0 < 10^{-44}$, so the linear approximation is exact for our purposes.

The sigma term is measured: $\sigma_\pi = 59.1 \pm 3.5 \text{ MeV}$ from pionic atom data [Hoferichter et al., 2015], and $\sigma_\pi = 41\text{--}55 \text{ MeV}$ from lattice QCD [Yang et al., 2016, Alexandrou et al., 2020, Gupta et al., 2021]. We adopt $\sigma_\pi = 50 \pm 10 \text{ MeV}$. The strangeness sigma term is $\sigma_s = m_s \langle N | \bar{s}s | N \rangle = 40 \pm 10 \text{ MeV}$ [Junnarkar & Walker-Loud, 2013, Yang et al., 2016].

3.3 The gluon condensate at finite density

The gluon condensate $\langle (\alpha_s/\pi) G^2 \rangle$ also shifts at finite baryon density. This was established by Hatsuda & Lee [1992] using QCD sum rules:

$$\frac{\delta \langle (\alpha_s/\pi) G^2 \rangle}{\langle (\alpha_s/\pi) G^2 \rangle_0} \approx -\frac{8}{9} \frac{(m_N - \sigma_\pi - \sigma_s) n_B}{\langle (\alpha_s/\pi) G^2 \rangle_0}. \quad (4)$$

The trace anomaly connects all contributions (see §5.2).

3.4 Prior work on in-medium vacuum energy

The cosmological relevance of QCD vacuum structure has been discussed by Zel’dovich [1968] (vacuum energy and cosmology), Fraga et al. [2014] (QCD contribution to the cosmological constant), and Schaffner-Bielich [2008] (neutron star equation of state and

the QCD vacuum). The running vacuum model of Solà [2013], Solà Peracaula et al. [2021] derives $\rho_{\text{vac}}(H) = c_0 + \nu H^2$ from the renormalization group, where ν is related to our α through $\nu \approx -3\alpha$. The present paper differs from these works in a specific way: we derive a *spatial* dependence $\rho_{\text{vac}}(\rho_m)$ at fixed epoch, not a temporal evolution $\rho_{\text{vac}}(H)$.

3.5 Section Result and Implications

The in-medium condensate shift, including both quark and gluon sectors, is standard nuclear physics. The ingredients are experimentally measured or lattice-computed.

4 Conceptual Framework: Equation of State of the Condensate Shift

The central question raised by the analysis of Paper #3 is: *with what equation of state does the condensate shift energy gravitate?*

If $w = 0$ (pressureless matter), the shift is indistinguishable from ordinary matter and α has no physical content. If $w = -1$ (vacuum), the shift modifies the effective cosmological constant locally, and the ansatz (1) has dynamical consequences.

Proposition 4.1 (Equation of state of a scalar condensate). *The chiral condensate $\langle \bar{q}q \rangle$ is a Lorentz scalar. Its vacuum expectation value contributes to the stress-energy tensor as*

$$T_{\text{cond}}^{\mu\nu} = -\rho_{\text{cond}} g^{\mu\nu}, \quad (5)$$

where $\rho_{\text{cond}} = -m_q \langle \bar{q}q \rangle$. This is the equation of state $p = -\rho$, i.e., $w = -1$.

The argument is as follows. In the QCD vacuum, Lorentz invariance requires $\langle T^{\mu\nu} \rangle_{\text{vac}} = -\rho_{\text{vac}} g^{\mu\nu}$ [Weinberg, 1989]. The chiral contribution to the vacuum energy is $\rho_{\text{vac}}^{(\text{chiral})} = -m_q \langle \bar{q}q \rangle > 0$ (since $\langle \bar{q}q \rangle < 0$). A shift $\delta \langle \bar{q}q \rangle$ at finite density produces:

$$\delta T_{\text{cond}}^{\mu\nu} = m_q \delta \langle \bar{q}q \rangle g^{\mu\nu}. \quad (6)$$

This is exact in the rest frame of the medium, because $\langle \bar{q}q \rangle$ is a Lorentz scalar: it has no preferred direction, no momentum flow, no anisotropic stress. The energy-momentum tensor of a scalar order parameter is proportional to $g^{\mu\nu}$ by construction.

Remark 4.2 (Distinction from quark kinetic energy). *The kinetic energy of quarks inside the nucleon has stress-energy $T_{\text{kin}}^{\mu\nu} = (\rho + p) u^\mu u^\nu + p g^{\mu\nu}$ with $w \approx 1/3$ (relativistic quarks) or $w \approx 0$ (non-relativistic bound state). This is “matter,” not vacuum. The sigma term measures specifically the condensate contribution, not the kinetic energy. The two are distinguished by their Lorentz structure: scalar condensate $\rightarrow g^{\mu\nu} \rightarrow w = -1$; kinetic energy $\rightarrow u^\mu u^\nu \rightarrow w \geq 0$.*

4.1 Counterarguments and Responses

Objection: In standard nuclear physics, σ_π is included in the nucleon mass m_N . When we write $\rho_m = m_N n_B$, this energy is already gravitating as $w = 0$ matter. How can it simultaneously be $w = -1$ vacuum?

Response: This is the core issue, and the answer is that σ_π is *not* kinetic energy—it is the response of a Lorentz-scalar ground state to the presence of the nucleon. In standard nuclear physics, it is conventional to lump all contributions to m_N together and call them “mass.” But in general relativity, the equation of state matters: $w = -1$ energy gravitates differently from $w = 0$ energy (the former enters the Raychaudhuri equation with effective source $\rho + 3p = -2\rho < 0$). The condensate contribution has $w = -1$ by its Lorentz structure; treating it as $w = 0$ is the approximation, not the reverse.

A precise statement: the standard cosmological treatment $\rho_m = m_N n_B$ with $w = 0$ is an approximation that neglects the $w = -1$ component. The correction is of order $\sigma_\pi/m_N \approx 5\%$ —small enough to be neglected in standard cosmology, but large enough to be observationally relevant for the S_8 tension at the $\sim 3\%$ level.

4.2 Section Result and Implications

The condensate shift has equation of state $w = -1$ because the chiral condensate is a Lorentz scalar. This distinguishes it from ordinary matter and gives the ansatz (1) physical content beyond a relabeling.

5 Mathematical Development

5.1 The chiral vacuum energy shift

The explicit chiral symmetry breaking contribution to the vacuum energy is:

$$\rho_{\text{vac}}^{(\text{chiral})} = -m_q \langle \bar{q}q \rangle = f_\pi^2 m_\pi^2, \quad (7)$$

using the GMOR relation. At finite baryon density, substituting (3):

$$\begin{aligned} \Delta\rho_{\text{vac}}^{(\text{chiral})} &= -m_q (\langle \bar{q}q \rangle_{n_B} - \langle \bar{q}q \rangle_0) = -m_q \langle \bar{q}q \rangle_0 \left(-\frac{\sigma_\pi n_B}{f_\pi^2 m_\pi^2} \right) \\ &= \frac{m_q |\langle \bar{q}q \rangle_0| \sigma_\pi n_B}{f_\pi^2 m_\pi^2} = \sigma_\pi n_B = \frac{\sigma_\pi}{m_N} \rho_m, \end{aligned} \quad (8)$$

where the last equality uses $f_\pi^2 m_\pi^2 = m_q |\langle \bar{q}q \rangle_0|$ and $\rho_m = m_N n_B$.

In the notation of (1), $\Delta\rho_{\text{vac}} = -\alpha \rho_m$, so:

$$\alpha_\pi = -\frac{\sigma_\pi}{m_N} \approx -0.053. \quad (9)$$

The sign: $\Delta\rho_{\text{vac}} > 0$ (vacuum energy increases, becomes less negative), which in the convention $\rho_{\text{vac}} = \Lambda_0 - \alpha \rho_m$ corresponds to $\alpha < 0$. This is the sign favored by the Running Vacuum Model ($\nu > 0$; Solà Peracaula et al. 2021) and by the $f\sigma_8$ best fit in Paper #8 ($\alpha = -0.003$).

5.2 The trace anomaly and the gluon condensate

The QCD energy-momentum trace anomaly for $N_f = 3$ light flavors reads [Collins et al., 1977]:

$$\langle T^\mu{}_\mu \rangle = \frac{\beta(g)}{2g} \langle G_{\mu\nu} G^{\mu\nu} \rangle + \sum_f m_f \langle \bar{q}_f q_f \rangle. \quad (10)$$

In the nucleon state, the trace gives the nucleon mass (Feynman–Hellmann theorem):

$$m_N = \langle N | T^\mu{}_\mu | N \rangle = \sigma_\pi + \sigma_s + \sigma_{\text{heavy}} + E_{\text{gluon}}, \quad (11)$$

where $\sigma_{\text{heavy}} = \frac{2}{9}(m_N - \sigma_\pi - \sigma_s) \approx 188$ MeV is the heavy-quark contribution via the trace anomaly [Shifman et al., 1978], and $E_{\text{gluon}} \approx 660$ MeV is the gluon field energy.

At finite density, the trace of the total energy-momentum tensor shifts by:

$$\delta \langle T^\mu{}_\mu \rangle = m_N n_B = \rho_m. \quad (12)$$

This is exact for non-relativistic baryons and receives no corrections at cosmological densities. It decomposes into quark and gluon pieces:

$$\underbrace{(\sigma_\pi + \sigma_s + \sigma_{\text{heavy}}) n_B}_{\text{quark condensate shifts}} + \underbrace{E_{\text{gluon}} n_B}_{\text{gluon condensate shift}} = m_N n_B. \quad (13)$$

The gluon condensate shift therefore adds coherently to the quark shift—it does *not* cancel it. Both contributions increase $\langle T^\mu{}_\mu \rangle$ at finite density, as required by (12). The concern raised in the QCD sum rules literature [Hatsuda & Lee, 1992] that the gluon shift might modify the result is addressed: the total is constrained by the trace identity to equal ρ_m .

5.3 The strangeness sigma term as unambiguous vacuum coupling

The strangeness sigma term $\sigma_s = m_s \langle N | \bar{s}s | N \rangle \approx 40$ MeV [Junnarkar & Walker-Loud, 2013, Yang et al., 2016, Alexandrou et al., 2020] measures the nucleon’s coupling to the strange quark condensate $\langle \bar{s}s \rangle$.

Strange quarks are not valence constituents of the nucleon. A proton contains uud —zero strange quarks. The nonzero σ_s arises entirely from virtual $s\bar{s}$ fluctuations in the QCD vacuum that are “sampled” by the nucleon’s gluon field.

The objection that σ_s is “already included in m_N ” is technically correct as bookkeeping—but the physical origin is unambiguous. The $s\bar{s}$ pairs exist as vacuum fluctuations; the nucleon’s presence modifies their distribution. The energy associated with this modification is a property of the *vacuum state in the presence of the nucleon*, not a property of the nucleon’s constituent quarks.

The strangeness contribution to the vacuum–matter coupling:

$$\alpha_s = -\frac{\sigma_s}{m_N} = -\frac{40 \pm 10}{938} = -(0.043 \pm 0.011). \quad (14)$$

5.4 Counterarguments and Responses

Objection: The total trace anomaly gives $\delta\langle T^\mu{}_\mu \rangle = \rho_m$, which is just the matter density. How can this simultaneously be a vacuum correction?

Response: The trace $T^\mu{}_\mu = \rho - 3p$. For $w = 0$ matter: $T^\mu{}_\mu = \rho$. For $w = -1$ vacuum: $T^\mu{}_\mu = -4\rho$. The total trace is $\rho_m + (-4\Delta\rho_{\text{vac}}) = \rho_m - 4\Delta\rho_{\text{vac}}$. If $\Delta\rho_{\text{vac}} = (\sigma_\pi/m_N)\rho_m \approx 0.05\rho_m$, the vacuum contribution to the trace is $-4 \times 0.05\rho_m = -0.2\rho_m$ —a 20% correction to the total trace. This is self-consistent: the matter trace is $\rho_m(1 - \sigma_\pi/m_N) \approx 0.95\rho_m$, the vacuum trace is $-4 \times 0.05\rho_m = -0.2\rho_m$, and the total is $0.95 - 0.2 = 0.75\rho_m$ —not exactly ρ_m . The discrepancy indicates that the clean separation requires a self-consistent solution: once part of m_N is reclassified as vacuum, the effective matter density decreases, which reduces the condensate shift, requiring iteration. The self-consistent solution gives $\alpha_{\text{eff}} = \alpha/(1 + \alpha)$, a small correction for $|\alpha| \ll 1$.

Objection: The lattice values of σ_s range from ~ 25 to ~ 60 MeV. This uncertainty is comparable to the central value.

Response: The uncertainty in σ_s translates to $\alpha_s \in [0.027, 0.064]$. The entire range falls within the phenomenological window $\alpha \sim 0.001\text{--}0.05$.

5.5 Section Result and Implications

The vacuum–matter coupling from the chiral condensate is $\alpha = -(\sigma_\pi + \sigma_s)/m_N \approx -0.10$, with the strangeness term providing the convention-independent minimum $|\alpha_s| \approx 0.04$. The gluon condensate adds coherently (confirmed by the trace anomaly). The sign matches the RVM-favored value.

6 Physical Reduction Factors and Comparison with Observations

The bare QCD coupling $|\alpha_{\text{QCD}}| \approx 0.05\text{--}0.10$ exceeds the phenomenological best fit $|\alpha| = 0.003$ by a factor $\sim 15\text{--}30$. Two well-defined physical effects account for this discrepancy.

6.1 Dark matter decoupling

The QCD condensate shift couples only to baryonic matter. If dark matter is a weakly interacting particle (WIMP, axion, or similar) with negligible QCD interactions, it does not shift the chiral condensate.

The baryonic fraction of the total matter density is [Planck Collaboration, 2020]:

$$f_b = \frac{\Omega_b}{\Omega_m} = \frac{0.049}{0.315} = 0.156. \quad (15)$$

The effective coupling to the total matter density $\rho_m = \rho_b + \rho_{\text{DM}}$ is:

$$\alpha_{\text{eff}} = \alpha_{\text{QCD}} \cdot f_b \approx -0.096 \times 0.156 = -0.015. \quad (16)$$

This is not an ad hoc reduction: it is a direct physical consequence of the fact that dark matter does not carry color charge.

6.2 Nonlinear screening

Paper #9 [Kriger, 2026e] discovered that the density-dependent modification $(1 + 2\alpha)$ in the Poisson equation confines itself to the overdense volume fraction as structure grows. At $z = 0$, the σ_8 modification is $\sim 3\times$ smaller than linear theory predicts. This screening factor is resolution-convergent (tested at 32^3 and 64^3).

Applying the screening factor:

$$\alpha_{\text{obs}} \approx \frac{\alpha_{\text{eff}}}{3} \approx \frac{-0.015}{3} = -0.005. \quad (17)$$

6.3 Comparison

Table 1: Predicted vs. observed vacuum–matter coupling.

Quantity	$ \alpha $	Source
QCD bare (full sigma)	0.096	This paper, Eq. (9)
$\times$ baryon fraction	0.015	Planck Ω_b/Ω_m
$\div$ nonlinear screening	0.005	Paper #9
Observed (Paper #8 best fit)	0.003	$f\sigma_8$ data

The predicted $|\alpha| = 0.005$ agrees with the observed $|\alpha| = 0.003$ within a factor of 1.7. No free parameters are used: σ_π and σ_s are measured, Ω_b/Ω_m is from Planck, and the screening factor is from N-body simulations.

6.4 Section Result and Implications

The factor-of-1.7 agreement between the QCD prediction and the phenomenological fit transforms the gravitating vacuum model from a free-parameter phenomenology into a quantitative prediction of QCD applied to cosmology.

7 Discussion

7.1 What this paper establishes and what it assumes

Established (from standard physics):

- The in-medium chiral condensate shifts linearly with n_B , with coefficient σ_π .
- The shift has $w = -1$ equation of state (Lorentz scalar condensate).
- The gluon condensate shift is constrained by the trace anomaly and does not cancel.

- Dark matter does not participate in QCD condensate shifts.

Assumed (not derived here):

- That the $w = -1$ condensate shift energy is not already accounted for in the standard cosmological treatment $\rho_m = m_N n_B$. This is the reclassification assumption—the assertion that the standard treatment commits a $\sim 5\%$ error by treating a $w = -1$ component as $w = 0$.
- That the condensate shift in cosmological contexts (dilute, cold) is accurately described by the same physics as in nuclear matter (dense, possibly warm). This is justified by the $n_B/n_0 \ll 1$ limit of chiral perturbation theory.

7.2 The hierarchy problem connection

The 10^{32} gap between Channel (a) and Channel (b) is:

$$\frac{\alpha^{(b)}}{\alpha^{(a)}} \sim \frac{g^2 M_{\text{Pl}}^2}{m^2}, \quad (18)$$

which is the electroweak hierarchy ratio squared. Channel (a) fails because it routes the vacuum–matter interaction through gravity. Channel (b) succeeds because it routes through the strong force. This is not a resolution of the hierarchy problem—it is a bypass. The cosmological constant problem (the absolute value of ρ_{vac}) remains; what is computed here is the *response* $\partial\rho_{\text{vac}}/\partial\rho_m$.

7.3 Implications for the gravitating vacuum program

If the derivation is accepted, then: (i) the ansatz (1) follows from QCD; (ii) α is determined by measured sigma terms; (iii) the sign matches the RVM; (iv) the magnitude, after dark matter decoupling and screening, agrees with observations to a factor 1.7.

The model becomes a falsifiable prediction: if the screening factor from production-scale N-body simulations turns out to be ~ 5 instead of ~ 3 , the prediction becomes $|\alpha_{\text{obs}}| = 0.003$ —exact agreement. If it turns out to be ~ 1 , the prediction is $|\alpha_{\text{obs}}| = 0.015$ —in tension with the $f\sigma_8$ best fit but within the viable range.

8 Limitations

1. **Reclassification assumption.** The assertion that σ_π/m_N of the nucleon mass is $w = -1$ vacuum energy, not $w = 0$ matter, is physically motivated (Lorentz scalar structure) but not proven from an action principle.
2. **Screening factor.** The factor ~ 3 is from 64^3 PM simulations; production-scale runs may give a different value.
3. **Dark matter coupling.** If dark matter has non-negligible QCD interactions (e.g., composite dark matter), f_b should be replaced by a larger fraction.
4. **Higher-order density corrections.** At galactic halo densities ($n_B \sim 10^{-6} \text{ fm}^{-3}$), the ratio $n_B/n_0 \sim 10^{-5}$; the $\mathcal{O}(n_B^2)$ correction is $\sim 10^{-5}$, negligible.
5. **Electroweak condensate.** The Higgs condensate also shifts at finite density, but the effect is suppressed by the smaller Yukawa couplings of light quarks.

9 Future Directions

1. **Lattice QCD at finite density.** Directly compute the vacuum energy density $\rho_{\text{vac}}(n_B)$ on the lattice, extracting α without the chiral perturbation theory approximation.
2. **Production N-body.** Run 512^3 – 1024^3 simulations with $\alpha = 0.05$ (bare QCD value) to determine the screening factor at percent precision.
3. **CLASS/CAMB implementation.** Use the predicted $\alpha_{\text{obs}} \approx 0.005$ in Boltzmann codes for MCMC comparison with Planck+DESI data.
4. **Dark matter sigma term.** If dark matter is an axion, compute the axion–condensate coupling to determine whether f_b should be modified.

10 Recommended Journal for Publication

This article is recommended for submission to **Foundations of Physics**. The paper bridges finite-density QCD (nuclear physics), the equation of state of scalar condensates (QFT), and the vacuum energy problem (cosmology). This interdisciplinary character—connecting the nucleon sigma term to the S_8 tension via the Lorentz structure of the chiral condensate—fits the journal’s focus on foundational questions in physics.

References

- Alexandrou, C., et al. 2020, Phys. Rev. D, 101, 034519
- Bazavov, A., et al. 2012, Phys. Rev. D, 85, 054503
- Le Bellac, M. 2011, Thermal Field Theory (Cambridge University Press)
- Borsanyi, S., et al. 2012, JHEP, 2012(08), 053
- Cohen, T. D., Furnstahl, R. J., Griegel, D. K. 1992, Phys. Rev. C, 45, 1881
- Collins, J. C., Duncan, A., Joglekar, S. D. 1977, Phys. Rev. D, 16, 438
- Drukarev, E. G., Levin, E. M. 1991, Nucl. Phys. A, 532, 695
- Fraga, E. S., Kurkela, A., Vuorinen, A. 2014, ApJ Lett., 781, L25
- Gasser, J., Leutwyler, H., Sainio, M. E. 1991, Phys. Lett. B, 253, 252
- Gell-Mann, M., Oakes, R. J., Renner, B. 1968, Phys. Rev., 175, 2195
- Gupta, R., et al. 2021, Phys. Rev. Lett., 127, 242002
- Hatsuda, T., Lee, S. H. 1992, Phys. Rev. C, 46, R34

- Hoferichter, M., Ruiz de Elvira, J., Kubis, B., Meißner, U.-G. 2015, Phys. Rev. Lett., 115, 092301
- Ji, X. 1995, Phys. Rev. Lett., 74, 1071
- Junnarkar, P., Walker-Loud, A. 2013, Phys. Rev. D, 87, 114510
- Kapusta, J. I., Gale, C. 2006, Finite-Temperature Field Theory, 2nd ed. (Cambridge University Press)
- Kruger, B. 2026a, “What If the Vacuum Gravitates Locally?,” Preprint
- Kruger, B. 2026b, “Theoretical Separation of Quantum Vacuum Energy from the Cosmological Constant,” Preprint
- Kruger, B. 2026c, “Matter-Dependent Vacuum Energy Density,” Preprint. doi:10.5281/zenodo.18896536
- Kruger, B. 2026d, “Structure Growth in the Gravitating Vacuum Model,” Preprint
- Kruger, B. 2026e, “N-body Simulations with a Gravitating Vacuum,” Preprint
- Planck Collaboration 2020, A&A, 641, A6
- Schaffner-Bielich, J. 2008, Nucl. Phys. A, 804, 309
- Shifman, M. A., Vainshtein, A. I., Zakharov, V. I. 1978, Nucl. Phys. B, 147, 385
- Solà, J. 2013, J. Phys. Conf. Ser., 453, 012015
- Solà Peracaula, J., et al. 2021, Eur. Phys. J. C, 81, 515
- Weinberg, S. 1989, Rev. Mod. Phys., 61, 1
- Yang, Y.-B., et al. 2016, Phys. Rev. D, 94, 054503
- Yang, Y.-B., et al. 2018, Phys. Rev. Lett., 121, 212001
- Zel’dovich, Ya. B. 1968, Sov. Phys. Usp., 11, 381

Peer Reviews and Revision History

First Round

Reviewer recommendation: Major Revision.

Major Issue 1 (gluon condensate): The reviewer asked whether tracking only the chiral piece misses a canceling contribution from the gluon condensate. Response: Section 5.2 now shows via the QCD trace anomaly that the gluon shift adds coherently—the total is constrained to equal $m_N n_B$ by Eq. (12).

Major Issue 2 (separation ambiguity): The reviewer argued that σ_s is already in m_N and that reclassifying it as vacuum requires a physical principle. Response: Section 4 provides the principle: the chiral condensate is a Lorentz scalar, so its energy has $T^{\mu\nu} \propto g^{\mu\nu}$, i.e., $w = -1$.

Major Issue 3 (equation of state): The reviewer asked what w the condensate shift carries. Response: Proposition 4.1 and the subsequent discussion show $w = -1$ from Lorentz structure.

Major Issue 4 (ad hoc reduction factors): The reviewer noted that bridging the factor ~ 30 with qualitative arguments is weak. Response: Section 6 shows that two quantitative factors (baryon fraction $\times 0.156$; screening $\div 3$) give $\alpha_{\text{obs}} = 0.005$ vs. observed 0.003—a factor 1.7, not 30.

Major Issue 5 (hierarchy claim): Overstated. Response: Section 7 now states explicitly that the finite-density QCD derivation *bypasses* the hierarchy, not resolves it.

Moderate Issue 6 (sign): Clarified in Section 5: $\alpha < 0$ throughout, matching RVM and Paper #8 best fit.

Moderate Issue 7 (dark matter): Promoted from limitation to Section 6: the baryon fraction is a key ingredient in the prediction, not a damage factor.

Moderate Issue 8 (prior literature): References to Zel'dovich (1968), Hatsuda & Lee (1992), Fraga et al. (2014), Schaffner-Bielich (2008) added in Section 3.

Moderate Issue 9 (Theorem/Proof): Removed. Derivations presented as sequential calculations with explicit steps.

Second Round

Placeholder.

Further Clarifications

Placeholder.