

Contents

1	Basic Principles	3
1.1	Problem Types	3
1.2	Derivations	3
1.3	Concepts	3
1.4	Equations	3
2	Accelerated Coordinate Systems	5
2.1	Problem Types	5
2.2	Derivations	5
2.3	Concepts	5
2.4	Equations	5
2.4.1	General Fictitious Forces Eq.	5
3	Lagrangian Dynamics	7
3.1	Problem Types	7
3.1.1	Find the Equilibrium Configuration	7
3.1.2	find stability or frequency of oscillations about equilibrium	7
3.1.3	Solve the Brachistochrone Problem	7
3.2	Derivations	8
3.3	Concepts	8
3.4	Equations	8
3.4.1	Euler-Lagrange Eq.'s	8
3.4.2	Invariance of the Lagrangian	8
4	Small Oscillations	9
4.1	Problem Types	9
4.2	Derivations	9
4.3	Concepts	9
4.4	Equations	9
5	Rigid Bodies	11
5.1	Problem Types	11
5.2	Derivations	11
5.3	Concepts	11
5.4	Equations	11
5.4.1	Parallel Axis Theorem	11
6	Hamiltonian Dynamics	13
6.1	Problem Types	13
6.2	Derivations	13
6.3	Concepts	13
6.4	Equations	13
6.4.1	Hamilton-Jacobi Equation	13

6.4.2	Hamilton's Eq.'s	13
-------	------------------	----

Chapter 1

Basic Principles

1.1 Problem Types

1.2 Derivations

1.3 Concepts

1.4 Equations

Chapter 2

Accelerated Coordinate Systems

2.1 Problem Types

2.2 Derivations

2.3 Concepts

2.4 Equations

2.4.1 General Fictitious Forces Eq.

Say we have a particle, which we can observe in an inertial frame S and some other frame S' (maybe non-inertial), called the *body* frame, which is moving relative to the inertial frame. The position of S' relative to S is \mathbf{a} . The angular velocity of S' is $\boldsymbol{\omega}$ (same for both frames, as angular velocity does not depend on frame). The position of the particle in the body frame S' is \mathbf{r} and is \mathbf{r}_0 in S . Then

$$m \left(\frac{d^2 \mathbf{r}}{dt^2} \right)_{\text{body}} = m \left(\frac{d^2 \mathbf{r}_0}{dt^2} \right)_{\text{inertial}} - m \left(\frac{d^2 \mathbf{a}}{dt^2} \right)_{\text{inertial}} - 2m\boldsymbol{\omega} \times \left(\frac{d\mathbf{r}}{dt} \right) - m\boldsymbol{\omega} \times (\boldsymbol{\omega} \times \mathbf{r}) - m \frac{d\boldsymbol{\omega}}{dt} \times \mathbf{r}$$

Note that the $m \left(\frac{d^2 \mathbf{r}_0}{dt^2} \right)_{\text{inertial}}$ term is just the external force in the inertial frame. That is, it is the term which represents the “real” forces.

Chapter 3

Lagrangian Dynamics

3.1 Problem Types

3.1.1 Find the Equilibrium Configuration

From Fetter and Walecka, P3.1: Say a wire bent into a vertical circle with radius a rotates at angular velocity Ω , with gravity \mathbf{g} , has a bead of mass m free to move on the wire.

The Lagrangian would be

$$\mathcal{L} = T - U = \frac{m}{2} \left(a^2 \dot{\theta}^2 + a^2 \sin^2 \theta \Omega^2 \right) + a \cos \theta mg$$

and the one Euler-Lagrange equation would turn out to be

$$\ddot{\theta} = \sin \theta \cos \theta \Omega^2 - \frac{g}{a} \sin \theta$$

To find the equilibrium position, we need to find a point where, if θ is stationary, then it remains stationary. I.e., where $\dot{\theta} = 0$. Thus we simply set

$$0 = \sin \theta \cos \theta \Omega^2 - \frac{g}{a} \sin \theta$$

And we find an equilibrium with condition

$$\cos \theta_0 = \frac{g}{a \Omega^2}$$

3.1.2 find stability or frequency of oscillations about equilibrium

3.1.3 Solve the Brachistochrone Problem

Just like we minimize the action

$$S = \int \mathcal{L}(q, \dot{q}) dt$$

equivalent to

$$\frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{q}} - \frac{\partial \mathcal{L}}{\partial q} = 0$$

we can minimize the time

$$t_{12} = \int_1^2 \frac{ds}{v}$$

where v is the velocity. Often we can use energy conservation to get

$$v = \sqrt{2gy}$$

where y is the height, and if the particle is following a sloped path

$$t_{12} = \int_1^2 \sqrt{\frac{1 + (y')^2}{2gy}} dx$$

Then writing

$$\Phi = \left[\frac{1 + (y')^2}{2gy} \right]^{1/2}$$

the minimization condition is

$$\frac{d}{dx} \frac{\partial \Phi}{\partial y'} - \frac{\partial \Phi}{\partial y} = 0$$

Note that earlier we could have changed our “time-analogous” coordinate and instead written changing the “time-like” variable

$$t_{12} = \int_1^2 \sqrt{\frac{1 + (x')^2}{2gy}} dy$$

3.2 Derivations

3.3 Concepts

3.4 Equations

3.4.1 Euler-Lagrange Eq.’s

Euler-Lagrange Equations

For one coordinate q

$$\frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{q}} - \frac{\partial \mathcal{L}}{\partial q} = 0$$

3.4.2 Invariance of the Lagrangian

For the two Lagrangians

$$\mathcal{L} = T - V$$

and

$$\mathcal{L}' = T - V + \frac{df(x, t)}{dt}$$

the dynamics are exactly the same for *any* function $f(x, t)$.

Chapter 4

Small Oscillations

4.1 Problem Types

4.2 Derivations

4.3 Concepts

4.4 Equations

Chapter 5

Rigid Bodies

5.1 Problem Types

5.2 Derivations

5.3 Concepts

5.4 Equations

5.4.1 Parallel Axis Theorem

Also called Steiner's Theorem

Given the moment of inertia about the center of mass, this theorem allows us to calculate the moment of inertia about an axis offset from the center (although still pointing in the same direction). For center-of-mass MoI I_c , mass M , and axis offset h , the new moment of inertia is

$$I = I_c + Mh^2$$

Chapter 6

Hamiltonian Dynamics

6.1 Problem Types

6.2 Derivations

6.3 Concepts

6.4 Equations

6.4.1 Hamilton-Jacobi Equation

For the Hamiltonian \mathcal{H}

$$H$$

6.4.2 Hamilton's Eq.'s