

Math 110BH Homework 7

Topic(s): Direct sums, free modules, Noetherian modules

February 24, 2023

For all of this assignment R denotes an arbitrary ring unless stated otherwise.

Problem 1*

Let M and M_1, \dots, M_n be R -modules. Prove that $M \cong \bigoplus_{j=1}^n M_j$ if and only if there exist R -linear maps $\iota_j : M_j \rightarrow M$ and $p_j : M \rightarrow M_j$ such that

- i) $p_j \circ \iota_j$ is the identity map on M_j for all j ,
- ii) $p_j \circ \iota_k = 0$ for all $j \neq k$, and
- iii) $\iota_1 \circ p_1 + \dots + \iota_n \circ p_n$ is the identity on M .

Problem 2

Let M, N be free R -modules with bases α, β , respectively. Use the universal property of free R -modules to prove that if there exists a bijection $g : \alpha \rightarrow \beta$, then there exists an R -module isomorphism $f : M \rightarrow N$ such that $f(a) = g(a)$ for all $a \in \alpha$.

Problem 3

Let M be a R -module. Prove that all of the following conditions are equivalent:

- i) All submodules of M are finitely generated.
- ii) M satisfies the ascending chain condition (ACC) for submodules. That is, whenever

$$M_1 \subseteq M_2 \subseteq M_3 \subseteq \dots$$

is an increasing chain of submodules, then there is a positive integer N such that for all $k \geq N$, $M_k = M_N$.

- iii) Every nonempty set of submodules of M contains a maximal element under inclusion.

*Note: If M satisfies any (so then all) of the above, then we say M is a **Noetherian module**.*

Problem 4*

Prove that if M and N are Noetherian modules, then the direct sum $M \oplus N$ is Noetherian.

Problem 5*

Let M be a Noetherian R -module and $f : M \rightarrow M$ be a surjective module homomorphism.

- a) Prove that there exists an n such that $\ker(f^n) = \ker(f^{n+k})$ for all $k \geq 0$. Then prove for such n that $\ker(f^n) \cap \operatorname{im}(f^n) = \{0\}$.
- b) Prove that f is an R -module isomorphism.