

# Lab: Arrays

Problems for exercises and homework for the [“Programming Fundamentals” course @ SoftUni](#).

You can check your solutions here: <https://judge.softuni.bg/Contests/172/Arrays-Lab>.

## 1. Day of Week

Enter a **day number** [1...7] and print the **day name** (in English) or “Invalid Day!”. Use an **array of strings**.

### Examples

Input	Output
1	Monday
2	Tuesday
7	Sunday
0	Invalid Day!

## 2. Reverse an Array of Integers

Write a program to read an **array of integers**, **reverse** it and **print** its elements. The input consists of a **number n** (the number of elements) + **n** integers, each as a separate line. Print the output on a single line (space separated).

### Examples

Input	Output
3 10 20 30	30 20 10
4 -1 20 99 5	5 99 20 -1

## 3. Last K Numbers Sums Sequence

Enter two integers **n** and **k**. Generate and print the following sequence of **n** elements:

- The first element is: **1**
- All other elements = sum of the previous **k** elements (if less than **k** are available, sum all of them)
- Example:  $n = 9, k = 5 \rightarrow 120 = 4 + 8 + 16 + 31 + 61$

### Examples

Input	Output
6 3	1 1 2 4 7 13
8 2	1 1 2 3 5 8 13 21



9	1 1 2 4 8 16 31 61 120
5	

## 4. Triple Sum

Write a program to read **an array of integers** and find all triples of elements **a**, **b** and **c**, such that **a + b == c** (where **a** stays to the left from **b**). Print **"No"** if no such triples exist.

### Examples


Input	Output
1 1 1 1	No
4 2 8 6	4 + 2 == 6 2 + 6 == 8
2 7 5 0	2 + 5 == 7 2 + 0 == 2 7 + 0 == 7 5 + 0 == 5
3 1 5 6 1 2	3 + 2 == 5 1 + 5 == 6 1 + 1 == 2 1 + 2 == 3 5 + 1 == 6 1 + 2 == 3

## 5. Rounding Numbers Away from Zero

Write a program to read **an array of real numbers** (space separated values), **round** them to the nearest integer in **"away from 0"** style and **print** the output as in the examples below.

Rounding in **"away from zero"** style means:

- To round to the nearest integer, e.g.  $2.9 \rightarrow 3$ ;  $-1.75 \rightarrow -2$
- In case the number is exactly in the middle of two integers (midpoint value), round it to the next integer which is away from the 0:



### Examples

Input	Output
0.9 1.5 2.4 2.5 3.14	0.9 => 1 1.5 => 2 2.4 => 2 2.5 => 3 3.14 => 3
-5.01 -1.599 -2.5 -1.50 0	-5.01 => -5 -1.599 => -2 -2.5 => -3 -1.50 => -2

$0 \Rightarrow 0$
-------------------

## 6. Reverse an Array of Strings

Write a program to read **an array of strings**, **reverse** it and **print** its elements. The input consists of a sequence of space separated strings. Print the output on a single line (space separated).

### Examples

Input	Output
a b c d e	e d c b a
-1 hi ho w	w ho hi -1

## 7. Sum Arrays

Write a program that reads two **arrays of integers** and sums them. When the arrays are not of the same size, duplicate the smaller array a few times.

### Examples

Input	Output	Comments
1 2 3 4 2 3 4 5	3 5 7 9	1 2 3 4 + 2 3 4 5 = 3 5 7 9
1 2 3 4 5 2 3	3 5 5 7 7	1 2 3 4 5 + 2 3 2 3 2 = 3 5 5 7 7
5 4 3 2 3 1 4	7 7 4 9	5 4 3 5 + 2 3 1 4 + 7 7 4 9

## 8. Condense Array to Number

Write a program to read **an array of integers** and **condense** them by **summing** adjacent couples of elements until a **single integer** is obtained. For example, if we have 3 elements {2, 10, 3}, we sum the first two and the second two elements and obtain {2+10, 10+3} = {12, 13}, then we sum again all adjacent elements and obtain {12+13} = {25}.

### Examples

Input	Output	Comments
2 10 3	25	2 10 3 → 2+10 10+3 → 12 13 → 12 + 13 → 25
5 0 4 1 2	35	5 0 4 1 2 → 5+0 0+4 4+1 1+2 → 5 4 5 3 → 5+4 4+5 5+3 → 9 9 8 → 9+9 9+8 → 18 17 → 18+17 → 35
1	1	1 is already condensed to number

## 9. Extract Middle 1, 2 or 3 Elements

Write a method to extract the **middle 1, 2 or 3 elements** from array of **n** integers and **print** them.

- **n = 1** -> **1** element
- **even n** -> **2** elements

- odd **n** -> **3** elements

Create a program that reads an **array of integers** (space separated values) and prints the middle elements in the format shown in the examples.

## Examples

Input	Output
5	{ 5 }
2 3 <b>8</b> <b>1</b> 7 4	{ 8, 1 }
1 2 3 4 5 6 7	{ 3, 4, 5 }
10 20 30 <b>40</b> <b>50</b> 60 70 80	{ 40, 50 }