

NYU CS Bridge to Tandon Course: Homework #8

Winter 2022

Jae Butler

Jiyeon Jung

Problem 7

a. Exercise 6.1.5

- (b) What is the probability that the hand is a three of a kind? A three of a kind has 3 cards of the same rank. The other two cards do not have the same rank as each other and do not have the same rank as the three with the same rank.

$$\begin{aligned}
 P(E) &= \frac{|E|}{|S|} \\
 |E| &= \binom{13}{1} \binom{4}{3} \binom{12}{2} \binom{4}{1} \binom{4}{1} \\
 |S| &= \binom{52}{5} \\
 \frac{|E|}{|S|} &= \frac{\binom{13}{1} \binom{4}{3} \binom{12}{2} \binom{4}{1} \binom{4}{1}}{\binom{52}{5}} \\
 \text{Solution} &= \frac{88}{4165}
 \end{aligned}$$

- (c) What is the probability that all 5 cards have the same suit?

$$\begin{aligned}
 P(E) &= \frac{|E|}{|S|} \\
 |E| &= \binom{13}{5} \binom{4}{1} \\
 |S| &= \binom{52}{5} \\
 \frac{|E|}{|S|} &= \frac{\binom{13}{5} \binom{4}{1}}{\binom{52}{5}} \\
 \text{Solution} &= \frac{33}{16660}
 \end{aligned}$$

- (d) What is the probability that the hand is a two of a kind? A two of a kind has two cards of the same rank (called the pair). Among the remaining three cards, not in the pair, no two have the same rank and none of them have the same rank as the pair

$$\begin{aligned}
 P(E) &= \frac{|E|}{|S|} \\
 |E| &= \binom{4}{2} \binom{13}{1} \binom{12}{3} \binom{4}{1} \binom{4}{1} \binom{4}{1} \\
 |S| &= \binom{52}{5} \\
 \frac{|E|}{|S|} &= \frac{\binom{4}{2} \binom{13}{1} \binom{12}{3} \binom{4}{1} \binom{4}{1} \binom{4}{1}}{\binom{52}{5}} \\
 \text{Solution} &= \frac{352}{833}
 \end{aligned}$$

b. Exercise 6.2.4

- (a) The hand has at least one club.

$$\begin{aligned}
 \overline{E} &= \text{The hand has no club} \\
 E &= \text{The hand has atleast one club} \\
 |S| &= \binom{52}{5} \\
 P(\overline{E}) &= \frac{\binom{39}{5}}{\binom{52}{5}} \\
 \text{Solution} &= \frac{7411}{9520}
 \end{aligned}$$

- (b) The hand has at least two cards with the same rank.

$$\begin{aligned}
 E &= \text{The hand has at least two cards with the same rank} \\
 |S| &= \binom{52}{5} \\
 P(E) &= 1 - P(0\text{card}) - P(1\text{card}) \\
 P(E) &= 1 - \frac{\binom{13}{0} \binom{39}{5}}{\binom{52}{5}} - \frac{\binom{13}{1} \binom{39}{4}}{\binom{52}{5}} \\
 \text{Solution} &= 1 - 0.2215 - .4114 = 0.3671
 \end{aligned}$$

(c) The hand has exactly one club or exactly one spade.

$$C = \text{Exactly 1 club} = \binom{13}{1} * \binom{39}{4}$$

$$S = \text{Exactly 1 Spade} = \binom{13}{1} * \binom{39}{4}$$

$$|S'| = \text{Sample Space} = \binom{52}{5}$$

$$P(C \cup S) = \binom{13}{1} * \binom{39}{4} + \binom{13}{1} * \binom{39}{4}$$

$$P(C \cup S) = \frac{2 * 13 * \binom{39}{4}}{\binom{52}{5}}$$

$$\text{Solution} = 26 * \frac{82251}{2598960} = 0.8228$$

(d) The hand has at least one club or at least one spade.

$$P(\overline{C} \cup \overline{S}) = \text{No club or no space}$$

$$P(C \cup S) = \text{at least one club and spades}$$

$$P(\overline{C} \cup \overline{S}) = \binom{26}{5} \text{no spade and no club deck}$$

$$|S'| = \text{Sample Space} = \binom{52}{5}$$

$$P(C \cup S) = 1 - \frac{\binom{26}{5}}{\binom{52}{5}}$$

$$\text{Solution} = 1 - \frac{65780}{2598960} = 1 - 0.0253 = 0.97468$$

Problem 8

a. Exercise 6.3.2 The letters {a, b, c, d, e, f, g} are put in a random order. Each permutation is equally likely. Define the following events:

A: The letter b falls in the middle (with three before it and three after it)

B: The letter c appears to the right of b, although c is not necessarily immediately to the right of b. For example, "agbdcef" would be an outcome in this event.

C: The letters "def" occur together in that order (e.g. "gdefbca")

(a) Calculate the probability of each individual event. That is, calculate $p(A)$, $p(B)$, and $p(C)$.

$$P(A) = \frac{6!}{7!} = \frac{1}{7}$$

$$P(B) = \frac{\frac{7!}{2!}}{7!} = \frac{1}{2}$$

$$P(C) = \frac{5!}{7!} = \frac{1}{42}$$

(b) What is $p(A|C)$?

$$P(A|C) = \frac{P|A \cap C|}{|C|} = \frac{2! * 3!}{5!} = \frac{1}{10}$$

(c) What is $p(B|C)$?

$$P(B|C) = \frac{P|B \cap C|}{|C|} = \frac{\frac{5!}{2!}}{5!} = \frac{1}{2}$$

(d) What is $p(A|B)$?

$$P(A|B) = \frac{P(A \cap B)}{|B|} = \frac{\frac{5!}{3!} * 3! * 3!}{\frac{7!}{2!}} = \frac{1}{7}$$

(e) Which pairs of events among A, B, and C are independent?

$P(A \cap B)$ and $P(B \cap C)$ is independent

$$P(A) = \frac{6!}{7!} = \frac{1}{7}$$

$$P(B) = \frac{\frac{7!}{2!}}{7!} = \frac{1}{2}$$

$$P(C) = \frac{5!}{7!} = \frac{1}{42}$$

$$P(A \cap B) = \frac{6!}{2} * \frac{1}{7!}$$

$$P(B \cap C) = \frac{5!}{2} * \frac{1}{7!}$$

b. Exercise 6.3.6 A biased coin is flipped 10 times. In a single flip of the coin, the probability of heads is $1/3$ and the probability of tails is $2/3$. The outcomes of the coin flips are mutually independent. What is the probability of each event?

$$Head = \frac{1}{3}$$

$$Tail = \frac{2}{3}$$

$$N = 10$$

b. The first 5 flips come up heads. The last 5 flips come up tails.

$$Solution = \binom{1}{3}^5 * \binom{2}{3}^5$$

(b)

(c) The first flip comes up heads. The rest of the flips come up tails.

$$Solution = \binom{1}{3} * \binom{2}{3}^9$$

c. Exercise 6.4.2 Assume that you have two dice, one of which is fair, and the other is biased toward landing on six, so that 0.25 of the time it lands on six, and 0.15 of the time it lands on each of 1, 2, 3, 4 and 5. You choose a die at random, and roll it six times, getting the values 4, 3, 6, 6, 5, 5. What is the probability that the die you chose is the fair die? The outcomes of the rolls are mutually independent.

(a) Find the probability of Fair die and biased die

Let F denote of the event that fair die is chosen and let X be the outcome of the event (values 4, 3, 6, 6, 5, 5)

$$\text{Fair die} = p(X_1) = p(X_2) \dots p(X_6) = \frac{1}{6}$$

$$\text{Biased die} = p(X_1) = p(X_2) \dots p(X_5) = 0.15 \quad p(X_6) = 0.25$$

$$P(X|F) = \left(\frac{1}{6}\right)^6$$

$$P(X|\bar{F}) = (0.15)^4 * (.025)^2$$

$$\text{BayesTheorem} = p(F|X) = \frac{p(X|F) * p(F)}{P(X|F) * p(F) + P(X|\bar{F}) * p(\bar{F})}$$

$$\text{Solution} = p(F|X) = \frac{\left(\frac{1}{6}\right)^6 * \frac{1}{2}}{\left(\frac{1}{6}\right)^6 * \frac{1}{2} + (0.15)^4 * (0.25)^2 * \frac{1}{2}} = .4038$$

Problem 9

a. Exercise 6.5.2

A hand of 5 cards is dealt from a perfectly shuffled deck of playing cards. Let the random variable A denote the number of aces in the hand.

(a) What is the range of A?

$$\{0, 1, 2, 3, 4\}$$

(b) Give the distribution over the random variable A.

$$|A| = 0 \frac{\binom{48}{5}}{\binom{52}{5}}, 1 \frac{\binom{4}{1} \binom{48}{4}}{\binom{52}{5}}, 2 \frac{\binom{4}{2} \binom{48}{3}}{\binom{52}{5}}, 3 \frac{\binom{4}{3} \binom{48}{2}}{\binom{52}{5}}, 4 \frac{\binom{4}{4} \binom{48}{1}}{\binom{52}{5}},$$

b. Exercise 6.6.1 Two student council representatives are chosen at random from a group of 7 girls and 3 boys. Let G be the random variable denoting the number of girls chosen. What is $E[G]$?

(a) G is a random variable denoting the number of girls chosen,

$$G = 0, 1, 2$$

$$\text{Probability of } (G = 0) = \frac{3}{45}$$

$$\text{Probability of } (G = 1) = \frac{21}{45}$$

$$\text{Probability of } (G = 2) = \frac{21}{45}$$

$$E[G] = (0 * \frac{3}{45}) + (1 * \frac{21}{45}) + (2 * \frac{21}{45})$$

$$E[G] = \frac{7}{5} = 1.4$$

b. Exercise 6.6.4 Expected values of squares.

- (a) A fair die is rolled once. Let X be the random variable that denotes the square of the number that shows up on the die. For example, if the die comes up 5, then $X = 25$. What is $E[X]$?

$$X = x^2$$

$$E[X] = (1^2 * \frac{1}{6}) + (2^2 * \frac{1}{6}) + (3^2 * \frac{1}{6}) + (4^2 * \frac{1}{6}) + (5^2 * \frac{1}{6}) + (6^2 * \frac{1}{6})$$

$$E[X] = \frac{91}{6} = 15.166$$

- (b) A fair coin is tossed three times. Let Y be the random variable that denotes the square of the number of heads. For example, in the outcome HTH, there are two heads and $Y = 4$. What is $E[Y]$?

$$Y = x^2$$

$$Y = 0, 1, 4, 9$$

$$E[Y] = (0^2 * \frac{1}{8}) + (1^2 * \frac{3}{8}) + (2^2 * \frac{3}{8}) + (3^2 * \frac{1}{8})$$

$$E[Y] = 3$$

b. Exercise 6.7.4 A class of 10 students hang up their coats when they arrive at school. Just before recess, the teacher hands one coat selected at random to each child. What is the expected number of children who get his or her own coat?

- (a) Let C denote the random variable children get his or her own coat
if children get his or her coat then $C_i = 1$ otherwise $C_i = 0$

$$E \sum_{i=1}^{10} C_i$$

$$E[C] = E[C_1] + E[C_2] + E[C_3] \dots E[C_{10}]$$

$$E[C] = 10 * E[\frac{1}{10}]$$

$$E[C_1] = (1 * \frac{1}{10}) + (0 * \frac{9}{10}) = \frac{1}{10}$$

$$E[C] = 10 * \frac{1}{10} = 1$$

Problem 10

b. Exercise 6.8.1 The probability that a circuit board produced by a particular manufacturer has a defect is 1 percent. You can assume that errors are independent, so the event that one circuit board has a defect is independent of whether a different circuit board has a defect.

- (a) What is the probability that out of 100 circuit boards made exactly 2 have defects?

$$\text{Binomial distribution} = b\binom{n}{k}P^kQ^{n-k}$$

$$N = 100(\text{Trials})$$

$$K = 2 \text{ (total defect)}$$

$$P = 1 - .01 = .99$$

$$b(2; 100, 0.01) = \binom{100}{2} * (0.01)^2(0.99)^{98} = 0.1849$$

- (b) What is the probability that out of 100 circuit boards made at least 2 have defects?

$$p = b(0; 100, 0.01) - b(1; 100, 0.01)$$

$$1 - \binom{100}{0} * (0.01)^0(0.99)^{100} - \binom{100}{1} * (0.01)^1(0.99)^{99} = 0.2642$$

- (c) What is the expected number of circuit boards with defects out of the 100 made?

$$E[C] = 100 * 0.01 = 1$$

- (d) Now suppose that the circuit boards are made in batches of two. Either both circuit boards in a batch have a defect or they are both free of defects. The probability that a batch has a defect is 1 percent. What is the probability that out of 100 circuit boards (50 batches) at least 2 have defects? What is the expected number of circuit boards with defects out of the 100 made? How do your answers compare to the situation in which each circuit board is made separately?

Probability of Zero defect out of 1 batch(50)

$$\text{Binomial distribution} = b\left(\begin{matrix} n \\ k \end{matrix}\right) P^k Q^{n-k}$$

$$N = 50$$

$$K = 0 \text{ (total defect)}$$

$$P = .01$$

$$b(0; 50, 0.01) = \binom{50}{0} * (0.01)^0 (0.99)^{50} = 0.60501$$

Probability of 1 defect out of 1 batch(50)

$$\text{Binomial distribution} = b\left(\begin{matrix} n \\ k \end{matrix}\right) P^k Q^{n-k}$$

$$N = 50$$

$$K = 1 \text{ (total defect)}$$

$$P = .01$$

$$b(1; 50, 0.01) = \binom{50}{1} * (0.01)^1 (0.99)^{49} = 0.30556$$

Combine probability of 0 and 1 defect from batch to get at least 2 batch with defects

$$P(0 \text{ Defect}) + P(1 \text{ defect}) = 0.6050 + 0.3055 = .9105$$

$$1 - P(0 \text{ Defect}) + P(1 \text{ defect}) = 1 - 0.9105 = 0.0895$$

Expected number of circuit board out of 100

$$E[C] = 50 * 0.01 = 0.5$$

When circuit boards are made in batches the probability of defect reduce to half compare to made with whole

b. Exercise 6.8.3 A gambler has a coin which is either fair (equal probability heads or tails) or is biased with a probability of heads equal to 0.3. Without knowing which coin he is using, you ask him to flip the coin 10 times. If the number of heads is at least 4, you conclude that the coin is fair. If the number of heads is less than 4, you conclude that the coin is biased.

- (a) What is the probability that you reach an incorrect conclusion if the coin is biased?

Solution: The incorrect conclusion is reached if there is three or more heads

$$1 - \left((0.7)^{10} + 10 * (0.3)^1 * (0.7)^9 + \binom{10}{2} * (0.3)^2 * (0.7)^8 + \binom{10}{3} * (0.3)^3 * (0.7)^7 \right)$$

$$\text{Solution : } 1 - (0.0824 * 7.888) = 0.35$$