

Engineering Mathematics

Complete MCQ-Oriented Notes

IBPS SO (IT Officer) Examination - Complete Preparation Series

100% Syllabus Coverage - MCQ-Oriented - Zero Filler

Compiled by Asabhya Maanav

Table of Contents

Engineering Mathematics - CIL MT Systems Complete Notes	6
1. Mathematical Logic	6
1.1 Propositional Logic and Connectives	6
1.2 Truth Tables; Tautology, Contradiction, Contingency	6
1.3 Logical Equivalences; De Morgan's Laws	7
1.4 Normal Forms: CNF and DNF	7
1.5 Rules of Inference; Validity of Arguments	7
1.6 First-Order (Predicate) Logic: Predicates and Quantifiers	8
1.7 Nested Quantifiers and Negation of Quantified Statements	8
2. Set Theory and Relations	9
2.1 Sets, Subsets, Power Set, Operations	9
2.2 Venn Diagrams; Inclusion-Exclusion Principle	9
2.3 Cartesian Product	9
2.4 Relations: Properties	9
2.5 Equivalence Relations and Equivalence Classes	10
2.6 Partial Order Relations; Hasse Diagrams	10
2.7 Functions: Injective, Surjective, Bijective	10
2.8 Countable and Uncountable Sets	10
3. Partial Orders and Lattices	11
3.1 POSET; Upper/Lower Bounds, LUB, GLB	11
3.2 Lattices; Bounded, Distributive, Complemented	11
3.3 Hasse Diagram Interpretation	11
4. Algebraic Structures	12
4.1 Binary Operations and Properties	12
4.2 Semigroup and Monoid	12
4.3 Groups; Abelian Groups; Subgroups	12
4.4 Order of a Group/Element; Cyclic Groups	12
5. Graph Theory	13
5.1 Graph Terminology	13
5.2 Handshaking Lemma	13
5.3 Types: Simple, Complete, Bipartite, Regular, Planar	13
5.4 Connectivity; Components	13

5.5 Eulerian and Hamiltonian Graphs	13
5.6 Trees and Spanning Trees	14
5.7 Graph Coloring; Chromatic Number	14
5.8 Matching	14
5.9 Planar Graphs; Euler's Formula	14
6. Combinatorics	15
6.1 Counting Principles	15
6.2 Permutations and Combinations	15
6.3 Pigeonhole Principle	15
6.4 Inclusion-Exclusion (Counting)	15
6.5 Binomial Theorem and Coefficients	15
6.6 Recurrence Relations	15
6.7 Generating Functions	16
7. Matrices and Determinants	17
7.1 Types of Matrices	17
7.2 Matrix Operations	17
7.3 Determinant: Properties and Computation	17
7.4 Inverse of a Matrix; Adjoint	17
7.5 Rank of a Matrix	18
7.6 Elementary Row Operations	18
8. System of Linear Equations	19
8.1 Homogeneous vs Non-Homogeneous	19
8.2 Consistency and Number of Solutions (Rank Method)	19
8.3 Gaussian Elimination	19
8.4 Cramer's Rule	19
9. Eigenvalues and Eigenvectors	20
9.1 Characteristic Equation	20
9.2 Computation	20
9.3 Properties	20
9.4 Cayley-Hamilton Theorem	20
9.5 Diagonalization	20
10. LU Decomposition	21
10.1 LU Factorization Concept and Method	21
10.2 Solving Systems Using LU	21

11. Limits, Continuity, Differentiability	22
11.1 Limits and Standard Limit Results	22
11.2 Continuity	22
11.3 Differentiability; Relation to Continuity	22
11.4 Derivative Rules	22
12. Applications of Derivatives	23
12.1 Maxima and Minima	23
12.2 Critical Points; First and Second Derivative Tests	23
12.3 Mean Value Theorems	23
13. Integration	24
13.1 Indefinite Integrals; Standard Integrals	24
13.2 Methods: Substitution, By Parts, Partial Fractions	24
13.3 Definite Integrals and Properties	24
13.4 Area Under a Curve	24
14. Probability	25
14.1 Sample Space, Events; Axioms	25
14.2 Conditional Probability; Multiplication Rule	25
14.3 Independent Events	25
14.4 Bayes' Theorem and Total Probability	25
14.5 Random Variables: Discrete and Continuous	25
14.6 PMF/PDF; CDF	25
14.7 Expectation, Variance, Standard Deviation	25
15. Probability Distributions	27
15.1 Uniform Distribution	27
15.2 Binomial Distribution	27
15.3 Poisson Distribution	27
15.4 Normal Distribution	27
15.5 Exponential Distribution	27
16. Statistics	28
16.1 Measures of Central Tendency	28
16.2 Measures of Dispersion	28
16.3 Correlation and Regression	28
17. Numerical Solutions of Equations	29
17.1 Non-linear Equations	29

17.2 Linear Systems	29
17.3 Convergence and Error	29
18. Numerical Integration	30
18.1 Trapezoidal Rule	30
18.2 Simpson's 1/3 Rule	30
18.3 Error in Numerical Integration	30
19. Numerical Solution of ODEs	31
19.1 Single-Step Methods	31
19.2 Multi-Step Methods	31
20. Exam Focus / Practice Checklist	32
20.1 Propositional Logic Equivalence / Validity	32
20.2 Relation/Function Property Identification	32
20.3 Lattice / Hasse Diagram Questions	32
20.4 Counting	32
20.5 Recurrence Solving	32
20.6 Matrix Rank / Determinant / Inverse	32
20.7 Eigenvalue Computation and Properties	32
20.8 Probability	32
20.9 Newton-Raphson / Trapezoidal / Simpson	32
20.10 Graph Theory	32

Engineering Mathematics - CIL MT Systems Complete Notes

- This guide covers the COMPLETE Engineering Mathematics syllabus for the **Coal India Limited (CIL) Management Trainee** Systems discipline objective CBT.
- Difficulty target: **conceptual-to-moderate** (direct formula application), but depth is kept at **GATE-level** for safety.
- Five pillars: **Discrete Mathematics, Linear Algebra, Calculus, Probability & Statistics, and Numerical Methods.**
- How to use: read the strict definition, memorize the red formulas, then absorb each **CBT Trap** which names the exact confusions examiners exploit.

1. Mathematical Logic

1.1 Propositional Logic and Connectives

- A **proposition** is a declarative statement that is either **true (T)** or **false (F)**, but never both.
- A **connective** combines propositions into compound propositions; the five standard ones are below.
- **Negation** (NOT, symbol $\neg p$ or $\sim p$): reverses truth value.
- **Conjunction** (AND, symbol $p \wedge q$): true only when **both** are true.
- **Disjunction** (OR, symbol $p \vee q$): true when **at least one** is true (inclusive OR).
- **Implication** (IF-THEN, symbol $p \rightarrow q$): false **ONLY** when **p is true and q is false**.
- **Biconditional** (IFF, symbol $p \leftrightarrow q$): true when **p and q have the same truth value**.

Connective	Symbol	True when
NOT	$\neg p$	p is false
AND	$p \wedge q$	both true
OR	$p \vee q$	at least one true
Implication	$p \rightarrow q$	except T $\rightarrow$ F
Biconditional	$p \leftrightarrow q$	same values

- For **n** propositional variables a truth table has **2^n** rows.
- Implication equivalents: $p \rightarrow q \equiv \neg p \vee q$.
- Contrapositive of $p \rightarrow q$ is $\neg q \rightarrow \neg p$ (logically equivalent).
- Converse of $p \rightarrow q$ is $q \rightarrow p$ (NOT equivalent).
- Inverse of $p \rightarrow q$ is $\neg p \rightarrow \neg q$ (NOT equivalent, but equivalent to converse).

Memory Trick

- "Implication is only a **liar when premise is true but conclusion false**" — remember the single F row of $p \rightarrow q$.

CBT Trap

- **Converse and inverse are NOT equivalent to the original**; only the **contrapositive** is.
- OR in logic is **inclusive** by default; XOR (**exclusive or**) is true when **exactly one** operand is true.
- A vacuous implication with a **false premise** is always **true**.

1.2 Truth Tables; Tautology, Contradiction, Contingency

- A **tautology** is a compound proposition that is **always true** for every assignment (e.g., $p \vee \neg p$).
- A **contradiction** is **always false** (e.g., $p \wedge \neg p$).
- A **contingency** is neither — its value **depends on inputs**.

- $p \rightarrow q$ is a tautology iff p logically implies q .

CBT Trap

- **Tautology vs satisfiable**: a tautology is true in ALL rows; satisfiable needs only **at least one** true row.

1.3 Logical Equivalences; De Morgan's Laws

- Two propositions are **logically equivalent** ($\equiv$) when they have **identical truth tables**.
- De Morgan: $\neg(p \wedge q) \equiv \neg p \vee \neg q$ and $\neg(p \vee q) \equiv \neg p \wedge \neg q$.
- Identity: $p \wedge T \equiv p$, $p \vee F \equiv p$.
- Domination: $p \vee T \equiv T$, $p \wedge F \equiv F$.
- Idempotent: $p \vee p \equiv p$, $p \wedge p \equiv p$.
- Distributive: $p \vee (q \wedge r) \equiv (p \vee q) \wedge (p \vee r)$.
- Absorption: $p \vee (p \wedge q) \equiv p$.

Law	Form
Commutative	$p \wedge q \equiv q \wedge p$
Associative	$(p \wedge q) \wedge r \equiv p \wedge (q \wedge r)$
De Morgan	$\neg(p \wedge q) \equiv \neg p \vee \neg q$
Distributive	$p \wedge (q \vee r) \equiv (p \wedge q) \vee (p \wedge r)$

CBT Trap

- When negating, De Morgan **flips AND to OR** and vice versa AND negates each term — students forget to flip the connective.

1.4 Normal Forms: CNF and DNF

- **DNF (Disjunctive Normal Form)**: an **OR of ANDs** — disjunction of conjunctive clauses (minterms).
- **CNF (Conjunctive Normal Form)**: an **AND of ORs** — conjunction of disjunctive clauses (maxterms).
- Every proposition can be written in both **CNF** and **DNF**.

Memory Trick

- **DNF = D** for "Disjunction outside"; **CNF = C** for "Conjunction outside".

CBT Trap

- Do not confuse **CNF (product of sums)** with **DNF (sum of products)**; SOP = DNF, POS = CNF.

1.5 Rules of Inference; Validity of Arguments

- **Modus Ponens**: from p and $p \rightarrow q$ infer q .
- **Modus Tollens**: from $\neg q$ and $p \rightarrow q$ infer $\neg p$.
- **Hypothetical Syllogism**: from $p \rightarrow q$ and $q \rightarrow r$ infer $p \rightarrow r$.
- **Disjunctive Syllogism**: from $p \vee q$ and $\neg p$ infer q .
- **Addition**: from p infer $p \vee q$.
- **Simplification**: from $p \wedge q$ infer p .
- An argument is **valid** when the conclusion is true whenever all premises are true.

CBT Trap

- **Affirming the consequent** (from q and $p \rightarrow q$ conclude p) and **denying the antecedent** are **fallacies**, not valid rules.

1.6 First-Order (Predicate) Logic: Predicates and Quantifiers

- A **predicate** $P(x)$ becomes a proposition once x is assigned a value or quantified.
- **Universal quantifier** $\forall x P(x)$: "for all x , $P(x)$ " — true if P holds for **every** element.
- **Existential quantifier** $\exists x P(x)$: "there exists x " — true if P holds for **at least one** element.

1.7 Nested Quantifiers and Negation of Quantified Statements

- Negation rule: $\neg \forall x P(x) \equiv \exists x \neg P(x)$ and $\neg \exists x P(x) \equiv \forall x \neg P(x)$.
- $\forall x \exists y$ differs from $\exists y \forall x$: order of quantifiers changes meaning.

CBT Trap

- $\forall x \exists y P(x,y)$ (each x has some y) is **weaker** than $\exists y \forall x P(x,y)$ (one y works for all x); swapping order is a classic trap.
- To negate, **swap** $\forall \leftrightarrow \exists$ and negate the inner predicate.

2. Set Theory and Relations

2.1 Sets, Subsets, Power Set, Operations

- A **set** is an unordered collection of distinct objects (elements).
- $A \subseteq B$ (subset): every element of A is in B; **proper subset** $A \subset B$ excludes equality.
- Power set** $P(A)$: set of all subsets; for $|A| = n$, $|P(A)| = 2^n$.
- Union** $A \cup B$: elements in A or B.
- Intersection** $A \cap B$: elements in both.
- Difference** $A - B$: in A but not B.
- Complement** A' : elements in universal set U not in A.

Operation	Notation	Definition
Union	$A \cup B$	in A or B
Intersection	$A \cap B$	in both
Difference	$A - B$	in A not B
Symmetric diff	$A \Delta B$	in exactly one

- Number of subsets of size r: $C(n, r)$; total subsets 2^n ; proper subsets $2^n - 1$.

CBT Trap

- Empty set** $\emptyset$ is a subset of EVERY set; $\emptyset \neq \{\emptyset\}$ ($|\{\emptyset\}| = 1$).
- Power set of empty set has **1** element, not 0.

2.2 Venn Diagrams; Inclusion-Exclusion Principle

- Two-set: $|A \cup B| = |A| + |B| - |A \cap B|$.
- Three-set: $|A \cup B \cup C| = |A| + |B| + |C| - |A \cap B| - |A \cap C| - |B \cap C| + |A \cap B \cap C|$.

Memory Trick

- Inclusion-exclusion alternates signs: **add singles, subtract pairs, add triples**.

2.3 Cartesian Product

- $A \times B$ = set of ordered pairs (a, b); $|A \times B| = |A| \times |B|$.
- $A \times B \neq B \times A$ in general (order matters).

2.4 Relations: Properties

- A **relation** R on set A is a subset of $A \times A$.
- Reflexive**: $(a, a) \in R$ for all a.
- Symmetric**: $(a, b) \in R \Rightarrow (b, a) \in R$.
- Antisymmetric**: $(a, b) \in R$ and $(b, a) \in R \Rightarrow a = b$.
- Transitive**: $(a, b), (b, c) \in R \Rightarrow (a, c) \in R$.
- On a set of n elements there are $2^{(n^2)}$ relations.

Property	Condition
Reflexive	$(a, a) \in R \forall a$
Symmetric	$aRb \Rightarrow bRa$

Property	Condition
Antisymmetric	$aRb \wedge bRa \Rightarrow a=b$
Transitive	$aRb \wedge bRc \Rightarrow aRc$

CBT Trap

- **Antisymmetric is NOT "not symmetric"**; a relation can be both symmetric and antisymmetric (only the diagonal).
- Number of **reflexive** relations = 2^{n^2-n} ; symmetric = $2^{n(n+1)/2}$.

2.5 Equivalence Relations and Equivalence Classes

- An **equivalence relation** is **reflexive + symmetric + transitive** (RST).
- It **partitions** the set into disjoint **equivalence classes**.
- Number of equivalence relations on n elements = **Bell number $B(n)$** .

CBT Trap

- A **partition** corresponds one-to-one with an **equivalence relation**; partition blocks are the classes.

2.6 Partial Order Relations; Hasse Diagrams

- A **partial order** (POSET relation) is **reflexive + antisymmetric + transitive**.
- A **Hasse diagram** draws the partial order with **transitive and reflexive edges removed** and direction implied upward.

CBT Trap

- **Equivalence vs partial order**: equivalence is symmetric; partial order is **antisymmetric** — opposite middle property.

2.7 Functions: Injective, Surjective, Bijective

- **Injective (one-one)**: distinct inputs map to distinct outputs.
- **Surjective (onto)**: every codomain element is hit.
- **Bijective**: both injective and surjective; has an **inverse**.
- Composition $(f \circ g)(x) = f(g(x))$.
- From a set of size m to size n : total functions n^m ; injective functions $n!/(n-m)!$ ($m \leq n$); surjective via inclusion-exclusion.

CBT Trap

- **Onto requires codomain = range**; counting onto functions uses **inclusion-exclusion**, not a simple factorial.
- $(f \circ g)$ means apply **g first**, then f .

2.8 Countable and Uncountable Sets

- A set is **countable** if it is finite or has a bijection with $\mathbb{N}$ (natural numbers).
- $\mathbb{Z}$ and $\mathbb{Q}$ are countable; $\mathbb{R}$ is **uncountable** (Cantor's diagonal argument).

CBT Trap

- **Rationals are countable** despite being dense; **reals are uncountable**.

3. Partial Orders and Lattices

3.1 POSET; Upper/Lower Bounds, LUB, GLB

- A **POSET** is a set with a partial order (**reflexive, antisymmetric, transitive**).
- **Upper bound** of subset S: an element $\geq$ every element of S.
- **LUB (supremum/join)**: the **least** upper bound.
- **GLB (infimum/meet)**: the **greatest** lower bound.

3.2 Lattices; Bounded, Distributive, Complemented

- A **lattice** is a POSET in which **every pair** of elements has both a **LUB (join, $\vee$)** and **GLB (meet, $\wedge$)**.
- **Bounded lattice**: has a **greatest element 1** and **least element 0**.
- **Distributive lattice**: meet distributes over join and vice versa.
- **Complemented lattice**: every element has a complement.

Lattice type	Key property
Bounded	has 0 and 1
Distributive	$a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$
Complemented	every element has complement
Boolean algebra	bounded + distributive + complemented

CBT Trap

- Every **chain (totally ordered set)** is a **distributive lattice**.
- A **Boolean algebra** is a **complemented distributive** lattice.

3.3 Hasse Diagram Interpretation

- Top node = **greatest element**; bottom = **least element**.
- An element covers another if directly connected with no intermediate.

CBT Trap

- Not every POSET is a lattice; check that **both LUB and GLB exist for all pairs**.

4. Algebraic Structures

4.1 Binary Operations and Properties

- A **binary operation** $*$ on set S maps $S \times S \rightarrow S$.
- **Closure**: $a*b \in S$ for all a, b .
- **Associativity**: $(a*b)*c = a*(b*c)$.
- **Identity e** : $a*e = e*a = a$.
- **Inverse**: $a*a^{-1} = e$.

4.2 Semigroup and Monoid

- **Semigroup** = closure + associativity.
- **Monoid** = semigroup + **identity element**.

4.3 Groups; Abelian Groups; Subgroups

- **Group** = monoid + **inverse** for every element (closure, associativity, identity, inverse).
- **Abelian (commutative) group**: additionally $a*b = b*a$.
- **Subgroup**: a subset that is itself a group under the same operation.

Structure	Properties
Semigroup	closure + associative
Monoid	+ identity
Group	+ inverse
Abelian group	+ commutative

Memory Trick

- Climb the ladder: **Semigroup** $\rightarrow$ **Monoid** $\rightarrow$ **Group** $\rightarrow$ **Abelian** adds **associativity-already, identity, inverse, commutativity**.

4.4 Order of a Group/Element; Cyclic Groups

- **Order of a group** = number of elements; **order of an element a** = smallest k with $a^k = e$.
- A **cyclic group** is generated by a single element; every cyclic group is **abelian**.
- **Lagrange's theorem**: order of a subgroup **divides** the order of the group.

CBT Trap

- **Group needs ALL four** axioms; missing inverse $\rightarrow$ only a monoid. Commutativity is NOT required for a group.

5. Graph Theory

5.1 Graph Terminology

- A **graph** $G = (V, E)$ has a vertex set V and edge set E .
- Degree** of a vertex = number of incident edges.
- A **path** is a walk with no repeated vertices; a **cycle** returns to start.

5.2 Handshaking Lemma

- Sum of degrees** = $2 \times \text{number of edges}$: $\sum \deg(v) = 2|E|$.
- The number of **odd-degree vertices** is always **even**.

CBT Trap

- Each edge contributes **2** to the degree sum; a common error is forgetting the factor of 2.

5.3 Types: Simple, Complete, Bipartite, Regular, Planar

- Simple graph**: no loops or multiple edges.
- Complete graph K_n** : every pair connected; edges $n(n-1)/2$.
- Bipartite**: vertices split into two sets with edges only across; **no odd cycles**.
- Regular**: every vertex has equal degree.
- Complete bipartite $K(m,n)$** : edges $m \times n$.

Graph	Edges
Complete K_n	$n(n-1)/2$
Complete bipartite $K_{m,n}$	$m \cdot n$
Tree on n vertices	$n-1$
Cycle C_n	n

CBT Trap

- A graph is **bipartite iff it has NO odd-length cycle**.

5.4 Connectivity; Components

- A graph is **connected** if there is a path between every pair.
- A **connected component** is a maximal connected subgraph.

5.5 Eulerian and Hamiltonian Graphs

- Eulerian circuit** (uses every edge once, returns to start) exists iff connected and **every vertex has even degree**.
- Eulerian path** (not closed) exists iff **exactly 0 or 2 vertices have odd degree**.
- Hamiltonian cycle** visits every **vertex** exactly once; **no simple necessary-and-sufficient condition** (NP-hard).

Concept	Touches	Condition
Eulerian	every edge	even degrees
Hamiltonian	every vertex	no simple condition

CBT Trap

- **Euler = edges, Hamilton = vertices**; Euler has an easy degree condition, Hamilton does NOT.

5.6 Trees and Spanning Trees

- A **tree** is a connected acyclic graph with **$n-1$** edges on n vertices.
- A tree has **exactly one path** between any two vertices.
- A **spanning tree** of a connected graph includes all vertices with **$n-1$** edges.
- Number of spanning trees of **$K_n = n^{(n-2)}$** (Cayley's formula).

CBT Trap

- Adding any edge to a tree creates **exactly one cycle**; removing any edge **disconnects** it.

5.7 Graph Coloring; Chromatic Number

- **Chromatic number $\chi(G)$** = minimum colors to color vertices so adjacent ones differ.
- **$\chi(K_n) = n$; $\chi(\text{bipartite}) = 2$** (if it has edges); **$\chi(\text{cycle } C_n) = 2$** if n even, **3** if n odd.
- **Four Color Theorem**: every planar graph is **4-colorable**.

CBT Trap

- **$\chi(\text{odd cycle}) = 3$** , not 2; bipartite needs exactly 2.

5.8 Matching

- A **matching** is a set of edges with no shared vertices.
- A **perfect matching** covers every vertex.

5.9 Planar Graphs; Euler's Formula

- A **planar graph** can be drawn with no edge crossings.
- **Euler's formula: $V - E + F = 2$** for connected planar graphs.
- Necessary condition: **$E \leq 3V - 6$** (simple, $V \geq 3$); for bipartite **$E \leq 2V - 4$** .
- **K_5 and $K_{3,3}$** are the basic **non-planar** graphs (Kuratowski).

CBT Trap

- **Euler's formula $V-E+F=2$** counts the **outer (unbounded) face** as one of the F faces.

6. Combinatorics

6.1 Counting Principles

- **Sum rule:** if tasks are mutually exclusive, total ways = $n_1 + n_2$.
- **Product rule:** for sequential independent choices, total = $n_1 \times n_2$.

6.2 Permutations and Combinations

- **Permutation** $P(n,r) = n!/(n-r)!$ — order matters.
- **Combination** $C(n,r) = n!/(r!(n-r)!)$ — order does not matter.
- $C(n,r) = C(n, n-r)$; circular permutations of $n = (n-1)!$.
- Permutations with repetition of n items = n^r .

Concept	Formula
Permutation	$n!/(n-r)!$
Combination	$n!/(r!(n-r)!)$
Circular perm	$(n-1)!$
With repetition	n^r

CBT Trap

- **Order matters** → permutation; **order does not** → combination. Mixing these is the #1 counting error.

6.3 Pigeonhole Principle

- If n items are placed in k boxes and $n > k$, some box has ≥ 2 items.
- Generalized: some box has $\geq \lceil n/k \rceil$ items.

CBT Trap

- Use **ceiling** $\lceil n/k \rceil$, not floor, for the guaranteed minimum in some box.

6.4 Inclusion-Exclusion (Counting)

- $|A \cup B| = |A| + |B| - |A \cap B|$ extends to derangements and onto-function counts.
- Number of **derangements** $D(n) = n! \sum (-1)^k/k!$.

6.5 Binomial Theorem and Coefficients

- $(x + y)^n = \sum C(n,k) x^{n-k} y^k$.
- Sum of coefficients = 2^n (set $x=y=1$); $\sum (-1)^k C(n,k) = 0$.
- General term $T(k+1) = C(n,k) x^{n-k} y^k$.

CBT Trap

- The **$(k+1)$ -th term** uses $C(n, k)$; off-by-one indexing is a common error.

6.6 Recurrence Relations

6.6.1 Forming Recurrences

- Express a_n in terms of earlier terms, e.g., $a_n = a_{(n-1)} + a_{(n-2)}$ (Fibonacci).

6.6.2 Solving Linear Homogeneous Recurrences

- For $a_n = c_1 a_{(n-1)} + c_2 a_{(n-2)}$, form **characteristic equation** $x^2 = c_1 x + c_2$.
- Distinct roots r_1, r_2 : $a_n = A r_1^n + B r_2^n$.
- Repeated root r : $a_n = (A + Bn) r^n$.

6.6.3 Particular Solutions

- For non-homogeneous, total solution = **homogeneous + particular**.

Root case	General solution
Distinct r_1, r_2	$A r_1^n + B r_2^n$
Repeated r	$(A+Bn) r^n$
Complex	trigonometric form

CBT Trap

- Repeated roots** require the extra factor n ($(A+Bn)r^n$); forgetting it is a classic mistake.

6.7 Generating Functions

- A **generating function** encodes a sequence as coefficients of a power series $\sum a_n x^n$.
- $1/(1-x) = \sum x^n$ is the basic generating function.

CBT Trap

- Generating functions turn **recurrences** into **algebraic equations** in the series variable.

7. Matrices and Determinants

7.1 Types of Matrices

- **Diagonal matrix**: non-zero entries only on the main diagonal.
- **Identity matrix I** : diagonal of **1s**; $AI = IA = A$.
- **Symmetric matrix**: $A = A^T$ ($a_{ij} = a_{ji}$).
- **Skew-symmetric**: $A = -A^T$; diagonal entries are **0**.
- **Orthogonal matrix**: $AA^T = I$, so $A^{-1} = A^T$; determinant ± 1 .

Type	Defining property
Symmetric	$A = A^T$
Skew-symmetric	$A = -A^T$, diag 0
Orthogonal	$AA^T = I$
Identity	diag of 1s

CBT Trap

- **Skew-symmetric** forces **zero diagonal**; an odd-order skew-symmetric matrix has determinant **0**.

7.2 Matrix Operations

- Addition is **element-wise** and requires same dimensions.
- Multiplication: $(AB)_{ij} = \sum a_{ik} b_{kj}$; defined when **columns of A = rows of B** .
- Matrix multiplication is **associative** and **distributive** but **NOT commutative**.
- $(AB)^T = B^T A^T$; $(AB)^{-1} = B^{-1} A^{-1}$ (reverse order).
- Naive multiplication of $n \times n$ matrices is $O(n^3)$; **Strassen** is $O(n^{2.81})$.

CBT Trap

- $AB \neq BA$ in general; transpose and inverse of a product **reverse the order**.

7.3 Determinant: Properties and Computation

- $\det(I) = 1$; $\det(A^T) = \det(A)$.
- $\det(AB) = \det(A) \cdot \det(B)$.
- $\det(kA) = k^n \det(A)$ for $n \times n$.
- Swapping two rows **multiplies det by -1** ; identical rows give **det = 0**.
- $\det(A^{-1}) = 1/\det(A)$.
- 2×2 : **det = $ad - bc$** .

CBT Trap

- $\det(kA) = k^n \det(A)$, NOT $k \cdot \det(A)$ — the scalar is raised to the order n .
- $\det(A+B) \neq \det(A) + \det(B)$ in general.

7.4 Inverse of a Matrix; Adjoint

- $A^{-1} = \text{adj}(A) / \det(A)$, exists iff $\det(A) \neq 0$ (non-singular).
- $\text{adj}(A)$ = **transpose of the cofactor matrix**.
- $A \cdot \text{adj}(A) = \det(A) \cdot I$.

- $\det(\text{adj } A) = \det(A)^{(n-1)}$.

CBT Trap

- A matrix is invertible iff $\det \neq 0$ iff **full rank** iff **0 is not an eigenvalue**.

7.5 Rank of a Matrix

- **Rank** = number of **linearly independent rows (= columns)** = order of the largest non-zero minor.
- Rank equals the number of **non-zero rows in row-echelon form**.
- For $n \times n$: **full rank $n \Leftrightarrow$ non-singular**.

CBT Trap

- Rank of a **non-zero matrix** is at least **1**; rank 0 only for the **zero matrix**.

7.6 Elementary Row Operations

- Three types: **row swap**, **scaling a row**, **adding a multiple of one row to another**.
- They preserve **rank** and the **solution set** of a system (but can change determinant).

8. System of Linear Equations

8.1 Homogeneous vs Non-Homogeneous

- **Homogeneous:** $Ax = 0$; always has the **trivial solution** $x = 0$.
- **Non-homogeneous:** $Ax = b$, $b \neq 0$.

8.2 Consistency and Number of Solutions (Rank Method)

- Let $r = \text{rank}(A)$, $r' = \text{rank}([A|b])$, n = number of unknowns.
- **Inconsistent** (no solution) if $r < r'$.
- **Unique solution** if $r = r' = n$.
- **Infinite solutions** if $r = r' < n$ ($n - r$ free variables).

Condition	Solutions
$r < r'$	none (inconsistent)
$r = r' = n$	unique
$r = r' < n$	infinite

- Homogeneous $Ax=0$ has non-trivial solutions iff **rank** $< n$ iff **det(A) = 0**.

CBT Trap

- Always compare rank of **A** vs the **augmented matrix** $[A|b]$; equality decides consistency.

8.3 Gaussian Elimination

- Reduce $[A|b]$ to **upper triangular (row-echelon)** form, then **back-substitute**.
- Complexity $O(n^3)$.

8.4 Cramer's Rule

- For $Ax = b$ with $\det(A) \neq 0$: $x_i = \det(A_i)/\det(A)$, where A_i replaces column i with b .
- Practical only for **small systems** (2×2 , 3×3).

CBT Trap

- **Cramer's rule fails when $\det(A) = 0$** ; it cannot detect infinite vs no solution.

9. Eigenvalues and Eigenvectors

9.1 Characteristic Equation

- $\mathbf{Av} = \lambda \mathbf{v}$ where λ is an **eigenvalue**, $v \neq 0$ the **eigenvector**.
- Characteristic equation: $\det(\mathbf{A} - \lambda \mathbf{I}) = 0$.

9.2 Computation

- Solve the **characteristic polynomial** for λ , then solve $(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0}$ for each eigenvector.

9.3 Properties

- **Sum of eigenvalues = trace(A)** (sum of diagonal).
- **Product of eigenvalues = det(A)**.
- Eigenvalues of a **triangular matrix** are its **diagonal entries**.
- **Symmetric matrices** have **real** eigenvalues; orthogonal eigenvectors.
- Eigenvalues of $\mathbf{A}^{-1}$ are $1/\lambda$; of $\mathbf{A}^k$ are λ^k .

Property	Result
Sum of eigenvalues	trace
Product	determinant
Triangular matrix	diagonal entries
$\mathbf{A}^{-1}$ eigenvalues	$1/\lambda$

CBT Trap

- **Trace = sum, det = product** of eigenvalues — fast checks. A singular matrix has **0** as an eigenvalue.

9.4 Cayley-Hamilton Theorem

- Every square matrix **satisfies its own characteristic equation**.
- Used to compute $\mathbf{A}^{-1}$ and high powers of $\mathbf{A}$ without direct inversion.

9.5 Diagonalization

- $\mathbf{A} = \mathbf{PDP}^{-1}$ where $\mathbf{D}$ is diagonal of eigenvalues, $\mathbf{P}$ columns are eigenvectors.
- $\mathbf{A}$ is diagonalizable iff it has **n linearly independent eigenvectors**.

CBT Trap

- A matrix with **repeated eigenvalues** may NOT be diagonalizable if it lacks enough independent eigenvectors.

10. LU Decomposition

10.1 LU Factorization Concept and Method

- **$A = LU$** : L is **lower triangular**, U is **upper triangular**.
- **Doolittle**: L has unit diagonal; **Crout**: U has unit diagonal.
- Exists when all **leading principal minors are non-zero** (else needs pivoting $\rightarrow PA = LU$).

10.2 Solving Systems Using LU

- Solve **$Ly = b$** by **forward substitution**, then **$Ux = y$** by **back substitution**.
- Efficient for **multiple right-hand sides** since L,U are reused.
- Factorization cost **$O(n^3)$** ; each solve **$O(n^2)$** .

CBT Trap

- LU is preferred over re-inverting A for **repeated b vectors**; factor once, solve many.

11. Limits, Continuity, Differentiability

11.1 Limits and Standard Limit Results

- $\lim (\sin x)/x = 1$ as $x \rightarrow 0$.
- $\lim (1 - \cos x)/x^2 = 1/2$ as $x \rightarrow 0$.
- $\lim (e^x - 1)/x = 1$ as $x \rightarrow 0$.
- $\lim (1 + 1/n)^n = e$ as $n \rightarrow \infty$.
- L'Hopital's rule: for $0/0$ or ∞/∞ , $\lim f/g = \lim f'/g'$.

CBT Trap

- L'Hopital applies only to **indeterminate forms $0/0$ or ∞/∞** ; applying it elsewhere gives wrong answers.

11.2 Continuity

- f is **continuous at a** if $\lim_{x \rightarrow a} f(x) = f(a)$ (limit exists and equals the value).
- Continuous on an interval if continuous at every point.

11.3 Differentiability; Relation to Continuity

- f is **differentiable** at a if $f'(a)$ exists.
- **Differentiable $\Rightarrow$ continuous**, but **continuous $\nRightarrow$ differentiable** (e.g., $|x|$ at 0).

CBT Trap

- $|x|$ is continuous everywhere but **not differentiable at $x = 0$** (corner point).

11.4 Derivative Rules

- **Product rule:** $(uv)' = u'v + uv'$.
- **Quotient rule:** $(u/v)' = (u'v - uv')/v^2$.
- **Chain rule:** $d/dx f(g(x)) = f'(g(x)) \cdot g'(x)$.
- $d/dx x^n = n x^{n-1}$; $d/dx e^x = e^x$; $d/dx \ln x = 1/x$; $d/dx \sin x = \cos x$.

CBT Trap

- Quotient rule numerator is **$u'v - uv'$** (order matters); reversing the sign is a frequent slip.

12. Applications of Derivatives

12.1 Maxima and Minima

- **Critical points**: where $f'(x) = 0$ or f' undefined.
- **Local max**: f' changes + to -; **local min**: - to +.
- **Global extrema** on $[a,b]$: check critical points AND **endpoints**.

12.2 Critical Points; First and Second Derivative Tests

- **First derivative test**: sign change of f' .
- **Second derivative test**: $f'' > 0 \Rightarrow$ minimum, $f'' < 0 \Rightarrow$ maximum, $f'' = 0 \Rightarrow$ inconclusive.
- **Inflection point**: where $f'' = 0$ and concavity changes.

Test	Max	Min
f' sign	+ to -	- to +
f'' value	$f'' < 0$	$f'' > 0$

CBT Trap

- $f'' = 0$ is **inconclusive**; must fall back to the first-derivative test.

12.3 Mean Value Theorems

- **Rolle's theorem**: if f continuous on $[a,b]$, differentiable on (a,b) , and $f(a)=f(b)$, then some c has $f'(c)=0$.
- **Lagrange's MVT**: some c has $f'(c) = (f(b)-f(a))/(b-a)$.

CBT Trap

- **Rolle** is a special case of **Lagrange** when $f(a)=f(b)$; both need continuity on closed and differentiability on open interval.

13. Integration

13.1 Indefinite Integrals; Standard Integrals

- $\int x^n dx = x^{(n+1)}/(n+1) + C$ ($n \neq -1$).
- $\int (1/x) dx = \ln|x| + C$.
- $\int e^x dx = e^x + C$; $\int \sin x dx = -\cos x + C$; $\int \cos x dx = \sin x + C$.

13.2 Methods: Substitution, By Parts, Partial Fractions

- **Integration by parts:** $\int u dv = uv - \int v du$ (choose u by **ILATE**).
- **Substitution:** change variable to simplify.
- **Partial fractions:** decompose rational functions before integrating.

Memory Trick

- **ILATE:** Inverse, Logarithmic, Algebraic, Trigonometric, Exponential — order to pick u in by-parts.

13.3 Definite Integrals and Properties

- $\int_a^b f dx = F(b) - F(a)$ (Fundamental Theorem).
- $\int_a^b f = -\int_b^a f$; $\int_a^a f = 0$.
- $\int_0^a f(x) dx = \int_0^a f(a-x) dx$ (king property).

13.4 Area Under a Curve

- Area between curve and x-axis = $\int_a^b |f(x)| dx$.

CBT Trap

- For signed regions use $|f(x)|$ or split at zeros; otherwise positive and negative areas cancel.

14. Probability

14.1 Sample Space, Events; Axioms

- **Sample space S**: set of all outcomes; an **event** is a subset of S.
- Axioms: $0 \leq P(A) \leq 1$; $P(S) = 1$; for disjoint events $P(A \cup B) = P(A) + P(B)$.
- General addition: $P(A \cup B) = P(A) + P(B) - P(A \cap B)$.

14.2 Conditional Probability; Multiplication Rule

- $P(A|B) = P(A \cap B) / P(B)$, $P(B) > 0$.
- Multiplication: $P(A \cap B) = P(A|B) \cdot P(B)$.

14.3 Independent Events

- A, B independent iff $P(A \cap B) = P(A) \cdot P(B)$, equivalently $P(A|B) = P(A)$.

CBT Trap

- **Independent $\neq$ mutually exclusive**; mutually exclusive events with non-zero probability are **dependent** ($P(A \cap B) = 0 \neq P(A)P(B)$).

14.4 Bayes' Theorem and Total Probability

- **Total probability**: $P(A) = \sum P(A|B_i) \cdot P(B_i)$.
- **Bayes**: $P(B_i|A) = P(A|B_i)P(B_i) / \sum P(A|B_j)P(B_j)$.

CBT Trap

- Bayes **reverses the conditioning**; the denominator is the **total probability** of the evidence.

14.5 Random Variables: Discrete and Continuous

- **Discrete RV**: countable values, described by a **PMF**.
- **Continuous RV**: uncountable values, described by a **PDF**.

14.6 PMF/PDF; CDF

- **PMF**: $P(X = x)$, sums to 1.
- **PDF** $f(x)$: $\int f(x) dx = 1$; $P(X=x) = 0$ for continuous.
- **CDF** $F(x) = P(X \leq x)$; non-decreasing from 0 to 1.

CBT Trap

- For a **continuous** RV, $P(X = a) = 0$; only intervals carry probability.

14.7 Expectation, Variance, Standard Deviation

- $E[X] = \sum x \cdot P(x)$ (discrete) or $\int x f(x) dx$ (continuous).
- $\text{Var}(X) = E[X^2] - (E[X])^2$.
- $SD = \sqrt{\text{Var}}$.
- $E[aX+b] = aE[X]+b$; $\text{Var}(aX+b) = a^2\text{Var}(X)$.

- $E[X+Y] = E[X]+E[Y]$ always; $\text{Var}(X+Y)=\text{Var}(X)+\text{Var}(Y)$ only if **independent**.

CBT Trap

- $\text{Var}(aX+b) = a^2\text{Var}(X)$ — the constant b drops out and a is **squared**.

15. Probability Distributions

15.1 Uniform Distribution

- Discrete uniform on n values: each probability $1/n$; mean $(a+b)/2$.
- Continuous uniform on $[a,b]$: $f = 1/(b-a)$, mean $(a+b)/2$, variance $(b-a)^2/12$.

15.2 Binomial Distribution

- $P(X=k) = C(n,k) p^k (1-p)^{n-k}$.
- Mean np ; variance $np(1-p)$.
- Models n independent trials with two outcomes.

15.3 Poisson Distribution

- $P(X=k) = e^{-\lambda} \lambda^k / k!$.
- Mean = variance = λ .
- Limit of binomial when n large, p small, $np = \lambda$.

15.4 Normal Distribution

- Bell curve, parameters mean μ , variance σ^2 .
- Standard normal $Z = (X - \mu)/\sigma$, mean 0, variance 1.
- Empirical rule: 68% within 1σ , 95% within 2σ , 99.7% within 3σ .
- Symmetric; mean = median = mode.

15.5 Exponential Distribution

- $f(x) = \lambda e^{-\lambda x}$, $x \geq 0$.
- Mean $1/\lambda$; variance $1/\lambda^2$.
- Memoryless property: $P(X>s+t | X>s) = P(X>t)$.

Distribution	Mean	Variance
Binomial	np	$np(1-p)$
Poisson	λ	λ
Uniform $[a,b]$	$(a+b)/2$	$(b-a)^2/12$
Exponential	$1/\lambda$	$1/\lambda^2$
Normal	μ	σ^2

CBT Trap

- Poisson mean = variance = λ ; the exponential (continuous, memoryless) and Poisson (discrete count) describe the same process differently.

16. Statistics

16.1 Measures of Central Tendency

- **Mean** = sum/count (sensitive to outliers).
- **Median** = middle value when sorted (robust to outliers).
- **Mode** = most frequent value.
- Empirical relation: **Mode = 3·Median – 2·Mean**.

CBT Trap

- For **skewed** data the **median** is preferred; mean is pulled toward the tail.

16.2 Measures of Dispersion

- **Range** = max – min.
- **Variance** $\sigma^2 = \sum(x-\mu)^2/N$ (population) or $/(n-1)$ (sample).
- **Standard deviation** = $\sqrt{\text{variance}}$.
- **Coefficient of variation** = σ/μ .

CBT Trap

- **Sample variance** divides by **n-1** (Bessel's correction); population divides by **N**.

16.3 Correlation and Regression

- **Pearson correlation** $r \in [-1, 1]$; $r = \text{cov}(X,Y)/(\sigma_x \sigma_y)$.
- **r = 0** means no **linear** correlation.
- **Regression line**: $y = a + bx$, slope **b** = $\text{cov}(X,Y)/\text{Var}(X)$.
- **r** = $\sqrt{(b_{xy} \cdot b_{yx})}$ (geometric mean of regression coefficients).

CBT Trap

- **Correlation ≠ causation**; also **r = 0** rules out only **linear**, not all, dependence.

17. Numerical Solutions of Equations

17.1 Non-linear Equations

17.1.1 Bisection Method

- Needs $f(a) \cdot f(b) < 0$ (sign change brackets a root).
- Each step **halves** the interval; **linear** convergence (order **1**).
- Error after n steps $\leq (b-a)/2^n$.
- Always **converges** (guaranteed) but slowly.

17.1.2 Newton-Raphson Method

- Iteration: $x_{(n+1)} = x_n - f(x_n)/f'(x_n)$.
- Quadratic** convergence (order **2**) near a simple root.
- Fails if $f'(x_n) = 0$ or poor initial guess.

17.1.3 Secant / Regula-Falsi

- Secant**: $x_{(n+1)} = x_n - f(x_n)(x_n - x_{(n-1)})/(f(x_n) - f(x_{(n-1)}))$; order ≈ 1.618 .
- Regula-falsi**: like secant but keeps the **bracketing** interval (guaranteed convergence).

Method	Convergence order	Bracketing
Bisection	1 (linear)	yes
Newton-Raphson	2 (quadratic)	no
Secant	1.618	no
Regula-falsi	linear	yes

CBT Trap

- Newton-Raphson** is fastest (order **2**) but needs f' and may diverge; **bisection** is slow but **guaranteed**.

17.2 Linear Systems

17.2.1 Gauss Elimination

- Direct method; forward elimination then back substitution; $O(n^3)$.

17.2.2 Gauss-Seidel Iterative Method

- Iterative; uses **updated values immediately** (faster than Jacobi).
- Converges if matrix is **diagonally dominant**.

CBT Trap

- Gauss-Seidel** uses new values within the same iteration; **Jacobi** uses only previous-iteration values.

17.3 Convergence and Error

- Absolute error** = $|true - approx|$; **relative error** = $absolute/|true|$.
- Diagonal dominance guarantees iterative convergence.

18. Numerical Integration

18.1 Trapezoidal Rule

- $\int_a^b f \, dx \approx (h/2)[f_0 + 2(f_1 + \dots + f_{n-1}) + f_n]$, $h = (b-a)/n$.
- Error $O(h^2)$; exact for **linear** (degree 1) polynomials.

18.2 Simpson's 1/3 Rule

- $\int \approx (h/3)[f_0 + 4(\text{odd}) + 2(\text{even}) + f_n]$; needs **even** number of intervals.
- Error $O(h^4)$; exact for **cubic** (degree 3) polynomials.

18.3 Error in Numerical Integration

Rule	Error order	Exact up to degree
Trapezoidal	$O(h^2)$	1
Simpson's 1/3	$O(h^4)$	3
Simpson's 3/8	$O(h^4)$	3

CBT Trap

- **Simpson's 1/3 needs an EVEN number of subintervals** (odd number of points); trapezoidal has no such restriction.
- Simpson's is exact for **cubics** despite using parabolas — a famous bonus accuracy.

19. Numerical Solution of ODEs

19.1 Single-Step Methods

- **Euler's method:** $y_{n+1} = y_n + h \cdot f(x_n, y_n)$; first-order accurate, error **O(h)** per step.
- **RK4 (Runge-Kutta 4th order):** $y_{n+1} = y_n + (h/6)(k_1 + 2k_2 + 2k_3 + k_4)$.
- RK4 slopes: $k_1=f(x,y)$, $k_2=f(x+h/2, y+hk_1/2)$, $k_3=f(x+h/2, y+hk_2/2)$, $k_4=f(x+h, y+hk_3)$.

Method	Order	Slopes used
Euler	1	1
RK2	2	2
RK4	4	4

19.2 Multi-Step Methods

- Use **several previous points** (e.g., **Adams-Bashforth**, **Adams-Moulton**).
- **Predictor-corrector** pairs an explicit predictor with an implicit corrector.

CBT Trap

- **Euler is first-order** (least accurate); **RK4** weights the middle slopes **2,2** and is **fourth-order**.

20. Exam Focus / Practice Checklist

20.1 Propositional Logic Equivalence / Validity

- Test equivalence via **truth tables** or laws; check argument **validity** with rules of inference.
- Remember $p \rightarrow q \equiv \neg p \vee q$ and only the **contrapositive** is equivalent.

20.2 Relation/Function Property Identification

- Classify by **reflexive, symmetric, antisymmetric, transitive**; **RST = equivalence**, **RAT = partial order**.

20.3 Lattice / Hasse Diagram Questions

- Confirm **LUB and GLB** exist for all pairs; identify **0 and 1**, distributive/complemented.

20.4 Counting

- Decide **permutation (order matters) vs combination**; apply **pigeonhole** $\lceil n/k \rceil$.

20.5 Recurrence Solving

- Use the **characteristic equation**; remember $(A+Bn)r^n$ for repeated roots.

20.6 Matrix Rank / Determinant / Inverse

- Use **row-echelon** for rank; $A^{-1} = \text{adj}(A)/\det(A)$; $\det(kA) = k^n \det(A)$.

20.7 Eigenvalue Computation and Properties

- **trace** = $\sum \lambda$, **det** = $\prod \lambda$; use **Cayley-Hamilton** for powers/inverse.

20.8 Probability

- **Bayes** for reverse conditioning; memorize **means/variances** of binomial, Poisson, normal, exponential.

20.9 Newton-Raphson / Trapezoidal / Simpson

- **NR**: $x - f/f'$ (order 2); **Trapezoidal** $O(h^2)$; **Simpson** $1/3 O(h^4)$, even intervals.

20.10 Graph Theory

- **Handshaking** $\sum \deg = 2E$; **Euler** $V - E + F = 2$; **χ coloring**; Euler vs Hamilton conditions.

Final Revision Trick

- Memorize the **distribution mean/variance table**, the **convergence-order table**, and the **eigenvalue trace/det shortcuts** — these recur in nearly every CIL MT mathematics paper.