

Learning Notes for Algebraic Geometry

Typed by
Bowen Liu¹

¹liubw22@mails.tsinghua.edu.cn

CONTENTS

Part 1. Preliminaries	3
1. Category theory	3
1.1. Category	3
1.2. Abelian category	4
2. Sheaf and cohomology	6
2.1. Sheaves	6
2.2. Derived functor formulation of sheaf cohomology	12
2.3. Acyclic resolution	14
2.4. Examples about acyclic sheaf	15
2.5. Proof of de Rham theorem using sheaf cohomology	20
2.6. Hypercohomology	21
Part 2. Schemes	23
3. Schemes and Morphisms	23
3.1. Schemes	23
3.2. Proj construction	25
4. Properties of schemes	27
4.1. Quasi-compact, irreducible, Noetherian topological space	27
4.2. Reduced, irreducible and integral scheme	28
4.3. Affine criterion	28
4.4. Noetherian scheme	29
5. Properties of morphisms	31
5.1. Quasi-compact, affine, finite type and finite	31
5.2. Birational morphism	33
5.3. Open immersion and closed immersion	34
5.4. Fibred product	35
5.5. Separated morphisms	36
5.6. Proper and projective morphisms	38
6. Coherent sheaves	40
6.1. $\mathcal{O}_X$ -modules	40
6.2. Coherent sheaves	41
6.3. Sheaf of ideals	43
6.4. Coherent sheaves on Proj	44
Part 3. Homework	47
7. Homework	47
7.1. Homework-1	47
7.2. Homework-2	52
7.3. Homework-3	58
7.4. Homework-4	60
7.5. Homework-5	61
7.6. Homework-6	65
7.7. Homework-7	67

7.8. Homework-8	69
7.9. Homework-9	71
7.10. Homework-10	78
7.11. Homework-11	80
7.12. Homework-12	83
7.13. Homework-13	86
References	89

Part 1. Preliminaries

1. CATEGORY THEORY

1.1. Category.

1.1.1. Category and Functors.

1.1.2. Morphisms.

Definition 1.1.1. A morphism $f: A \rightarrow B$ in $\mathcal{C}$ is called a **monomorphism** if for any two morphisms $\alpha, \beta: C \rightarrow A$ satisfying $f \circ \alpha = f \circ \beta$, we have $\alpha = \beta$.

Definition 1.1.2. A morphism $f: A \rightarrow B$ in $\mathcal{C}$ is called an **epimorphism** if for any two morphisms $\alpha, \beta: A \rightarrow C$ satisfying $\alpha \circ f = \beta \circ f$, we have $\alpha = \beta$.

Definition 1.1.3. A morphism is called a **bimorphism** if it is both a monomorphism and an epimorphism.

Definition 1.1.4. A morphism is called an **isomorphism** if it admits a two-sided inverse.

Remark 1.1.1. Every isomorphism is a bimorphism, but in general a bimorphism need not be an isomorphism. For example, in the category of rings, the canonical morphism $\mathbb{Z} \rightarrow \mathbb{Q}$ is both a monomorphism and an epimorphism, but it is not an isomorphism.

1.1.3. Categorical objects.

Definition 1.1.5. Let $\{A_i\}_{i \in I}$ be a family of objects in the category $\mathcal{C}$. The **direct product** of the family $\{A_i\}_{i \in I}$ is a tuple $(\prod_{i \in I} A_i, p_i)$, where $\prod_{i \in I} A_i$ is an object in $\mathcal{C}$, and $p_i: \prod_{i \in I} A_i \rightarrow A_i$ is a family of morphisms called the projections, satisfying the following universal property: For any object C and any family of morphisms $f_i: C \rightarrow A_i$, there exists a unique morphism $f: C \rightarrow \prod_{i \in I} A_i$ such that $p_i \circ f = f_i$ for all $i \in I$.

$$\begin{array}{ccc} \prod_{i \in I} A_i & \xleftarrow{\quad f \quad} & C \\ p_i \downarrow & \swarrow f_i & \\ A_i & & \end{array}$$

Definition 1.1.6. Let $\{A_i\}_{i \in I}$ be a family of objects in the category $\mathcal{C}$. The **direct sum** of the family $\{A_i\}_{i \in I}$ is a tuple $(\oplus_{i \in I} A_i, k_i)$, where $\oplus_{i \in I} A_i$ is an object in $\mathcal{C}$, and $k_i: A_i \rightarrow \oplus_{i \in I} A_i$ is a family of morphisms called the canonical injections, satisfying the following universal property: For any object C and any family of morphisms $f_i: A_i \rightarrow C$, there exists a unique morphism $f: \oplus_{i \in I} A_i \rightarrow C$ such that $f \circ k_i = f_i$ for all $i \in I$.

$$\begin{array}{ccc} \oplus_{i \in I} A_i & \xrightarrow{\quad f \quad} & C \\ k_i \uparrow & \searrow f_i & \\ A_i & & \end{array}$$

1.2. Abelian category.

1.2.1. Additive category.

Definition 1.2.1. A category $\mathcal{C}$ is called an **additive category** if it satisfies the following conditions for any objects A, B, C in $\mathcal{C}$:

- (1) the **direct product** of A and B exists;
- (2) $\text{Hom}(A, B)$ is an abelian group, and $0 \in \text{Hom}(A, B)$ is called the **zero morphism**;
- (3) there exists a **zero object** $0 \in \mathcal{C}$, which is both initial and terminal;
- (4) the map

$$\begin{aligned} \text{Hom}(A, B) \times \text{Hom}(B, C) &\rightarrow \text{Hom}(A, C) \\ (f, g) &\mapsto g \circ f \end{aligned}$$

is bilinear.

Remark 1.2.1.

- (1) $\mathcal{C}$ has all finite products and coproducts, and they coincide.
- (2) the **zero object** is both an initial object and a terminal object.


Definition 1.2.2. Let $\mathcal{C}$ and $\mathcal{D}$ be two additive categories. A functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is called an **additive functor** if

$$\text{Hom}(A, B) \rightarrow \text{Hom}(F(A), F(B))$$

is a group homomorphism for all $A, B \in \text{Ob}(\mathcal{C})$.

Definition 1.2.3. Let $\mathcal{C}$ be a category, let $X, Y \in \text{Ob}(\mathcal{C})$, and let $f, g \in \text{Hom}(X, Y)$. The **equalizer** of f and g is an object E , together with a morphism $e: E \rightarrow X$ such that

- (1) $f \circ e = g \circ e$.
- (2) If there exists $e': E' \rightarrow X$ such that $f \circ e' = g \circ e'$, then there exists a unique $\theta: E' \rightarrow E$ such that $e' = e \circ \theta$.



Definition 1.2.4. Let $\mathcal{C}$ be an additive category and $f: A \rightarrow B$ be a morphism in $\mathcal{C}$.

- (1) The **kernel** of f is the **equalizer** of f and $0: A \rightarrow B$.
- (2) The **cokernel** of f is the **coequalizer** of f and the zero morphism $0: A \rightarrow B$.

Remark 1.2.2.

- (1) The kernel and cokernel may not exist in an additive category in general. For example, $\mathcal{O} \rightarrow \mathcal{O}(1)$ has no cokernel in the category of vector bundles.

- (2) A kernel is necessarily a monomorphism, and a cokernel is necessarily an epimorphism.
- (3) There is a natural morphism $\text{coim } f \rightarrow \text{im } f$ induced by the universal property

$$\begin{array}{ccccccc}
 \ker f & \longrightarrow & A & \xrightarrow{f} & B & \longrightarrow & \text{coker } f \\
 & & \downarrow & & \uparrow & & \\
 & & \text{coim } f & \dashrightarrow & \text{im } f & &
 \end{array}$$

1.2.2. Abelian category.

Definition 1.2.5. An abelian category $\mathcal{C}$ is an additive category with zero objects such that for every morphism f in $\mathcal{C}$, the kernel and the **cokernel** of f exist, and the canonical morphism $\text{coim } f \rightarrow \text{im } f$ is an isomorphism.

Proposition 1.2.1. In an abelian category, a bimorphism is an isomorphism.

Definition 1.2.6. In an abelian category, a sequence of morphisms

$$A \xrightarrow{u} B \xrightarrow{v} C$$

is called **exact** if $v \circ u = 0$ and the canonical morphism from $\text{coim } u \rightarrow \ker v$ is an isomorphism.

Definition 1.2.7. An exact sequence of the form

$$0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$$

is called a **short exact sequence**.

Definition 1.2.8. A short exact sequence $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ is called **split** if it is isomorphic to

$$0 \rightarrow A \rightarrow A \oplus C \rightarrow C \rightarrow 0,$$

where $A \rightarrow A \oplus C$ and $A \oplus C \rightarrow C$ are the canonical morphisms.

2. SHEAF AND COHOMOLOGY

2.1. Sheaves. Along this section, X denotes a topological space.

2.1.1. Definitions and Examples.

Definition 2.1.1. A presheaf of abelian groups $\mathcal{F}$ on X consists of the following data:

- (1) For any open subset U of X , $\mathcal{F}(U)$ is an abelian group.
- (2) If $U \subseteq V$ are two open subsets of X , then there is a group homomorphism $r_{VU}: \mathcal{F}(V) \rightarrow \mathcal{F}(U)$. Moreover, above data satisfy
 - I $\mathcal{F}(\emptyset) = 0$.
 - II $r_{UU} = \text{id}$.
 - III If $W \subseteq U \subseteq V$ are open subsets of X , then $r_{VW} = r_{UW} \circ r_{VU}$.

Moreover, $\mathcal{F}$ is called a **sheaf** if it satisfies the following extra conditions

- IV Let $\{V_i\}_{i \in I}$ be an open covering of open subset $U \subseteq X$ and $s \in \mathcal{F}(U)$. If $s|_{V_i} := r_{UV_i}(s) = 0$ for all $i \in I$, then $s = 0$.
- V Let $\{V_i\}_{i \in I}$ be an open covering of open subset $U \subseteq X$ and $s_i \in \mathcal{F}(V_i)$. If $s_i|_{V_i \cap V_j} = s_j|_{V_i \cap V_j}$ for all $i, j \in I$, then there exists $s \in \mathcal{F}(U)$ such that $s|_{V_i} = s_i$ for all $i \in I$.

Example 2.1.1 (constant presheaf). For an abelian group G , the constant presheaf assigns to each open subset U the group G itself, but in general it is not a sheaf.

Definition 2.1.2. A morphism $\varphi: \mathcal{F} \rightarrow \mathcal{G}$ between presheaves consists of the following data:

- (1) For any open subset U of X , there is a group homomorphism $\varphi(U): \mathcal{F}(U) \rightarrow \mathcal{G}(U)$.
- (2) If $U \subseteq V$ are two open subsets of X , then the following diagram commutes

$$\begin{array}{ccc} \mathcal{F}(V) & \xrightarrow{\varphi(V)} & \mathcal{G}(V) \\ \downarrow r_{VU} & & \downarrow r_{VU} \\ \mathcal{F}(U) & \xrightarrow{\varphi(U)} & \mathcal{G}(U) \end{array}$$

Notation 2.1.1. For convenience, for $s \in \mathcal{F}(U)$, we often write $\varphi(s)$ instead of $\varphi(U)(s)$.

Remark 2.1.1. The morphisms between sheaves are defined as morphisms of presheaves.

Definition 2.1.3. A morphism of presheaves $\varphi: \mathcal{F} \rightarrow \mathcal{G}$ is called an **isomorphism** if it has a two-sided inverse, that is, there exists a morphism of presheaves $\psi: \mathcal{G} \rightarrow \mathcal{F}$ such that $\psi \circ \varphi = \text{id}_{\mathcal{F}}$ and $\varphi \circ \psi = \text{id}_{\mathcal{G}}$.

Remark 2.1.2. A morphism of presheaves $\varphi: \mathcal{F} \rightarrow \mathcal{G}$ is an isomorphism if and only if for every open subset $U \subseteq X$, $\varphi(U): \mathcal{F}(U) \rightarrow \mathcal{G}(U)$ is an isomorphism of abelian groups.

2.1.2. Stalks.

Definition 2.1.4. For a presheaf $\mathcal{F}$ and $p \in X$, the stalk at p is defined as

$$\mathcal{F}_p = \varinjlim_{p \in U} \mathcal{F}(U)$$

Remark 2.1.3 (alternative definition). To avoid the language of direct limits, we give an equivalent description of the stalk. For $p \in U \cap V$, sections $s_U \in \mathcal{F}(U)$ and $s_V \in \mathcal{F}(V)$ are equivalent if there exists an open neighborhood $p \in W \subseteq U \cap V$ such that $s_U|_W = s_V|_W$. An element $s_p \in \mathcal{F}_p$, called a germ, is an equivalence class $[s_U]$; for $s \in \mathcal{F}(U)$, the germ defined by s is denoted by $s|_p$.

Notation 2.1.2.

- (1) For $s \in \mathcal{F}(U)$ and $p \in U$, $s|_p$ denotes the equivalence class it gives.
- (2) For $s_p \in \mathcal{F}_p$, a section $s \in \mathcal{F}(U)$ with $s|_p = s_p$ is called a representative of the germ; it is not unique.

Definition 2.1.5. Given a morphism of sheaves $\varphi: \mathcal{F} \rightarrow \mathcal{G}$, it induces a morphism of abelian groups $\varphi_p: \mathcal{F}_p \rightarrow \mathcal{G}_p$ as follows:

$$\begin{aligned} \varphi_p: \mathcal{F}_p &\rightarrow \mathcal{G}_p \\ s_p &\mapsto \varphi(s)|_p. \end{aligned}$$

Remark 2.1.4. It is necessary to check the φ_p is well-defined since there are different choices s such that $s|_p = s_p$.

Proposition 2.1.1. Let $\varphi: \mathcal{F} \rightarrow \mathcal{G}$ be a morphism between sheaves. Then φ is an isomorphism if and only if the induced map $\varphi_p: \mathcal{F}_p \rightarrow \mathcal{G}_p$ is an isomorphism for every $p \in X$.

Proof. It follows if φ is an isomorphism between sheaves, then it induces an isomorphism on stalks. Conversely, it suffices to show $\varphi(U): \mathcal{F}(U) \rightarrow \mathcal{G}(U)$ is an isomorphism for every open subset $U \subseteq X$.

- (1) Injectivity: For $s, s' \in \mathcal{F}(U)$ such that $\varphi(s) = \varphi(s')$, by passing to stalks one has $\varphi_p(s|_p) = \varphi_p(s'|_p)$ for every $p \in U$, and thus $s|_p = s'|_p$ since φ_p is an isomorphism. By definition of stalks there exists an open subset $V_p \subseteq U$ containing p such that s agrees with s' on V_p . Then it gives an open covering $\{V_p\}$ of U , and by axiom (IV) one has $s = s'$ on U .
- (2) Surjectivity: For $t \in \mathcal{G}(U)$, by passing to stalks there exists $s_p \in \mathcal{F}_p$ such that $\varphi_p(s_p) = t|_p$ for every $p \in U$ since φ_p is surjective. By definition of stalks there exists an open subset $V_p \subseteq U$ containing p and $s \in \mathcal{F}(V_p)$ such that $\varphi(s) = t$ on V_p . This gives a collection of sections defined on an open covering $\{V_p\}$ of U , and by injectivity we proved above one has these sections agree with each other on the intersections. Then by axiom (V) there exists a section $s \in \mathcal{F}(U)$ such that $\varphi(s) = t$.

□

2.1.3. Sheafification. In Example 2.1.1, we come across a presheaf that is not a sheaf. To obtain a sheaf from a presheaf, we require a process known as sheafification. One approach to defining sheafification is through its universal property.

Definition 2.1.6. Given a presheaf $\mathcal{F}$ there is a sheaf $\mathcal{F}^+$ and a morphism $\theta: \mathcal{F} \rightarrow \mathcal{F}^+$ with the property that for any sheaf $\mathcal{G}$ and any morphism $\varphi: \mathcal{F} \rightarrow \mathcal{G}$ there is a unique morphism $\bar{\varphi}: \mathcal{F}^+ \rightarrow \mathcal{G}$ such that the following diagram commutes:

$$\begin{array}{ccc} \mathcal{F} & \xrightarrow{\varphi} & \mathcal{G} \\ \downarrow \theta & \nearrow \bar{\varphi} & \\ \mathcal{F}^+ & & \end{array}$$

The universal property shows that if the sheafification exists, then it is unique up to a unique isomorphism. One way to give an explicit construction of sheafification is to glue stalks together in a suitable way. Let $\mathcal{F}^+(U)$ be a set of functions

$$f: U \rightarrow \coprod_{p \in U} \mathcal{F}_p$$

such that $f(p) \in \mathcal{F}_p$ and for every $p \in U$ there is an open subset $V_p \subseteq U$ containing p and $t \in \mathcal{F}(V_p)$ such that $t|_q = f(q)$ for all $q \in V_p$.

Proposition 2.1.2. $\mathcal{F}^+$ is the **sheafification** of $\mathcal{F}$.

Proof. First, we show $\mathcal{F}^+$ is a sheaf: It follows $\mathcal{F}^+$ is a presheaf, so it suffices to check conditions (IV) and (V) in the definition. Let $U \subseteq X$ be an open subset and $\{V_i\}$ be an open covering of U .

- (1) If $s \in \mathcal{F}^+(U)$ such that $s|_{V_i} = 0$ for all i , then s must be zero: It suffices to show $s(p) = 0$ for all $p \in U$. For any $p \in U$, then there exists an open subset V_i contains p , hence $s(p) = s|_{V_i}(p) = 0$.
- (2) Suppose there exists a collection of sections $\{s_i \in \mathcal{F}^+(V_i)\}_{i \in I}$ such that

$$s_i|_{V_i \cap V_j} = s_j|_{V_i \cap V_j}$$

holds for all $i, j \in I$. Now we construct $s \in \mathcal{F}^+(U)$ as follows: For $p \in U$ and V_i containing p , we define $s(p) = s_i(p)$. This is well-defined since s_i agree on the intersections, so it remains to show $s \in \mathcal{F}^+(U)$. It follows $s(p) \in \mathcal{F}_p$. For $p \in U$, there exists V_i containing p , and thus there exists $W_i \subseteq V_i$ containing p and $t \in \mathcal{F}(W_i)$ such that $t|_q = s_i(q) = s(q)$ for all $q \in W_i$.

There is a canonical morphism $\theta: \mathcal{F} \rightarrow \mathcal{F}^+$ as follows: For open subset $U \subseteq X$, and $s \in \mathcal{F}(U)$, $\theta(s)$ is defined by

$$\begin{aligned} \theta(s): U &\rightarrow \coprod_{p \in U} \mathcal{F}_p \\ p &\mapsto s|_p. \end{aligned}$$

Note that if $\mathcal{F}$ is a sheaf, the canonical morphism $\theta: \mathcal{F} \rightarrow \mathcal{F}^+$ is an isomorphism.

- (1) Injectivity: If $s \in \mathcal{F}(U)$ such that $s|_p = 0$ for all $p \in U$, then there exists an open covering $\{V_i\}_{i \in I}$ of U such that $s|_{V_i} = 0$, by axiom (IV) of sheaf one has $s = 0$.
- (2) Surjectivity: For $f \in \mathcal{F}^+(U)$ and every $p \in U$, the definition of $\mathcal{F}^+$ gives a neighborhood $p \in V_p \subseteq U$ and a section $t_p \in \mathcal{F}(V_p)$ such that $f(q) = t_p|_q$ for every $q \in V_p$. On overlaps, t_p and $t_{p'}$ have the same germ at every point, hence agree as sections. The sheaf axiom glues them to a section $s \in \mathcal{F}(U)$ satisfying $\theta(s) = f$.

Finally, we show $\mathcal{F}^+$ satisfies the universal property of sheafification. A morphism of presheaves $\varphi: \mathcal{F} \rightarrow \mathcal{G}$ induces a map on stalks

$$\varphi_p: \mathcal{F}_p \rightarrow \mathcal{G}_p.$$

For $f \in \mathcal{F}^+(U)$, the composite of f with the map

$$\coprod_{p \in U} \varphi_p: \coprod_{p \in U} \mathcal{F}_p \rightarrow \coprod_{p \in U} \mathcal{G}_p$$

gives a map $\tilde{\varphi}(f): U \rightarrow \coprod_{p \in U} \mathcal{G}_p$, and in fact $\tilde{\varphi}(f) \in \mathcal{G}^+(U)$: For $p \in U$, $\tilde{\varphi}(f)(p) \in \mathcal{G}_p$ since $f(p) \in \mathcal{F}_p$ and $\varphi_p: \mathcal{F}_p \rightarrow \mathcal{G}_p$. If for all $q \in V_p$ we have $t|_q = f(q)$, then

$$\tilde{\varphi}(f)(q) = \varphi_q(f(q)) = \varphi_q(t|_q) = \varphi(t)|_q.$$

Since $\mathcal{G}$ is a sheaf, the canonical morphism $\theta': \mathcal{G} \rightarrow \mathcal{G}^+$ is an isomorphism, and so we can define $\bar{\varphi} := \theta'^{-1} \circ \tilde{\varphi}$. We now show $\varphi = \bar{\varphi} \circ \theta = \theta'^{-1} \circ \tilde{\varphi} \circ \theta$. It is easy to show they coincide on each stalk since $\varphi_p = \theta_p'^{-1} \circ \tilde{\varphi}_p \circ \theta_p$, and thus $\varphi = \bar{\varphi} \circ \theta$ by Proposition 2.1.1. Furthermore, uniqueness follows from the fact that $\bar{\varphi}_p$ is uniquely determined by φ_p . $\square$

Remark 2.1.5. From the construction, one can see the **stalk** of $\mathcal{F}^+$ at p is exactly $\mathcal{F}_p$.

Remark 2.1.6. The sheafification can be described categorically. Let $\text{Sh}(X, \mathbf{Ab})$ and $\text{PSh}(X, \mathbf{Ab})$ denote the categories of sheaves and presheaves of abelian groups on X . The natural inclusion $\iota: \text{Sh}(X, \mathbf{Ab}) \hookrightarrow \text{PSh}(X, \mathbf{Ab})$ has sheafification as its left adjoint.

Example 2.1.2 (constant sheaf). For an abelian group G , the associated constant sheaf $\underline{G}$ is the **sheafification** of the constant presheaf. By the construction of sheafification, $\underline{G}$ can be explicitly expressed as

$$\underline{G}(U) = \{\text{locally constant function } f: U \rightarrow G\}$$

2.1.4. Exact sequence of sheaf. Given a morphism $\varphi: \mathcal{F} \rightarrow \mathcal{G}$ between sheaves of abelian groups, there are the following presheaves

$$U \mapsto \ker \varphi(U)$$

$$U \mapsto \text{im } \varphi(U)$$

$$U \mapsto \text{coker } \varphi(U),$$

since $\varphi(U): \mathcal{F}(U) \rightarrow \mathcal{G}(U)$ is a group homomorphism.

Proposition 2.1.3. Kernel of a morphism between sheaves is a sheaf.

Proof. Let $\{V_i\}_{i \in I}$ be an open covering of U .

- (1) For $s \in \ker \varphi(U)$, if $s|_{V_i} = 0$, then $s = 0$ since s is also in $\mathcal{F}(U)$.
- (2) If there exists $s_i \in \ker \varphi(V_i)$ such that $s_i|_{V_i \cap V_j} = s_j|_{V_i \cap V_j}$, then they glue together to get $s \in \mathcal{F}(U)$. Note that

$$\varphi(U)(s)|_{V_i} = \varphi(V_i)(s|_{V_i}) = \varphi(V_i)(s_i) = 0$$

Then $s \in \ker \varphi(U)$.

□

But the image of a morphism may not be a sheaf. Although we can prove the first requirement in the same way, the proof for the second requirement fails: Suppose $s_i \in \text{im } \varphi(V_i)$, and suppose that we can glue them together to get a $s \in \mathcal{G}(U)$, but s may not be the image of some $t \in \mathcal{F}(U)$. The cokernel fails to be a sheaf for the same reason.

Definition 2.1.7. Let $\varphi: \mathcal{F} \rightarrow \mathcal{G}$ be a morphism between sheaves of abelian groups. Then the image and the **cokernel** of φ are defined to be the **sheafification** of the following presheaves

$$U \mapsto \text{im } \varphi(U)$$

$$U \mapsto \text{coker } \varphi(U)$$

respectively.

Definition 2.1.8. For a sequence of sheaves:

$$\dots \rightarrow \mathcal{F}^{i-1} \xrightarrow{\varphi^{i-1}} \mathcal{F}^i \xrightarrow{\varphi^i} \mathcal{F}^{i+1} \rightarrow \dots$$

It is called **exact** at $\mathcal{F}^i$, if $\ker \varphi^i = \text{im } \varphi^{i-1}$. If a sequence is exact everywhere, then it is an exact sequence of sheaves.

Definition 2.1.9. An exact sequence of sheaves is called a **short exact sequence** if it looks like

$$0 \rightarrow \mathcal{F} \xrightarrow{\varphi} \mathcal{G} \xrightarrow{\psi} \mathcal{H} \rightarrow 0$$

Proposition 2.1.4. Let $\varphi: \mathcal{F} \rightarrow \mathcal{G}$ be a morphism between sheaves of abelian groups. Then for any $p \in X$, one has

$$(\ker \varphi)_p = \ker \varphi_p$$

$$(\text{im } \varphi)_p = \text{im } \varphi_p.$$

Proof. For (1), it follows $(\ker \varphi)_p \subseteq \ker \varphi_p$. Conversely, if $s_p \in \ker \varphi_p$, then $\varphi_p(s_p) = 0 \in \mathcal{G}_p$. In other words, there exists an open subset U containing p and $s \in \mathcal{F}(U)$ such that $s|_p = s_p$ and $\varphi(s)|_p = 0$, which implies there is another open subset V containing p such that $\varphi(s)|_V = 0$. Hence $\varphi(s|_V) = 0$, that is, $s|_V \in \ker \varphi(V)$. Thus $s_p = (s|_V)|_p \in (\ker \varphi)_p$.

For (2), It follows $(\text{im } \varphi)_p \subseteq \text{im } \varphi_p$ since the sheafification does not change stalk. Conversely, if $s_p \in \text{im } \varphi_p$, then there exists $t_p \in \mathcal{F}_p$ such that $\varphi_p(t_p) = s_p$. Suppose $t \in \mathcal{F}(U)$ is a section of some open subset U containing p such that $t|_p = t_p$. Then $\varphi(t)|_p = \varphi_p(t_p) = s_p$. In other words, s_p is in the **stalk** of

the image presheaf at p , but the sheafification does not change stalk, so we have $s_p \in (\text{im } \varphi)_p$. $\square$

Corollary 2.1.1. The sequence of sheaves

$$\dots \rightarrow \mathcal{F}^{i-1} \xrightarrow{\varphi^{i-1}} \mathcal{F}^i \xrightarrow{\varphi^i} \mathcal{F}^{i+1} \rightarrow \dots$$

is exact if and only if the sequence of abelian groups is exact

$$\dots \rightarrow \mathcal{F}_p^{i-1} \xrightarrow{\varphi_p^{i-1}} \mathcal{F}_p^i \xrightarrow{\varphi_p^i} \mathcal{F}_p^{i+1} \rightarrow \dots$$

for all $p \in X$.

Corollary 2.1.2. The sequence of sheaves

$$0 \rightarrow \mathcal{F} \rightarrow \mathcal{G}$$

is exact if and only if for any open subset U , the following sequence of abelian groups is exact

$$0 \rightarrow \mathcal{F}(U) \rightarrow \mathcal{G}(U).$$

Method one. For any open subset $U \subseteq X$, one has

$$\varphi(U): \mathcal{F}(U) \rightarrow \mathcal{G}(U)$$

is injective, since by definition we have for any open subset $U \subseteq X$, $\ker \varphi(U) = 0$, that is injectivity. $\square$

Method two. Or from another point of view, for each $p \in U$, we have

$$\varphi_p: \mathcal{F}_p \rightarrow \mathcal{G}_p$$

is injective. That is $\ker \varphi_p = 0$. So we obtain $(\ker \varphi(U))_p = 0$ for all $p \in U$. But for a section $s \in \mathcal{F}(U)$ if we have $s|_p = 0$, then we must have $s = 0$. So we obtain $\ker \varphi(U) = 0$. $\square$

Example 2.1.3 (exponential sequence). Let X be a complex manifold and $\mathcal{O}_X$ be its holomorphic function sheaf. Then

$$0 \rightarrow 2\pi\sqrt{-1}\mathbb{Z} \rightarrow \mathcal{O}_X \xrightarrow{\exp} \mathcal{O}_X^* \rightarrow 0$$

is an exact sequence of sheaves, called exponential sequence.

Proof. The difficulty is to show $\exp$ is surjective on stalks at $p \in X$. That is we need to construct logarithms of functions $g \in \mathcal{O}_X^*(U)$ for U , a neighborhood of p . We may choose U is simply-connected, then define

$$\log g(q) = \log g(p) + \int_{\gamma_q} \frac{dg}{g}$$

for $q \in U$, where γ_q is a path from p to q in U , and the definition is independent of the choice of γ_q since U is simply-connected. $\square$

Remark 2.1.7. In fact, U is simply-connected is crucial for constructing logarithm. If we consider $X = \mathbb{C}$ and $U = \mathbb{C} \setminus \{0\}$, then

$$\exp: \mathcal{O}_X(U) \rightarrow \mathcal{O}_X^*(U)$$

cannot be surjective.

2.2. Derived functor formulation of sheaf cohomology. The category $\underline{Ab}_X$: sheaves of abelian groups on X . In this section we will introduce sheaf cohomology by considering it as a derived functor.

Given an exact sequence of sheaf as follows

$$0 \rightarrow \mathcal{F}' \xrightarrow{\phi} \mathcal{F} \xrightarrow{\psi} \mathcal{F}''.$$

By taking sections over an open subset U , we obtain a sequence of abelian groups

$$0 \rightarrow \mathcal{F}'(U) \xrightarrow{\phi(U)} \mathcal{F}(U) \xrightarrow{\psi(U)} \mathcal{F}''(U).$$

Above sequence is not only exact at $\mathcal{F}'(U)$, but also is exact at $\mathcal{F}(U)$. In other words, the functor given by taking section over open subset is a left exact functor.

- (1) First, we show $\ker \psi(U) \supseteq \operatorname{im} \phi(U)$. For $s \in \mathcal{F}'(U)$, if we want to show $\psi \circ \phi(s) = 0$, it suffices to show $(\psi \circ \phi(s))|_p = 0$ for all $p \in U$ since $\mathcal{F}''$ is a sheaf. For any $p \in U$, by considering stalk at p we obtain an exact sequence of abelian groups

$$0 \rightarrow \mathcal{F}'_p \xrightarrow{\phi_p} \mathcal{F}_p \xrightarrow{\psi_p} \mathcal{F}''_p.$$

Then we obtain $\psi_p \circ \phi_p(s|_p) = 0$, which implies $(\psi \circ \phi(s))|_p = 0$.

- (2) Conversely, Given $s \in \ker \psi(U)$, we have $s|_p \in \ker \psi_p$ for any $p \in U$. By exactness of stalks, there exists $t_p \in \mathcal{F}'_p$ such that $\phi_p(t_p) = s|_p$. Thus there exists an open subset V_i containing p and $t_i \in \mathcal{F}'(V_i)$ such that $\phi(t_i) = s|_{V_i}$. Now it suffices to show these t_i can be glued together to obtain $t \in \mathcal{F}'(U)$, and since $\mathcal{F}$ is a sheaf, it suffices to check these t_i agree on intersections $V_i \cap V_j$. Note that $\phi(t_i|_{V_i \cap V_j} - t_j|_{V_i \cap V_j}) = s|_{V_i \cap V_j} - s|_{V_i \cap V_j} = 0$, then these t_i agree on intersections since ϕ is injective.

Remark 2.2.1. From the above argument, we can see that

$$0 \rightarrow \mathcal{F}' \xrightarrow{\phi} \mathcal{F} \xrightarrow{\psi} \mathcal{F}''$$

is exact if and only if for any open subset $U \subseteq X$

$$0 \rightarrow \mathcal{F}'(U) \xrightarrow{\phi(U)} \mathcal{F}(U) \xrightarrow{\psi(U)} \mathcal{F}''(U)$$

is exact.

In homological algebra, we always consider the **derived functor** of a left or right-exact functor. In particular, the global section functor is left exact, and its right derived functor defines the cohomology of a sheaf. Before giving the definition of derived functors, first, we define the **injective resolution** of a sheaf.

Definition 2.2.1. A sheaf $\mathcal{I}$ is injective if $\operatorname{Hom}(-, \mathcal{I})$ is an exact functor.

Definition 2.2.2. Let $\mathcal{F}$ be a sheaf. An injective resolution of $\mathcal{F}$ is an exact sequence

$$0 \rightarrow \mathcal{F} \rightarrow \mathcal{I}^0 \rightarrow \mathcal{I}^1 \rightarrow \mathcal{I}^2 \rightarrow \dots$$

where $\mathcal{I}^i$ are injective for all i .

Theorem 2.2.1. Every sheaf admits an injective resolution.

Theorem 2.2.2. Let $\mathcal{F} \rightarrow \mathcal{I}^\bullet$ and $\mathcal{G} \rightarrow \mathcal{G}^\bullet$ be two resolutions, and let $\phi: \mathcal{F} \rightarrow \mathcal{G}$ be a morphism of sheaves. Then there exists a morphism $\tilde{\phi}: \mathcal{I}^\bullet \rightarrow \mathcal{G}^\bullet$ lifting ϕ ; such a lift is unique up to homotopy.

Definition 2.2.3. Let $\mathcal{F}$ be a sheaf of abelian groups. Then

$$H^p(X, \mathcal{F}) := H^p(\mathcal{I}^\bullet(X)).$$

Remark 2.2.2. The theorem 2.2.2 shows that the definition of sheaf cohomology is independent of the choice of injective resolution.

Example 2.2.1. By definition, the 0-th cohomology is exactly the global sections

$$H^0(X, \mathcal{F}) := \ker \{\mathcal{I}^0(X) \rightarrow \mathcal{I}^1(X)\}.$$

Thus $H^0(X, \mathcal{F}) = \mathcal{F}(X)$, the global sections of sheaf.

Example 2.2.2. If $\mathcal{F}$ is an injective sheaf, then $H^i(X, \mathcal{F}) = 0$ for all $i > 0$, since the **sheaf cohomology** of an injective sheaf can be computed by using the following special injective resolution

$$0 \rightarrow \mathcal{F} \xrightarrow{\text{id}} \mathcal{F} \rightarrow 0 \rightarrow \dots$$

Theorem 2.2.3 (zig-zag). If

$$0 \rightarrow \mathcal{F} \rightarrow \mathcal{G} \rightarrow \mathcal{H} \rightarrow 0$$

is a short exact sequence of sheaves, then there is an induced long exact sequence of abelian groups

$$0 \rightarrow H^0(X, \mathcal{F}) \rightarrow H^0(X, \mathcal{G}) \rightarrow H^0(X, \mathcal{H}) \rightarrow H^1(X, \mathcal{F}) \rightarrow H^1(X, \mathcal{G}) \rightarrow \dots$$

Definition 2.2.4. Let $f: X \rightarrow Y$ be a continuous map between topological spaces and $\mathcal{F}$ be a sheaf of abelian groups on X . The **direct image** of $\mathcal{F}$, denoted by $f_*\mathcal{F}$, is a sheaf on Y defined by

$$f_*\mathcal{F}(U) := \mathcal{F}(f^{-1}(U)).$$

Proposition 2.2.1. $f_*: \underline{Ab}_X \rightarrow \underline{Ab}_Y$ is a left exact functor.

Proof. Given an exact sequence of sheaves on X

$$0 \rightarrow \mathcal{F}' \rightarrow \mathcal{F} \rightarrow \mathcal{F}''.$$

Then we need to show

$$0 \rightarrow f_*\mathcal{F}' \rightarrow f_*\mathcal{F} \rightarrow f_*\mathcal{F}''$$

is also an exact sequence of sheaves on Y . By Remark 2.2.1 it suffices to show that for any open subset $V \subseteq Y$, we obtain the following exact sequence

$$0 \rightarrow f_*\mathcal{F}'(V) \rightarrow f_*\mathcal{F}(V) \rightarrow f_*\mathcal{F}''(V),$$

and that is exactly

$$0 \rightarrow \mathcal{F}'(f^{-1}(V)) \rightarrow \mathcal{F}(f^{-1}(V)) \rightarrow \mathcal{F}''(f^{-1}(V)).$$

Since f is continuous, $f^{-1}(V)$ is an open subset in X , and thus above sequence of abelian is exact since $0 \rightarrow \mathcal{F}' \rightarrow \mathcal{F} \rightarrow \mathcal{F}''$ is exact. $\square$

2.3. Acyclic resolution. In practice it may be difficult for us to choose an injective resolution, so we usually use other resolutions to compute sheaf cohomology.

Definition 2.3.1. A sheaf $\mathcal{F}$ is acyclic if $H^i(X, \mathcal{F}) = 0$ for all $i > 0$.

Example 2.3.1. Injective sheaf is acyclic.

Definition 2.3.2. Let $\mathcal{F}$ be a sheaf. An acyclic resolution of $\mathcal{F}$ is an exact sequence

$$0 \rightarrow \mathcal{F} \rightarrow \mathcal{A}^0 \rightarrow \mathcal{A}^1 \rightarrow \mathcal{A}^2 \rightarrow \dots$$

where $\mathcal{A}^i$ is acyclic for all i .

Proposition 2.3.1. The cohomology of the sheaf $\mathcal{F}$ can be computed using an acyclic resolution.

In fact, it is a standard homological technique, called dimension shifting, so we will state this technique in the language of homological algebra. We first consider a simple version of it.

Example 2.3.2. Let $\mathcal{F}$ be a left exact functor and $0 \rightarrow A \rightarrow M_1 \rightarrow B \rightarrow 0$ be an exact sequence with M_1 is $\mathcal{F}$ -acyclic. Then $R^{i+1}\mathcal{F}(A) \cong R^i\mathcal{F}(B)$ for $i > 0$, and $R^1\mathcal{F}(A)$ is the **cokernel** of $\mathcal{F}(M_1) \rightarrow \mathcal{F}(B)$.

Proof. By considering the long exact sequence induced by $0 \rightarrow A \rightarrow M^1 \rightarrow B \rightarrow 0$, one has

$$R^i\mathcal{F}(M^1) \rightarrow R^i\mathcal{F}(B) \rightarrow R^{i+1}\mathcal{F}(A) \rightarrow R^{i+1}\mathcal{F}(M^1)$$

(1) If $i > 0$, then $R^i\mathcal{F}(M^1) = R^{i+1}\mathcal{F}(M^1) = 0$ since M^1 is $\mathcal{F}$ -acyclic, and thus $R^{i+1}\mathcal{F}(A) \cong R^i\mathcal{F}(B)$ for $i > 0$.

(2) If $i = 0$, then

$$0 \rightarrow \mathcal{F}(M^1) \rightarrow \mathcal{F}(B) \rightarrow R^1\mathcal{F}(A) \rightarrow 0$$

implies $R^1\mathcal{F}(A) = \text{coker}\{\mathcal{F}(M^1) \rightarrow \mathcal{F}(B)\}$. $\square$

We now prove dimension shifting in a general setting.

Lemma 2.3.1 (dimension shifting). If

$$0 \rightarrow A \rightarrow M^1 \rightarrow M^2 \rightarrow \dots \rightarrow M^m \rightarrow B \rightarrow 0$$

is exact with M^i is $\mathcal{F}$ -acyclic, then $R^{i+m}\mathcal{F}(A) \cong R^i\mathcal{F}(B)$ for $i > 0$, and $R^m\mathcal{F}(A)$ is the **cokernel** of $\mathcal{F}(M^m) \rightarrow \mathcal{F}(B)$.

Proof. Prove it by induction on m . For $m = 1$, we already see it in Example 2.3.2. Assume it holds for $m < k$, then for $m = k$, we split $0 \rightarrow A \rightarrow M^1 \rightarrow M^2 \rightarrow \dots \rightarrow M^k \xrightarrow{d_k} B \rightarrow 0$ into two exact sequences

$$0 \rightarrow A \rightarrow M^1 \rightarrow M^2 \rightarrow \dots \rightarrow M^{k-1} \rightarrow \ker d_k \rightarrow 0$$

$$0 \rightarrow \ker d_k \rightarrow M^k \xrightarrow{d_k} B \rightarrow 0.$$

Then by induction hypothesis, for $i > 0$ we have

$$\begin{aligned} R^{i+k-1}\mathcal{F}(A) &\cong R^i\mathcal{F}(\ker d_k) \\ R^{i+1}\mathcal{F}(\ker d_k) &\cong R^i\mathcal{F}(B). \end{aligned}$$

Combine these two isomorphisms together we obtain $R^{i+k}\mathcal{F}(A) \cong R^i\mathcal{F}(B)$ for $i > 0$, as desired. For $i = 0$, it suffices to let $i = 1$ in $R^{i+k-1}\mathcal{F}(A) \cong R^i\mathcal{F}(\ker d_k)$, then we obtain

$$R^k\mathcal{F}(A) = R^1\mathcal{F}(\ker d_k) = \operatorname{coker}\{\mathcal{F}(M^k) \rightarrow \mathcal{F}(B)\}.$$

This completes the proof. $\square$

Corollary 2.3.1. If $0 \rightarrow A \rightarrow M^\bullet$ is a $\mathcal{F}$ -acyclic resolution, then $R^i\mathcal{F}(A) = H^i(\mathcal{F}(M^\bullet))$.

Proof. Truncate the resolution as

$$\begin{aligned} 0 \rightarrow A \rightarrow M^0 \rightarrow M^1 \rightarrow \dots \rightarrow M^{i-1} \rightarrow B \rightarrow 0 \\ 0 \rightarrow B \rightarrow M^i \rightarrow M^{i+1} \rightarrow \dots \end{aligned}$$

Since we already have $R^i\mathcal{F}(A) = \operatorname{coker}\{\mathcal{F}(M^{i-1}) \rightarrow \mathcal{F}(B)\}$, and $\mathcal{F}$ is left exact, one has

$$\mathcal{F}(B) = \ker\{\mathcal{F}(M^i) \rightarrow \mathcal{F}(M^{i+1})\}.$$

Thus we obtain

$$R^i\mathcal{F}(A) = \operatorname{coker}\{\mathcal{F}(M^{i-1}) \rightarrow \ker\{\mathcal{F}(M^i) \rightarrow \mathcal{F}(M^{i+1})\}\} = H^i(\mathcal{F}(M^\bullet)).$$

$\square$

2.4. Examples about acyclic sheaf.

2.4.1. *Flabby sheaf.* First kind of acyclic sheaf is flabby² sheaf.

Definition 2.4.1. A sheaf $\mathcal{F}$ is flabby if for all open $U \subseteq V$, the restriction map $\mathcal{F}(V) \rightarrow \mathcal{F}(U)$ is surjective.

We now see some examples about flabby sheaves.

Example 2.4.1. A constant sheaf on an irreducible topological space is flabby.

Proof. Note that the constant presheaf on an irreducible topological space is a sheaf in fact, and one checks directly this constant presheaf is flabby. $\square$

In particular, we have

²Some authors also call this flasque.

Example 2.4.2. Let X be an algebraic variety. Then constant sheaf $\mathbb{Z}_X$ is flabby.

Example 2.4.3. If $\mathcal{F}$ is a flabby sheaf on X , and $f: X \rightarrow Y$ is a continuous map, then $f_*\mathcal{F}$ is a flabby sheaf on Y .

Proof. For $V \subseteq W$ in Y , it suffices to show $f_*\mathcal{F}(W) \rightarrow f_*\mathcal{F}(V)$ is surjective, and that is

$$\mathcal{F}(f^{-1}W) \rightarrow \mathcal{F}(f^{-1}V)$$

it is surjective since $\mathcal{F}$ is flabby. $\square$

Example 2.4.4. An injective sheaf is flabby.

Proof. Let $\mathcal{I}$ be an injective sheaf and $V \subseteq U$ be open subsets. Now we define the sheaf $\underline{\mathbb{Z}}_U$ on X by

$$\underline{\mathbb{Z}}_U := \begin{cases} \underline{\mathbb{Z}}(W) & W \subseteq U \\ 0 & \text{otherwise} \end{cases}$$

where $\underline{\mathbb{Z}}$ is constant sheaf valued in $\mathbb{Z}$, and similarly we define sheaf $\underline{\mathbb{Z}}_V$. By construction one has $\underline{\mathbb{Z}}_U(W) = \underline{\mathbb{Z}}_V(W)$ unless $W \subseteq U$ and $W \not\subseteq V$. Thus we obtain an exact sequence

$$0 \rightarrow \underline{\mathbb{Z}}_V \rightarrow \underline{\mathbb{Z}}_U.$$

Applying the functor $\text{Hom}(-, \mathcal{I})$, which is exact, we obtain an exact sequence

$$\text{Hom}(\underline{\mathbb{Z}}_U, \mathcal{I}) \rightarrow \text{Hom}(\underline{\mathbb{Z}}_V, \mathcal{I}) \rightarrow 0.$$

We now explain why we need such a weird sheaf $\underline{\mathbb{Z}}_U$. In fact, we will prove $\text{Hom}(\underline{\mathbb{Z}}_U, \mathcal{I}) = \mathcal{I}(U)$. Indeed since $\varphi: \underline{\mathbb{Z}}_U \rightarrow \mathcal{I}$ is a sheaf morphism. Then if $W \not\subseteq U$, then $\varphi(U)$ must be zero. If $W = U$, then the group of sections of $\underline{\mathbb{Z}}_U(U)$ over any connected component is simply $\mathbb{Z}$ and hence $\varphi(U)$ on this connected component is determined by the image of $1 \in \mathbb{Z}$. Thus $\varphi(U)$ can be thought of as an element of $\mathcal{I}(U)$. Now on any proper open subset of U , φ is determined by restriction maps. Hence $\text{Hom}(\underline{\mathbb{Z}}_U, \mathcal{I}) = \mathcal{I}(U)$, as desired. The same argument shows $\text{Hom}(\underline{\mathbb{Z}}_U, \mathcal{I}) = \mathcal{I}(V)$, and thus we obtain an exact sequence

$$\mathcal{I}(U) \rightarrow \mathcal{I}(V) \rightarrow 0,$$

which shows $\mathcal{I}$ is flabby. $\square$

Our goal is to prove a flabby sheaf is acyclic, but we still need some property of flabby sheaves.

Proposition 2.4.1. If $0 \rightarrow \mathcal{F}' \xrightarrow{\phi} \mathcal{F} \xrightarrow{\psi} \mathcal{F}'' \rightarrow 0$ is an exact sequence of sheaves, and $\mathcal{F}'$ is flabby, then for any open subset U , the sequence

$$0 \rightarrow \mathcal{F}'(U) \xrightarrow{\phi(U)} \mathcal{F}(U) \xrightarrow{\psi(U)} \mathcal{F}''(U) \rightarrow 0$$

is exact.

Proof. It suffices to show $\mathcal{F}(U) \rightarrow \mathcal{F}''(U) \rightarrow 0$ is exact. Here we only gives a sketch of the proof. Since we have exact sequence on stalks for each $p \in U$ as follows

$$0 \rightarrow \mathcal{F}'_p \xrightarrow{\phi_p} \mathcal{F}_p \xrightarrow{\psi_p} \mathcal{F}''_p \rightarrow 0$$

Then for each $s \in \mathcal{F}''(U)$, there exists $t_p \in \mathcal{F}_p$ such that $\psi_p(t_p) = s|_p$, so there exists open subset $V_p \subseteq U$ containing p and $t \in \mathcal{F}(V_p)$ such that $\psi(t) = s|_{V_p}$. If we can glue these t together then we get a section in $\mathcal{F}(U)$ and is mapped to s , which completes the proof. However, they may not equal on the intersection. However, this can be handled as follows, consider another point q and $t' \in \mathcal{F}(V_q)$ such that $\psi(t') = s|_{V_q}$, $(t' - t)|_{V_p \cap V_q} \in \ker \psi(V_p \cap V_q) = \text{im } \phi(V_p \cap V_q)$. So there exists $t'' \in \mathcal{F}'(V_p \cap V_q)$ such that

$$\phi(t'') = (t' - t)|_{V_p \cap V_q}$$

Now since $\mathcal{F}'$ is flabby, then there exists $t''' \in \mathcal{F}(V_p)$ such that $t'''|_{V_p \cap V_q} = t''$. And consider $t + \phi(t''') \in \mathcal{F}(V_p)$, which will coincide with t' on $V_p \cap V_q$. After above corrections, we can glue t after correction together. $\square$

Proposition 2.4.2. If $0 \rightarrow \mathcal{F}' \rightarrow \mathcal{F} \rightarrow \mathcal{F}'' \rightarrow 0$ is an exact sequence of sheaves, and if $\mathcal{F}'$ and $\mathcal{F}$ are flabby, then $\mathcal{F}''$ is flabby.

Proof. Take $V \subseteq U$ and consider the following diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathcal{F}'(U) & \longrightarrow & \mathcal{F}(U) & \longrightarrow & \mathcal{F}''(U) & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathcal{F}'(V) & \longrightarrow & \mathcal{F}(V) & \longrightarrow & \mathcal{F}''(V) & \longrightarrow & 0 \end{array}$$

Then the desired result follows from five lemma. $\square$

Proposition 2.4.3. A flabby sheaf is acyclic.

Proof. Let $\mathcal{F}$ be a flabby sheaf. Since there are enough injective objects in the category of sheaves of abelian groups, there is an exact sequence

$$0 \rightarrow \mathcal{F} \rightarrow \mathcal{I} \rightarrow \mathcal{Q} \rightarrow 0$$

where $\mathcal{I}$ is injective. By Example 2.4.4 we have that $\mathcal{I}$ is flabby, and thus by Proposition 2.4.2 we have $\mathcal{Q}$ is flabby. Consider the long exact sequence induced from the above short exact sequence

$$\mathcal{F}(X) \rightarrow \mathcal{I}(X) \rightarrow \mathcal{Q}(X) \rightarrow H^1(X, \mathcal{F}) \rightarrow H^1(X, \mathcal{I}) \rightarrow \dots$$

Note that $H^1(X, \mathcal{I}) = 0$ since $\mathcal{I}$ is injective, and thus acyclic. Then $H^1(X, \mathcal{F}) = \text{coker}\{\mathcal{I}(X) \rightarrow \mathcal{Q}(X)\}$. But Proposition 2.4.1 shows that $\mathcal{I}(X) \rightarrow \mathcal{Q}(X)$ is surjective since $\mathcal{F}$ is flabby, so $H^1(X, \mathcal{F}) = 0$.

We now prove $H^k(X, \mathcal{F}) = 0$ for $k > 0$ by induction on k , and above argument shows it is true for $k = 1$. Assume this holds for $k < n$, and consider

$$\dots \rightarrow H^{n-1}(X, \mathcal{Q}) \rightarrow H^n(X, \mathcal{F}) \rightarrow H^n(X, \mathcal{I}) \rightarrow H^n(X, \mathcal{Q}) \rightarrow \dots$$

By induction hypothesis, we can reduce above sequence to

$$\dots \rightarrow 0 \rightarrow H^n(X, \mathcal{F}) \rightarrow 0 \rightarrow H^n(X, \mathcal{Q}) \rightarrow \dots$$

which implies $H^n(X, \mathcal{F}) = 0$. This completes the proof. $\square$

2.4.2. Soft sheaf. The second kind of acyclic sheaves is called **soft** sheaves, which is quite similar to flabby.

Definition 2.4.2. A sheaf $\mathcal{F}$ over X is soft if for any closed subset $S \subseteq X$ the restriction map $\mathcal{F}(X) \rightarrow \mathcal{F}(S)$ is surjective.

Remark 2.4.1. For closed subset S , the section over it is defined by

$$\mathcal{F}(S) := \varinjlim_{S \subseteq U} \mathcal{F}(U)$$

Parallel to Proposition 2.4.1 and Proposition 2.4.2, soft sheaf has the following properties:

Proposition 2.4.4. If $0 \rightarrow \mathcal{F}' \xrightarrow{\phi} \mathcal{F} \xrightarrow{\psi} \mathcal{F}'' \rightarrow 0$ is an exact sequence of sheaves, and $\mathcal{F}'$ is soft, then the following sequence

$$0 \rightarrow \mathcal{F}'(X) \xrightarrow{\phi(X)} \mathcal{F}(X) \xrightarrow{\psi(X)} \mathcal{F}''(X) \rightarrow 0$$

is exact.

Proposition 2.4.5. If $0 \rightarrow \mathcal{F}' \rightarrow \mathcal{F} \rightarrow \mathcal{F}'' \rightarrow 0$ is an exact sequence of sheaves, and if $\mathcal{F}'$ and $\mathcal{F}$ are soft, then $\mathcal{F}''$ is soft.

Proposition 2.4.6. A soft sheaf is acyclic.

So you may wonder, what is the difference between flabby and soft since the definitions are quite similar, and both of them are acyclic. Directly by definition of sections over a closed subset, we know that every flabby sheaf is soft, but converse fails

Example 2.4.5. The **sheaf** of smooth functions on a smooth manifold is soft but not flabby.

Lemma 2.4.1. If $\mathcal{M}$ is a sheaf of modules over a soft sheaf of rings $\mathcal{R}$, then $\mathcal{M}$ is a soft sheaf.

Proof. Let $s \in \mathcal{M}(K)$ for some closed subset $K \subseteq X$. Then s extends to some open neighborhood U of K . Let $\rho \in \mathcal{R}(K \cup (X \setminus U))$ be defined by

$$\rho = \begin{cases} 1, & \text{on } K \\ 0, & \text{on } X \setminus U \end{cases}$$

Since $\mathcal{R}$ is soft, ρ extends to a section over X , then $\rho \circ s$ is the desired extension of s . $\square$

2.4.3. Fine sheaf. Another important kind of acyclic sheaves, which behaves like sheaf of differential forms Ω_X^k is called **fine sheaf**. Recall what is a partition of unity: Let $U = \{U_i\}_{i \in I}$ be a locally finite open covering of topological space X . A partition of unity subordinate to U is a collection of continuous functions $f_i : U_i \rightarrow [0, 1]$ for each $i \in I$ such that its support lies in U_i , and for any $x \in X$

$$\sum_{i \in I} f_i(x) = 1.$$

Definition 2.4.3. A fine sheaf $\mathcal{F}$ on X is a sheaf of $\mathcal{A}$ -modules, where $\mathcal{A}$ is a sheaf of rings such that for every locally finite open covering $\{U_i\}_{i \in I}$ of X , there is a partition of unity

$$\sum_{i \in I} \rho_i = 1$$

where $\rho_i \in \mathcal{A}(X)$ and $\text{supp}(\rho_i) \subseteq U_i$.

Remark 2.4.2. For a sheaf $\mathcal{F}$ on X and a section $s \in \mathcal{F}(X)$, its support is defined as

$$\text{supp}(s) := \overline{\{x \in X : s|_x \neq 0\}}.$$

Proposition 2.4.7. A fine sheaf is acyclic.

Proof. Let $\mathcal{F}$ be a sheaf of $\mathcal{A}$ -modules and a fine sheaf. Choose an injective resolution

$$0 \rightarrow \mathcal{F} \xrightarrow{d} \mathcal{I}^0 \xrightarrow{d} \mathcal{I}^1 \xrightarrow{d} \mathcal{I}^2 \xrightarrow{d} \dots$$

such that $\mathcal{I}^i$ are injective sheaves of $\mathcal{A}$ -modules. Let $s \in \mathcal{I}^p(X)$ such that $ds = 0$. Then by exactness of injective resolution we have X is covered by open subsets U_i such that for each i there is an element $t_i \in \mathcal{I}^{p-1}(U_i)$ such that $dt_i = s|_{U_i}$. By passing to a refinement we may assume that the cover $\{U_i\}$ is locally finite. Let $\{\rho_i\}$ be a partition of unity subordinate to $\{U_i\}$. Then we obtain $t = \sum \rho_i t_i \in \mathcal{I}^{p-1}(X)$ such that $dt = s$. This completes the proof. $\square$

Example 2.4.6. Let M be a smooth manifold and $\pi : E \rightarrow M$ be a vector bundle. Then the **sheaf** of smooth sections of E is a $C^\infty(M)$ -module sheaf, which is a fine sheaf. In particular, the **sheaf** of tangent bundle, sheaf of differential forms Ω_M and k -forms Ω_M^k are fine sheaves.

Remark 2.4.3. As a consequence, it is meaningless to compute cohomology of sheaf of differential k -forms, or any other vector bundle over a smooth manifold. But in complex version, something interesting happens: Let $(X, \mathcal{O}_X)$ be a complex manifold and $\pi : E \rightarrow X$ be a holomorphic vector bundle. Then the **sheaf** of holomorphic sections of E is not a fine sheaf since there is no partition of unity may not be holomorphic, so the cohomology of holomorphic vector bundle is not trivial, and that is what Dolbeault cohomology computes.

For fine sheaf and soft sheaf, we have

Lemma 2.4.2. Fine sheaf is soft.

Proof. Let $\mathcal{F}$ be a fine sheaf, $S \subseteq X$ closed and $s \in \mathcal{F}(S)$. Let $\{U_i\}$ be an open covering of S and $s_i \in \mathcal{F}(U_i)$ such that

$$s_i|_{S \cap U_i} = s|_{S \cap U_i}.$$


Let $U_0 = X - S$, and $s_0 = 0$. Then $\{U_i\} \sqcup \{U_0\}$ is an open covering of X . without loss of generality, we assume this open covering is locally finite and choose a partition of unity $\{\rho_i\}$ subordinate to it. Then

$$\bar{s} := \sum_i \rho_i(s_i)$$

is a section in $\mathcal{F}(X)$ which extends s . □

Remark 2.4.4. Until now, we have shown that soft, fine and flabby sheaves are acyclic. Lemma 2.4.2 shows fine sheaf is soft, and by definition a flabby sheaf is soft. The Example 2.4.5 shows that soft sheaf may not be flabby, and constant sheaf on an irreducible space is flabby but not fine. In a summary, we obtain the following relations:

Acyclic



2.5. Proof of de Rham theorem using sheaf cohomology. As we already know, for constant sheaf $\underline{\mathbb{R}}$ over a smooth manifold M , we obtain the following fine resolution

$$0 \rightarrow \underline{\mathbb{R}} \xrightarrow{i} \Omega_M^0 \xrightarrow{d} \Omega_M^1 \xrightarrow{d} \Omega_M^2 \xrightarrow{d} \dots$$

And de Rham cohomology computes the **sheaf cohomology** of $\underline{\mathbb{R}}$. de Rham theorem implies that de Rham cohomology equals to the singular cohomology with real coefficients. So if we can give constant sheaf another resolution using singular cochains, we may derive the de Rham cohomology.

We state this in a general setting: Let X be a topological manifold, and a constant sheaf $\underline{G}$ over X , where G is an abelian group. Let $S^p(U, G)$ be the group of singular cochains in U with coefficients in G , and let δ denote the coboundary operator.

Let $S^p(G)$ be the sheaf over X generated by the presheaf $U \mapsto S^p(U, G)$, with induced differential mapping $S^p(G) \xrightarrow{\delta} S^{p+1}(G)$.

Similar to Poincaré lemma, we have for a unit ball U in Euclidean space, we obtain the following sequence

$$\dots \rightarrow S^{p-1}(U, G) \xrightarrow{\delta} S^p(U, G) \xrightarrow{\delta} S^{p+1}(U, G) \rightarrow \dots$$

is exact. So we obtain the following resolution of the constant sheaf $\underline{G}$

$$0 \rightarrow \underline{G} \rightarrow S^0(G) \xrightarrow{\delta} S^1(G) \xrightarrow{\delta} S^2(G) \rightarrow \dots$$

Remark 2.5.1. If M is a smooth manifold, then we can consider smooth chains, that is $f: \Delta^p \rightarrow U$, where f is a smooth function. The corresponding results above still hold, and we have a resolution by smooth cochains with coefficients in G :

$$0 \rightarrow \underline{G} \rightarrow S_\infty^\bullet(G)$$

So if we choose $G = \mathbb{R}$, then it suffices to show $0 \rightarrow \underline{\mathbb{R}} \rightarrow S_\infty^\bullet(\mathbb{R})$ is an acyclic resolution, then we obtain de Rham theorem.

First, note that S_∞^p is a S_∞^0 -module, given by cup product on open subsets. Then by Lemma 2.4.1 and the fact S_∞^0 is soft we know that it is a soft resolution. This completes the proof.

2.6. Hypercohomology. In homological algebra, the **hypercohomology** is a generalization of cohomology functor which takes as input not objects in abelian category but instead chain complexes of objects.

One of the motivations for hypercohomology is to generalize the zig-zag lemma, that is, the **short exact sequence** of sheaves induces a long exact sequence of cohomology groups. It turns out hypercohomology gives techniques for constructing a similar cohomological associated long exact sequence from an arbitrary long exact sequence

$$0 \rightarrow \mathcal{F}_1 \rightarrow \mathcal{F}_2 \rightarrow \dots \rightarrow \mathcal{F}_k \rightarrow 0$$

We now give the definition of hypercohomology: Let $\mathcal{F}^\bullet: \dots \rightarrow \mathcal{F}^{i-1} \rightarrow \mathcal{F}^i \rightarrow \mathcal{F}^{i+1} \rightarrow \dots$ be a complex of sheaves of abelian groups, which is bounded from below, that is, $\mathcal{F}^n = 0$ for $n \ll 0$. Then $\mathcal{F}^\bullet$ admits an injective resolution $\mathcal{F}^\bullet \rightarrow \mathcal{I}^\bullet$. In other words,

$$\begin{array}{ccccccc} \dots & \longrightarrow & \mathcal{F}^{i-1} & \longrightarrow & \mathcal{F}^i & \longrightarrow & \mathcal{F}^{i+1} \longrightarrow \dots \\ & & \downarrow & & \downarrow & & \downarrow \\ \dots & \longrightarrow & \mathcal{I}^{i-1} & \longrightarrow & \mathcal{I}^i & \longrightarrow & \mathcal{I}^{i+1} \longrightarrow \dots \end{array}$$

such that

- (1) All $\mathcal{I}^i$ are injective sheaves.
- (2) The induced homomorphism $H^i(\mathcal{F}^\bullet) \rightarrow H^i(\mathcal{I}^\bullet)$ is an isomorphism.

The **hypercohomology** of $\mathcal{F}^\bullet$ is defined by

$$H^i(X, \mathcal{F}^\bullet) := H^i(\Gamma(X, \mathcal{I}^\bullet))$$

Definition 2.6.1. For a sheaf $\mathcal{F}$, $\mathcal{F}^\bullet[n]$ is a sheaf of complex defined by

$$(\mathcal{F}^\bullet[n])^i = \begin{cases} \mathcal{F} & i = n \\ 0 & \text{otherwise.} \end{cases}$$

Example 2.6.1. Let $\mathcal{F}$ be a sheaf and $0 \rightarrow \mathcal{F} \rightarrow \mathcal{I}^0 \rightarrow \mathcal{I}^1 \rightarrow \dots$ be an injective resolution of $\mathcal{F}$. Then

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{F} & \longrightarrow & 0 & \longrightarrow & 0 \longrightarrow \dots \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathcal{I}^0 & \longrightarrow & \mathcal{I}^1 & \longrightarrow & \mathcal{I}^2 \longrightarrow \dots \end{array}$$

is an injective resolution of $\mathcal{F}^\bullet[0]$. Indeed, $\mathcal{I}^i$ are injective for all $i \geq 0$, and

$$H^i(\mathcal{I}^\bullet) = \begin{cases} \mathcal{F}, & n = 0 \\ 0, & \text{otherwise} \end{cases} = H^i(\mathcal{F}^\bullet[0])$$

So by definition of hypercohomology, we have $H^i(X, \mathcal{F}^\bullet[0]) = H^i(\Gamma(X, \mathcal{I}^\bullet)) = H^i(X, \mathcal{F}^\bullet)$. In general, one has

$$H^i(X, \mathcal{F}^\bullet[n]) \cong H^{i+n}(X, \mathcal{F}).$$

Theorem 2.6.1 (zig-zag). Let $0 \rightarrow \mathcal{F}^\bullet \rightarrow \mathcal{G}^\bullet \rightarrow \mathcal{H}^\bullet \rightarrow 0$ be a short exact sequence of complexes of sheaves which are bounded from below. Then there is an induced long exact sequence

$$\dots \rightarrow H^{i-1}(X, \mathcal{H}^\bullet) \rightarrow H^i(X, \mathcal{F}^\bullet) \rightarrow H^i(X, \mathcal{G}^\bullet) \rightarrow H^i(X, \mathcal{H}^\bullet) \rightarrow H^{i+1}(X, \mathcal{F}^\bullet) \rightarrow \dots$$

Part 2. Schemes

3. SCHEMES AND MORPHISMS

3.1. Schemes. Throughout this lecture, all rings are assumed to be commutative with identity element, and all homomorphisms of rings are assumed to map 1 to 1.

3.1.1. Ringed space.

Definition 3.1.1. A ringed space is a pair $(X, \mathcal{O}_X)$ consisting of a topological space X and a sheaf of rings $\mathcal{O}_X$ on X .

Definition 3.1.2. A ringed space $(X, \mathcal{O}_X)$ is a locally ringed space if for every $p \in X$, the stalk $\mathcal{O}_{X,p}$ is a local ring.

Definition 3.1.3. Let $(X, \mathcal{O}_X)$ and $(Y, \mathcal{O}_Y)$ be two ringed spaces. A morphism from $(X, \mathcal{O}_X)$ to $(Y, \mathcal{O}_Y)$ is a pair $(f, f^\sharp)$ consisting of a continuous map $f: X \rightarrow Y$ and a morphism of sheaves $f^\sharp: \mathcal{O}_Y \rightarrow f_*\mathcal{O}_X$.

Remark 3.1.1. Let $(f, f^\sharp)$ be a morphism between ringed spaces $(X, \mathcal{O}_X)$ and $(Y, \mathcal{O}_Y)$. For every point $p \in X$, there is a homomorphism $(f_*\mathcal{O}_X)_{f(p)} \rightarrow \mathcal{O}_{X,p}$ defined by

$$(f_*\mathcal{O}_X)_{f(p)} = \varinjlim_{f(p) \in V} (f_*\mathcal{O}_X)(V) = \varinjlim_{p \in f^{-1}(V)} \mathcal{O}_X(f^{-1}(V)) \rightarrow \varinjlim_{p \in U} \mathcal{O}_X(U) = \mathcal{O}_{X,p}.$$

On the other hand, the morphism of sheaves $f^\sharp: \mathcal{O}_Y \rightarrow f_*\mathcal{O}_X$ induces a homomorphism on stalks

$$(f^\sharp)_p: \mathcal{O}_{Y,f(p)} \rightarrow (f_*\mathcal{O}_X)_{f(p)}.$$

By composing above two homomorphisms, there is a homomorphism

$$f_p^\sharp: \mathcal{O}_{Y,f(p)} \rightarrow \mathcal{O}_{X,p}.$$

Definition 3.1.4. A morphism $(f, f^\sharp)$ between locally ringed spaces $(X, \mathcal{O}_X)$ and $(Y, \mathcal{O}_Y)$ is a morphism between ringed spaces, and for each $p \in X$, the morphism

$$f_p^\sharp: \mathcal{O}_{Y,f(p)} \rightarrow \mathcal{O}_{X,p}$$

is a local homomorphism of local rings.

Definition 3.1.5. An isomorphism of locally ringed spaces is a morphism with a two-sided inverse.

3.1.2. Affine schemes.

Definition 3.1.6. Let A be a ring. The spectrum of A is a locally ringed space, consisting of the following data:

- (1) A topological space $\text{Spec} A$, which is the set of all prime ideals of A , equipped with the Zariski topology, that is, all closed subsets of $\text{Spec} A$ are of the form $V(\mathfrak{a}) = \{\mathfrak{p} \in \text{Spec} A \mid \mathfrak{a} \subseteq \mathfrak{p}\}$.

- (2) A structure sheaf $\mathcal{O}_{\text{Spec}A}$, which is defined as follows: For every open subset U of $\text{Spec}A$, $\mathcal{O}_{\text{Spec}A}(U)$ consists of mappings $s: U \rightarrow \coprod_{\mathfrak{p} \in \text{Spec}A} A_{\mathfrak{p}}$ satisfying the following two conditions:
- (a) For every $\mathfrak{p} \in U$, one has $s(\mathfrak{p}) \in A_{\mathfrak{p}}$.
 - (b) For every $\mathfrak{p} \in U$, there exists a neighborhood $U_{\mathfrak{p}}$ of $\mathfrak{p}$ contained in U and $a, f \in A$ such that for every $\mathfrak{q} \in U_{\mathfrak{p}}$, one has $f \notin \mathfrak{q}$ and $s(\mathfrak{q}) = a/f$ in $A_{\mathfrak{q}}$.

Definition 3.1.7. A locally ringed space that is isomorphic to $(\text{Spec}A, \mathcal{O}_{\text{Spec}A})$ for some ring A is called an **affine scheme**.

Definition 3.1.8. Let A be a ring and $f \in A$. The distinguished open subset $D(f)$ is defined by $\text{Spec}A \setminus V(f)$.

Proposition 3.1.1. The distinguished open subsets form a topology basis of the Zariski topology.

Corollary 3.1.1. The **affine scheme** is quasi-compact.

Proposition 3.1.2. Let A be a ring.

- (1) For every $\mathfrak{p} \in \text{Spec}A$, there is a canonical isomorphism $\mathcal{O}_{\mathfrak{p}} \cong A_{\mathfrak{p}}$.
- (2) For every $f \in A$, there is a canonical isomorphism $\mathcal{O}(D(f)) \cong A_f$. In particular, $\mathcal{O}(\text{Spec}A) \cong A$.

Proof. See Proposition 1.2.4 in [Fu06]. □

Proposition 3.1.3.

- (1) Let $\phi: A \rightarrow B$ be a homomorphism of rings. Then ϕ induces a canonical morphism of locally ringed spaces

$$(f, f^{\#}): (\text{Spec}B, \mathcal{O}_{\text{Spec}B}) \rightarrow (\text{Spec}A, \mathcal{O}_{\text{Spec}A}).$$

- (2) Any morphism $(f, f^{\#})$ between $(\text{Spec}B, \mathcal{O}_{\text{Spec}B})$ and $(\text{Spec}A, \mathcal{O}_{\text{Spec}A})$ is obtained this way.

Proof. See Proposition 1.2.5 in [Fu06]. □

Proposition 3.1.4. For any $f \in A$, there is a canonical isomorphism of locally ringed spaces

$$(D(f), \mathcal{O}_{\text{Spec}A}|_{D(f)}) \cong (\text{Spec}A_f, \mathcal{O}_{\text{Spec}A_f}).$$

Proof. See Proposition 1.2.6 in [Fu06]. □

3.1.3. Schemes.

Definition 3.1.9. A scheme $(X, \mathcal{O}_X)$ is a locally ringed space for which there exists an open covering $\{U_i\}_{i \in I}$ of X such that each $(U_i, \mathcal{O}_X|_{U_i})$ is an affine scheme.

Notation 3.1.1. For convenience, if there is no ambiguity of the underlying topological space, we will simply write $\mathcal{O}$ instead of $\mathcal{O}_X$.

Proposition 3.1.5. Let $(X, \mathcal{O}_X)$ be a scheme and U be an open subset of X . Then $(U, \mathcal{O}_X|_U)$ is a scheme.

Proof. It follows from Proposition 3.1.4. $\square$

Proposition 3.1.6. Let X be a scheme and A be a ring. Then there is a one-to-one correspondence between the set of morphisms of schemes from X to $\text{Spec} A$ and the set of homomorphisms of rings from A to $\mathcal{O}_X(X)$.

Proof. See Proposition 1.2.8 in [Fu06]. $\square$

3.2. Proj construction. In this section we fix a graded ring S with decomposition $S = \bigoplus_{d=0}^{\infty} S_d$.

Proposition 3.2.1. An ideal of S is called a **homogeneous ideal** if it satisfies one of the following equivalent conditions:

- (1) $\mathfrak{a} = \bigoplus_d (\mathfrak{a} \cap S_d)$.
- (2) If $a \in \mathfrak{a}$ and $a = \sum_d a_d$ with $a_d \in S_d$, then $a_d \in \mathfrak{a}$.
- (3) $\mathfrak{a}$ is generated by homogeneous elements as an additive subgroup of S .
- (4) $\mathfrak{a}$ is generated by homogeneous elements an ideal of S .

Proposition 3.2.2. Let $\mathfrak{a}$ be a proper homogeneous ideal of S . If for any homogeneous elements f and g in S such that $fg \in \mathfrak{a}$, one has either $f \in \mathfrak{a}$ or $g \in \mathfrak{a}$, then $\mathfrak{a}$ is a prime ideal.

Proposition 3.2.3. Let $\mathfrak{a}, \mathfrak{b}$ be homogeneous ideals of S . Then

- (1) $\mathfrak{a} + \mathfrak{b}, \mathfrak{a}\mathfrak{b}, \mathfrak{a} \cap \mathfrak{b}$ are homogeneous ideals.
- (2) $\sqrt{\mathfrak{a}}$ is a homogeneous ideal.

Proposition 3.2.4 (Proj). Let $S_+ = \bigoplus_{d=1}^{\infty} S_d$ and $\text{Proj } S$ be the set of all homogeneous prime ideals of S not containing S_+ . For any homogeneous ideal $\mathfrak{a}$ of S , define

$$V_+(\mathfrak{a}) = \{\mathfrak{p} \in \text{Proj } S \mid \mathfrak{a} \subseteq \mathfrak{p}\}.$$

Then

- (1) $V_+(0) = \text{Proj } S$ and $V_+(S) = \emptyset$.
- (2) $\bigcap_{i \in I} V_+(\mathfrak{a}_i) = V_+(\sum_{i \in I} \mathfrak{a}_i)$ for any family of homogeneous ideals $\{\mathfrak{a}_i\}_{i \in I}$ of S .
- (3) $V_+(\mathfrak{a}) \cap V_+(\mathfrak{b}) = V_+(\mathfrak{a}\mathfrak{b}) = V_+(\mathfrak{a} \cap \mathfrak{b})$ for any homogeneous ideals $\mathfrak{a}$ and $\mathfrak{b}$ of S .

In particular, $\text{Proj } S$ is a topological space so that closed subsets are of the form $V_+(\mathfrak{a})$ for homogeneous ideals $\mathfrak{a}$ of S . This topology is called the Zariski topology of $\text{Proj } S$.

We now define the structure sheaf on $\text{Proj } S$. For any homogeneous prime ideal $\mathfrak{p} \in \text{Proj } S$, consider the ring

$$S_{(\mathfrak{p})} = \left\{ \frac{a}{t} \in S_{\mathfrak{p}} \mid a \text{ and } t \text{ are homogeneous of the same degree} \right\}.$$

For open subset $U \subseteq \text{Proj } S$, $\mathcal{O}_{\text{Proj } S}(U)$ is defined to be the set of functions $s: U \rightarrow \prod_{\mathfrak{p} \in U} S_{(\mathfrak{p})}$ such that

- (1) For every $\mathfrak{p} \in U$, one has $s(\mathfrak{p}) \in S_{(\mathfrak{p})}$.

- (2) For every $\mathfrak{p} \in U$, there exists a neighborhood $U_{\mathfrak{p}}$ of $\mathfrak{p}$ contained in U and homogeneous elements $a, f \in S$ of the same degree such that for every $\mathfrak{q} \in U_{\mathfrak{p}}$, one has $f \notin \mathfrak{q}$ and $s(\mathfrak{q}) = a/f$ in $S_{(\mathfrak{q})}$.

Proposition 3.2.5.

- (1) For every $\mathfrak{p} \in \text{Proj } S$, there is a canonical isomorphism $\mathcal{O}_{\text{Proj } S, \mathfrak{p}} \cong S_{(\mathfrak{p})}$.
(2) For every homogeneous element $f \in S_+$, let

$$D_+(f) = \text{Proj } S \setminus V_+(f) = \{\mathfrak{p} \in \text{Proj } S \mid f \notin \mathfrak{p}\}.$$

Then $D_+(f)$ is open in $\text{Proj } S$, and open subsets of this type form a basis for the topology of $\text{Proj } S$. Moreover, there is an isomorphism of locally ringed spaces

$$(D_+(f), \mathcal{O}_{\text{Proj } S}|_{D_+(f)}) \cong (\text{Spec } S_{(f)}, \mathcal{O}_{\text{Spec } S_{(f)}}).$$

In particular, $(\text{Proj } S, \mathcal{O}_{\text{Proj } S})$ is a scheme.

Proof. See Proposition 1.2.10 in [Fu06]. □

4. PROPERTIES OF SCHEMES

4.1. Quasi-compact, irreducible, Noetherian topological space.

4.1.1. Quasi-compact scheme.

Definition 4.1.1. A scheme $(X, \mathcal{O}_X)$ is called **quasi-compact** if any open covering of X admits a finite subcovering.

Remark 4.1.1. A scheme is quasi-compact if and only if it is a finite union of affine schemes.

4.1.2. Irreducible topological space.

Definition 4.1.2. A topological space X is called irreducible if X is not the union of two proper closed subsets. A subset $Y \subseteq X$ is called irreducible if Y is irreducible with the induced topology.

Proposition 4.1.1. Let X be a topological space and $Y \subseteq X$ be a subset equipped with induced topological. If Y is irreducible, then $\overline{Y}$ is also irreducible.

Proof. Suppose $\overline{Y}$ is not irreducible, so we have closed subsets S, T in X such that $\overline{Y}$ is not contained in either S or T but $\overline{Y} \subset T \cup S$. If $Y \subseteq S$, then $\overline{Y} \subseteq \overline{S} = S$, a contradiction, so Y is not contained in S . Similarly Y is not contained in T . But $Y \subseteq \overline{Y} \subseteq S \cup T$, thus Y is not irreducible, a contradiction. $\square$

Proposition 4.1.2. Let A be a ring. A closed subset of $\text{Spec} A$ is irreducible if and only if it is of the form $V(\mathfrak{p})$ for some prime ideal $\mathfrak{p}$ of A .

Proof. See Proposition 1.3.2 in [Fu06]. $\square$

Proposition 4.1.3. Let X be a scheme. For any irreducible closed subset Y of X , there exists a unique point $y \in Y$ such that $Y = \overline{\{y\}}$

Proof. See Proposition 1.3.3 in [Fu06]. $\square$

4.1.3. Noetherian topological space.

Definition 4.1.3. A topological space X is called a **Noetherian topological space** if the family of closed subsets of X satisfies the descending chain conditions.

Example 4.1.1. If A is a Noetherian ring, then $\text{Spec} A$ is a Noetherian topological space.

Proposition 4.1.4. Suppose X is a Noetherian topological space.

- (1) For every closed subset Y of X , there is a decomposition $Y = Y_1 \cup \cdots \cup Y_n$ into closed irreducible subsets Y_i such that $Y_i \not\subseteq Y_j$ whenever $i \neq j$, where Y_i are called irreducible component of Y .
- (2) An irreducible closed subset Y is an irreducible component of X if and only if Y is maximal among the family of irreducible closed subset of X .

Proof. See Proposition 1.3.4 in [Fu06]. $\square$

Definition 4.1.4. Let A be a ring. A prime ideal of A is called a **minimal prime ideal** if it contains no prime ideal other than itself.

Corollary 4.1.1. Let A be a Noetherian ring. Then there is a one to one correspondence between the family of irreducible components of $\text{Spec} A$ and the family of minimal prime ideals of A .

4.2. Reduced, irreducible and integral scheme.

Definition 4.2.1. Let $(X, \mathcal{O}_X)$ be a scheme. Then it is

- (1) connected if X is connected.
- (2) irreducible if X is irreducible.
- (3) reduced if for every open subset U of X , $\mathcal{O}_X(U)$ is reduced.
- (4) integral if for every open subset U of X , $\mathcal{O}_X(U)$ is an integral domain.
- (5) locally integral if $\mathcal{O}_{X,p}$ is an integral domain for every $p \in X$.

Proposition 4.2.1. A scheme $(X, \mathcal{O}_X)$ is integral if and only if it is irreducible and reduced.

Proof. See Proposition 1.3.6 in [Fu06]. □

Proposition 4.2.2. Let $(X, \mathcal{O}_X)$ be an integral scheme and ξ be its generic point. Then $\mathcal{O}_{X,\xi}$ is a field.

Proposition 4.2.3. A scheme $(X, \mathcal{O}_X)$ is reduced if and only if $\mathcal{O}_{X,p}$ is reduced for every $p \in X$.

Proof. Suppose $(X, \mathcal{O}_X)$ is reduced and $s \in \mathcal{O}_{X,p}$ such that $s^n = 0$. Then there exists an open neighborhood U of p such that $s^n = 0$ in $\mathcal{O}_X(U)$, and thus $s = 0$ since $\mathcal{O}_X(U)$ is reduced. Conversely, suppose $\mathcal{O}_{X,p}$ is reduced for every $p \in X$ and $s \in \mathcal{O}_X(U)$ such that $s^n = 0$. Then by passing to the stalks one has $s_p = 0$ for every $p \in U$, and thus $s = 0$. □

Proposition 4.2.4. Let $(X, \mathcal{O}_X)$ be a scheme such that X is a Noetherian topological space. Then $(X, \mathcal{O}_X)$ is locally integral if and only if it is reduced and its irreducible component are disjoint.

Proof. See Proposition 1.3.8 in [Fu06]. □

4.3. Affine criterion.

Definition 4.3.1. Let $(X, \mathcal{O}_X)$ be a scheme. For any section $f \in \mathcal{O}_X(X)$, X_f is defined to be the subset of X consisting of those $p \in X$ such that the germ of f at p is a unit in $\mathcal{O}_{X,p}$.

Proposition 4.3.1. Let $(X, \mathcal{O}_X)$ be a scheme.

- (1) For every $f \in \mathcal{O}_X(X)$, X_f is open. It is empty if and only if there exists an open covering $\{U_i\}_{i \in I}$ of X such that each $f|_{U_i}$ is nilpotent.
- (2) For any $f, g \in \mathcal{O}_X(X)$, we have $X_f \cap X_g = X_{fg}$.
- (3) Let $(\varphi, \varphi^\#): (X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$ be a morphism of schemes and $f \in \mathcal{O}_Y(Y)$. Then $\varphi^{-1}(Y_f) = X_{\varphi^\#(f)}$.

- (4) Suppose X can be covered by finitely many affine open subschemes $\{U_i\}_{i \in I}$ such that $U_i \cap U_j$ can be covered by finitely many affine open subschemes for all $i, j \in I$. Let $A = \mathcal{O}_X(X)$. Then for any $f \in A$, we have $\mathcal{O}_X(X_f) = A_f$.

Proof. Here we only give the proof of (1), and for others one can refer to the Proposition 1.3.9 in [Fu06]. Suppose $\{U_i = \text{Spec } A_i\}_{i \in I}$ is an affine covering of X and define $f_i = f|_{U_i}$. Then it follows that $X_f = \bigcup_{i \in I} D(f_i)$, and thus it is open.

We now prove the half part:

- (a) Suppose there exists an open covering $\{U_i\}_{i \in I}$ such that each $f_i := f|_{U_i}$ is nilpotent. Then without loss of generality we may assume this open covering consists of affine open subsets (by taking refinement when necessary). By previous result one has $X_f = \bigcup_{i \in I} D(f_i) = \emptyset$ since $D(f_i) = \emptyset$ if f_i is nilpotent.
- (b) Conversely, if X_f is empty, then there exists an affine open covering $\{U_i = \text{Spec } A_i\}_{i \in I}$ such that $D(f_i) = \emptyset$ for each $i \in I$, and thus f_i is nilpotent. $\square$

Proposition 4.3.2. A scheme $(X, \mathcal{O}_X)$ is affine if and only if there exist finitely many sections $f_1, \dots, f_n \in \mathcal{O}_X(X)$ generating the unit ideal of $\mathcal{O}_X(X)$ such that each open subscheme $(X_{f_i}, \mathcal{O}_X|_{X_{f_i}})$ is affine.

Proof. See Proposition 1.3.10 in [Fu06]. $\square$

4.4. Noetherian scheme.

Definition 4.4.1. A scheme $(X, \mathcal{O}_X)$ is called locally Noetherian if it can be covered by affine open subschemes $\{U_i = \text{Spec } A_i\}_{i \in I}$ such that each A_i is Noetherian, and it is called Noetherian if it is quasi-compact and locally Noetherian.

Proposition 4.4.1. Let $(X, \mathcal{O}_X)$ be a Noetherian scheme. Then every open subset of X is quasi-compact.

Proof. If X is Noetherian, then it admits a finite affine open covering $\{U_i = \text{Spec } A_i\}_{i \in I}$ such that each A_i is Noetherian. In particular, X is a Noetherian topological space, and thus every open subset of X is quasi-compact. $\square$

Proposition 4.4.2. Let $(X, \mathcal{O}_X)$ be a locally Noetherian scheme. Then for any affine open subscheme $U = \text{Spec } A$ of X , A is Noetherian. In particular, an affine scheme $(\text{Spec } A, \mathcal{O}_{\text{Spec } A})$ is locally Noetherian (and thus Noetherian) if and only if A is Noetherian.

Proof. See Proposition 1.3.11 in [Fu06]. $\square$

Proposition 4.4.3. If $(X, \mathcal{O}_X)$ is a Noetherian scheme, then X is a Noetherian topological space, but the converse is not true.

Proof. Recall that if a ring A is Noetherian, then $\text{Spec } A$ is a Noetherian topological space. Now since $(X, \mathcal{O}_X)$ is a Noetherian scheme, then there exists a

finite affine open covering $\{U_i = \operatorname{Spec} A_i\}_{i=1}^n$ such that each A_i is a Noetherian ring, and it follows that a finite union of Noetherian topological space is also Noetherian.

Conversely, consider the ring $A = k[x_1, x_2, \dots]/(x_1^2, x_2^2, \dots)$, where k is a field. Since each variable x_n is nilpotent, every prime must contain $I = (x_1, x_2, \dots)$. But A/I is just k , so I is already a maximal ideal, and thus I is the only prime ideal. This shows $\operatorname{Spec} A$ has only one point, and it is obviously Noetherian. But I is not finitely generated, so A is not Noetherian. In particular, $\operatorname{Spec} A$ is not a Noetherian scheme. $\square$

5. PROPERTIES OF MORPHISMS

5.1. Quasi-compact, affine, finite type and finite.

Definition 5.1.1. Let $\pi: X \rightarrow Y$ be a morphism of schemes. It is called

- (1) **quasi-compact** if there exists a covering of Y by affine open subschemes $\{V_i\}_{i \in I}$ such that each $\pi^{-1}(V_i)$ is quasi-compact.
- (2) **affine** if there exists a covering of Y by affine open subschemes $\{V_i\}_{i \in I}$ such that each $\pi^{-1}(V_i)$ is affine.
- (3) **locally of finite type** if there exists a covering of Y by affine open subschemes $\{V_i = \text{Spec} B_i\}_{i \in I}$ such that each $\pi^{-1}(V_i)$ can be covered by affine open subschemes $\{U_{ij} = \text{Spec} A_{ij}\}_{j \in J_i}$ for some finitely generated B_i -algebra A_{ij} .
- (4) **finite type** if it is quasi-compact and locally of finite type.
- (5) **finite** if there exists a covering of Y by affine open subschemes $\{V_i = \text{Spec} B_i\}_{i \in I}$ such that each $\pi^{-1}(V_i) = \text{Spec} A_i$ for some finitely generated B_i -module A_i .

Definition 5.1.2. Let $(X, \mathcal{O}_X)$ be a scheme and $\{\text{Spec} A_i\}_{i \in I}$ be an open covering of X . A property $\mathcal{P}$ is called **affine-local**, if every $\text{Spec} A_i$ has property $\mathcal{P}$ implies any affine open subset of X has property $\mathcal{P}$.

Lemma 5.1.1. Suppose $\text{Spec} A$ and $\text{Spec} B$ are affine open subschemes of a scheme X . Then $\text{Spec} A \cap \text{Spec} B$ is the union of open subsets that are simultaneously distinguished open subschemes of $\text{Spec} A$ and $\text{Spec} B$.

Proof given in [Vak17]. It suffices to show for any $\mathfrak{p} \in \text{Spec} A \cap \text{Spec} B$, there exists an open neighborhood of $\mathfrak{p}$ in $\text{Spec} A \cap \text{Spec} B$ that is simultaneously distinguished in both $\text{Spec} A$ and $\text{Spec} B$. Let $\text{Spec} A_f$ be a distinguished open subset of $\text{Spec} A$ contained in $\text{Spec} A \cap \text{Spec} B$ and containing $\mathfrak{p}$ and $\text{Spec} B_g$ be a distinguished open subset of $\text{Spec} B$ contained in $\text{Spec} A_f$ and containing $\mathfrak{p}$. Then $g \in \mathcal{O}_X(\text{Spec} B) = B$ restricts to an element $g' \in \mathcal{O}_X(\text{Spec} A_f) = A_f$. Note that

$$\text{Spec} B_g = \text{Spec} A_f \setminus \{\mathfrak{q} \in \text{Spec} A_f \mid g' \in \mathfrak{q}\} = (\text{Spec} A_f)_{g'}.$$

If $g' = g''/f^n$ in A_f , where $g'' \in A$, then $(\text{Spec} A_f)_{g'} = \text{Spec} A_{fg''}$, and this completes the proof. $\square$

Lemma 5.1.2 (affine communication lemma). Let $\mathcal{P}$ be a property enjoyed by some affine open subsets of a scheme X , such that

- (1) if an affine open subset $\text{Spec} A \hookrightarrow X$ has property $\mathcal{P}$, then for any $f \in A$, $\text{Spec} A_f \hookrightarrow X$ does too.
- (2) If $(f_1, \dots, f_n) = A$, and $\text{Spec} A_{f_i} \hookrightarrow X$ has property $\mathcal{P}$ for each i , then so does $\text{Spec} A \hookrightarrow X$.

Then the property $\mathcal{P}$ is affine-local.

Proof given in [Vak17]. Suppose $\{\text{Spec} A_i\}$ be an affine open covering of the scheme X enjoying the property $\mathcal{P}$. For any affine open subscheme $\text{Spec} A$ of X , by Lemma 5.1.1 we may cover $\text{Spec} A$ with a finite number of distinguished

open subsets $\text{Spec} A_{g_i}$, each of which is distinguished in some $\text{Spec} A_i$. By (1) one has each $\text{Spec} A_{g_i}$ has property $\mathcal{P}$, and thus $\text{Spec} A$ has the property $\mathcal{P}$ by (2). $\square$

Proposition 5.1.1. Let $\pi: X \rightarrow Y$ be a morphism of schemes.

- (1) π is **quasi-compact** if and only if for every affine open subscheme V of Y , $f^{-1}(V)$ is quasi-compact.
- (2) π is **affine** if and only if for every affine open subscheme V of Y , $f^{-1}(V)$ is affine.
- (3) π is of **locally finite type** if and only if for every affine open subscheme $V = \text{Spec} B$ of Y , $f^{-1}(V)$ can be covered by affine open subschemes $\{U_j = \text{Spec} A_j\}_{j \in J}$ such that each A_j is a finitely generated B -algebra.
- (4) π is of **finite type** if and only if for every affine open subscheme $V = \text{Spec} B$ of Y , $f^{-1}(V)$ can be covered by finitely many affine open subschemes $\{U_j = \text{Spec} A_j\}_{j \in J}$ such that each A_j is a finitely generated B -algebra.
- (5) π is **finite** if and only if for every affine open subscheme $V = \text{Spec} B$ of Y , $f^{-1}(V) = \text{Spec} A$ for some finitely generated B -module A .

Proof. Here we only give the proof of (1), and proofs for others are similar. By affine communication lemma, it suffices to show the following two statements

- (a) If $\text{Spec} A \subseteq Y$ is an affine open subset such that $\pi^{-1}(\text{Spec} A)$ is quasi-compact, then for any $f \in A$ one has $\pi^{-1}(\text{Spec} A_f)$ is quasi-compact.
- (b) If $(f_1, \dots, f_n) = A$ and $\pi^{-1}(\text{Spec} A_{f_i})$ is quasi-compact for each i , then $\pi^{-1}(\text{Spec} A)$ is quasi-compact.

The statement (b) is easy since the finite union of quasi-compact subsets is still quasi-compact, and now we prove statement (a). Suppose $\text{Spec} A \subseteq Y$ is an affine open subset such that $\pi^{-1}(\text{Spec} A)$ is quasi-compact. Then for any $f \in A$, one has

$$\pi^{-1}(\text{Spec} A_f) = (\pi^{-1}(\text{Spec} A))_{\pi^\sharp(f)},$$

where $\pi^\sharp(f) \in \mathcal{O}_X(\pi^{-1}(\text{Spec} A))$. On the other hand, since $\pi^{-1}(\text{Spec} A)$ is quasi-compact, there exists a finite affine open covering $\mathcal{U}$ of it, and by (1) of Proposition 4.3.1 one has the intersection of $(\pi^{-1}(\text{Spec} A))_{\pi^\sharp(f)}$ with any affine open subset in $\mathcal{U}$ is still affine. This shows $(\pi^{-1}(\text{Spec} A))_{\pi^\sharp(f)}$ is quasi-compact since it is a finite union of affine open subsets. $\square$

Corollary 5.1.1. Any affine morphism is quasi-compact.

Corollary 5.1.2. If the source of a morphism is a Noetherian scheme, then the morphism is quasi-compact.

Proof. It follows from Proposition 4.4.1. $\square$

Proposition 5.1.2. Suppose $f: X \rightarrow Y$ is a morphism of schemes. Then f is locally of finite type if and only if for every affine open subscheme $V = \text{Spec} B$ of Y and every affine open subscheme $U = \text{Spec} A$ of X such that $f(U) \subseteq V$, the B -algebra A is finitely generated.

Proof. See Exercise 7.7.1. $\square$

5.2. Birational morphism.

Proposition 5.2.1. Let $X \rightarrow S$ and $Y \rightarrow S$ be two morphisms and assume $Y \rightarrow S$ is locally of finite type.

- (1) Let $f, g: X \rightarrow Y$ be two morphisms making the following diagram commutes

$$\begin{array}{ccc} X & \xrightarrow{\quad} & Y \\ & \searrow & \swarrow \\ & S & \end{array}$$

Let $x \in X$ such that $f(x) = g(x) = y$ and such that $f_x^\# = g_x^\#$. Then there exists an open neighborhood U of x in X such that $f|_U = g|_U$.

- (2) Suppose S is locally Noetherian. Let $x \in X$ and $y \in Y$ be two points such that their images in S are the same point $s \in S$, and let $\phi: \mathcal{O}_{Y,y} \rightarrow \mathcal{O}_{X,x}$ be a homomorphism making the following diagram commutes

$$\begin{array}{ccc} \mathcal{O}_{X,x} & \xleftarrow{\quad \phi \quad} & \mathcal{O}_{Y,y} \\ & \nwarrow \quad \nearrow & \\ & \mathcal{O}_{S,s} & \end{array}$$

Then there exists an open neighborhood U of x in X and a morphism $f: U \rightarrow Y$ such that $f(x) = y$, $f_x^\# = \phi$ and the following diagram commutes

$$\begin{array}{ccc} U & \xrightarrow{\quad f \quad} & Y \\ & \searrow & \swarrow \\ & S & \end{array}$$

- (3) Suppose S is locally Noetherian, $X \rightarrow S$ is also locally of finite type, and $f: X \rightarrow Y$ is a morphism making the diagram in (1) commutes. Assume $f(x) = y$ and $f_x^\#: \mathcal{O}_{Y,y} \rightarrow \mathcal{O}_{X,x}$ is an isomorphism. Then there exist open neighborhoods U of x in X and V of y in Y such that f induces an isomorphism from U to V .

Proof. See Proposition 1.3.13 of [Fu06]. □

Definition 5.2.1. Let X and Y be integral schemes.

- (1) A morphism $f: X \rightarrow Y$ is dominant if $\overline{f(X)} = Y$.
(2) A dominant morphism $f: X \rightarrow Y$ is called birational if $f_\xi^\#: \mathcal{O}_{Y,\eta} \rightarrow \mathcal{O}_{X,\xi}$ is an isomorphism, where ξ, η are generic points of X and Y respectively.

Corollary 5.2.1. Let S be a locally Noetherian scheme and let X and Y be two integral schemes. Suppose we obtain the following commutative diagram

$$\begin{array}{ccc} X & \xrightarrow{\quad f \quad} & Y \\ & \searrow & \swarrow \\ & S & \end{array}$$

such that $X \rightarrow S$ and $Y \rightarrow S$ are locally of finite type and f is a birational morphism. Then there exists non-empty open subsets $U \subseteq X$ and $V \subseteq Y$ such that f induces an isomorphism from U to V .

5.3. Open immersion and closed immersion.

Definition 5.3.1. A morphism $(f, f^\sharp): (Z, \mathcal{O}_Z) \rightarrow (X, \mathcal{O}_X)$ is called an **open immersion** if it induces an isomorphism of $(Z, \mathcal{O}_Z)$ with an open subscheme of $(X, \mathcal{O}_X)$.

Definition 5.3.2. A morphism $(f, f^\sharp): (Z, \mathcal{O}_Z) \rightarrow (X, \mathcal{O}_X)$ is called a **closed immersion** if it induces a homeomorphism of Z with a closed subset of X , and $f^\sharp: \mathcal{O}_X \rightarrow f_*\mathcal{O}_Z$ is surjective.

Definition 5.3.3. A morphism $Z \rightarrow X$ is called an **immersion** if it can be written as a composite $Z \rightarrow U \rightarrow X$ such that $U \rightarrow X$ is an open immersion and $Z \rightarrow U$ is a closed immersion.

Definition 5.3.4. A subset Z of X is called locally closed if it is the intersection of an open subset with a closed subset.

Remark 5.3.1. In other words, a subset Z is locally closed if and only if Z is open in $\overline{Z}$.

Proposition 5.3.1. Let $(f, f^\sharp): (Z, \mathcal{O}_Z) \rightarrow (X, \mathcal{O}_X)$ be a morphism of schemes.

- (1) $(f, f^\sharp): (Z, \mathcal{O}_Z) \rightarrow (X, \mathcal{O}_X)$ is an open immersion if and only if f induces a homeomorphism of Z with an open subset of X and $f_p^\sharp: \mathcal{O}_{X, f(p)} \rightarrow \mathcal{O}_{Z, p}$ is an isomorphism for every $p \in Z$.
- (2) $(f, f^\sharp): (Z, \mathcal{O}_Z) \rightarrow (X, \mathcal{O}_X)$ is an immersion if and only if f induces a homeomorphism of Z with a locally closed subset of X and $f_p^\sharp: \mathcal{O}_{X, f(p)} \rightarrow \mathcal{O}_{Z, p}$ is surjective for every $p \in Z$.
- (3) The immersions are monomorphisms in the category of schemes. Moreover, the composite of immersions is an immersion, so are open immersion and closed immersion.

Proof. See Exercise 7.8.1. □

Proposition 5.3.2. Let A be a ring.

- (1) For every ideal $\mathfrak{a}$ of A , the morphism $\varphi: \operatorname{Spec} A/\mathfrak{a} \rightarrow \operatorname{Spec} A$ induced by the canonical homomorphism $\phi: A \rightarrow A/\mathfrak{a}$ is a closed immersion.
- (2) Every closed immersion into $\operatorname{Spec} A$ is isomorphic to $\operatorname{Spec} A/\mathfrak{a} \rightarrow \operatorname{Spec} A$ for some ideal $\mathfrak{a}$ of A .

Proof. See Proposition 1.3.16 of [Fu06]. □

Corollary 5.3.1. A closed immersion is affine, and thus quasi-compact.

Proposition 5.3.3 (reduced closed subscheme structure). Let $(X, \mathcal{O}_X)$ be a scheme and Y be a closed subset of X . Then there exists a unique reduced scheme structure $(Y, \mathcal{O}_Y)$ on Y which makes Y a closed subscheme of X . If $(Z, \mathcal{O}_Z) \rightarrow (X, \mathcal{O}_X)$ is a closed immersion such that its image in X contains Y ,

then there exists a unique morphism $(Y, \mathcal{O}_Y) \rightarrow (Z, \mathcal{O}_Z)$ such that the following diagram commutes

$$\begin{array}{ccc} Y & \longrightarrow & X \\ \downarrow & \nearrow & \\ Z & & \end{array}$$

Proof. See Proposition 1.3.18 of [Fu06]. □

5.4. Fibred product. In this section S always is a scheme.

Definition 5.4.1.

- (1) An S -scheme is a scheme X together with a morphism $X \rightarrow S$.
- (2) An S -morphism from an S -scheme X to an S -scheme Y is a morphism $X \rightarrow Y$ such that the diagram

$$\begin{array}{ccc} X & \longrightarrow & Y \\ & \searrow & \swarrow \\ & S & \end{array}$$

commutes.

Remark 5.4.1. For any scheme X , there is a unique morphism $X \rightarrow \operatorname{Spec} \mathbb{Z}$, so the category of schemes coincides with the category of $\operatorname{Spec} \mathbb{Z}$ -schemes.

Definition 5.4.2. Let X and Y be S -schemes. The product in the category of S -schemes is called the **fibred product** of X and Y over S , which is a S -scheme denoted by $X \times_S Y$.

Proposition 5.4.1. For S -schemes X and Y , their fibred product over S exists and is unique up to unique isomorphism.

Proof. See Proposition 1.3.20 of [Fu06]. □

Proposition 5.4.2. Let $f: X \rightarrow Y$ be a morphism of schemes and y be a point in Y with residue field $k(y)$. The projection

$$X \times_Y \operatorname{Spec} k(y) \rightarrow X$$

induces a homeomorphism from $X \times_Y \operatorname{Spec} k(y)$ with $f^{-1}(y)$ on the underlying topological spaces.

Proof. See Proposition 1.3.21 of [Fu06]. □

Proposition 5.4.3. Let

$$\begin{array}{ccc} X & & Y \\ & \searrow & \swarrow \\ & T & \\ & \downarrow & \\ & S & \end{array}$$

be morphisms of schemes. Then the diagram

$$\begin{array}{ccc}
X \times_T Y & \longrightarrow & X \times_S Y \\
\downarrow & & \downarrow \\
T & \longrightarrow & T \times_S T
\end{array}$$

is Cartesian, where the top horizontal arrow is the unique morphism whose composites with the two projections of $X \times_S Y$ to its factors are the two projections of $X \times_T Y$ to its factors, and the right vertical arrow is the unique morphism whose compositions with the two projections of $T \times_S T$ to its factors are the morphisms $X \rightarrow T$ and $Y \rightarrow T$ respectively.

Proof. See Exercise 7.9.4. □

5.4.1. Base change.

Definition 5.4.3. Let $f: X \rightarrow Y$ be a morphism of schemes. A property $\mathcal{P}$ of f is called **stable under base change**, if for any morphism $Y' \rightarrow Y$, the base change $f': X \times_Y Y' \rightarrow Y'$ also has property $\mathcal{P}$.

Proposition 5.4.4.

- (1) The quasi-compact morphism is stable under base change.
- (2) The finite type morphism is stable under base change.
- (3) The finite morphism is stable under base change.
- (4) Immersions (including closed and open immersions) are stable under base change.

Proof. Here we only give the proof of (1), and the proofs for others are similar.

Let $f: X \rightarrow Y$ be a quasi-compact morphism and $g: Y' \rightarrow Y$ be a morphism. For any affine open subset $U = \text{Spec } A$ of Y , we choose an affine open covering $\{V_i = \text{Spec } B_i\}$ of $g^{-1}(U)$, where the B_i are A -algebras. Then

$$f'^{-1}(V_i) = X \times_Y V_i = f^{-1}(U) \otimes_U V_i.$$

On the other hand, since f is quasi-compact, there exists a finite affine open covering $\{W_j = \text{Spec } C_j\}_{j=1}^n$ of $f^{-1}(U)$, where the C_j are A -algebras. Then

$$f'^{-1}(V_i) = f^{-1}(U) \otimes_U V_i = \bigcup_{j=1}^n \text{Spec } C_j \otimes_A B_i.$$

This shows $f'^{-1}(V_i)$ is quasi-compact. By varying U and thus $g^{-1}(U)$, we obtain an open covering $\{V_i\}_{i \in I}$ of Y' such that $f'^{-1}(V_i)$ is quasi-compact for each $i \in I$. This shows f' is quasi-compact. □

5.5. Separated morphisms.

Definition 5.5.1. Let $f: X \rightarrow Y$ be a morphism of schemes. The diagonal morphism $\Delta_{X/Y}: X \rightarrow X \times_Y X$ is the unique morphism satisfying

$$p \circ \Delta_{X/Y} = q \circ \Delta_{X/Y} = \text{id}_X.$$

Definition 5.5.2. Let $f: X \rightarrow Y$ be a morphism of schemes. It is called **separated** if $\Delta_{X/Y}$ is a closed immersion.

Definition 5.5.3. A scheme X is called separated if the canonical morphism $X \rightarrow \operatorname{Spec} \mathbb{Z}$ is separated.

Definition 5.5.4.

- (1) A morphism $f: X \rightarrow Y$ of schemes is called **quasi-separated** if the **diagonal morphism** is quasi-compact.
- (2) A scheme X is quasi-separated if the canonical morphism $X \rightarrow \operatorname{Spec} \mathbb{Z}$ is quasi-separated.

Proposition 5.5.1. Let $f: \operatorname{Spec} B \rightarrow \operatorname{Spec} A$ be a morphism of affine schemes. Then f is separated.

Proof. See Proposition 1.3.22 of [Fu06]. □

Proposition 5.5.2. Let $f: X \rightarrow Y$ be a morphism of schemes.

- (1) The diagonal morphism $\Delta: X \rightarrow X \times_Y X$ is an immersion.
- (2) $f: X \rightarrow Y$ is separated if and only if $\Delta_{X/Y}(X)$ is a closed subset of $X \times_Y X$.

Proof. See Proposition 1.3.23 of [Fu06]. □

Proposition 5.5.3.

- (1) A morphism $f: X \rightarrow Y$ of schemes is separated if and only if there exists an open covering $\{V_i\}_{i \in I}$ of Y such that $f^{-1}(V_i) \rightarrow V_i$ is separated.
- (2) The **immersion** is separated.
- (3) The composite of two separated morphisms is separated.
- (4) The **separated morphism** is stable under base change.
- (5) Let $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ be morphisms of schemes. If $g \circ f$ is separated, then f is separated.

Proof. Here we only give the proof of (1), and for others one can refer to the Proposition 1.3.26 in [Fu06].

Let $\{V_i\}_{i \in I}$ be an open covering of Y and $U_i = f^{-1}(V_i)$ such that $U_i \rightarrow V_i$ is separated. By (2) of Proposition 5.5.2 it suffices to show $\Delta_{X/Y}(X)$ is a closed subset of $X \times_Y X$, and it suffices to find an open covering $\mathcal{U}$ such that the preimage of each open subset in $\mathcal{U}$ under $\Delta_{X/Y}$ is closed. Suppose $g: X \times_Y X \rightarrow Y$ is the natural morphism. Then $\{g^{-1}(V_i)\}$ gives an open covering of $X \times_Y X$ and by the property of fibred product one has $g^{-1}(V_i) = U_i \times_{V_i} U_i$. It follows $\Delta_{X/Y}^{-1}(U_i \times_{V_i} U_i) = U_i$, and $U_i \rightarrow U_i \times_{V_i} U_i$ is closed since $U_i \rightarrow V_i$ is separated.

Conversely, suppose $f: X \rightarrow Y$ is separated. Then for any open covering $\{V_i\}_{i \in I}$ of Y , $f^{-1}(V_i) \rightarrow V_i$ is the base change of f under $V_i \hookrightarrow Y$, and thus it is separated since the **separated morphism** is stable under base change. □

Proposition 5.5.4. Let X be a S -scheme, where S is an affine scheme.

- (1) If X is separated over S , then the intersection of any two affine open subschemes is affine, and this fails in general if X is not separated.
- (2) If X is quasi-separated over S , then the intersection of any two affine open subschemes is a finite union of affine subschemes,

Proof. For (1), Let $U = \text{Spec} A$ and $V = \text{Spec} B$ be two affine open subschemes of X . Then by the property of fibred product one has $U \times_S V$ is affine. On the other hand, by Proposition 5.4.3 one has the following diagram is Cartesian

$$\begin{array}{ccc} U \times_X V & \longrightarrow & U \times_S V \\ \downarrow & & \downarrow \\ X & \longrightarrow & X \times_S X \end{array}$$

Thus $U \cap V = U \times_X V \rightarrow U \times_S V$ is a closed immersion since $X \rightarrow X \times_S X$ is a closed immersion, and the **closed immersion** is stable under base change. As a consequence, $U \cap V$ is affine since every closed immersion into an affine scheme is affine, and the proof of (2) is similar.

In general, consider the affine plane with double origin, that is, two copies U and V of affine plane $\mathbb{A}_k^2$ over a field k glued along the identity morphism on the open subscheme $\mathbb{A}_k^2 \setminus \{0\}$. Then the intersection $U \cap V \cong \mathbb{A}_k^2 \setminus \{0\}$, which is not affine. $\square$

5.6. Proper and projective morphisms.

5.6.1. Proper morphisms.

Definition 5.6.1. A morphism $f: X \rightarrow Y$ of schemes is called **universally closed**, if for any morphism $Y' \rightarrow Y$, the base change $f': X \times_Y Y' \rightarrow Y'$ of f is a closed map on the underlying topological spaces.

Definition 5.6.2. A morphism $f: X \rightarrow Y$ of schemes is proper if it is of finite type, separated, and universally closed.

Proposition 5.6.1.

- (1) A morphism $f: X \rightarrow Y$ of schemes is proper if and only if there exists an open covering $\{V_i\}_{i \in I}$ of Y such that $f^{-1}(V_i) \rightarrow V_i$ is proper.
- (2) The **closed immersion** is proper.
- (3) The composite of two proper morphisms is proper.
- (4) The proper morphism is stable under base change.
- (5) Let $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ be morphisms of schemes. If $g \circ f$ is proper and g is separated, then f is proper.

Proposition 5.6.2. The finite morphism is proper.

5.6.2. Projective morphisms.

Definition 5.6.3. The projective space $\mathbb{P}_{\mathbb{Z}}^n$ is defined by $\text{Proj } \mathbb{Z}[x_0, \dots, x_n]$, and for any scheme Y , projective n -space over Y is the Y -scheme $\mathbb{P}_Y^n := \mathbb{P}_{\mathbb{Z}}^n \times_{\text{Spec } \mathbb{Z}} Y$.

Definition 5.6.4. A morphism $f: X \rightarrow Y$ of schemes is projective if f can be factorized as a composite

$$X \rightarrow \mathbb{P}_Y^n \rightarrow Y$$

such that $X \rightarrow \mathbb{P}_Y^n$ is a closed immersion and $\mathbb{P}_Y^n \rightarrow Y$ is the projection. It is called quasi-projective if it can be factorized as above with $X \rightarrow \mathbb{P}_Y^n$ being an immersion.

Definition 5.6.5. A S -scheme X is called **projective** over S , if $X \rightarrow S$ is a projective morphism.

Lemma 5.6.1. For any finitely generated A -module M , one has

$$\{\mathfrak{p} \in \operatorname{Spec} A \mid M_{\mathfrak{p}} = 0\} = \operatorname{Spec} A \setminus V(\operatorname{ann}_A(M)).$$

Proposition 5.6.3. The projective morphism is proper.

Proposition 5.6.4.

- (1) The **closed immersion** is projective.
- (2) The composite of projective morphisms is projective.
- (3) The projective morphism is stable under base change.
- (4) Let $f: X \rightarrow Y$ and $f': X' \rightarrow Y'$ be projective S -morphisms between S -schemes. Then $f \times f': X \times_S X' \rightarrow Y \times_S Y'$ is projective.
- (5) Let $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ be morphism of schemes. If $g \circ f$ is projective and g is separated, then f is projective.

Proposition 5.6.5 (Segre embedding). There exists a closed immersion

$$\mathbb{P}_S^m \times_S \mathbb{P}_S^n \rightarrow \mathbb{P}_S^{(m+1)(n+1)-1},$$

which is an S -morphism.

6. COHERENT SHEAVES

6.1. $\mathcal{O}_X$ -**modules.** Let $(X, \mathcal{O}_X)$ be a ringed space.

Definition 6.1.1. A sheaf of $\mathcal{O}_X$ -module (or $\mathcal{O}_X$ -module) is a sheaf $\mathcal{F}$ such that

- (1) every open subset $U \subseteq X$, $\mathcal{F}(U)$ is an $\mathcal{O}_X(U)$ -module;
- (2) for every inclusion of open subsets $V \subseteq U$, the restriction $\mathcal{F}(U) \rightarrow \mathcal{F}(V)$ is compatible with the module structure via the ring homomorphism $\mathcal{O}_X(U) \rightarrow \mathcal{O}_X(V)$.

Definition 6.1.2. Let $\mathcal{F}$ and $\mathcal{G}$ be $\mathcal{O}_X$ -modules. A morphism of $\mathcal{O}_X$ -modules is a morphism of sheaves $\phi: \mathcal{F} \rightarrow \mathcal{G}$ such that for each open subset $U \subseteq X$, $\phi(U): \mathcal{F}(U) \rightarrow \mathcal{G}(U)$ is a homomorphism of $\mathcal{O}_X(U)$ -modules.

Notation 6.1.1. The set of morphisms between $\mathcal{O}_X$ -modules $\mathcal{F}$ and $\mathcal{G}$ is denoted by $\text{Hom}_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G})$, which is a $\mathcal{O}_X(X)$ -module.

Example 6.1.1. Let $\mathcal{F}$ and $\mathcal{G}$ be $\mathcal{O}_X$ -modules.

- (1) The sheaf $\text{hom}_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G})$ is the $\mathcal{O}_X$ -module

$$U \mapsto \text{Hom}_{\mathcal{O}_X|_U}(\mathcal{F}|_U, \mathcal{G}|_U).$$

- (2) The tensor $\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{G}$ is the $\mathcal{O}_X$ -module associated to the presheaf

$$U \mapsto \mathcal{F}(U) \otimes_{\mathcal{O}_X(U)} \mathcal{G}(U).$$

Remark 6.1.1. For any $p \in X$, one has

$$(\mathcal{F} \otimes \mathcal{G})_p = \mathcal{F}_p \otimes \mathcal{G}_p.$$

Example 6.1.2. Let $\{\mathcal{F}_i\}_{i \in I}$ be a family of $\mathcal{O}_X$ -modules.

- (1) The direct sum in the category of $\mathcal{O}_X$ -module is the sheaf associated to the presheaf

$$U \mapsto \bigoplus_{i \in I} \mathcal{F}_i(U).$$

- (2) The direct product in the category of $\mathcal{O}_X$ -module is the sheaf associated to the presheaf

$$U \mapsto \prod_{i \in I} \mathcal{F}_i(U).$$

Example 6.1.3. Let I be a direct set.

- (1) For a direct system $(\mathcal{F}_i, \phi_{ij})$ of $\mathcal{O}_X$ -modules, its direct limit in the category of $\mathcal{O}_X$ -modules is the sheaf associated to the presheaf

$$U \mapsto \varinjlim_i \mathcal{F}_i(U).$$

- (2) For an inverse system $(\mathcal{F}_i, \phi_{ij})$ of $\mathcal{O}_X$ -modules, its direct limit in the category of $\mathcal{O}_X$ -modules is the sheaf associated to the presheaf

$$U \mapsto \varprojlim_i \mathcal{F}_i(U).$$

Definition 6.1.3. Let $(f, f^\sharp): (X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$ be a morphism of ringed spaces.

- (1) Let $\mathcal{F}$ be an $\mathcal{O}_X$ -module. The direct image $f_*\mathcal{F}$ is an $f_*\mathcal{O}_X$ -module, and it becomes a $\mathcal{O}_Y$ -module via the morphism $f^\sharp: \mathcal{O}_Y \rightarrow f_*\mathcal{O}_X$.
- (2) Let $\mathcal{G}$ be an $\mathcal{O}_Y$ -module. The inverse image of $\mathcal{G}$ is defined to be $\mathcal{O}_X$ -module

$$f^*\mathcal{G} = \mathcal{O}_X \otimes_{f^{-1}\mathcal{O}_Y} f^{-1}\mathcal{G}.$$

Definition 6.1.4. An $\mathcal{O}_X$ -module $\mathcal{F}$ is called of finite presentation if there exists an open covering $\{U_i\}_{i \in I}$ of X such that on each U_i , there is an exact sequence of the form

$$\mathcal{O}_{U_i}^{\oplus m_i} \rightarrow \mathcal{O}_{U_i}^{\oplus n_i} \rightarrow \mathcal{F}|_{U_i} \rightarrow 0.$$

Definition 6.1.5. An $\mathcal{O}_X$ -module $\mathcal{F}$ is called of finite type if there exists an open covering $\{U_i\}_{i \in I}$ of X such that on each U_i , there is an exact sequence of the form

$$\mathcal{O}_{U_i}^{\oplus n_i} \rightarrow \mathcal{F}|_{U_i} \rightarrow 0.$$

Proposition 6.1.1. Let $(X, \mathcal{O}_X)$ be a ringed space and $\mathcal{F}, \mathcal{G}$ be $\mathcal{O}_X$ -modules. If $\mathcal{F}$ is of finite representation, then for every $p \in X$, one has

$$(\mathcal{H}om_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G}))_p \cong \text{Hom}_{\mathcal{O}_{X,p}}(\mathcal{F}_p, \mathcal{G}_p).$$

Proof. See Proposition 1.4.1 in [Fu06]. □

6.2. Coherent sheaves.

6.2.1. Sheaf associated to a module over affine space.

Definition 6.2.1. Let A be a ring and M be a A -module. The $\mathcal{O}_{\text{Spec} A}$ -module associated to M , denoted by $M^\sim$ is defined as follows: For every open subset $U \subseteq \text{Spec} A$, $M^\sim(U)$ consists of those mappings $s: U \rightarrow \coprod_{p \in \text{Spec} A} M_p$ satisfying the following two conditions

- (1) For every $p \in U$, one has $s(p) \in M_p$.
- (2) For every $p \in U$, there exists an open neighborhood U_p of p , $m \in M$ and $f \in A$, such that for every $q \in U_p$, one has $f \notin q$ and $s(q) = m/f$ in M_q .

For every inclusion of open subsets $V \subseteq U$, $M^\sim(U) \rightarrow M^\sim(V)$ is defined to be the restriction of mappings.

Example 6.2.1. Let A be a ring. Then $A^\sim = \mathcal{O}_{\text{Spec} A}$.

Proposition 6.2.1. Let A be a ring and M be a A -module.

- (1) $(M^\sim)_p \cong M_p$ for every $p \in \text{Spec} A$.
- (2) $(M^\sim)(D(f)) \cong M_f$ for every $f \in A$.
- (3) A sequence of A -modules

$$M' \rightarrow M \rightarrow M''$$

is exact if and only if the sequence of $\mathcal{O}_{\text{Spec} A}$ -modules

$$M'^\sim \rightarrow M^\sim \rightarrow M''^\sim$$

is exact.

(4) For any A -modules M and N , one has

$$\begin{aligned}\mathrm{Hom}_{\mathcal{O}_{\mathrm{Spec}A}}(M^\sim, N^\sim) &\cong \mathrm{Hom}_A(M, N) \\ M^\sim \otimes_{\mathcal{O}_{\mathrm{Spec}A}} N^\sim &\cong (M \otimes_A N)^\sim.\end{aligned}$$

If M is an A -module with finite presentation, then

$$\mathcal{H}om_{\mathcal{O}_{\mathrm{Spec}A}}(M^\sim, N^\sim) \cong (\mathrm{Hom}_A(M, N))^\sim.$$

(5) For a family $\{M_i\}_{i \in I}$ of A -modules, one has

$$\bigoplus_i M_i^\sim \cong (\bigoplus_i M_i)^\sim.$$

(6) For a direct system $(M_i, \phi_{ij})_{i \in I}$ of A -modules, one has

$$\varinjlim_i M_i^\sim \cong (\varinjlim_i M_i)^\sim.$$

Proof. Here we only give the proofs of (3), (5) and (6), since (1),(2) can be proved by a similar argument as shown in the structure sheaf case, and the proof of (4) is shown in Proposition 1.4.2 in [Fu06].

For (3), Note that a sequence of A -modules $M' \rightarrow M \rightarrow M''$ is exact if and only if

$$M'_p \rightarrow M_p \rightarrow M''_p$$

is exact for every $p \in \mathrm{Spec}A$, and thus it is equivalent to the sequence of sheaves $M'^\sim \rightarrow M^\sim \rightarrow M''^\sim$ is exact by (1).

For (5), It suffices to note that for each $p \in \mathrm{Spec}A$, one has

$$(\bigoplus_i M_i^\sim)_p \cong \bigoplus_i (M_i^\sim)_p \cong \bigoplus_i M_{i,p} \cong (\bigoplus_i M_i)_p^\sim,$$

since taking stalk commutes with direct sum, and by the same argument one can prove (6), since both taking stalk and direct limit are colimits, and colimits commute with colimits. $\square$

Proposition 6.2.2. Let $\phi: A \rightarrow B$ be a homomorphism of rings and $f: \mathrm{Spec}B \rightarrow \mathrm{Spec}A$ be the corresponding morphism.

- (1) For every B -module N , one has $f_* N^\sim \cong N^\sim$, where on the right hand N is regarded as an A -module.
- (2) For every A -module M , one has $f^* M^\sim \cong (B \otimes_A M)^\sim$.

Proof. See Proposition 1.4.3 in [Fu06]. $\square$

6.2.2. Coherent sheaves on Noetherian scheme.

Definition 6.2.2. Let $(X, \mathcal{O}_X)$ be a scheme. A sheaf of $\mathcal{O}_X$ -module $\mathcal{F}$ is called **quasi-coherent** if X can be covered with affine open subschemes $U_i = \mathrm{Spec}A_i$ such that $\mathcal{F}|_{U_i} \cong M_i^\sim$ for some A_i -module M_i .

Definition 6.2.3. Let $(X, \mathcal{O}_X)$ be a Noetherian scheme. A sheaf of $\mathcal{O}_X$ -module $\mathcal{F}$ is called **quasi-coherent** if X can be covered with affine open subschemes $U_i = \mathrm{Spec}A_i$ such that $\mathcal{F}|_{U_i} \cong M_i^\sim$ for some finitely generated A_i -module M_i .

Proposition 6.2.3. Let X be a scheme and $\mathcal{F}$ be a quasi-coherent sheaf on X . Then for any affine open subscheme $U = \text{Spec } A$ of X , there exists an A -module M such that $\mathcal{F}|_U = M^\sim$. If X is Noetherian and $\mathcal{F}$ is coherent, then M is finitely generated.

Proof. See Proposition 1.4.5 in [Fu06]. $\square$

Corollary 6.2.1. Let X be a scheme and $\mathcal{F}, \mathcal{G}$ be quasi-coherent $\mathcal{O}_X$ -modules.

- (1) The tensor product $\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{G}$ is quasi-coherent, and if $\mathcal{F}$ is of finite presentation, then $\mathcal{H}om_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G})$ is quasi-coherent.
- (2) Let $\phi: \mathcal{F} \rightarrow \mathcal{G}$ be a morphism. Then $\ker \phi, \text{coker } \phi$ and $\text{im } \phi$ are quasi-coherent.

If X is Noetherian, then the same statements hold for coherent sheaf.

Proof. See Corollary 1.4.6 in [Fu06]. $\square$

Proposition 6.2.4. Let $f: X \rightarrow Y$ be a morphism of schemes.

- (1) If $\mathcal{G}$ is a quasi-coherent $\mathcal{O}_Y$ -module, then $f^*\mathcal{G}$ is a quasi-coherent $\mathcal{O}_X$ -module.
- (2) If X and Y are Noetherian and $\mathcal{G}$ is coherent, then $f^*\mathcal{G}$ is coherent.
- (3) If f is quasi-compact and quasi-separated, and $\mathcal{F}$ is a quasi-coherent $\mathcal{O}_X$ -module, then $f_*\mathcal{F}$ is a quasi-coherent $\mathcal{O}_Y$ -module.

Proof. See Proposition 1.4.9 in [Fu06]. $\square$

6.2.3. Coherent sheaves on general scheme.

Definition 6.2.4. Let $(X, \mathcal{O}_X)$ be a ringed space. An $\mathcal{O}_X$ -module $\mathcal{F}$ is called **coherent** if $\mathcal{F}$ is of finite type and for every open subset U of X and every homomorphism $u: \mathcal{O}_U^{\oplus n} \rightarrow \mathcal{F}|_U$, the **kernel** of u is of finite type.

Proposition 6.2.5. Let $(X, \mathcal{O}_X)$ be a ringed space.

- (1) Suppose $\mathcal{O}_X$ is coherent. Prove that an $\mathcal{O}_X$ -module $\mathcal{F}$ is coherent if and only if $\mathcal{F}$ is of finite presentation.
- (2) Prove that Definition 6.2.4 of coherence coincides with the one in Definition 6.2.3 for Noetherian schemes.

Proof. See Exercise 7.11.2. $\square$

Lemma 6.2.1 (Oka). Let X be a complex manifold and $\mathcal{O}_X$ be the **sheaf** of holomorphic functions (and thus $(X, \mathcal{O}_X)$ gives a locally ringed space). Then $\mathcal{O}_X$ is coherent.

6.3. Sheaf of ideals.

Definition 6.3.1. Let $(X, \mathcal{O}_X)$ be a ringed space. A sheaf of ideals $\mathcal{I}$ of $\mathcal{O}_X$ is an $\mathcal{O}_X$ -submodule of $\mathcal{O}_X$.


Definition 6.3.2. Let $(X, \mathcal{O}_X)$ be a ringed space and $\mathcal{F}$ be an $\mathcal{O}_X$ -module. The **support** of $\mathcal{F}$ is defined to be the set

$$\text{supp } \mathcal{F} = \{p \in X \mid \mathcal{F}_p \neq 0\}.$$

Proposition 6.3.1. Let X be a scheme and $\mathcal{F}$ be a quasi-coherent $\mathcal{O}_X$ -module of finite type. Then $\text{supp } \mathcal{F}$ is a closed subset of X . Suppose furthermore that X is Noetherian. Then for every coherent sheaf of ideals $\mathcal{I}$ of $\mathcal{O}_X$ such that $\text{supp } \mathcal{F} \subseteq \text{supp}(\mathcal{O}_X/\mathcal{I})$, there exists $n \in \mathbb{N}$ such that $\mathcal{I}^n \mathcal{F} = 0$.

Proof. See Proposition 1.4.14 of [Fu06]. $\square$

Definition 6.3.3. Let $f: X \rightarrow Y$ be a morphism between schemes. A closed immersion $i: Z \rightarrow Y$ such that $f = i \circ g$ for a morphism $g: X \rightarrow Z$ is called a **scheme theoretic image**, if i satisfies the following universal property: For every factorization $f = i' \circ g'$ such that $i': Z' \rightarrow Y$ is a closed immersion and $g': X \rightarrow Z'$ is a morphism, there exists a unique morphism $j: Z \rightarrow Z'$ such that $i' \circ j = i$ and $j \circ g = g'$.




Proposition 6.3.2. Let $f: X \rightarrow Y$ be a quasi-compact and quasi-separated morphism between schemes. Then $\overline{f(X)}$ is its scheme theoretic image.

Corollary 6.3.1. Let $f: Z \rightarrow X$ be an immersion. If f is quasi-compact, then f can be factorized as $Z \rightarrow F \rightarrow X$ such that the first arrow is an open immersion and the second arrow is a closed immersion.

Proof. See Corollary 1.4.17 of [Fu06]. $\square$

Proposition 6.3.3 (Chow's lemma). Let S be a Noetherian scheme and $f: X \rightarrow S$ be a proper morphism. Then there exists a projective morphism $g: X' \rightarrow X$ such that $f \circ g$ is projective, and g induces an isomorphism $g^{-1}(U) \rightarrow U$ for some dense open subset U of X .



Proof. See Proposition 1.4.18 of [Fu06]. $\square$

6.4. Coherent sheaves on Proj. Let $S = \bigoplus_{d=0}^{\infty} S_d$ be a graded ring and M is a graded S -module.

Definition 6.4.1. $M(n)$ is the graded S -module whose underlying S -module is M and whose grading is defined by $(M(n))_d = M_{d+n}$.

Definition 6.4.2. For every homogeneous prime ideal $\mathfrak{p}$ of S , define $M_{(\mathfrak{p})}$ to be the $S_{(\mathfrak{p})}$ -module

$$M_{(\mathfrak{p})} = \left\{ \frac{m}{t} \mid m \in M \text{ and } t \in S \setminus \mathfrak{p} \text{ are homogeneous of the same degree} \right\}.$$

Definition 6.4.3. For every homogeneous element f in S , define $M_{(f)}$ to be the $S_{(f)}$ -module

$$M_{(f)} = \left\{ \frac{m}{f^k} \mid k \in \mathbb{N} \text{ and } m \in M \text{ is homogeneous of degree } k \deg f \right\}.$$

Definition 6.4.4. The $\mathcal{O}_{\text{Proj } S}$ -module associated to M , denoted by $M^\sim$ is defined as follows: For every open subset $U \subseteq \text{Spec } A$, $M^\sim(U)$ consists of those mappings $s : U \rightarrow \coprod_{p \in \text{Spec } A} M_{(p)}$ satisfying the following two conditions

- (1) For every $p \in U$, one has $s(p) \in M_{(p)}$.
- (2) For every $p \in U$, there exists an open neighborhood U_p of p , $m \in M$ and $f \in S$ of the same degree, such that for every $q \in U_p$, one has $f \notin q$ and $s(q) = m/f$ in $M_{(q)}$.

For every inclusion of open subsets $V \subseteq U$, $M^\sim(U) \rightarrow M^\sim(V)$ is defined to be the restriction of mappings.

Proposition 6.4.1.

- (1) For every $p \in \text{Proj } S$, one has $(M^\sim)_p \cong M_p$.
- (2) For every homogeneous element f in S of positive degree, one has $M^\sim|_{D(f)} \cong (M_{(f)})^\sim$.
- (3) $M^\sim$ is quasi-coherent. If S is Noetherian and M is finitely generated S -module, then $\text{Proj } S$ is Noetherian and $M^\sim$ is coherent.

Definition 6.4.5. For any integer n , $\mathcal{O}_{\text{Proj } S}(n)$ is defined by $S(n)^\sim$ and $\mathcal{O}_{\text{Proj } S}(1)$ is called the **twisting** sheaf. Moreover, for any $\mathcal{O}_{\text{Proj } S}$ -module $\mathcal{F}$, define

$$\mathcal{F}(n) = \mathcal{F} \otimes_{\mathcal{O}_{\text{Proj } S}} \mathcal{O}_{\text{Proj } S}(n).$$

Proposition 6.4.2. Let S be a graded ring which is generated by S_1 as a S_0 -algebra.

- (1) $\mathcal{O}_{\text{Proj } S}(n)$ is an invertible sheaf for every n .
- (2) Let M and N be S -modules. Then

$$(M \otimes_S N)^\sim \cong M^\sim \otimes_{\mathcal{O}_{\text{Proj } S}} N^\sim.$$

Proposition 6.4.3. Let S be a graded rings which are generated by S_1 as S_0 -algebra and so is T . Let $\phi : T \rightarrow S$ be a homomorphism with the property $\phi(S_d) \subseteq T_d$ for all d , and

$$U = \{q \in \text{Proj } T \mid S_+ \not\subseteq \phi^{-1}(q)\}.$$

Then U is an open subset of $\text{Proj } T$ and ϕ induces a canonical morphism $f : U \rightarrow \text{Proj } S$.

- (1) For every graded S -module M , one has

$$f^* M^\sim \cong (T \otimes_S M)^\sim|_U.$$

In particular, one has

$$f^* \mathcal{O}_{\text{Proj } S}(n) \cong \mathcal{O}_{\text{Proj } T}(n)|_U.$$

(2) For every graded T -module N , one has

$$f_*(N^\sim|_U) \cong N^\sim,$$

where on the right hand side, N is regarded as an S -module.

6.4.1. Graded S -module associated to $\mathcal{O}_{\text{Proj } S}$ -module. Let S be a graded ring which is generated by S_1 as S_0 -algebra and $X = \text{Proj } S$.

Definition 6.4.6. For any $\mathcal{O}_X$ -module $\mathcal{F}$, the graded S -module associated to $\mathcal{F}$ is defined by

$$\Gamma_*(\mathcal{F}) = \bigoplus_{n=-\infty}^{\infty} \mathcal{F}(n)(X),$$

where the S acts on $\Gamma_*(\mathcal{F})$ by the following composite of canonical homomorphisms

$$S_d \otimes \mathcal{F}(n)(X) \rightarrow \mathcal{O}_{\text{Proj } S}(d)(X) \otimes \mathcal{F}(n)(X) \rightarrow \mathcal{F}(d+n)(X).$$

Proposition 6.4.4. Let $\mathcal{F}$ be a quasi-coherent $\mathcal{O}_{\text{Proj } S}$ -module. Then $\Gamma_*(\mathcal{F})^\sim \cong \mathcal{F}$.

Corollary 6.4.1.

- (1) Let X be a closed subscheme of $\mathbb{P}_A^m = \text{Proj } A[x_0, \dots, x_m]$, where A is a ring. Then X is isomorphic to $\text{Proj } A[x_0, \dots, x_m]/\mathfrak{a}$ for some homogeneous ideal $\mathfrak{a}$ of $A[x_0, \dots, x_m]$.
- (2) A scheme X over $\text{Spec } A$ is projective if and only if it is isomorphic to $\text{Proj } S$ for some graded ring S such that $S_0 = A$ and S is finitely generated by S_1 as S_0 -algebra.

6.4.2. Ampleness and globally generated.

Definition 6.4.7. Let X be a S -scheme. An invertible $\mathcal{O}_X$ -module $\mathcal{L}$ is called **very ample** over S if there exists an immersion $i: X \rightarrow \mathbb{P}_S^m$ which is an S -morphism such that

$$\mathcal{L} \cong i^*(\mathcal{O}_{\mathbb{P}_S^m}(1)).$$

Definition 6.4.8. An $\mathcal{O}_X$ -module $\mathcal{F}$ is generated by global sections if there exists a family of sections $s_i \in \mathcal{F}(X)$ such that for every $p \in X$, $\mathcal{F}_p$ is generated by the germs $(s_i)_p$ as an $\mathcal{O}_{X,p}$ -module.

Theorem 6.4.1 (Serre). Let X be a scheme proper over $\text{Spec } A$ for a Noetherian ring A , let $\mathcal{O}_X(1)$ be an invertible $\mathcal{O}_X$ -module very ample over $\text{Spec } A$, and let $\mathcal{F}$ be a coherent $\mathcal{O}_X$ -module. There exists an integer N such that for every $n \geq N$, the $\mathcal{O}_X$ -module $\mathcal{F}(n) = \mathcal{F} \otimes \mathcal{O}_X \mathcal{O}_X(1)^{\otimes n}$ is generated by finitely many global sections.

Part 3. Homework

7. HOMEWORK

7.1. Homework-1.

Exercise 7.1. 1. A filtered abelian group is a pair $(A, F^\bullet A)$ such that A is an abelian group and

$$\dots \supset F^i A \supset F^{i+1} A \supset \dots$$

is a decreasing family of subgroups of A with indices $i \in \mathbb{Z}$. A homomorphism $f: (A, F^\bullet A) \rightarrow (B, F^\bullet B)$ of filtered abelian groups is a homomorphism $f: A \rightarrow B$ of abelian groups such that $f(F^i A) \subset F^i B$ for all $i \in \mathbb{Z}$.

- (1) Prove that filtered abelian groups form an additive category with zero objects and every morphism has kernel and cokernel.
- (2) Given an example of a morphism f such that the canonical morphism $\text{coim } f \rightarrow \text{im } f$ is not an isomorphism.

Proof. For (1), Suppose $(A, F^\bullet A)$ and $(B, F^\bullet B)$ are filtered abelian groups. The **direct product** of $(A, F^\bullet A)$ and $(B, F^\bullet B)$ is given by $(A \oplus B, F^\bullet(A \oplus B))$, where the filtration of $A \oplus B$ is given by $F^i(A \oplus B) = F^i A \oplus F^i B$, and it follows morphisms between $(A, F^\bullet A)$ and $(B, F^\bullet B)$ form an abelian group such that the composition is bilinear. This shows the category of filtered abelian groups is additive, and the zero object in this category is the zero group with trivial filtration.

Suppose $f: (A, F^\bullet A) \rightarrow (B, F^\bullet B)$ is a morphism between filtered abelian groups. Since f is also a group homomorphism between abelian groups, it has kernel and cokernel in the category of abelian groups. More precisely, $\ker f \subset A$ and $\text{coker } f = B/\text{im } f$. Then the filtrations on A and B induce filtrations on $\ker f$ and $\text{coker } f$ respectively, and thus it gives kernel and cokernel in the category of filtered abelian groups.

For (2), Suppose $A = \mathbb{Z} \oplus \mathbb{Z}$ with filtration $\mathbb{Z} \oplus \mathbb{Z} \supset \mathbb{Z} \supset \{0\}$ and $B = \mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}$ with filtration $\mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z} \supset \mathbb{Z} \oplus \mathbb{Z} \supset \{0\}$. For homomorphism given by

$$\begin{aligned} A &\rightarrow B \\ (a, b) &\mapsto (a, b, 0), \end{aligned}$$

the coimage is exactly A with filtration $\mathbb{Z} \oplus \mathbb{Z} \supset \mathbb{Z} \supset \{0\}$, but the image is $\mathbb{Z} \oplus \mathbb{Z}$ with filtration $\mathbb{Z} \oplus \mathbb{Z} \supset \mathbb{Z} \oplus \mathbb{Z} \supset \{0\}$. $\square$

Exercise 7.2. Let $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ be a sequence of morphisms in an abelian category. Prove the following statements are equivalent:

- (1) The sequence is a short exact sequence.
- (2) $B \rightarrow C$ is an epimorphism and $A \rightarrow B$ is its kernel.
- (3) $A \rightarrow B$ is a monomorphism and $B \rightarrow C$ is its cokernel.

Proof. First, we show the following lemma:

Lemma 7.1.1. Suppose $B \xrightarrow{v} C \rightarrow 0$ is a sequence of morphisms in an abelian category. Then the following statements are equivalent:

- (a) $B \rightarrow C \rightarrow 0$ is exact.
 (b) the **cokernel** of v is $C \rightarrow 0$.
 (c) v is an epimorphism.

Proof.

(a) to (b): If $B \rightarrow C \rightarrow 0$ is exact, then $\text{coim } v = \text{im } v$ is the **kernel** of $C \rightarrow 0$, that is the $\text{im } v = C \rightarrow C$. On the other hand, $\text{im } v$ is the **kernel** of $\text{cokernel } v$. Thus the **cokernel** of v is $C \rightarrow 0$.

(b) to (a): If the **cokernel** of v is $C \rightarrow 0$, then $\text{coim } v = \text{im } v = \ker(\text{coker } v) = \ker\{C \rightarrow 0\}$, that is $B \rightarrow C \rightarrow 0$ is exact.

(b) to (c): If the **cokernel** of v is $C \rightarrow 0$ and $\alpha, \beta: C \rightarrow D$ are morphisms such that $\alpha \circ v = \beta \circ v$. Then $(\alpha - \beta) \circ v = 0$, and thus by the universal property of **cokernel** there exists the following commutative diagram

$$\begin{array}{ccccc} B & \xrightarrow{v} & C & \longrightarrow & 0 \\ & & \downarrow \alpha - \beta & \swarrow & \\ & & D & & \end{array}$$

This shows $\alpha = \beta$, that is, v is an epimorphism.

(c) to (b): If v is an epimorphism and $f: C \rightarrow D$ is a morphism such that $f \circ v = 0$, then $f = 0$ since v is an epimorphism, and thus every morphism f such that $f \circ v = 0$ factors through $C \rightarrow 0$, that is, the **cokernel** of v is $C \rightarrow 0$. $\square$

Remark 7.1.1. By the same argument one can see a sequence of morphisms $0 \rightarrow A \xrightarrow{u} B$ in abelian category is exact if and only if u is a monomorphism, also if and only if $0 \rightarrow A$ is the **kernel** of u .

Now suppose $0 \rightarrow A \xrightarrow{u} B \xrightarrow{v} C \rightarrow 0$ is an exact sequence in abelian category. Then we claim u is the **kernel** of v : Since $v \circ u = 0$, by the universal property of **kernel** there exists the following diagram

$$\begin{array}{ccccc} 0 & \longrightarrow & A & \xrightarrow{u} & B \\ & & \downarrow & \searrow \bar{u} & \uparrow \\ & & \text{coim}\{0 \rightarrow A\} & \longrightarrow & \ker v \end{array}$$

Note that $\bar{u}$ is an epimorphism, since $A \rightarrow \text{coim}\{0 \rightarrow A\}$ is an epimorphism and $\text{coim}\{0 \rightarrow A\} \rightarrow \ker v$ is an isomorphism. Moreover, $\bar{u}$ is a monomorphism since u is a monomorphism: If $\alpha, \beta: D \rightarrow A$ such that $\bar{u} \circ \alpha = \bar{u} \circ \beta$, then we compose them with $\ker v \rightarrow B$, one has $u \circ \alpha = u \circ \beta$, and thus $\alpha = \beta$. Then $\bar{u}$ is both monomorphism and epimorphism, and since the category is abelian, one has $\bar{u}$ is an isomorphism, and thus u is the **kernel** of v . By the same argument, one checks directly v is the **cokernel** of u .

In a summary, above arguments show that (1) implies (2) and (3). To see (2) implies (1), it suffices to show $0 \rightarrow A \rightarrow B \rightarrow C$ is exact, since $v: B \rightarrow C$ is epimorphism already implies $B \rightarrow C \rightarrow 0$ is exact. Firstly, since u is the **kernel** of v , then it is monomorphism, and thus $0 \rightarrow A \rightarrow B$ is exact. By

previous lemma one has $0 \rightarrow A$ is the **kernel** of u , and thus $\text{coim } u = A \rightarrow A$. On the other hand, kernel of v is u . This shows the coimage of u is exactly the **kernel** of v , that is $A \rightarrow B \rightarrow C$ is exact.

$$\begin{array}{ccccccc} 0 & \longrightarrow & A & \xrightarrow{u} & B & \xrightarrow{v} & C \\ & & \downarrow \cong & & \uparrow u & & \\ & & \text{coim } u = A & \dashrightarrow & \ker v = A & & \end{array}$$

□

Exercise 7.3. Let A and B be objects in an abelian category. Prove that the canonical sequence

$$0 \rightarrow A \xrightarrow{i_1} A \oplus B \xrightarrow{p_2} B \rightarrow 0$$

is exact.

Proof. By Exercise 7.2 it suffices to show i_1 is a monomorphism and cokernel of i_1 is p_2 . By definition there exists $p_1: A \oplus B \rightarrow A$ such that $p_1 \circ i_1 = \text{id}_A$ and $i_2: B \rightarrow A \oplus B$ such that $p_2 \circ i_2 = \text{id}_B$. Moreover, $p_2 \circ i_1 = p_1 \circ i_2 = 0$ and $i_1 \circ p_1 + i_2 \circ p_2 = \text{id}_{A \oplus B}$.

- (1) Suppose $\alpha, \beta: C \rightarrow A$ are morphisms such that $i_1 \circ \alpha = i_1 \circ \beta$. Then $p_1 \circ i_1 \circ \alpha = p_1 \circ i_1 \circ \beta$ implies $\alpha = \beta$, and thus i_1 is a monomorphism.
- (2) Suppose $\alpha: C \rightarrow A \oplus B$ is a morphism such that $p_2 \circ \alpha = 0$. Then

$$i_1 \circ p_1 \circ \alpha = (i_1 \circ p_1 + i_2 \circ p_2) \circ \alpha = \alpha.$$

Thus we obtain the following commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & A & \xrightarrow{i_1} & A \oplus B & \xrightarrow{p_2} & B \longrightarrow 0 \\ & & \uparrow p_1 \circ \alpha & \nearrow \alpha & & & \\ & & C & & & & \end{array}$$

This shows $i_1: A \rightarrow A \oplus B$ satisfies the universal property of kernel.

□

Exercise 7.4. Let I be a category whose objects form a set, and let F be a covariant functor from I to the category of Abelian groups. For each $i \in I$, let $k_i: F(i) \rightarrow \bigoplus_{i \in I} F(i)$ be the canonical monomorphism. Let H be the subgroup of $\bigoplus_{i \in I} F(i)$ generated by

$$k_i(x_i) - k_j(F(i \rightarrow j)(x_i)),$$

where $i \rightarrow j$ goes over all morphisms in I , and x_i goes over all elements $F(i)$. Set

$$\varinjlim_{i \in I} F(i) = \left(\bigoplus_{i \in I} F(i) \right) / H.$$

Let $\phi_i: F(i) \rightarrow \varinjlim_{i \in I} F(i)$ be the composite of k_i with the projection $\bigoplus_{i \in I} F(i) \rightarrow (\bigoplus_{i \in I} F(i))/H$. Then we obtain $\phi_j \circ F(i \rightarrow j) = \phi_i$ for every morphism $i \rightarrow j$ in I . If A is an abelian group and $\psi_i: F(i) \rightarrow A$ ($i \in I$) is a family of homomorphisms such that $\psi_j \circ F(i \rightarrow j) = \psi_i$ for all morphisms $i \rightarrow j$ in I , then there exists one

and only one homomorphism $\psi: \varinjlim_{i \in I} F(i) \rightarrow A$ such that $\psi \circ \phi_i = \psi_i$ for all i .

Proof. First, we show the existence: Note that by the universal property of direct sum, there exists a morphism $\phi: \bigoplus_i F(i) \rightarrow A$, such that $\psi_i = \phi \circ k_i$, where $k_i: F(i) \rightarrow \bigoplus_i F(i)$ is the canonical inclusion. Moreover, for any element $k_i(x_i) - k_j(F(i \rightarrow j)(x_i)) \in H$, one has

$$\phi(k_i(x_i) - k_j(F(i \rightarrow j)(x_i))) = \psi_i(x_i) - \psi_j \circ F(i \rightarrow j)(x_i) = 0.$$

This shows $H \subseteq \ker \phi$, and thus we obtain a morphism $\psi: \varinjlim_{i \in I} F(i) \rightarrow A$ induced by ϕ , and it follows $\psi_i = \psi \circ \phi_i$.

$$\begin{array}{ccccc} & F(i) & & & \\ & \downarrow & \searrow \psi_i & & \\ \phi_i \swarrow & \bigoplus_i F(i) & \xrightarrow{\phi} & A & \\ & \downarrow & \searrow \psi & & \\ & \varinjlim_{i \in I} F(i) & & & \end{array}$$

Before we begin to prove the uniqueness, we claim any element of $\varinjlim_{i \in I} F(i)$ can be written in the form $\phi_i(x_i)$ for some $i \in I$ and some $x_i \in F(i)$: For any element $x \in \varinjlim_{i \in I} F(i) = \bigoplus_{i \in I} F(i)/H$, we write it as

$$x = \sum_{j=1}^n \phi_i(x_j), \quad x_j \in F(j).$$

It suffices to check the case of $n = 2$: Since I is a directed set, there exists $k \in I$ such that $k \geq 1, k \geq 2$. Then

$$\phi_1(x_1) + \phi_2(x_2) = \phi_k \circ F(1 \rightarrow k)(x_1) + \phi_k \circ F(2 \rightarrow k)(x_2).$$

Then x can be written as $\phi_k(F(1 \rightarrow k)(x_1) + F(2 \rightarrow k)(x_2))$ as desired.

Let us show the uniqueness: If $\psi': \varinjlim_{i \in I} F(i) \rightarrow A$ is another morphism such that $\psi_i = \psi' \circ \phi_i$ for all $i \in I$. By above claim, we know each element can be written as $\phi_i(x_i)$ for $x_i \in F(i)$. So it suffices to check $\psi(\phi_i(x_i)) = \psi'(\phi_i(x_i))$, which is clear

$$\psi(\phi_i(x_i)) = \psi_i(x_i) = \psi'(\phi_i(x_i)).$$

□

Exercise 7.5. Let $(A_i, \phi_{ji})_{i \in \mathbb{N}}$ be an inverse system of abelian groups over the direct set $(\mathbb{N}, \leq)$ of natural numbers. Consider the homomorphism

$$f: \prod_{i \in \mathbb{N}} A_i \rightarrow \prod_{i \in \mathbb{N}} A_i, \quad (a_i) \mapsto (a_i - \phi_{i+1,i}(a_{i+1})).$$

Define $\varprojlim_i^1 A_i = \text{coker } f$. Let $u: (A'_i, \phi_{ji})_{i \in \mathbb{N}} \rightarrow (A_i, \phi_{ji})_{i \in \mathbb{N}}$ and $v: (A_i, \phi_{ji})_{i \in \mathbb{N}} \rightarrow (A''_i, \phi_{ji})_{i \in \mathbb{N}}$ be morphisms of inverse systems of abelian groups such that the sequences

$$0 \rightarrow A'_i \xrightarrow{u_i} A_i \xrightarrow{v_i} A''_i \rightarrow 0$$

are exact for all i .

Prove that we have an exact sequence $0 \rightarrow \varprojlim_i A'_i \rightarrow \varprojlim_i A_i \rightarrow \varprojlim_i A''_i \rightarrow \varprojlim_i^1 A'_i \rightarrow \varprojlim_i^1 A_i \rightarrow \varprojlim_i^1 A''_i \rightarrow 0$.

Proof. Consider the following commutative diagram consisting of exact sequences

$$\begin{array}{ccccccc}
 & \ker f' & & \ker f & & \ker f'' & \\
 & \downarrow & & \downarrow & & \downarrow & \\
 0 & \longrightarrow & \prod_{i \in I} A'_i & \xrightarrow{u} & \prod_{i \in I} A_i & \xrightarrow{v} & \prod_{i \in I} A''_i \longrightarrow 0 \\
 & & \downarrow f' & & \downarrow f & & \downarrow f'' \\
 0 & \longrightarrow & \prod_{i \in I} A'_i & \xrightarrow{u} & \prod_{i \in I} A_i & \xrightarrow{v} & \prod_{i \in I} A''_i \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & \operatorname{coker} f' & & \operatorname{coker} f & & \operatorname{coker} f''
 \end{array}$$

Since $\ker f \cong \varprojlim_i A_i$, the snake lemma yields the desired result. $\square$

7.2. Homework-2.

7.2.1. *Part I.* In the following, we work with morphisms in an abelian category $\mathcal{C}$.

Exercise 7.6. Let $f: A \rightarrow B$ and $g: B \rightarrow C$ be morphisms.

(1) Suppose f and g are monomorphisms. Prove $g \circ f$ is a monomorphism.

(2) Suppose $g \circ f$ is a monomorphism. Prove f is a monomorphism.

Proof. For (1), Suppose $\alpha, \beta: D \rightarrow A$ are arbitrary morphisms such that $g \circ f \circ \alpha = g \circ f \circ \beta$. Then $f \circ \alpha = f \circ \beta$ since g is a monomorphism, and thus $\alpha = \beta$ since f is also a monomorphism.

For (2), Suppose $\alpha, \beta: D \rightarrow A$ are arbitrary morphisms such that $f \circ \alpha = f \circ \beta$. By composing g one has

$$g \circ f \circ \alpha = g \circ f \circ \beta,$$

and thus $\alpha = \beta$ since $g \circ f$ is a monomorphism. $\square$

Exercise 7.7. Let $f: A \rightarrow B$ be a morphism in $\mathcal{C}$. Recall that we have a commutative diagram

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ \downarrow & & \uparrow \\ \text{coim } f & \xrightarrow{\cong} & \text{im } f \end{array}$$

Moreover $A \rightarrow \text{coim } f$ is an epimorphism and $\text{im } f \hookrightarrow B$ is a monomorphism. Suppose we have a commutative diagram

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ \phi \downarrow & & \uparrow \psi \\ C & \xrightarrow{\cong} & D \end{array}$$

such that $\phi: A \rightarrow C$ is an epimorphism, $\psi: D \hookrightarrow B$ is a monomorphism, and $C \cong D$ is an isomorphism. Prove that there exist isomorphisms $\text{coim } f \xrightarrow{\cong} C$ and $D \xrightarrow{\cong} \text{im } f$ such that the following diagram commutes:

$$\begin{array}{ccccccc} A & \xrightarrow{\phi} & C & \xrightarrow{\cong} & D & \xrightarrow{\psi} & B \\ & \searrow & \uparrow \cong & & \uparrow \cong & \nearrow & \\ & & \text{coim } f & \xrightarrow{\cong} & \text{im } f & & \end{array}$$

Thus $\phi: A \rightarrow C$ can be identified with $\phi: A \rightarrow \text{coim } f$, and $\psi: D \hookrightarrow B$ can be identified with $\text{im } f \hookrightarrow B$.

Proof. For convenience we denote the **kernel** of f by $\tau: \ker f \rightarrow A$, denote the isomorphism between C and D by g , and denote canonical morphism from A to $\text{coim } f$ by u .

Note that $\psi \circ g \circ \phi \circ \tau = f \circ \tau = 0$. Then $\phi \circ \tau = 0$ since ψ is a monomorphism and g is an isomorphism. By universal property of cokernel there is a morphism from $\text{coim } f \rightarrow C$, denoted by α . Since $\alpha \circ u = \phi$ and both ϕ and u are

epimorphisms, one has α is an epimorphism. By the same argument one can see there exists a morphism $\beta: \text{im } f \rightarrow D$ which is a monomorphism. Since $\mathcal{C}$ is an abelian category, there is canonical isomorphism between $\text{coim } f$ and $\text{im } f$, and thus α is a monomorphism and β is an epimorphism. This shows both α and β are isomorphisms in $\mathcal{C}$, since $\mathcal{C}$ is an abelian category.

$$\begin{array}{ccccccccc}
 0 & \longrightarrow & \ker f & \xrightarrow{\tau} & A & \xrightarrow{f} & B & \longrightarrow & \text{coker } f & \longrightarrow & 0 \\
 & & & \searrow u & \downarrow \phi & & \uparrow \psi & & \swarrow v & & \\
 & & \text{coim } f & \xrightarrow{\alpha} & C & \xrightarrow{g} & D & \xleftarrow{\beta} & \text{im } f & & \\
 & & & & & & & & & \searrow & \\
 & & & & & & & & & \cong &
 \end{array}$$

□

Exercise 7.8. Define the opposite category $\mathcal{C}^\circ$ of $\mathcal{C}$ as follows:

- (a) $\mathcal{C}^\circ$ has the same objects as $\mathcal{C}$. For any object A in $\mathcal{C}$, we denote the corresponding object in $\mathcal{C}^\circ$ by A° .
- (b) For any objects A and B in $\mathcal{C}$, we define

$$\text{Hom}_{\mathcal{C}^\circ}(A^\circ, B^\circ) = \text{Hom}_{\mathcal{C}}(B, A).$$

For any morphism $\phi: A \rightarrow B$ in $\mathcal{C}$, we denote by $\phi^\circ: B^\circ \rightarrow A^\circ$ the corresponding morphism in $\mathcal{C}^\circ$.

Then

- (1) Prove that $\mathcal{C}^\circ$ is an abelian category.
- (2) Suppose

$$A \xrightarrow{\phi} B \xrightarrow{\psi} C$$

is an exact sequence in $\mathcal{C}$. Prove that

$$C^\circ \xrightarrow{\psi^\circ} B^\circ \xrightarrow{\phi^\circ} A^\circ$$

is an exact sequence in $\mathcal{C}^\circ$.

Proof. For (1), Firstly, let us see $\mathcal{C}$ is an additive category. For objects A°, B° and C° of $\mathcal{C}^\circ$, by definition $\text{Hom}_{\mathcal{C}^\circ}(A^\circ, B^\circ) = \text{Hom}_{\mathcal{C}}(B, A)$ is an abelian group, and the composition

$$\text{Hom}_{\mathcal{C}^\circ}(A^\circ, B^\circ) \times \text{Hom}_{\mathcal{C}^\circ}(B^\circ, C^\circ) \rightarrow \text{Hom}_{\mathcal{C}^\circ}(A^\circ, C^\circ)$$

is bilinear. Moreover, the **direct sum** of A°, B° in $\mathcal{C}^\circ$ is the product of A, B in $\mathcal{C}$, which also exists. Second, we show $\mathcal{C}^\circ$ is an abelian category. For morphism $f^\circ: B^\circ \rightarrow A^\circ$ in $\mathcal{C}^\circ$ corresponding to $f: A \rightarrow B$ in $\mathcal{C}$, we will show the **kernel** of f° is the **cokernel** of f . For arbitrary morphism $\alpha^\circ: C^\circ \rightarrow B^\circ$ such that $f^\circ \circ \alpha^\circ = 0$, by the universal property of kernel, there exists the following commutative diagram

$$\begin{array}{ccccc}
\ker f^\circ & \longrightarrow & B^\circ & \xrightarrow{f^\circ} & A^\circ \\
\uparrow & \nearrow \alpha^\circ & & & \\
C^\circ & & & &
\end{array}$$

This corresponds to the following commutative diagram in the category $\mathcal{C}$

$$\begin{array}{ccccc}
\ker f^\circ & \longleftarrow & B & \xleftarrow{f} & A \\
\downarrow & \nwarrow \alpha & & & \\
C & & & &
\end{array}$$

Then by uniqueness of cokernel, one has $\ker f^\circ$ is exactly the **cokernel** of f . Similarly one can show the **cokernel** of f° is exactly the **kernel** of f . This shows for any morphism $f^\circ: B^\circ \rightarrow A^\circ$, it has kernel and cokernel since $\mathcal{C}$ is an abelian category. Moreover, by the same argument one checks directly $\text{coim } f^\circ$ is isomorphic to $\text{im } f$, and $\text{im } f^\circ$ is isomorphic to $\text{coim } f$, and thus

$$\text{coim } f^\circ \cong \text{im } f^\circ,$$

since $\mathcal{C}$ is an abelian category.

For (2), Note that $A \xrightarrow{\phi} B \xrightarrow{\psi} C$ is exact if and only if $\ker \psi = \text{coim } \phi$, and since $\mathcal{C}$ is an abelian category, it is equivalent to $\ker \psi = \text{im } \phi$. By arguments in the proof of (1) it is equivalent to $\text{coker } \psi^\circ = \text{coim } \phi^\circ$. $\square$

7.2.2. Part II.

Exercise 7.9. Let X be a topological space, A an abelian group endowed with the discrete topology, and $\mathcal{F}$ the sheaf so that $\mathcal{F}(U)$ is the group of continuous maps from U to A for every open subset U of X . Prove that $\mathcal{F}$ is isomorphic to the sheaf associated to the constant presheaf $U \mapsto A$.

Proof. Firstly note that if A is equipped with discrete topology, then continuous map f from U to A is locally constant since every point $a \in A$ is an open subset, and thus its preimage $f^{-1}(a)$ is an open subset in U . On the other hand, by the construction of constant sheaf associated to the constant presheaf, the sections of it over U are also locally constant maps from U to A . This shows $\mathcal{F}$ is exactly the **sheafification** of constant presheaf. $\square$

Exercise 7.10. For every open subset U of the complex plane $\mathbb{C}$, let $\mathcal{O}(U)$ be the ring of holomorphic functions on U , and let $\mathcal{O}^*(U)$ be the group of units in $\mathcal{O}(U)$. Prove that the morphism $\mathcal{O} \rightarrow \mathcal{O}^*$ defined by

$$\begin{aligned}
\mathcal{O}(U) &\rightarrow \mathcal{O}^*(U) \\
f &\mapsto e^{2\pi\sqrt{-1}f}
\end{aligned}$$

is an epimorphism in the category of sheaves of abelian groups, but not an epimorphism in the category of presheaves. Here we regard $\mathcal{O}$ as a sheaf of abelian groups with respect to addition of functions. Prove that the **kernel** of

this morphism is isomorphic to the sheaf associated to the constant presheaf $U \mapsto \mathbb{Z}$.

Proof. For the first part, if we want to show $\mathcal{O} \xrightarrow{\exp} \mathcal{O}^* \rightarrow 0$ is an exact sequence in the category of sheaves of abelian groups, it suffices to check for each $x \in \mathbb{C}$, the following sequence of stalks is exact

$$\mathcal{O}_x \xrightarrow{\exp} \mathcal{O}_x^* \rightarrow 0.$$

It holds since for any non-vanishing holomorphic function f , $\log f$ is well-defined on a sufficiently small neighborhood of x , which proves the surjectivity. On the other hand, $\mathcal{O} \xrightarrow{\exp} \mathcal{O}^* \rightarrow 0$ is not an exact sequence in the category of presheaves of abelian groups, since

$$\mathcal{O}(\mathbb{C}^*) \xrightarrow{\exp} \mathcal{O}^*(\mathbb{C}^*) \rightarrow 0$$

fails to be an exact sequence.

For the half part, we need to prove

$$0 \rightarrow \underline{\mathbb{Z}} \xrightarrow{2\pi\sqrt{-1}} \mathcal{O} \xrightarrow{\exp} \mathcal{O}^*$$

is an exact sequence in the category of sheaves of abelian groups. It suffices to show for any open subset $U \subseteq \mathbb{C}$, the following sequence of abelian groups is exact

$$0 \rightarrow \underline{\mathbb{Z}}(U) \xrightarrow{2\pi\sqrt{-1}} \mathcal{O}(U) \xrightarrow{\exp} \mathcal{O}^*(U).$$

If $u: U \rightarrow \mathbb{Z}$ is a locally constant function, then it follows $\exp(2\pi\sqrt{-1}u) = 0$. Conversely, if $v: U \rightarrow \mathbb{C}$ is a holomorphic function such that $\exp v = 0$. Then for each $x \in U$, $v(x) = 2\pi\sqrt{-1}u(x)$, where $u: U \rightarrow \mathbb{Z}$ is a continuous function since v is continuous, and thus $v \in 2\pi\sqrt{-1}\underline{\mathbb{Z}}(U)$, since continuous integral-valued function is locally constant. $\square$

Exercise 7.11. Let $\mathcal{C}$ be a category. For any object $X \in \text{ob } \mathcal{C}$, let $\tilde{X}: \mathcal{C} \rightarrow (\text{Sets})$ be the contravariant functor from $\mathcal{C}$ to the category of sets defined by

$$\tilde{X}(Y) = \text{Hom}(Y, X).$$

A functor from $\mathcal{C}$ to the category of sets is called representable by X if it is isomorphic to $\tilde{X}$. For any contravariant functor $G: \mathcal{C} \rightarrow (\text{Sets})$, prove that we have a one-to-one correspondence

$$\begin{aligned} \text{Hom}(\tilde{X}, G) &\rightarrow G(X) \\ \alpha &\mapsto \alpha_X(\text{id}_X), \end{aligned}$$

where $\text{Hom}(\tilde{X}, G)$ is the set of natural transformations from the functor $\tilde{X}$ to the functor G . Prove the same result for covariant functors.

Proof. Let us first check this correspondence is surjective: For an object $s \in G(X)$, we define $\alpha = \alpha(s): \tilde{X} \rightarrow G$ as follows: For $X' \in \mathcal{C}$, let $\alpha_{X'}: \tilde{X}(X') \rightarrow G(X')$ be the morphism of set which sends $f: X' \rightarrow X$ to $G(f)(s)$. We now show $\alpha: \tilde{X} \rightarrow G$ is a natural transformation: For any morphism $g: X'' \rightarrow X'$ in $\mathcal{C}$, it suffices to show the following diagram commutes

$$\begin{array}{ccc}
\tilde{X}(X') & \xrightarrow{\alpha_{X'}} & G(X') \\
\downarrow \tilde{X}(g) & & \downarrow G(g) \\
\tilde{X}(X'') & \xrightarrow{\alpha_{X''}} & G(X'')
\end{array}$$

For any element $f \in \tilde{X}(X')$, that is, a morphism $f: X' \rightarrow X$, one has

$$G(f \circ g)(s) = G(g) \circ G(f)(s).$$

This shows above diagram commutes by the construction of α . Moreover, it follows

$$\alpha_C(\text{id}_X) = G(\text{id}_X)(s) = s$$

as desired.

To see above correspondence is injective: If there are two natural transformation $\alpha, \eta: \tilde{X} \rightarrow G$ such that $\alpha_X(\text{id}_X) = \eta_X(\text{id}_X)$, we need to show $\alpha = \eta$. In other words, it suffices to show for any $X' \in \mathcal{C}$, we have $\alpha_{X'} = \eta_{X'}$. For any morphism $g: X' \rightarrow X$, as α is a natural transformation, we obtain the following commutative diagram

$$\begin{array}{ccc}
\tilde{X}(X) & \xrightarrow{\alpha_X} & G(X) \\
\downarrow \tilde{X}(g) & & \downarrow G(g) \\
\tilde{X}(X') & \xrightarrow{\alpha_{X'}} & G(X')
\end{array}$$

It follows that

$$G(g) \circ \alpha_X(\text{id}_X) = \alpha_{X'} \circ \tilde{X}(g)(\text{id}_X) = \alpha_{X'}(g).$$

Similarly as η is a natural transformation, one has $(G(g) \circ \eta_X)(\text{id}_X) = \eta_{X'}(g)$. Hence

$$\alpha_{X'}(g) = G(g) \circ \alpha_X(\text{id}_X) = G(g) \circ \eta_X(\text{id}_X) = \eta_{X'}(g).$$

By considering the opposite category, it follows the same result holds for co-variant functors. $\square$

Exercise 7.12. Let $u: \mathcal{C} \rightarrow \mathcal{D}$ be a functor. Suppose that for each object $D \in \text{ob } \mathcal{D}$, the functor

$$\begin{aligned}
\mathcal{C} &\rightarrow (\text{Sets}) \\
C &\mapsto \text{Hom}(u(C), D)
\end{aligned}$$

is representable by an object $v(D) \in \text{ob } \mathcal{C}$. Then $v: \mathcal{D} \rightarrow \mathcal{C}$ is a functor right adjoint to u .

Proof. In other words, for any objects $C \in \mathcal{C}, D \in \mathcal{D}$, there is an one-to-one correspondence

$$\text{Hom}(u(C), D) \cong \text{Hom}(C, v(D)).$$

Thus by definition v is a right adjoint to u . $\square$

Exercise 7.13. Let $u: \mathcal{C} \rightarrow \mathcal{D}$ be a functor.

- (1) We say u is faithful (resp. fully faithful) if for any objects $C_1, C_2 \in \text{ob } \mathcal{C}$, the map

$$\text{Hom}(C_1, C_2) \rightarrow \text{Hom}(u(C_1), u(C_2))$$

is injective (resp. bijective).

- (2) We say u is essentially surjective if for any object D in $\mathcal{D}$, there exists an object C in $\mathcal{C}$ such that we have an isomorphism $u(C) \cong D$.
- (3) We say u is an equivalence of categories if u is both fully faithful and essentially surjective.

Suppose u is an equivalence of categories. For any $D \in \text{ob } \mathcal{D}$, choose an object $v(D) \in \text{ob } \mathcal{C}$ such that $u \circ v(D) \cong D$. Prove that v is a functor that is both left and right adjoint to $u: \mathcal{D} \rightarrow \mathcal{C}$. It is called a quasi-inverse of u .

Proof. First, we show v is a functor: If $f: D_1 \rightarrow D_2$ is a morphism in $\mathcal{D}$, then consider the following commutative diagram

$$\begin{array}{ccccc} D_1 & \xrightarrow{v} & v(D_1) & \xrightarrow{u} & D_1 \\ \downarrow f & & \downarrow v(f) & & \downarrow f \\ D_2 & \xrightarrow{v} & v(D_2) & \xrightarrow{u} & D_2 \end{array}$$

Since u is an equivalence of categories and is therefore fully faithful, there exists a morphism $v(f): v(D_1) \rightarrow v(D_2)$ still making above diagram commutes, which shows v is a functor.

We now show v is the right adjoint of u , that is to show for any $C \in \mathcal{C}$ and $D \in \mathcal{D}$, there is a one-to-one correspondence $\text{Hom}(u(C), D) = \text{Hom}(C, v(D))$. Note that u is essentially surjective, so there exists C' such that $u(C') = D$, and thus

$$\text{Hom}(u(C), D) = \text{Hom}(u(C), u(C')) = \text{Hom}(C, C').$$

On the other hand, one has

$$\text{Hom}(C, v(D)) = \text{Hom}(C, v \circ u(C')) = \text{Hom}(u(C), u \circ v \circ u(C')) = \text{Hom}(u(C), u(C')) = \text{Hom}(C, C').$$

This shows v is the right adjoint of u , and by the same argument one can see v is the left adjoint of u . $\square$

7.3. Homework-3.

Exercise 7.14. Let A be a ring. For every open subset U of $\text{Spec} A$, let S_U be the multiplicative subset $S_U = \bigcap_{\mathfrak{p} \in U} (A - \mathfrak{p})$, and let $\mathcal{P}(U) = S_U^{-1}A$. For every inclusion $V \subseteq U$ of open subsets, we have $S_U \subseteq S_V$ and hence we have a canonical homomorphism $\mathcal{P}(U) \rightarrow \mathcal{P}(V)$. This makes $\mathcal{P}$ a presheaf of rings on $\text{Spec} A$. Prove that $\mathcal{O}_{\text{Spec} A} \cong \mathcal{P}^+$.

Proof. It suffices to show for each point $\mathfrak{p} \in \text{Spec} A$, one has

$$\mathcal{O}_{\text{Spec} A, \mathfrak{p}} \cong \mathcal{P}_{\mathfrak{p}}^+.$$

Note that $\mathcal{O}_{\text{Spec} A, \mathfrak{p}} \cong A_{\mathfrak{p}}$, so it suffices to show $\mathcal{P}_{\mathfrak{p}}^+ \cong A_{\mathfrak{p}}$. But by the construction of $\mathcal{P}$, one has

$$\mathcal{P}_{\mathfrak{p}}^+ = \mathcal{P}_{\mathfrak{p}} = \varinjlim_{\mathfrak{p} \in U} \mathcal{P}(U) = \varinjlim_{\mathfrak{p} \in U} S_U^{-1}A.$$

Now it suffices to show that $A_{\mathfrak{p}}$ satisfies the universal property of inverse limit $\varinjlim_{\mathfrak{p} \in U} S_U^{-1}A$, which follows from the universal property of localization. $\square$

Exercise 7.15. Let S be a multiplicative subset of a ring A . Prove that the canonical morphism $\text{Spec} S^{-1}A \rightarrow \text{Spec} A$ induces an embedding on the underlying topological spaces.

Proof. Recall that the prime ideals in $S^{-1}A$ are in one to one correspondence with prime ideals in A which do not intersect with S , and the correspondence is given by pullback. This shows the canonical morphism $\phi: \text{Spec} S^{-1}A \rightarrow \text{Spec} A$ is bijective, and it follows that ϕ is continuous, so it suffices to show that ϕ is closed.

Note that every ideal in $S^{-1}A$ is an extended ideal, that is, it is of the form $S^{-1}\mathfrak{a}$, where $\mathfrak{a} \subseteq A$ is an ideal. Then

$$\begin{aligned} \phi(V(S^{-1}\mathfrak{a})) &= \phi(\{S^{-1}\mathfrak{p} \mid S^{-1}\mathfrak{a} \subseteq S^{-1}\mathfrak{p}, \mathfrak{p} \text{ is prime}\}) \\ &= \{\mathfrak{p} \mid \mathfrak{a} \subseteq \mathfrak{p}, \mathfrak{p} \text{ is prime}\} \\ &= V(\mathfrak{a}). \end{aligned}$$

This completes the proof. $\square$

Exercise 7.16. Let x be a point in scheme X , and let $k(x) = \mathcal{O}_{X,x}/\mathfrak{m}_x$ be the residue field at x . Construct a natural morphism $i: \text{Spec} k(x) \rightarrow X$ with image x so that the homomorphism $\mathcal{O}_{X,x} \rightarrow k(x)$ induced by $i^\sharp$ is the canonical homomorphism.

Proof. If we want to construct a morphism from scheme $\text{Spec} k(x) \rightarrow X$, it suffices to construct a continuous map between topological spaces and a morphism between structure sheaves.

- (1) For the continuous map between topological spaces $\text{Spec} k(x)$ and X , we simply send $\text{Spec} k(x)$ to the point $x \in X$ since $\text{Spec} k(x)$ is just a single point.

- (2) For the morphism $i^\#$ between structure sheaves, it is defined as follows:
 For open subset $U \subseteq X$, $i^\#: \mathcal{O}_X(U) \rightarrow \mathcal{O}_{\text{Spec } k(x)}(i^{-1}(U)) = k(x)$ is defined by

$$\mathcal{O}_X(U) \xrightarrow{\alpha} \mathcal{O}_{X,x} \rightarrow k(x),$$

where α is given by taking limit if $x \in U$, otherwise α is zero map.

Then above data gives a morphism between schemes $\text{Spec } k(x)$ and X , and by definition the homomorphism $\mathcal{O}_{X,x} \rightarrow k(x)$ induced by $i^\#$ is canonical morphism. $\square$

7.4. Homework-4.

Exercise 7.17. Let X be a topological space and $\mathcal{C}_X$ be the **sheaf** of complex-valued functions on X . Prove that $(X, \mathcal{C}_X)$ is a locally ringed space. Moreover, for each $p \in X$, one has

$$\mathfrak{m}_p = \{f \in \mathcal{C}_{X,p} \mid f(p) = 0\},$$

and the residue field $k(p) = \mathbb{C}$.

Proof. To show $(X, \mathcal{C}_X)$ is a locally ringed space, it suffices to show that for each $p \in X$, every element in $\mathcal{C}_{X,p} \setminus \mathfrak{m}_p$ is a unit. Then $\mathfrak{m}_p$ is the unique maximal ideal and thus $\mathcal{C}_{X,p}$ is the local ring.

For $f \in \mathcal{C}_{X,p} \setminus \mathfrak{m}_p$, since $f(p) \neq 0$, we may construct a continuous function g defined on an open neighborhood U of p such that $g(p) = 1/f(p)$. Then g is an inverse of f in $\mathcal{C}_{X,p}$.

To see the residue field, it suffices to note that

$$0 \rightarrow \mathfrak{m}_p \rightarrow \mathcal{C}_{X,p} \rightarrow \mathbb{C} \rightarrow 0$$

is an exact sequence. □

Exercise 7.18. Let $f: U \rightarrow X$ be an embedding of topological spaces. Then for any sheaf $\mathcal{F}$ defined on U and $p \in U$, prove that

$$(f_* \mathcal{F})_{f(p)} \cong \mathcal{F}_p.$$

Proof. Since f is a topological embedding, without loss of generality we may assume $U \subseteq X$ equipped with subspace topology and f is the inclusion map $i: U \hookrightarrow X$. By definition one has

$$\begin{aligned} (i_* \mathcal{F})_p &= \varinjlim_{p \in V \subseteq X} i_* \mathcal{F}(V) \\ &= \varinjlim_{p \in V \subseteq X} \mathcal{F}(i^{-1}(V)) \\ &= \varinjlim_{p \in V \subseteq X} \mathcal{F}(V \cap U). \end{aligned}$$

On the other hand, since U is equipped with subspace topology, every open subset of U containing p is exactly of the form $V \cap U$, where $V \subseteq X$ is an open subset containing p . This shows

$$\varinjlim_{p \in V \subseteq X} \mathcal{F}(V \cap U) = \varinjlim_{p \in U} \mathcal{F}(U) = \mathcal{F}_p$$

□

7.5. Homework-5.

7.5.1. Part I.

Exercise 7.5.1. Let S be a graded ring.

(1) Let $\mathfrak{p}$ be a prime ideal of S , and let

$$\mathfrak{p}' = \bigoplus_d (\mathfrak{p} \cap S_d).$$

Prove that $\mathfrak{p}'$ is a homogeneous prime ideal of S .

(2) Let $\mathfrak{a}$ be a homogeneous ideal of S . Prove that $\sqrt{\mathfrak{a}}$ is the intersection of homogeneous prime ideals containing $\mathfrak{a}$.

Proof. For (1), Note that for any degree d , one has

$$\mathfrak{p}' \cap S_d = \mathfrak{p} \cap S_d.$$

This shows

$$\mathfrak{p}' = \bigoplus_d (\mathfrak{p} \cap S_d) = \bigoplus_d (\mathfrak{p}' \cap S_d),$$

and thus $\mathfrak{p}$ is a homogeneous ideal.

To see $\mathfrak{p}'$ is prime, it suffices to show if a, b are two homogeneous elements such that $ab \in \mathfrak{p}'$, then either a or b in $\mathfrak{p}$. Since both a and b are homogeneous, then ab is also homogeneous. If $ab \in \mathfrak{p} \cap S_d$, then either a or b in $\mathfrak{p}$ since $\mathfrak{p}$ is prime, and thus either a or b in some $\mathfrak{p} \cap S_{d'}$ since both a and b are homogeneous. This completes the proof of $\mathfrak{p}'$ is a prime homogeneous ideal.

For (2), Suppose I is the set of all homogeneous prime ideals of S containing $\mathfrak{a}$. Firstly one has $\sqrt{\mathfrak{a}} \subseteq \bigcap_{\mathfrak{p}' \in I} \mathfrak{p}'$ since $\sqrt{\mathfrak{a}}$ equals the intersection of all prime ideals containing $\mathfrak{a}$. On the other hand, for any prime ideal $\mathfrak{p}$ containing $\mathfrak{a}$, one has the homogeneous prime $\mathfrak{p}' = \bigoplus_d (\mathfrak{p} \cap S_d) \subseteq \mathfrak{p}$ also contains $\mathfrak{a}$, since

$$\mathfrak{a} = \bigoplus_d (\mathfrak{a} \cap S_d) \subseteq \bigoplus_p (\mathfrak{a} \cap S_d).$$

Thus $\sqrt{\mathfrak{a}} = \bigoplus_{\mathfrak{p}' \in I} \mathfrak{p}'$ as desired. $\square$

Exercise 7.5.2. Let $\phi: S \rightarrow T$ be a homomorphism of graded rings. Suppose there exists an integer $m \geq 1$ such that $\phi(S_d) \subset T_{md}$ for all d . Let I be the **homogeneous ideal** of T generated by $\phi(S_+)$, and let $U = \text{Proj } T - V(T)$. Construct a morphism $f: U \rightarrow \text{Proj } S$ of schemes so that $f(q) = \phi^{-1}(q)$ for any $q \in U$ and that $f_q^\sharp: \mathcal{O}_{\text{Proj } S, f(q)} \rightarrow \mathcal{O}_{\text{Proj } T, q}$ can be identified with the homomorphism $\phi_q: S_{(\phi^{-1}(q))} \rightarrow T_{(q)}$.

Proof. Firstly let us construct the continuous map between the base topological spaces U and $\text{Proj } S$. Since $\phi: S \rightarrow T$ is a homomorphism of graded rings, one has the pullback of a homogeneous ideal $q \subseteq T$ still is a homogeneous ideal of S . As a consequence, if $q \in U = \text{Proj } T \setminus V(T)$, one has $\phi^{-1}(q)$ is a homogeneous prime ideal, and $\phi^{-1}(q)$ does not contain S_+ , otherwise q will contain T , which is a contradiction. This shows $f(q) := \phi^{-1}(q)$ gives a well-defined map from U to $\text{Spec } S$, which is also continuous.

We now construct the morphism between structure sheaves $\mathcal{O}_{\text{Proj } S}$ and $f_*\mathcal{O}_U$, where $\mathcal{O}_U$ is $\mathcal{O}_{\text{Spec } T|U}$ in fact. For each open subset $V \subseteq \text{Spec } S$, it suffices to construct a homomorphism of rings

$$\mathcal{O}_{\text{Proj } S}(V) \rightarrow \mathcal{O}_U(f^{-1}(V)) = \mathcal{O}_{\text{Proj } T|U}(f^{-1}(V)),$$

which is compatible with the restriction map between different open subsets. Given $q \in \text{Proj } T \setminus V(T)$, we will construct a homomorphism $\phi_q: S_{\phi^{-1}(q)} \rightarrow T_{(q)}$ as follows: For any element $s/t \in S_{\phi^{-1}(q)}$, $\phi(s/t)$ is defined by $\phi(s)/\phi(t)$. Since $t \notin \phi^{-1}(q)$, one has $\phi(t) \notin q$, and thus $\phi(s/t) \in T_q$. On the other hand, since ϕ is a ring homomorphism between graded rings, it maps elements of degree zero to the one of degree zero, and thus $\phi(s/t) \in T_{(q)}$ as desired.

For any element $s \in \mathcal{O}_{\text{Proj } S}(V)$, it is a map $s: V \rightarrow \coprod_{\mathfrak{p}} S_{\mathfrak{p}}$ satisfying some properties. Given $q \in f^{-1}(V)$ with $\phi^{-1}(q) = \mathfrak{p}$, that is, $f(q) = \mathfrak{p} \in V$. By composing

$$q \xrightarrow{f} \mathfrak{p} \xrightarrow{s_{\mathfrak{p}}} S_{(\mathfrak{p})} \xrightarrow{\phi_q} T_{(q)},$$

one can construct a map $t: f^{-1}(V) \rightarrow \coprod_{q \in f^{-1}(V)} T_{(q)}$. A routine check shows that t gives an element of $\mathcal{O}_U(f^{-1}(V))$, and this correspondence gives a morphism between sheaves $\mathcal{O}_{\text{Proj } S} \rightarrow f_*\mathcal{O}_U$ such that the induced morphism on stalks is exactly $\phi_q: S_{(\phi^{-1}(q))} \rightarrow T_{(q)}$. $\square$

Exercise 7.5.3. Let A be a ring, I an ideal of A , and $S = \bigoplus_{d=0}^{\infty} I^d$. Then S is a graded ring. We call $\text{Proj } S$ the blowing-up of $\text{Spec } A$ along the ideal I . Prove that the inclusion $A = S_0 \hookrightarrow S$ induces a morphism of schemes $f: \text{Proj } S \rightarrow \text{Spec } A$ such that over the open subset $U = \text{Spec } A - V(I)$, f induces an isomorphism $f^{-1}(U) \xrightarrow{\cong} U$.

Proof. For convenience we denote the inclusion $i: A \hookrightarrow S$. First, we show the inclusion i gives a continuous map f between topological spaces between $\text{Proj } S$ and $\text{Spec } A$ with $\text{im } f \subseteq U$. Given $\mathfrak{p} \in \text{Proj } S$, one has $i^{-1}(\mathfrak{p})$ is a prime ideal of A , and $i^{-1}(\mathfrak{p})$ cannot contain the ideal I , otherwise $\mathfrak{p}$ contains all power of I , and thus it contains S_+ , a contradiction.

- (1) Note that the continuous map $f: \text{Proj } S \rightarrow U$ is surjective, since for any prime ideal $\mathfrak{p} \subseteq A$, automatically it is a homogeneous prime ideal in S with $f(\mathfrak{p}) = \mathfrak{p}$.
- (2) On the other hand, if homogeneous prime ideals $\mathfrak{p}, \mathfrak{q} \in S$ such that $i^{-1}(\mathfrak{p}) = i^{-1}(\mathfrak{q})$, then $\mathfrak{p} = i^{-1}(\mathfrak{p})S_+ = i^{-1}(\mathfrak{q})S_+ = \mathfrak{q}$. This shows f is injective.
- (3) Finally, we show f is a closed map. Suppose $V_+(\mathfrak{a})$ is a closed subset of $\text{Proj } S$, where $\mathfrak{a} \subseteq S$ is a homogeneous prime ideal. Then

$$\begin{aligned} f(V_+(\mathfrak{a})) &= f(\{\mathfrak{p} \in \text{Proj } S \mid \mathfrak{a} \subseteq \mathfrak{p}\}) \\ &= \{i^{-1}(\mathfrak{p}) \mid \mathfrak{p} \in \text{Proj } S, \mathfrak{a} \subseteq \mathfrak{p}\} \\ &= \{\mathfrak{p} \in \text{Spec } A \mid i^{-1}(\mathfrak{a}) \subseteq \mathfrak{p}\} \\ &= V(i^{-1}(\mathfrak{a})). \end{aligned}$$

Above arguments shows that $f^{-1}(U) \xrightarrow{\cong} U$ as topological spaces.

To see there exists an isomorphism between structure sheaves $\mathcal{O}_U$ and $f_*\mathcal{O}_{f^{-1}(U)}$, it suffices to show $\mathcal{O}_{U,f(p)} \cong \mathcal{O}_{f^{-1}(U),p}$ holds for each $p \in f^{-1}(U)$. On one hand, one has

$$\mathcal{O}_{U,f(p)} = A_{i^{-1}(p)}.$$

On the other hand, $\mathcal{O}_{f^{-1}(U),p} \cong S_{(p)}$. Note that

$$S_{(p)} = \left\{ \frac{s}{t} \mid s \in S, t \in p, s, t \text{ are homogeneous and of the same degree.} \right\}$$

It follows that there exists an inclusion $A_{i^{-1}(p)} \hookrightarrow S_{(p)}$. Conversely, for any $s/t \in S_{(p)}$, it suffices to construct an element $a/b \in A_{i^{-1}(p)}$ such that $a/b = s/t$ in $S_{(p)}$. $\square$

Exercise 7.5.4. Let $A = R[x_1, \dots, x_n]$ be a polynomial ring, let $I = (x_1, \dots, x_n)$ be the ideal of A generated by $x_1, \dots, x_n$, let $S = \bigoplus_{d=0}^{\infty} I^d$, let $T = A[y_1, \dots, y_n]$ be the graded ring so that T_d consists of homogeneous polynomials of degree d in the variables $y_1, \dots, y_n$ with coefficients in A , and let J be the **homogeneous ideal** of T generated by $x_i y_j - x_j y_i$ ($i, j = 1, \dots, n$). Consider the **epimorphism** of graded rings $\phi: T \rightarrow S$ so that $\phi(a) = a \in S_0$ for any $a \in A$ and $\phi(y_i) = x_i \in S_1$. Prove that ϕ induces an isomorphism of schemes $\text{Proj } S \xrightarrow{\cong} \text{Proj } A[y_1, \dots, y_n]/J$.

Proof. Here we only give a proof for $n = 2$, and cases for more variables are similar. Firstly note that the **kernel** of the **epimorphism** of graded rings

$$\phi: T \rightarrow S$$

contains the ideal J , since $\phi(x_1 y_2 - x_2 y_1) = 0$. Conversely, if $\alpha = a y_1 + b y_2$ is mapped to 0, then it must be in the ideal generated by $x_1 y_2 - x_2 y_1$ since x_1, x_2 are algebraically independent. This shows $\phi: T \rightarrow S$ induces an isomorphism between graded rings S and $A[y_1, \dots, y_n]/J$, that is, a ring isomorphism which preserves the degree. Thus it induces an isomorphism between schemes $\text{Proj } S$ and $\text{Proj } A[y_1, \dots, y_n]/J$. $\square$

7.5.2. Part II.

Exercise 7.5.5. Let S be a graded ring and p be a homogeneous prime ideal. Prove that

- (1) If $f \in S \setminus p$, then $S_{(f)} \rightarrow S_{(p)}$ is injective.
- (2) $S_{(p)}$ is a local ring.

Proof. For (1), Suppose $a/f^n = b/f^m$ in $S_{(p)}$. Then there exists a homogeneous element $s \in S \setminus p$ such that

$$s(a f^m - b f^n) = 0.$$

Then $a f^m - b f^n = 0$ since p is a prime ideal, and thus $a/f^n = b/f^m$ in $S_{(f)}$.

For (2), Consider

$$\mathfrak{m} = \left\{ \frac{s}{t} \mid s, t \in p, s, t \text{ are homogeneous and of the same degree} \right\}.$$

Note that any element outside of $\mathfrak{m}$ is invertible, and thus $\mathfrak{m}$ is the only maximal ideal of the local ring $S_{(\mathfrak{p})}$. $\square$

Exercise 7.5.6. Prove that if $\emptyset \neq U \subseteq \operatorname{Spec} A$, then $0 \neq 1 \in \mathcal{O}_{\operatorname{Spec} A}(U)$.

Proof. Since $\emptyset \neq U$, we may assume there exists a non-zero ideal $\mathfrak{p} \in U$. If $0 = 1 \in \mathcal{O}_{\operatorname{Spec} A}$, that is, $\mathcal{O}_{\operatorname{Spec} A}(U)$ is a zero ring, then for any open subset $V \subseteq U$, one has $\mathcal{O}_{\operatorname{Spec} A}(V)$ is also zero ring. In particular, $\varinjlim_{\mathfrak{p} \in U} \mathcal{O}_{\operatorname{Spec} A}(U) = A_{\mathfrak{p}}$ is a zero ring, a contradiction. $\square$

7.6. Homework-6.

Exercise 7.6.1. Let X be a Noetherian topological space and let $\mathcal{F}$ be a presheaf. Suppose that for every open subset U and every finite open covering $(U_i)_{i \in I}$, the following conditions hold:

- (1) For any sections $s, t \in \mathcal{F}(U)$ such that $s|_{U_i} = t|_{U_i}$ for every $i \in I$, then $s = t$.
- (2) Let $s_i \in \mathcal{F}(U_i)$ be sections such that $s_i|_{U_i \cap U_j} = s_j|_{U_i \cap U_j}$ for every pair $i, j \in I$. Then there exists a section $s \in \mathcal{F}(U)$ such that $s|_{U_i} = s_i$ for every $i \in I$.

Prove that $\mathcal{F}$ is a sheaf.

Proof. For a Noetherian topological space X , any open subset $U \subseteq X$ is quasi-compact, that is, any open covering of U admits a finite subcovering. Thus it suffices to check above two conditions for every finite open covering. $\square$

Exercise 7.6.2. Let $(\mathcal{F}_i, \phi_{ij})_{i \in I}$ be a direct system of sheaves of abelian groups on X , and let $\varinjlim_i \mathcal{F}_i$ be the sheaf associated to the presheaf $U \mapsto \varinjlim_i \mathcal{F}_i(U)$. Prove that $\varinjlim_i \mathcal{F}_i$ is the direct limit of $(\mathcal{F}_i, \phi_{ij})_{i \in I}$ in the category of sheaves, and for every $P \in X$, we have

$$(\varinjlim_i \mathcal{F}_i)_P \cong \varinjlim_i \mathcal{F}_{i,P}.$$

Suppose furthermore that X is a Noetherian topological space. Prove that the presheaf $U \mapsto \varinjlim_i \mathcal{F}_i(U)$ is a sheaf.

Proof. First, we show $\varinjlim_i \mathcal{F}_i$ satisfies the universal property of the direct limit of $(\mathcal{F}_i, \phi_{ij})_{i \in I}$. Suppose $\mathcal{C}$ is a sheaf and $\psi_i: \mathcal{F}_i \rightarrow \mathcal{C}$ are morphisms such that $\psi_j \phi_{ij} = \psi_i$. Then For any open subset $U \subseteq X$, by the universal property of $\varinjlim_i \mathcal{F}_i(U)$, one has

$$\begin{array}{ccc} \mathcal{F}_i(U) & \xrightarrow{\psi_i(U)} & \varinjlim_i \mathcal{F}_i(U) \dashrightarrow \mathcal{C}(U) \\ \downarrow \phi_{ij}(U) & \nearrow & \uparrow \psi_j(U) \\ \mathcal{F}_j(U) & & \end{array}$$

For convenience we denote $\psi(U): \varinjlim_i \mathcal{F}_i(U) \rightarrow \mathcal{C}(U)$. Since all of $\phi_{ij}, \psi_i, \psi_j$ are morphisms between (pre)sheaves, it follows that the collection of group homomorphisms $\{\psi(U)\}_{U \subseteq X}$ gives a morphism of presheaves $U \mapsto \varinjlim_i \mathcal{F}_i(U)$ and $\mathcal{C}$, and thus gives a morphism of sheaves $\varinjlim_i \mathcal{F}_i$ and $\mathcal{C}$.

Secondly, note that

$$\begin{array}{ccc}
\mathcal{F}_i(U) & \longrightarrow & \varinjlim_i \mathcal{F}_i(U) \\
\downarrow & \searrow & \uparrow \\
\mathcal{F}_{i,p} & & \mathcal{F}_j(U) \\
\downarrow & \swarrow & \downarrow \\
\varinjlim_i \mathcal{F}_{i,p} & \longleftarrow & \mathcal{F}_{j,p}
\end{array}$$

The morphism $\varinjlim_i \mathcal{F}_i(U) \rightarrow \varinjlim_i \mathcal{F}_{i,p}$ satisfies the universal property of taking stalk, and thus by the uniqueness one has

$$(\varinjlim_i \mathcal{F}_i)_p \cong \varinjlim_i \mathcal{F}_{i,p}.$$

Finally let us suppose X is a Noetherian topological space and prove that $U \mapsto \varinjlim_i \mathcal{F}_i(U)$ is a sheaf. For every finite open covering $\{U_\alpha\}_{\alpha \in \mathcal{A}}$ of U . If $s, t \in \varinjlim_i \mathcal{F}_i(U)$ such that $s|_{U_\alpha} = t|_{U_\alpha}$ for all $\alpha \in \mathcal{A}$, then for each $\alpha \in \mathcal{A}$, there exists N_α such that for all $i > N_\alpha$, one has $s|_{U_\alpha} = t|_{U_\alpha}$ in $\mathcal{F}_i(U_\alpha)$. Since $\mathcal{A}$ is a finite index set, we may take $N > \max\{N_\alpha \mid \alpha \in \mathcal{A}\}$. Then for all $i > N$ and $\alpha \in \mathcal{A}$, one has $s|_{U_\alpha} = t|_{U_\alpha}$ in $\mathcal{F}_i(U_\alpha)$, and since $\mathcal{F}_i$ is a sheaf, one has $s = t$ in $\mathcal{F}_i(U)$ for all $i > N$, and thus $s = t$ in $\varinjlim_i \mathcal{F}_i(U)$. Similarly, one can check the other condition for $\varinjlim_i \mathcal{F}_i$ to be a sheaf by the same argument. $\square$

Exercise 7.6.3. Let $(\mathcal{F}_i, \phi_{ij})_{i \in I}$ be an inverse system of sheaves of abelian groups on X . Prove that the presheaf $U \mapsto \varprojlim_i \mathcal{F}_i(U)$ is a sheaf and it is the inverse limit of $(\mathcal{F}_i, \phi_{ij})_{i \in I}$ in the category of sheaves.

Proof. By the same argument as above exercise one can show that $\varprojlim_i \mathcal{F}_i$ is the inverse limit in the category of sheaves, so here we only prove that $U \mapsto \varprojlim_i \mathcal{F}_i(U)$ is a sheaf.

Recall that a presheaf $\mathcal{F}$ is a sheaf if and only if for an open subset U and open covering $\{U_\alpha\}$ of U , the following sequence is exact

$$0 \rightarrow \mathcal{F}(U) \rightarrow \prod_{\alpha} \mathcal{F}(U_\alpha) \rightarrow \prod_{\alpha, \beta} \mathcal{F}(U_{\alpha\beta}).$$

Since the inverse limit is left exact, one has

$$0 \rightarrow \varprojlim_i \mathcal{F}_i(U) \rightarrow \prod_{\alpha} \varprojlim_i \mathcal{F}_i(U_\alpha) \rightarrow \prod_{\alpha, \beta} \varprojlim_i \mathcal{F}_i(U_{\alpha\beta}).$$

This completes the proof. $\square$

7.7. Homework-7.

Exercise 7.7.1. Let $f: X \rightarrow Y$ be a morphism of schemes which is locally of finite type. Let $V = \text{Spec} B$ be an affine open subscheme of Y and $U = \text{Spec} A$ be an affine open subscheme of X such that $f(U) \subseteq V$.

- (1) Prove that $f^{-1}(V)$ can be covered by affine open subschemes $\{U_\lambda = \text{Spec} A_\lambda\}_{\lambda \in \Lambda}$ such that A_λ are finitely generated B -algebras.
- (2) Prove that there exist finitely many $a_1, \dots, a_n \in A$ such that $D(a_1), \dots, D(a_n)$ cover U , and each A_{a_i} is a finitely generated B -algebra.
- (3) Prove that A is a finitely generated B -algebra.

Proof. For (1), Since $V \cap V_i$ is an open subset of $V_i = \text{Spec} B_i$, then $V \cap V_i$ is a union of distinguished open subsets $D(f_{ik}) = \text{Spec}(B_i)_{f_{ik}}$, and for each $f_{ik} \in B_i$, for convenience we still use f_{ik} to denote the image of f_{ik} under the morphism $B_i \rightarrow A_{ij}$. Then $(A_{ij})_{f_{ik}}$ is a finitely generated $(B_i)_{f_{ik}}$ -algebra. After relabelling the index, in fact we have shown that V is covered by affine schemes $\text{Spec} C_i$ such that each $f^{-1}(\text{Spec} C_i)$ is a union of affine schemes $\text{Spec} D_{ij}$, where D_{ij} is a finitely generated C_i -algebra.

For each point $p \in V$, suppose it lies in the affine scheme $\text{Spec} C_i$. Then there exists a distinguished open subset $\text{Spec} B_{f_p} \subseteq \text{Spec} C_i$ which contains p . For convenience we still use f_p to denote the image of f_p under the morphism $B \rightarrow C_i \rightarrow D_{ij}$. Then each $(D_{ij})_{f_p}$ is a finitely generated B_{f_p} -algebra, and thus a finitely generated B -algebra. This completes the proof.

For (2), Since $f^{-1}(V)$ is covered by affine schemes $\text{Spec} A_\lambda$, then $U = \text{Spec} A$ is also covered by the intersection of $\text{Spec} A \cap \text{Spec} A_\lambda$. Moreover, since both A and A_λ are affine, then by Lemma 5.1.1 one can pick a collection of open subset $U_{\lambda i}$ such that $U_{\lambda i}$ are simultaneously the distinguished open subsets of $\text{Spec} A$ and $\text{Spec} A_\lambda$. For convenience we write

$$U_{\lambda i} \cong \text{Spec} A_{f_{\lambda i}} \cong \text{Spec}(A_\lambda)_{g_{\lambda i}}.$$

Since A_λ is a finitely generated B -algebra, so is $(A_\lambda)_{g_{\lambda i}}$, and thus each $A_{f_{\lambda i}}$ can be realized as a finitely generated B -algebra, which completes the proof.

For (3), Note that $\text{Spec} A$ is covered by distinguished open subsets $D(a_1), \dots, D(a_n)$ if and only if $(a_1, \dots, a_n) = A$. Thus it reduces to the following lemma of commutative algebra.

Lemma 7.7.1. Let A be a B -algebra and $(a_1, \dots, a_n) = A$. If A_{a_i} is a finitely generated B -algebra for each i , then A is also a finitely generated B -algebra.

Proof. For each $x \in A$, since A_{a_i} is finitely generated B -algebra, its image in A_{a_i} is equal to some

$$F^i\left(\frac{x_1^i}{k_1^i}, \dots, \frac{x_{j_i}^i}{k_{j_i}^i}, \frac{x_i^i}{a_i^{j_i}}\right),$$

where $x_j^i/a_i^{k_j^i}$ are generators of A_{a_i} over B , and F^i is some polynomial with coefficients in B . After multiplying by a large power of a_i , there are n equations

in A which are of the form

$$a_i^N x = \tilde{F}^i(x_i^i, \dots, x_{j_i}^i, a_i).$$

On the other hand, since $(a_1, \dots, a_n) = 1$, there exists $m_1, \dots, m_n \in A$ such that

$$m_1 a_1 + \dots + m_n a_n = 1.$$

Exponentiate above equation to the nN -th power and multiply by x , one has

$$x = G(a_1, \dots, a_n, m_1, \dots, m_n, x_j^i),$$

where G is a polynomial with coefficients in B , since each monomial of m_i , there exists some a_j in the coefficients such that the power of a_j is $\geq N$, and thus $a_i^N x$ can be replaced by $\tilde{F}^i$. This shows any $x \in A$ can be expressed as a polynomial of m_i, a_i, x_j^i with coefficients in B , and thus A is a finitely generated B -algebra. □

□

7.8. Homework-8.

Exercise 7.8.1. Let $(f, f^\sharp): (Z, \mathcal{O}_Z) \rightarrow (X, \mathcal{O}_X)$ be a morphism of schemes.

- (1) $(f, f^\sharp): (Z, \mathcal{O}_Z) \rightarrow (X, \mathcal{O}_X)$ is an open immersion if and only if f induces a homeomorphism of Z with an open subset of X and $f_p^\sharp: \mathcal{O}_{X, f(p)} \rightarrow \mathcal{O}_{Z, p}$ is an isomorphism for every $p \in Z$.
- (2) $(f, f^\sharp): (Z, \mathcal{O}_Z) \rightarrow (X, \mathcal{O}_X)$ is an immersion if and only if f induces a homeomorphism of Z with a locally closed subset of X and $f_p^\sharp: \mathcal{O}_{X, f(p)} \rightarrow \mathcal{O}_{Z, p}$ is an epimorphism for every $p \in Z$.
- (3) The immersions are monomorphisms in the category of schemes. Moreover, the composite of immersions is an immersion, so are open immersions and closed immersions.

Proof. For (1), Note that by definition one has $(f, f^\sharp): (Z, \mathcal{O}_Z) \rightarrow (X, \mathcal{O}_X)$ is an open immersion if and only if it induces an isomorphism between $(Z, \mathcal{O}_Z)$ and an open subscheme of $(X, \mathcal{O}_X)$. Since f has already induced a homeomorphism of Z with an open subset of X , it suffices to show for every $p \in Z$, $f_p^\sharp: \mathcal{O}_{X, f(p)} \rightarrow \mathcal{O}_{Z, p}$ is an isomorphism if and only if $(f_* \mathcal{O}_Z)_{f(p)} \cong \mathcal{O}_{X, f(p)}$. In general it fails, but since f induces a homeomorphism, one has

$$(f_* \mathcal{O}_Z)_{f(p)} = \varinjlim_{p \in f^{-1}(V)} \mathcal{O}_Z(f^{-1}(V)) = \varinjlim_{p \in U} \mathcal{O}_Z(U) = \mathcal{O}_{Z, p}$$

is an isomorphism. On the other hand, one has the following commutative diagram

$$\begin{array}{ccc} f_p^\sharp: \mathcal{O}_{X, f(p)} & \xrightarrow{\quad} & \mathcal{O}_{Z, p} \\ & \searrow & \nearrow \\ & (f_* \mathcal{O}_Z)_{f(p)} & \end{array}$$

Thus $f_p^\sharp$ is an isomorphism if and only if $(f_* \mathcal{O}_Z)_{f(p)} \cong \mathcal{O}_{X, f(p)}$, and by the same argument one can show (2).

For (3), Since the composite of epimorphisms is an epimorphism, one has the composite of immersions is an immersion, so are open immersions and closed immersions.

We now show immersions are monomorphisms in the category of schemes. Suppose $\alpha, \beta: \tilde{Z} \rightarrow Z$ are two morphisms between schemes such that $f \circ \alpha = f \circ \beta$, where f is an immersion. Firstly $\alpha = \beta$ as morphisms between topological spaces since f induces homeomorphism between underlying topological spaces. Moreover, on each stalk one has

$$\alpha^\sharp \circ f^\sharp = \beta^\sharp \circ f^\sharp,$$

and thus one has $\alpha^\sharp = \beta^\sharp$ on each stalk since $f^\sharp$ is an epimorphism. □

Exercise 7.8.2. Let S be a graded ring and let $\mathfrak{a}$ be a homogeneous ideal of S . Prove that the canonical homomorphism $S \rightarrow S/\mathfrak{a}$ induces a closed immersion $\text{Proj } S/\mathfrak{a} \rightarrow \text{Proj } S$.

Proof. Firstly the canonical homomorphism $S \rightarrow S/\mathfrak{a}$ is a homomorphism of graded rings, and thus the pullback of homogeneous prime ideals are still homogeneous prime ideals. For any $\mathfrak{p}/\mathfrak{a} \in \text{Proj } S/\mathfrak{a}$, the pullback of $\mathfrak{p}/\mathfrak{a}$ is $\mathfrak{p}$, which is a homogeneous prime ideal which contains $\mathfrak{a}$ and cannot contain S_+ , otherwise $\mathfrak{p}/\mathfrak{a}$ contains $(S/\mathfrak{a})_+$, a contradiction. On the other hand, any homogeneous prime ideal $\mathfrak{p} \in V_+(\mathfrak{a})$ gives an element in $\text{Proj } S/\mathfrak{a}$, and these two constructions are inverse to each other. Thus one has $\text{Proj } S/\mathfrak{a} \cong V_+(\mathfrak{a})$ as sets. Moreover, it is also easy to show the canonical homomorphism $\text{Proj } S/\mathfrak{a} \rightarrow V_+(\mathfrak{a})$ is a closed map and thus it is a homeomorphism with respect to the Zariski topology.

Now it suffices to show for each $\mathfrak{p}/\mathfrak{a} \in \text{Proj } S/\mathfrak{a}$, the canonical homomorphism $\mathcal{O}_{\text{Proj } S, \mathfrak{p}} \rightarrow \mathcal{O}_{\text{Proj } S/\mathfrak{a}, \mathfrak{p}/\mathfrak{a}}$ is surjective. Note that

$$\begin{aligned}\mathcal{O}_{\text{Proj } S, \mathfrak{p}} &\cong S_{(\mathfrak{p})} \\ \mathcal{O}_{\text{Proj } S/\mathfrak{a}, \mathfrak{p}/\mathfrak{a}} &\cong (S/\mathfrak{a})_{(\mathfrak{p}/\mathfrak{a})},\end{aligned}$$

and the canonical homomorphism is given by projection, which is surjective. This completes the proof. $\square$

7.9. Homework-9.

7.9.1. Part I.

Exercise 7.9.1. Finish step 3 of the proof of Proposition 1.3.20 as follows.

(1) Let

$$\begin{array}{ccc} U_i \times_S Y & \xrightarrow{q_i} & Y \\ \downarrow p_i & & \downarrow \\ U_i & \longrightarrow & S \end{array}$$

be the **fibred product** of U_i and Y . Prove that there exists one and only one isomorphism $\phi_{ij}: p_i^{-1}(U_i \cap U_j) \rightarrow p_j^{-1}(U_i \cap U_j)$ such that the following diagrams commute:

$$\begin{array}{ccc} p_i^{-1}(U_i \cap U_j) & \xrightarrow{\phi_{ij}} & p_j^{-1}(U_i \cap U_j) \\ \searrow p_i & & \downarrow p_j \\ & & U_i \cap U_j \end{array} \quad \begin{array}{ccc} p_i^{-1}(U_i \cap U_j) & \xrightarrow{\phi_{ij}} & p_j^{-1}(U_i \cap U_j) \\ \searrow q_i & & \downarrow q_j \\ & & Y \end{array}$$

(2) Prove that $\phi_{ij} = \phi_{ji}^{-1}$ and $\phi_{jk} \circ \phi_{ij} = \phi_{ik}$ when restricted to $p_i^{-1}(U_i \cap U_j \cap U_k)$. So we can glue the schemes $U_i \times_S Y$ together to get a scheme Z .

(3) Suppose we have a commutative diagram

$$\begin{array}{ccc} Z & \xrightarrow{q} & Y \\ \downarrow p & & \downarrow \\ X & \longrightarrow & S \end{array}$$

and suppose X has an open covering $X = \bigcup_i U_i$ such that

$$\begin{array}{ccc} p^{-1}(U_i) & \xrightarrow{q} & Y \\ \downarrow p & & \downarrow \\ U_i & \longrightarrow & S \end{array}$$

are fibred product for all i . Prove that Z in the first diagram is the **fibred product** of X and Y over S .

Proof. For (1), By step 2 of the proof of Proposition 1.3.20, one has $p_i^{-1}(U_i \cap U_j)$ is the **fibred product** of $U_i \cap U_j$ and Y over S , so is $p_j^{-1}(U_i \cap U_j)$. Thus there exists one and only one isomorphism $\phi_{ij}: p_i^{-1}(U_i \cap U_j) \rightarrow p_j^{-1}(U_i \cap U_j)$ by the universal property of fibred product.

For (2), Note that $\phi_{ij} \circ \phi_{ji}$ is an isomorphism such that the following diagram commutes

$$\begin{array}{ccc} p_i^{-1}(U_i \cap U_j) & \xrightarrow{\phi_{ij} \circ \phi_{ji}} & p_i^{-1}(U_i \cap U_j) \\ \searrow p_i & & \downarrow p_i \\ & & U_i \cap U_j, \end{array}$$

so is the identity map. Then by the fact that the **fibred product** is unique up to a unique isomorphism, one has $\phi_{ij} \circ \phi_{ji} = \text{id}$, that is, $\phi_{ij} = \phi_{ji}^{-1}$ as desired. The same argument shows that $\phi_{jk} \circ \phi_{ij} = \phi_{ik}$.

For (3), In order to prove that Z satisfies the universal property of fibred product, we need to show for any commutative diagram

$$\begin{array}{ccc} W & \xrightarrow{\beta} & Y \\ \alpha \downarrow & & \downarrow \\ X & \longrightarrow & S, \end{array}$$

there exists a unique morphism $W \rightarrow Z$ such that the following diagram commutes

$$\begin{array}{ccccc} W & & & & \\ & \searrow \beta & & & \\ & & Z & \xrightarrow{q} & Y \\ & \searrow \alpha & \downarrow p & & \downarrow \\ & & X & \longrightarrow & S. \end{array}$$

Since $X = \bigcup_i U_i$, an observation is that the morphism $\alpha: W \rightarrow X$ is equivalent to a collection of morphisms $\{\alpha_i: W \rightarrow U_i\}$ which are compatible with each other. Thus for each i there exists a unique morphism $W \rightarrow p^{-1}(U_i)$ such that the following commutative diagram

$$\begin{array}{ccccc} W & & & & \\ & \searrow \beta & & & \\ & & p^{-1}(U_i) & \xrightarrow{q} & Y \\ & \searrow \alpha_i & \downarrow p & & \downarrow \\ & & U_i & \longrightarrow & S \end{array}$$

since $p^{-1}(U_i)$ is the **fibred product** of U_i and Y over S . By uniqueness the collection of morphisms $\{W \rightarrow p^{-1}(U_i)\}$ can be glued to a unique morphism from $W \rightarrow Z$, as desired. $\square$

Exercise 7.9.2. Prove the isomorphism $X_i \times_S Y \cong X_i \times_{S_i} Y_i$ in step 6 in the proof of Proposition 1.3.20.

Proof. It suffices to note that given morphisms $f: Z \rightarrow X$ and $g: Z \rightarrow Y$ over S , the image of g must land inside Y_i . $\square$

7.9.2. Part II.

Exercise 7.9.3. Let $\mathcal{C}$ be a category in which fibred product exists. Consider commutative diagrams

$$\begin{array}{ccc}
X'' & \xrightarrow{f'} & X' & \xrightarrow{g'} & X \\
\downarrow & & \downarrow & & \downarrow \\
S'' & \xrightarrow{f} & S' & \xrightarrow{g} & S
\end{array}
\quad
\begin{array}{ccc}
X'' & \xrightarrow{g' \circ f'} & X \\
\downarrow & & \downarrow \\
S'' & \xrightarrow{g \circ f} & S
\end{array}$$

Suppose the second square is Cartesian. Prove that the first square is Cartesian if and only if the third one is Cartesian.

Proof. Suppose the first square is Cartesian and consider the following commutative diagram

$$\begin{array}{ccc}
W & \xrightarrow{\beta} & X \\
\alpha \searrow & & \downarrow \\
& X'' & \xrightarrow{g' \circ f'} X \\
& \downarrow & \downarrow \\
& S'' & \xrightarrow{g \circ f} S
\end{array}$$

By composing α and f , one obtains the following commutative diagram


$$\begin{array}{ccc}
W & \xrightarrow{\beta} & X \\
\phi \searrow & & \downarrow \\
& X' & \xrightarrow{g'} X \\
f \circ \alpha \searrow & & \downarrow \\
& S' & \xrightarrow{g} S
\end{array}$$

where the morphism $\phi: W \rightarrow X'$ is induced by the assumption that the second square is Cartesian. On the other hand, consider the following commutative diagram


$$\begin{array}{ccc}
W & \xrightarrow{\phi} & X' \\
\psi \searrow & & \downarrow \\
& X'' & \xrightarrow{f'} X' \\
\alpha \searrow & & \downarrow \\
& S'' & \xrightarrow{f} S'
\end{array}$$

where the morphism $\psi: W \rightarrow X''$ is induced by the assumption that the first square is Cartesian, and that is exactly the morphism making the third square to be Cartesian.


Conversely, suppose the third square is Cartesian, and consider the following commutative diagram



Then by the assumption the third square is Cartesian, one has the following commutative diagram



where the induced morphism $W \rightarrow X''$ is denoted by ϕ . In order to show the first square is Cartesian, it suffices to show the following diagram commutes



which follows from the second square is Cartesian. $\square$

Exercise 7.9.4.

- (1) Let $f: X \rightarrow S$ and $g: Y \rightarrow S$ be maps of sets. Prove that their fibred product is $(X \times_S Y, p, q)$, where

$$X \times_S Y = \{(x, y) \mid x \in X, y \in Y, f(x) = g(y)\},$$

and $p: X \times_S Y \rightarrow X, q: X \times_S Y \rightarrow Y$ are the projections $p(x, y) = x$ and $q(x, y) = y$ respectively.

- (2) Use the description of the fibred product in (1) to prove Proposition 5.4.3 for the category of sets.
- (3) Let $\mathcal{C}$ be a category and let

$$\begin{array}{ccc} X' & \longrightarrow & X \\ \downarrow & & \downarrow \\ S' & \longrightarrow & S \end{array}$$

be a commutative diagram in $\mathcal{C}$. For every object $Z \in \text{Ob } \mathcal{C}$, it induces a commutative diagram

$$\begin{array}{ccc}
\mathrm{Hom}(Z, X') & \longrightarrow & \mathrm{Hom}(Z, X) \\
\downarrow & & \downarrow \\
\mathrm{Hom}(Z, S') & \longrightarrow & \mathrm{Hom}(Z, S)
\end{array}$$

in the category of sets. Prove that the first diagram is Cartesian in $\mathcal{C}$ if and only if the second diagram is Cartesian in the category of sets for every object $Z \in \mathrm{Ob}\mathcal{C}$.

- (4) Use (2) and (3) to prove Proposition 1.3.24 for every category $\mathcal{C}$ in which fibred product exists.

Proof. For (1), Suppose there exist morphisms (between sets) $\alpha: W \rightarrow X$ and $\beta: W \rightarrow Y$ such that the following diagram commutes

$$\begin{array}{ccc}
W & \xrightarrow{\beta} & Y \\
\alpha \downarrow & & \downarrow g \\
X & \xrightarrow{f} & S.
\end{array}$$

Then one can construct

$$\begin{aligned}
\varphi: W &\rightarrow X \times_S Y \\
w &\mapsto (\alpha(w), \beta(w))
\end{aligned}$$

such that the following diagram commutes

$$\begin{array}{ccccc}
W & & & & \\
\downarrow \alpha & \searrow \varphi & & \searrow \beta & \\
X & \xrightarrow{f} & X \times_S Y & \longrightarrow & Y \\
& & \downarrow & & \downarrow g \\
& & S & &
\end{array}$$

Moreover, any morphism from $W \rightarrow X \times_S Y$ such that above diagram commutes must be of this form. This shows $X \times_S Y$ satisfies the universal property of fibred product.

For (2), Given the morphisms between sets as follows

$$\begin{array}{ccc}
X & & Y \\
& \searrow f & \swarrow g \\
& T & \\
& \downarrow \phi & \\
& S. &
\end{array}$$

By using the description of the fibred product in (1), the following diagram commutes

$$\begin{array}{ccc}
X \times_T Y & \longrightarrow & X \times_S Y \\
\downarrow & & \downarrow \\
T & \longrightarrow & T \times_S T,
\end{array}$$

since the morphisms in above diagram can be described as follows:

- (a) $X \times_T Y \rightarrow X \times_S Y$ is given by $(x, y) \mapsto (x, y)$.
- (b) $X \times_S Y \rightarrow T \times_S T$ is given by $(x, y) \mapsto (f(x), g(y))$.
- (c) $T \rightarrow T \times_S T$ is given by $t \mapsto (t, t)$.
- (d) $X \times_T Y \rightarrow T$ is given by $(x, y) \mapsto f(x)$ or $(x, y) \mapsto g(y)$, since in this case $f(x) = g(y)$.

Moreover, given morphisms $\alpha: W \rightarrow T$ and $\beta: W \rightarrow X \times_S Y$ such that the following diagram commutes

$$\begin{array}{ccccc}
W & & & & \\
& \searrow \beta & & & \\
& & X \times_T Y & \longrightarrow & X \times_S Y \\
& \searrow \alpha & \downarrow & & \downarrow \\
& & T & \longrightarrow & T \times_S T,
\end{array}$$

an observation is that the image of β lies in $X \times_T Y$, and thus there is a unique morphism from W to $X \times_T Y$ such that above diagram commutes, which is exactly β itself. This shows $X \times_T Y$ satisfies the universal property of fibred product.

For (3), Suppose

$$\begin{array}{ccc}
X' & \longrightarrow & X \\
\downarrow & & \downarrow f \\
S' & \xrightarrow{g} & S
\end{array}$$

is Cartesian. For any $Z \in \text{Ob } \mathcal{C}$ and morphisms $\alpha: Z \rightarrow S'$ and $\beta: Z \rightarrow X$ such that $f \circ \beta = g \circ \alpha$, by the universal property there exists a unique $\phi: Z \rightarrow X'$. Conversely given morphisms $Z \rightarrow X'$, it is easy to construct morphisms $\alpha: Z \rightarrow S'$ and $\beta: Z \rightarrow X$ such that $f \circ \beta = g \circ \alpha$. This shows $\text{Hom}(Z, X') \cong \text{Hom}(Z, S') \times_{\text{Hom}(Z, S)} \text{Hom}(Z, X)$ by the description of fibred product in (1). This shows

$$\begin{array}{ccc}
\text{Hom}(Z, X') & \longrightarrow & \text{Hom}(Z, X) \\
\downarrow & & \downarrow \\
\text{Hom}(Z, S') & \longrightarrow & \text{Hom}(Z, S)
\end{array}$$

is Cartesian, and by the same argument one can prove the converse statement.

For (4), By (3), it suffices to show that for each $Z \in \text{Ob}\mathcal{C}$, the following diagram is Cartesian

$$\begin{array}{ccc} \text{Hom}(Z, X \times_T Y) & \longrightarrow & \text{Hom}(Z, X \times_S Y) \\ \downarrow & & \downarrow \\ \text{Hom}(Z, T) & \longrightarrow & \text{Hom}(Z, T \times_S T). \end{array}$$

Note that by the proof of (3), one can see

$$\begin{aligned} \text{Hom}(Z, T \times_S T) &= \text{Hom}(Z, T) \times_{\text{Hom}(Z, S)} \text{Hom}(Z, T) \\ \text{Hom}(Z, X \times_S Y) &= \text{Hom}(Z, X) \times_{\text{Hom}(Z, S)} \text{Hom}(Z, Y) \\ \text{Hom}(Z, X \times_T Y) &= \text{Hom}(Z, X) \times_{\text{Hom}(Z, T)} \text{Hom}(Z, Y), \end{aligned}$$

and thus the desired result follows from (2). $\square$

7.9.3. Part III.

Exercise 7.9.5. Let X, Y be S -schemes and $f, g: X \rightarrow Y$ be S -morphisms. Suppose $(f, g): X \rightarrow Y \times_S Y$ is the morphism such that $p \circ (f, g) = f, q \circ (f, g) = g$ and K is the **fibred product** of X and Y over $Y \times_S Y$, which can be seen as follows

$$\begin{array}{ccccccc} & & K & \xrightarrow{b} & Y & & \\ & \nearrow h' & \downarrow \iota & & \downarrow \Delta & & \\ Z & \xrightarrow{h} & X & \xrightarrow{(f, g)} & Y \times_S Y & \xrightarrow{p} & Y \\ & & & & \downarrow q & & \downarrow \\ & & & & Y & \longrightarrow & S. \end{array}$$

- (1) Prove that $\iota: K \rightarrow X$ is an immersion and $f \circ \iota = g \circ \iota$.
- (2) Let $h: Z \rightarrow X$ be a morphism such that $f \circ h = g \circ h$. Prove that there is a unique morphism $h': Z \rightarrow K$ such that $\iota \circ h' = h$.

Proof. For (1), If $\alpha, \beta: Z \rightarrow K$ are morphisms such that $\iota \circ \alpha = \iota \circ \beta$, then one also has $\Delta \circ b \circ \alpha = \Delta \circ b \circ \beta$. Then by the universal property of K one can see morphism from $Z \rightarrow K$ with such property must be unique, and thus $\alpha = \beta$. This shows ι is a monomorphism. For the half part, note that

$$\begin{aligned} p \circ (f, g) \circ \iota &= f \circ \iota \\ q \circ (f, g) \circ \iota &= g \circ \iota. \end{aligned}$$

On the other hand, $(f, g) \circ \iota = \Delta \circ b$ and $p \circ \Delta = q \circ \Delta$.

For (2), Since $h: Z \rightarrow X$ satisfies $f \circ h = g \circ h$, one has $p \circ (f, g) \circ h = q \circ (f, g) \circ h$, which gives a morphism $Z \rightarrow Y$ satisfying desired commutative property, and thus by the universal property of K as a fibred product, there exists a morphism $h': Z \rightarrow K$ such that $\iota \circ h' = h$. $\square$

7.10. Homework-10.

7.10.1. Part I.

Exercise 7.10.1. Let X be an S -scheme, $S' \rightarrow S$ a morphism, $X' = X \times_S S'$ the base change of X , $\Delta: X \rightarrow X \times_S X$ and $\Delta': X' \rightarrow X' \times_{S'} X'$ the diagonal morphisms. Use the result in Exercise 7 on page 58 of the textbook to prove the following diagram is Cartesian:

$$\begin{array}{ccc} X' & \xrightarrow{\Delta'} & X' \times_{S'} X' \\ \downarrow & & \downarrow \\ X & \xrightarrow{\Delta} & X \times_S X \end{array}$$

Proof. By (3) of Exercise 7 on page 58, it suffices to show for every scheme Z , the following diagram is Cartesian

$$\begin{array}{ccc} \mathrm{Hom}(Z, X') & \longrightarrow & \mathrm{Hom}(Z, X' \times_{S'} X') \\ \downarrow & & \downarrow \\ \mathrm{Hom}(Z, X) & \longrightarrow & \mathrm{Hom}(Z, X \times_S X). \end{array}$$

On the other hand, one has

$$\begin{aligned} \mathrm{Hom}(Z, X \times_S X) &= \mathrm{Hom}(Z, X) \times_{\mathrm{Hom}(Z, S)} \mathrm{Hom}(Z, X) \\ \mathrm{Hom}(Z, X' \times_{S'} X') &= \mathrm{Hom}(Z, X') \times_{\mathrm{Hom}(Z, S')} \mathrm{Hom}(Z, X'). \end{aligned}$$

Thus it suffices to show the following diagram is Cartesian

$$\begin{array}{ccc} \mathrm{Hom}(Z, X') & \longrightarrow & \mathrm{Hom}(Z, X') \times_{\mathrm{Hom}(Z, S')} \mathrm{Hom}(Z, X') \\ \downarrow & & \downarrow \\ \mathrm{Hom}(Z, X) & \longrightarrow & \mathrm{Hom}(Z, X) \times_{\mathrm{Hom}(Z, S)} \mathrm{Hom}(Z, X), \end{array}$$

which is clear by the description of fibred product in the category of sets. $\square$

Exercise 7.10.2. Let X and S be locally compact topological spaces, S is Hausdorff, and let $f: X \rightarrow S$ be a continuous map.

- (1) Prove that a proper map is a closed map.
- (2) Let S' be a locally compact topological space and let $g: S' \rightarrow S$ be a continuous map. For any proper map $f: X \rightarrow S$, prove the base change $f': X \times_S S' \rightarrow S'$ of f is proper.

Proof. For (1), Let $V \subseteq X$ be a closed subset. It suffices to show $S \setminus f(V)$ is open. For $s \in S \setminus f(V)$, there exists an open neighborhood of U of s with compact closure since S is locally compact. Then $f^{-1}(U)$ is compact since f is proper. Let $E = V \cap f^{-1}(U)$. Then E is compact since it is a closed subset of a compact set, and hence $f(E)$. Again by S is Hausdorff, one has $f(E)$ is closed in S . Then $U \setminus f(E)$ is an open neighborhood of s which is disjoint from $f(V)$. This shows $S \setminus f(V)$ is open.

For (2), Note that the fibred product in the category of topological spaces can be described as $X \times_S S' = \{(x, y) \in X \times S' \mid f(x) = g(y)\}$ and $f': X \times_S S' \rightarrow S'$ is given by $(x, y) \mapsto y$. Then for any compact subset $K \subseteq S'$, one has

$$(f')^{-1}(K) \subseteq f^{-1}g(K) \times K,$$

which is a compact subset since f is proper. $\square$

7.10.2. Part II.

Exercise 7.10.3. The set

$$I = \{(i_0, \dots, i_n) \mid i_0 + \dots + i_n = d, i_0, \dots, i_n \in \mathbb{Z}_{\geq 0}\}$$

has $\binom{d+n}{n}$ solutions. Prove that the homomorphism

$$\begin{aligned} \varphi: \mathbb{Z}[y_{i_0 \dots i_n}]_{(i_0, \dots, i_n) \in I} &\rightarrow \mathbb{Z}[x_0, \dots, x_n] \\ y_{i_0 \dots i_n} &\mapsto x_0^{i_0} \cdots x_n^{i_n} \end{aligned}$$

induces a morphism $\mathbb{P}_{\mathbb{Z}}^n \rightarrow \mathbb{P}_{\mathbb{Z}}^{\binom{d+n}{n}-1}$.

Proof. For convenience, we use $A = \bigoplus_{n \in \mathbb{Z}_{\geq 0}} A_n$ to denote the graded ring $\mathbb{Z}[y_{i_0 \dots i_n}]_{(i_0, \dots, i_n) \in I}$. Note that for each degree n , one has $\varphi(A_n) \subseteq \mathbb{Z}[x_0, \dots, x_n]_{nd}$. This shows ring homomorphism φ preserves the grade, and thus it induces a morphism between Proj. $\square$

7.11. Homework-11.

7.11.1. Part I.

Exercise 7.11.1. Let X be a topological space, $\mathcal{B}$ a basis of topology for X , $\mathcal{F}$ and $\mathcal{G}$ sheaves of abelian groups on X . Suppose for every member U in $\mathcal{B}$, we are provided with a homomorphism $\phi(U): \mathcal{F}(U) \rightarrow \mathcal{G}(U)$ such that for every pair $V \subseteq U$ in $\mathcal{B}$, the following diagram commutes,

$$\begin{array}{ccc} \mathcal{F}(U) & \xrightarrow{\phi(U)} & \mathcal{G}(U) \\ \downarrow & & \downarrow \\ \mathcal{F}(V) & \xrightarrow{\phi(V)} & \mathcal{G}(V) \end{array}$$

Prove that there exists a unique morphism of sheaves $\phi: \mathcal{F} \rightarrow \mathcal{G}$ such that for any member U in $\mathcal{B}$, one has $\phi(U): \mathcal{F}(U) \rightarrow \mathcal{G}(U)$ coincides with $\phi(U)$.

Proof. For convenience we denote $\mathcal{B} = \{U_\alpha\}_{\alpha \in A}$ and use ϕ_α to denote $\phi(U_\alpha)$. For any open subset $U \subseteq X$, we write it as $U = \bigcup_\alpha U_\alpha$, since $\mathcal{B}$ is a basis for topology. Then for any $x \in U$, if $x \in U_\alpha$, then we define $\phi(U)(x) = \phi_\alpha(x)$. This is well-defined since if $x \in U_\alpha \cap U_\beta$, then

$$\phi_\alpha(x) = \phi(U_\alpha \cap U_\beta)(x) = \phi_\beta(x).$$

This gives a unique morphism of sheaves $\phi: \mathcal{F} \rightarrow \mathcal{G}$ such that for any $U \in \mathcal{B}$, one has $\phi(U): \mathcal{F}(U) \rightarrow \mathcal{G}(U)$ coincides with $\phi(U)$. $\square$

7.11.2. Part II.

Exercise 7.11.2 (coherent sheaf on ringed space). Let $(X, \mathcal{O}_X)$ be a ringed space. An $\mathcal{O}_X$ -module $\mathcal{F}$ is called **coherent** if $\mathcal{F}$ is of finite type and for every open subset U of X and every homomorphism $u: \mathcal{O}_U^{\oplus n} \rightarrow \mathcal{F}|_U$, the **kernel** of u is of finite type.

- (1) Suppose $\mathcal{O}_X$ is coherent. Prove that an $\mathcal{O}_X$ -module $\mathcal{F}$ is coherent if and only if $\mathcal{F}$ is of finite presentation.
- (2) Prove that this definition of coherence coincides with the one in Definition 6.2.3 for Noetherian schemes.

Proof. For (1), First, we show if $\mathcal{F}$ is coherent, then $\mathcal{F}$ is of finite presentation (To prove this, we do not need the assumption $\mathcal{O}_X$ is coherent). Since $\mathcal{F}$ is of finite type, there exists an open covering $\{U_i\}$ of X such that

$$\mathcal{O}_{U_i}^{\oplus n_i} \xrightarrow{u_i} \mathcal{F}|_{U_i} \rightarrow 0$$

is exact, and the $\ker u_i$ is of finite type implies that there exists a refinement of $\{U_i\}$, denoted by $\{U_{ij}\}$ such that the following sequence is exact

$$\mathcal{O}_{U_{ij}}^{\oplus m_{ij}} \rightarrow \mathcal{O}_{U_{ij}}^{\oplus n_i} \rightarrow \mathcal{F}|_{U_{ij}} \rightarrow 0.$$

This shows $\mathcal{F}$ is of finite presentation. For the converse statement, firstly let us prove the following lemma.

Lemma 7.1. An $\mathcal{O}_X$ -module $\mathcal{F}$ is coherent if and only if there exists an open covering $\{U_i\}$ of X such that $\mathcal{F}|_{U_i}$ is coherent.

Proof. It follows that if $\mathcal{F}$ is coherent, then every $\mathcal{F}|_{U_i}$ is coherent. Conversely, suppose there exists an open covering $\{U_i\}$ of X such that $\mathcal{F}|_{U_i}$ is coherent. Firstly, since each $\mathcal{F}|_{U_i}$ is of finite type, there exists a refinement $\{U_{ij}\}$ of $\{U_i\}$ such that the following sequences are exact

$$\mathcal{O}|_{U_{ij}}^{\oplus n_{ij}} \rightarrow \mathcal{F}|_{U_{ij}} \rightarrow 0,$$

and thus $\mathcal{F}$ is of finite type. For every homomorphism $u: \mathcal{O}_U^{\oplus n} \rightarrow \mathcal{F}|_U$, its restriction gives the following homomorphism

$$u_i: \mathcal{O}_{U \cap U_i}^{\oplus n} \rightarrow \mathcal{F}|_{U \cap U_i},$$

and the $\ker u_i$ is of finite type since $\mathcal{F}|_{U \cap U_i}$ is coherent, and thus by the previous argument one has $\ker u$ is of finite type. $\square$

Now suppose $\mathcal{O}_X$ is coherent and $\mathcal{F}$ is of finite representation. Then there exists an open covering $\{U_i\}$ such that the following sequence

$$\mathcal{O}_{U_i}^{\oplus m_i} \rightarrow \mathcal{O}_{U_i}^{\oplus n_i} \rightarrow \mathcal{F}|_{U_i} \rightarrow 0$$

is exact, and thus $\mathcal{F}|_{U_i}$ is coherent, since the quotient of coherent sheaves is coherent³. Then by Lemma 7.1 one has $\mathcal{F}$ is coherent.

For (2), Suppose X is a Noetherian scheme. In this case, $\mathcal{O}_X$ is coherent, and thus it suffices to show the definition of coherence for Noetherian schemes in Definition 6.2.3 is equivalent to finite presentation.

(a) If $\mathcal{F}$ is of finite presentation, then without loss of generality we may assume there exists an affine open covering $\{U_i = \text{Spec } A_i\}$ such that

$$\mathcal{O}_{U_i}^{\oplus m_i} \rightarrow \mathcal{O}_{U_i}^{\oplus n_i} \rightarrow \mathcal{F}|_{U_i} \rightarrow 0$$

is exact. Note that $\mathcal{O}_{U_i}^{\oplus n_i}$ itself is of the form $M_i^\sim$, where $M_i = A_i^{\oplus n_i}$. This shows $\mathcal{F}$ is coherent (in the sense of Definition 6.2.3).

(b) If $\mathcal{F}$ is coherent (in the sense of Definition 6.2.3), then there exists an affine open covering $\{U_i = \text{Spec } A_i\}$ such that

$$M_i^\sim \xrightarrow{u_i} \mathcal{F}|_{U_i} \rightarrow 0$$

is exact, where M_i is a finitely generated A_i -module. Since X is Noetherian, one has A_i is a Noetherian ring, and thus M_i is also a Noetherian module since it is finitely generated. In particular, any submodule of M_i is finitely generated, and thus $\ker u_i$ is of finite type. $\square$

³However, it is also highly non-trivial with the definition of coherence given here, but I do not want to give a proof here. See Lemma 17.12.4 in [Sta23].

Exercise 7.11.3. Let X and Y be S -schemes, let $f, g: X \rightarrow Y$ be S -morphisms, and let U be a dense open subset of X such that $f|_U = g|_U$ as morphisms of schemes. Suppose Y is separated over S . Then $f = g$ as maps on topological spaces. Suppose furthermore that X is reduced. Then $f = g$ as morphisms of schemes.

Proof. Consider the following diagram

$$\begin{array}{ccc} X \times_{Y \times_S Y} Y & \longrightarrow & Y \\ \downarrow & & \downarrow \Delta_{Y/S} \\ X & \xrightarrow{h=f \times_S g} & Y \times_S Y. \end{array}$$

If Y is separated, then $\Delta_{Y/S}(Y) \subseteq Y \times_S Y$ is a closed subset, and thus $h^{-1}(\Delta_{Y/S}(Y))$ is also a closed subset since h is continuous. On the other hand, suppose $U \subseteq X$ is the dense open subset such that $f|_U = g|_U$. Then $U \subseteq h^{-1}(\Delta_{Y/S}(Y))$, and thus one has $h^{-1}(\Delta_{Y/S}(Y)) = X$ by taking closure of U . As a consequence, one has $f = g$ on X as maps on topological spaces.

Now it suffices to show $f^\sharp, g^\sharp$ induce the same morphisms between sheaves, and thus without loss of generality we may assume both $X = \operatorname{Spec} A$ and $Y = \operatorname{Spec} B$ are affine schemes. For convenience we write $f^\sharp: \mathcal{O}_{\operatorname{Spec} A} \rightarrow f^{-1}\mathcal{O}_{\operatorname{Spec} B}$ by using adjoint between f_* and f^{-1} . If $\ker(f^\sharp - g^\sharp) \subseteq \operatorname{Spec} A$ is of the form $V(\mathfrak{b})$, then one has $U \subseteq V(\mathfrak{b})$, and thus $V(\mathfrak{b}) = \operatorname{Spec} A$, since U is dense. As a consequence, $\mathfrak{b}$ is the radical of A , and thus $\mathfrak{b} = 0$ since A is reduced. $\square$

7.12. Homework-12.

7.12.1. Part I.

Exercise 7.12.1. 1. Let $f: X \rightarrow Y$ be a continuous map between topological spaces, let $\mathcal{F}$ and $\mathcal{G}$ be sheaves of abelian groups on X and Y , respectively, let $\delta: \mathcal{G} \rightarrow f_*\mathcal{F}$ be a morphism, and let $\psi: f^{-1}\mathcal{G} \rightarrow \mathcal{F}$ be the morphism induced by δ by adjunction. For any $x \in X$, and $\delta_x: \mathcal{G}_{f(x)} \rightarrow \mathcal{F}_x$ be the composite

$$\begin{aligned} \mathcal{G}_{f(x)} \xrightarrow{\delta_{f(x)}} (f_*\mathcal{F})_{f(x)} &= \varinjlim_{f(x) \in V} (f_*\mathcal{F})(V) \\ &= \varinjlim_{x \in f^{-1}(V)} \mathcal{F}(f^{-1}(V)) \rightarrow \varinjlim_{x \in U} \mathcal{F}(U) = \mathcal{F}_x, \end{aligned}$$

and let $\psi_x: \mathcal{G}_{f(x)} \rightarrow \mathcal{F}_x$ be the composite

$$\mathcal{G}_{f(x)} \cong (f^{-1}\mathcal{G})_x \xrightarrow{\psi_x} \mathcal{F}_x.$$

Prove that $\delta_x = \psi_x$.

Proof. For open subset $U \subseteq X$, recall that the adjoint morphism $\psi(U)$ is given by the following composite

$$f^{-1}\mathcal{G}(U) \xrightarrow{f^{-1}\delta(U)} f^{-1}f_*\mathcal{F}(U) \rightarrow \mathcal{F}(U).$$

By passing to stalks one has the following commutative diagram

$$\begin{array}{ccccc} \mathcal{G}_{f(x)} & \xrightarrow{\delta_{f(x)}} & (f_*\mathcal{F})_{f(x)} & \longrightarrow & \mathcal{F}_x \\ \cong \downarrow & & \downarrow \cong & & \downarrow \cong \\ (f^{-1}\mathcal{G})_x & \xrightarrow{(f^{-1}\delta)_x} & (f^{-1}f_*\mathcal{F})_x & \longrightarrow & \mathcal{F}_x \end{array}$$

This completes the proof. $\square$

Exercise 7.12.2. Let $(f, f^\sharp): (X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$ be a morphism of locally ringed spaces, $\mathcal{F}$ an $\mathcal{O}_X$ -module, $\mathcal{G}$ an $\mathcal{O}_Y$ -module, $\delta: \mathcal{G} \rightarrow f_*\mathcal{F}$ a homomorphism of $\mathcal{O}_Y$ -modules, and $\psi: f^*\mathcal{G} \rightarrow \mathcal{F}$ the morphism induced by δ by adjunction. For any $x \in X$, and $\delta_x: \mathcal{G}_{f(x)} \rightarrow \mathcal{F}_x$ be defined as in above exercise, and let $\psi_x: \mathcal{O}_{X,x} \otimes_{\mathcal{O}_{Y,f(x)}} \mathcal{G}_{f(x)} \rightarrow \mathcal{F}_x$ be the composite

$$\mathcal{O}_{X,x} \otimes_{\mathcal{O}_{Y,f(x)}} \mathcal{G}_{f(x)} \cong (f^*\mathcal{G})_x \xrightarrow{\psi_x} \mathcal{F}_x.$$

Prove that

$$\psi_x(r \otimes s) = r\delta_x(s)$$

for any $r \in \mathcal{O}_{X,x}$ and $s \in \mathcal{G}_{f(x)}$.

Proof. For open subset $U \subseteq X$, recall that the adjoint morphism $\psi(U)$ is given by the following composite

$$f^*\mathcal{G}(U) = (\mathcal{O}_X \otimes_{f^{-1}\mathcal{O}_Y} f^{-1}\mathcal{G})(U) \xrightarrow{\text{id} \otimes f^{-1}\delta(U)} (\mathcal{O}_X \otimes_{f^{-1}\mathcal{O}_Y} f^{-1}f_*\mathcal{F})(U) \rightarrow \mathcal{F}(U).$$

Similarly, by passing to stalks one has the following commutative diagram

$$\begin{array}{ccccc}
\mathcal{O}_{X,x} \otimes_{\mathcal{O}_{Y,f(x)}} \mathcal{G}_{f(x)} & \xrightarrow{\text{id} \otimes \delta_{f(x)}} & \mathcal{O}_{X,x} \otimes_{\mathcal{O}_{Y,f(x)}} (f_* \mathcal{F})_{f(x)} & \longrightarrow & \mathcal{F}_x \\
\downarrow \cong & & \downarrow \cong & & \downarrow \cong \\
(f^* \mathcal{G})_x & \xrightarrow{(f^* \delta)_x} & (f^* f_* \mathcal{F})_x & \longrightarrow & \mathcal{F}_x
\end{array}$$

This completes the proof. $\square$

Exercise 7.12.3. Let $(X, \mathcal{O}_X)$ be a scheme and $\mathcal{F}$ a quasi-coherent $\mathcal{O}_X$ -module. Suppose $\mathcal{F}$ is of finite presentation. Prove that for any affine open subset $U \cong \text{Spec } A$, we have $\mathcal{F}|_U \cong M^\sim$ for some A -module M with finite presentation.

Proof. Firstly, for any affine open subset $U \cong \text{Spec } A$, we may assume $\mathcal{F}|_U \cong M^\sim$ for some A -module M , so it suffices to show M is of finite presentation. Since $\mathcal{F}$ is of finite presentation, there exists an open covering of U , without loss of generality we may assume it is given by $\{U_i = \text{Spec } A_{f_i}\}_{i=1}^n$ with $(f_1, \dots, f_n) = A$, such that on each U_f one has the following exact sequence

$$\mathcal{O}_{U_i}^{\oplus m_i} \rightarrow \mathcal{O}_{U_i}^{\oplus n_i} \rightarrow \mathcal{F}|_{U_i} \rightarrow 0,$$

where $m_i, n_i \in \mathbb{Z}_{>0}$. In other words, the localization M_{f_i} is of finite presentation for each $i = 1, 2, \dots, n$, and thus M is of finite presentation. $\square$

7.12.2. Part II.

Exercise 7.12.4. Let X be a scheme, $\mathcal{L}$ be an invertible $\mathcal{O}_X$ -module with $f \in \mathcal{L}(X)$ and $\mathcal{F}$ be a quasi-coherent $\mathcal{O}_X$ -module.

- (1) Define X_f to be the subset of X consisting of those points $x \in X$ such that the germ of f at x does not lie in $\mathfrak{m}_x \mathcal{L}_x$, where $\mathfrak{m}_x$ is the maximal ideal of the local ring $\mathcal{O}_{X,x}$. Then X_f is open.
- (2) Suppose X is quasi-compact. If $s \in \mathcal{F}(X)$ is a section whose restriction to X_f vanishes, then there exists a natural number n such that the section $s \otimes f^{\otimes n} \in (\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{L}^{\otimes n})(X)$ vanishes.
- (3) Suppose X is quasi-compact and quasi-separated. Given a section t in $\mathcal{F}(X_f)$, there exists a natural number n such that the section $t \otimes f^{\otimes n}$ in $(\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{L}^{\otimes n})(X_f)$ can be extended to a section in $(\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{L}^{\otimes n})(X)$.

Proof. For (1), Since $\mathcal{L}$ is an invertible $\mathcal{O}_X$ -module, there exists an open covering $\{U_i\}$ of X such that $\mathcal{L}|_{U_i} \cong \mathcal{O}_{U_i}$. If we use f_i to denote $f|_{U_i}$ and use X_{f_i} to denote the subset of U_i consisting of those points $x \in U_i$ such that the germ of f_i at x does not lie in $\mathfrak{m}_x \mathcal{L}_x$, then $X_f = \bigcup_i X_{f_i}$, and by (1) of Proposition 4.3.1 one has each X_{f_i} is open.

For (2), Since X is quasi-compact, we may assume $\{U_i = \text{Spec } A_i\}_{i=1}^n$ is a finite affine open covering of X , and thus $X_f \cap \text{Spec } A_i = D(f_i)$, where $f_i := f|_{U_i}$. Since the restriction of s to X_f vanishes, one has the restriction of s to each $D(f_i) = \text{Spec}(A_i)_{f_i}$ vanishes for each $i = 1, \dots, n$, and thus there exists some $n_i \in \mathbb{Z}_{>0}$ such that $f_i^{n_i} s = 0$ in A_i . Now it suffices to take $n = \max_i \{n_i\}$, one has $s \otimes f^{\otimes n} \in (\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{L}^{\otimes n})(X)$ vanishes.

For (3), Since X is quasi-compact and quasi-separated, by (2) of Proposition 5.5.4 we may assume $\{U_i = \text{Spec} A_i\}_{i=1}^n$ is a finite affine open covering of X such that each $U_i \cap U_j$ is a union of finitely many affine open subschemes. For convenience we denote $t_i := t|_{U_i}$. Note that $t_i - t_j$ vanishes on $U_i \cap U_j$, and since $U_i \cap U_j$ is a union of finite affine open subschemes, then by (2) there exists some $n_{ij} \in \mathbb{N}$ such that $(t_i - t_j) \otimes f_{ij}^{n_{ij}} \in (\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{L}^{\otimes n_{ij}})(U_i \cap U_j)$ vanishes. On the other hand, for each $t_i \in \mathcal{F}(X_f \cap U_i)$, there exists some $m_i \in \mathbb{N}$ such that $t_i \otimes f^{m_i}$ extends to $(\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{L}^{\otimes m_i})(X_f \cap U_i)$. Since everything is finite, there exists a sufficiently large n such that $t \otimes f^n$ extends to a section in $(\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{L}^{\otimes n})(X)$. $\square$

7.13. Homework-13.

Exercise 7.13.1. Let S be a graded ring which is generated by S_1 as an S_0 -algebra.

- (1) Prove that for every $f \in S_1$ and every graded S -module M , we have an $S_{(f)}$ -module isomorphism

$$\bigoplus_{n=-\infty}^{\infty} M(n)_{(f)} \cong M_f.$$

In particular, taking $M = S$, we get an $S_{(f)}$ -module isomorphism

$$\bigoplus_{n=-\infty}^{\infty} S(n)_{(f)} \xrightarrow{\cong} S_f.$$

Prove that we have an S_f -module isomorphism

$$M_{(f)} \otimes_{S_{(f)}} S_f \cong M_f$$

- (2) Let M and N be graded S -modules. Construct an isomorphism

$$M_{(f)} \otimes_{S_{(f)}} N_{(f)} \xrightarrow{\cong} (M \otimes_S N)_{(f)},$$

so that after taking the tensor product $\otimes_{S_f} S_f$ and composing with the isomorphisms in (1), we get the canonical isomorphism

$$M_f \otimes_{S_f} N_f \cong (M \otimes_S N)_f.$$

- (3) Let $\mathfrak{p} \in \text{Proj } S$, and let T be the set of homogeneous elements in $S \setminus \mathfrak{p}$. Prove that we have an $S_{(\mathfrak{p})}$ -module isomorphism

$$\bigoplus_{n=-\infty}^{\infty} M(n)_{(\mathfrak{p})} \xrightarrow{\cong} T^{-1}M.$$

In particular, taking $M = S$, we get an $S_{(\mathfrak{p})}$ -module isomorphism

$$\bigoplus_{n=-\infty}^{\infty} S(n)_{(\mathfrak{p})} \xrightarrow{\cong} T^{-1}S.$$

Prove that we have a $T^{-1}S$ -module isomorphism

$$M_{(\mathfrak{p})} \otimes_{S_{(\mathfrak{p})}} T^{-1}S \xrightarrow{\cong} T^{-1}M.$$

- (4) Let M and N be graded S -modules. Construct an isomorphism

$$M_{(\mathfrak{p})} \otimes_{S_{(\mathfrak{p})}} N_{(\mathfrak{p})} \xrightarrow{\cong} (M \otimes_S N)_{(\mathfrak{p})}$$

so that after taking the tensor product $\otimes_{T^{-1}S} T^{-1}S$ and composing with the isomorphisms in (3), we get the canonical isomorphism

$$T^{-1}M \otimes_{T^{-1}S} T^{-1}N \xrightarrow{\cong} T^{-1}(M \otimes_S N).$$

Proof. For (1), Consider the following homomorphism

$$\begin{aligned} \phi: \bigoplus_{n=-\infty}^{\infty} M(n)_{(f)} &\rightarrow M_f \\ \left(\frac{m_n}{f^{k_n}}\right) &\mapsto \sum_{n=-\infty}^{\infty} \frac{m_n}{f^{k_n}}. \end{aligned}$$

It is well-defined since there are only finitely many $m_n/f^{k_n} \neq 0$.

(a) First, we show ϕ surjective: Since for any $m/f^k \in M_f$, we may write $m = \sum_i m_i$, where m_i are homogeneous elements of degree i . If $i = k + n$, then $m_i/f^k \in M(n)_{(f)}$, and thus

$$\phi\left(\frac{m_i}{f^k}\right) = \sum_i \frac{m_i}{f^k} = \frac{m}{f^k}.$$

(b) We now prove ϕ is injective: If $\sum_{n=-\infty}^{\infty} m_n/f^{k_n} = 0$ in M_f , then there exists some $N \in \mathbb{N}$ such that

$$f^N \sum_{n=-\infty}^{\infty} \frac{m_n}{f^{k_n}} = 0.$$

Since $\deg m_n = k_n + n$, then by degree reason $f^N m_n/f^{k_n} = 0$ for every $n \in \mathbb{Z}$. On the other hand, multiplying by f^N gives an isomorphism from $M(n)_{(f)}$ to $M(n+N)_{(f)}$ for each $n \in \mathbb{Z}$ since $f \in S_1$. This shows $m_n/f^{k_n} = 0$ for each $n \in \mathbb{Z}$, and thus ϕ is injective. In particular, one has

$$\bigoplus_{n=-\infty}^{\infty} S(n)_{(f)} \xrightarrow{\cong} S_f.$$

Moreover, since $M(n)_{(f)} \cong M_{(f)} \otimes_{S_{(f)}} S(n)_{(f)}$ as $S_{(f)}$ -modules, one has

$$\begin{aligned} M_f &\cong \bigoplus_{n=-\infty}^{\infty} M(n)_{(f)} \\ &\cong \bigoplus_{n=-\infty}^{\infty} (M_{(f)} \otimes_{S_{(f)}} S(n)_{(f)}) \\ &\cong M_{(f)} \otimes_{S_{(f)}} S_f. \end{aligned}$$

For (2), Consider the following homomorphism

$$\begin{aligned} \phi: M_{(f)} \otimes_{S_{(f)}} N_{(f)} &\rightarrow (M \otimes_S N)_{(f)} \\ \sum_i \frac{m_i}{f^{k_i}} \otimes \frac{n_i}{f^{l_i}} &\mapsto \sum_i \frac{m_i \otimes n_i}{f^{k_i+l_i}}. \end{aligned}$$

It follows that ϕ is surjective, so it suffices to check it is injective. If

$$\sum_i \frac{m_i \otimes n_i}{f^{k_i+l_i}} = \frac{1}{f^N} \sum_i f^{\alpha_i} m_i \otimes n_i = 0$$

in $(M \otimes_S N)_{(f)}$, where $N = \max_i \{k_i + l_i\}$ and $k_i + l_i + \alpha_i = N$ for all i , then $\sum_i f^{\alpha_i} m_i \otimes n_i = 0$ in $M \otimes_S N$, and thus

$$\sum_i \frac{m_i}{f^{k_i}} \otimes \frac{n_i}{f^{l_i}} = 0$$

in $M_{(f)} \otimes_{S_{(f)}} N_{(f)}$. This shows the injectivity. As a consequence, after taking the tensor product $\otimes_{S_{(f)}} S_f$ and composing with the isomorphisms in (1), we get the canonical isomorphism

$$M_f \otimes_{S_f} N_f \cong (M \otimes_S N)_f.$$

For (3), Note that

$$\begin{aligned} \bigoplus_{n=-\infty}^{\infty} M(n)_{(p)} &\stackrel{(a)}{\cong} \bigoplus_{n=-\infty}^{\infty} \varinjlim_{p \in D_+(f)} (M(n)_{(f)})^\sim \\ &\cong \varinjlim_{p \in D_+(f)} \bigoplus_{n=-\infty}^{\infty} (M(n)_{(f)})^\sim \\ &\cong \varinjlim_{p \in D_+(f)} \left(\bigoplus_{n=-\infty}^{\infty} M(n)_{(f)} \right)^\sim \\ &\cong \varinjlim_{p \in D_+(f)} (M_f)^\sim \\ &\stackrel{(b)}{\cong} T^{-1}M, \end{aligned}$$

where (a) and (b) hold from S is generated by S_1 as an S_0 -algebra, and by the same argument as (1) one can show

$$M_{(p)} \otimes_{S_{(p)}} T^{-1}S \xrightarrow{\cong} T^{-1}M.$$

For (4), It follows from (2) by using the same modification as (3). $\square$

Exercise 7.13.2. Let X be a scheme, and Y be a closed subscheme of X with ideal sheaf $\mathcal{I}$. Define the blowing-up $\tilde{X}$ of X along Y to be the scheme $\text{Proj}(\bigoplus_{n=0}^{\infty} \mathcal{I}^n)$ over X , where $\text{Proj}(\bigoplus_{n=0}^{\infty} \mathcal{I}^n)$ is obtained by gluing $\text{Proj}(\bigoplus_{n=0}^{\infty} \mathcal{I}^n(U))$ over open affine subschemes U of X .

- (1) Let $\pi: \tilde{X} \rightarrow X$ be the canonical morphism. Prove that π induces an isomorphism $\pi^{-1}(X \setminus Y) \xrightarrow{\cong} X \setminus Y$.
- (2) Let $f: X' \rightarrow X$ be a closed immersion, and let $Y' = X' \times_X Y$ be the **closed subscheme** of X' obtained from $Y \rightarrow X$ by base change. Prove that the blowing-up $\tilde{X}'$ of X' along Y' is isomorphic to the **scheme theoretic image** of the composite

$$X' \setminus Y' \xrightarrow{f} X \setminus Y \xrightarrow{\pi^{-1}} \pi^{-1}(X \setminus Y) \hookrightarrow \tilde{X}$$

Proof. For (1), Suppose $\{U_\alpha = \text{Spec } A_\alpha\}$ is an affine open covering of X . If we write $I_\alpha = \mathcal{I}(U_\alpha)$, then $\pi: \tilde{X} \rightarrow X$ on the affine piece U_α is induced by the natural inclusion

$$A_\alpha \hookrightarrow S_\alpha = \bigoplus_{d \geq 0} I_\alpha^d.$$

By previous homework (Exercise 7.5.3), one has $\pi|_{\pi^{-1}(U_\alpha)}: \pi^{-1}(U_\alpha \setminus V(I_\alpha)) \rightarrow U_\alpha \setminus V(I_\alpha)$ is an isomorphism. This shows $\pi: \pi^{-1}(X \setminus Y) \rightarrow X \setminus Y$ is an isomorphism. $\square$

REFERENCES

- [Fu06] L. Fu. *Algebraic Geometry*. Mathematics series for graduate students. Tsinghua University Press, 2006.
- [Sta23] The Stacks project authors. The stacks project. <https://stacks.math.columbia.edu>, 2023.
- [Vak17] Ravi Vakil. The rising sea: Foundations of algebraic geometry. *preprint*, 2017.