

Uncertainty, consistency and hypothesis testing


Thomas Leavitt

July 22, 2022

Variance of Difference-in-Means

Variance of Difference-in-Means estimator

- Yesterday we showed that Difference-in-Means estimator in “village heads” example is



- What is the variance of this estimator?
- Why do we care about variance of an estimator?

Variance of Difference-in-Means estimator

- Variance is average squared distance of estimator from its expected value:

$$\underbrace{E \left[\underbrace{\left(\hat{\tau}(Z, Y) - \underbrace{E[\hat{\tau}(Z, Y)]}_{\text{Expected value}} \right)^2}_{\text{Squared distance from expected value}} \right]}_{\text{Average (or expected) squared distance}}$$

Variance of Difference-in-Means estimator

- Variance is average squared distance of estimator from its expected value:

$$\underbrace{\mathbb{E} \left[\underbrace{\left(\hat{\tau}(\mathbf{Z}, \mathbf{Y}) - \underbrace{\mathbb{E}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]}_{\text{Expected value}} \right)^2}_{\text{Squared distance from expected value}} \right]}_{\text{Average (or expected) squared distance}}$$

- Diff-in-Means unbiased for τ , so write variance as

$$\mathbb{E} [(\hat{\tau}(\mathbf{Z}, \mathbf{Y}) - \tau)^2]$$

Variance of Difference-in-Means estimator

- Variance is average squared distance of estimator from its expected value:

$$\text{E} \left[\underbrace{\left(\underbrace{\hat{\tau}(\mathbf{Z}, \mathbf{Y}) - \underbrace{\text{E}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]}_{\text{Expected value}}}_{\text{Squared distance from expected value}} \right)^2}_{\text{Average (or expected) squared distance}} \right]$$

- Diff-in-Means unbiased for τ , so write variance as

$$\text{E} [(\hat{\tau}(\mathbf{Z}, \mathbf{Y}) - \tau)^2]$$

- In “village heads” example with 21 possible assignments

$$(\hat{\tau}(\mathbf{z}_1, \mathbf{y}_1) - \tau)^2 \Pr(\mathbf{Z} = \mathbf{z}_1) + \dots + (\hat{\tau}(\mathbf{z}_{21}, \mathbf{y}_{21}) - \tau)^2 \Pr(\mathbf{Z} = \mathbf{z}_{21})$$

Variance is ≈ 21.19

Variance of Difference-in-Means estimator

- Neyman (1923) derived expression for variance of Diff-in-Means under complete random assignment

$$\text{Var} [\hat{\tau} (\mathbf{Z}, \mathbf{Y})] = \frac{1}{N-1} \left(\frac{n_T \sigma_{y_C}^2}{n_C} + \frac{n_C \sigma_{y_T}^2}{n_T} + 2 \sigma_{y_C, y_T} \right)$$

where

$$\sigma_{y_C}^2 = \frac{1}{N} \sum_{i=1}^N \left(y_{Ci} - \frac{1}{N} \sum_{i=1}^N y_{Ci} \right)^2 \text{ is var of control POs}$$

$$\sigma_{y_T}^2 = \frac{1}{N} \sum_{i=1}^N \left(y_{Ti} - \frac{1}{N} \sum_{i=1}^N y_{Ti} \right)^2 \text{ is var of treated POs}$$

$$\sigma_{y_C, y_T} = \frac{1}{N} \sum_{i=1}^N \left(y_{Ci} - \frac{1}{N} \sum_{i=1}^N y_{Ci} \right) \left(y_{Ti} - \frac{1}{N} \sum_{i=1}^N y_{Ti} \right) \text{ is cov of POs}$$

Variance of Difference-in-Means estimator

- Neyman (1923) derived expression for variance of Diff-in-Means under complete random assignment

$$\text{Var} [\hat{\tau} (\mathbf{Z}, \mathbf{Y})] = \frac{1}{N-1} \left(\frac{n_T \sigma_{y_C}^2}{n_C} + \frac{n_C \sigma_{y_T}^2}{n_T} + 2\sigma_{y_C, y_T} \right)$$

where

$$\sigma_{y_C}^2 = \frac{1}{N} \sum_{i=1}^N \left(y_{Ci} - \frac{1}{N} \sum_{i=1}^N y_{Ci} \right)^2 \text{ is var of control POs}$$

$$\sigma_{y_T}^2 = \frac{1}{N} \sum_{i=1}^N \left(y_{Ti} - \frac{1}{N} \sum_{i=1}^N y_{Ti} \right)^2 \text{ is var of treated POs}$$

$$\sigma_{y_C, y_T} = \frac{1}{N} \sum_{i=1}^N \left(y_{Ci} - \frac{1}{N} \sum_{i=1}^N y_{Ci} \right) \left(y_{Ti} - \frac{1}{N} \sum_{i=1}^N y_{Ti} \right) \text{ is cov of POs}$$

★ Note that $\sigma_{y_C}^2$, $\sigma_{y_T}^2$ and σ_{y_C, y_T} depend on unknown potential outcomes

Variance of Difference-in-Means estimator

- Sometimes you might see equivalent expression (Imbens and Rubin 2015)

$$\frac{S_{y_C}}{n_C} + \frac{S_{y_T}}{n_T} - \frac{S_\tau}{N},$$

where

$$S_{y_C} = \frac{1}{N-1} \sum_{i=1}^N \left(y_{Ci} - \frac{1}{N} \sum_{i=1}^N y_{Ci} \right)^2$$

$$S_{y_T} = \frac{1}{N-1} \sum_{i=1}^N \left(y_{Ti} - \frac{1}{N} \sum_{i=1}^N y_{Ti} \right)^2$$

$$S_\tau = \frac{1}{N-1} \sum_{i=1}^N \left(\tau_i - \frac{1}{N} \sum_{i=1}^N \tau_i \right)^2$$

Variance of Difference-in-Means estimator

- Sometimes you might see equivalent expression (Imbens and Rubin 2015)

$$\frac{S_{y_C}}{n_C} + \frac{S_{y_T}}{n_T} - \frac{S_\tau}{N},$$

where

$$S_{y_C} = \frac{1}{N-1} \sum_{i=1}^N \left(y_{Ci} - \frac{1}{N} \sum_{i=1}^N y_{Ci} \right)^2$$

$$S_{y_T} = \frac{1}{N-1} \sum_{i=1}^N \left(y_{Ti} - \frac{1}{N} \sum_{i=1}^N y_{Ti} \right)^2$$

$$S_\tau = \frac{1}{N-1} \sum_{i=1}^N \left(\tau_i - \frac{1}{N} \sum_{i=1}^N \tau_i \right)^2$$

★ Note that S_{y_C} , S_{y_T} and S_τ depend on unknown potential outcomes

Variance of Difference-in-Means estimator

- Example: "Village heads" study (Gerber and Green 2012, Chapter 2):

Village	Budget share (%)		
	y_C	y_T	τ
1	10	15	5
2	15	15	0
3	20	30	10
4	20	15	-5
5	10	20	10
6	15	15	0
7	15	30	15
Average	15	20	5

Variance of Difference-in-Means estimator

- Example: “Village heads” study (Gerber and Green 2012, Chapter 2):

Village	Budget share (%)		
	y_C	y_T	τ
1	10	15	5
2	15	15	0
3	20	30	10
4	20	15	-5
5	10	20	10
6	15	15	0
7	15	30	15
Average	15	20	5

- With access to true POs, we can directly calculate $\text{Var} [\hat{\tau} (\mathbf{Z}, \mathbf{Y})]$:

$$\sigma_{y_C}^2 \approx 14.29, \sigma_{y_T}^2 \approx 42.86, \sigma_{y_C, y_T} \approx 7.14$$

- So, $\text{Var} [\hat{\tau} (\mathbf{Z}, \mathbf{Y})] = \frac{1}{N-1} \left(\frac{n_T \sigma_{y_C}^2}{n_C} + \frac{n_C \sigma_{y_T}^2}{n_T} + 2 \sigma_{y_C, y_T} \right) \approx 21.19$

Variance of Difference-in-Means estimator

- Example: “Village heads” study (Gerber and Green 2012, Chapter 2):

Village	Budget share (%)		
	y_C	y_T	τ
1	10	15	5
2	15	15	0
3	20	30	10
4	20	15	-5
5	10	20	10
6	15	15	0
7	15	30	15
Average	15	20	5

- With access to true POs, we can directly calculate $\text{Var}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]$:

$$\sigma_{y_C}^2 \approx 14.29, \sigma_{y_T}^2 \approx 42.86, \sigma_{y_C, y_T} \approx 7.14$$

- So, $\text{Var}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})] = \frac{1}{N-1} \left(\frac{n_T \sigma_{y_C}^2}{n_C} + \frac{n_C \sigma_{y_T}^2}{n_T} + 2\sigma_{y_C, y_T} \right) \approx 21.19$

- In practice, $\sigma_{y_C}^2$, $\sigma_{y_T}^2$ and σ_{y_C, y_T} unknown, so we estimate $\text{Var}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]$

Variance estimation

Variance estimation

We showed that the Difference-in-Means estimator's variance is

$$\text{Var} [\hat{\tau}(\mathbf{Z}, \mathbf{Y})] = \frac{1}{N-1} \left(\frac{n_T \sigma_{y_C}^2}{n_C} + \frac{n_C \sigma_{y_T}^2}{n_T} + 2\sigma_{y_C, y_T} \right)$$

- We have unbiased estimators for $\sigma_{y_C}^2$ and $\sigma_{y_T}^2$, but not σ_{y_C, y_T}
- So what do we do?

Variance estimation

We showed that the Difference-in-Means estimator's variance is

$$\text{Var} [\hat{\tau} (\mathbf{Z}, \mathbf{Y})] = \frac{1}{N-1} \left(\frac{n_T \sigma_{y_C}^2}{n_C} + \frac{n_C \sigma_{y_T}^2}{n_T} + 2 \sigma_{y_C, y_T} \right)$$

- We have unbiased estimators for $\sigma_{y_C}^2$ and $\sigma_{y_T}^2$, but not σ_{y_C, y_T}
- So what do we do?
- We use **conservative** “plug-in” estimator (Neyman 1923)
- **Conservative** means that

$$\mathbb{E} \left[\widehat{\text{Var}} [\hat{\tau} (\mathbf{Z}, \mathbf{Y})] \right] \geq \text{Var} [\hat{\tau} (\mathbf{Z}, \mathbf{Y})]$$

Variance estimation

We showed that the Difference-in-Means estimator's variance is

$$\text{Var} [\hat{\tau} (\mathbf{Z}, \mathbf{Y})] = \frac{1}{N-1} \left(\frac{n_T \sigma_{y_C}^2}{n_C} + \frac{n_C \sigma_{y_T}^2}{n_T} + 2 \sigma_{y_C, y_T} \right)$$

- We have unbiased estimators for $\sigma_{y_C}^2$ and $\sigma_{y_T}^2$, but not σ_{y_C, y_T}
- So what do we do?
- We use **conservative** “plug-in” estimator (Neyman 1923)
- **Conservative** means that

$$\mathbb{E} \left[\widehat{\text{Var}} [\hat{\tau} (\mathbf{Z}, \mathbf{Y})] \right] \geq \text{Var} [\hat{\tau} (\mathbf{Z}, \mathbf{Y})]$$

- When potential outcomes perfectly positively correlated

$$\mathbb{E} \left[\widehat{\text{Var}} [\hat{\tau} (\mathbf{Z}, \mathbf{Y})] \right] = \text{Var} [\hat{\tau} (\mathbf{Z}, \mathbf{Y})]$$

- Otherwise, $\widehat{\text{Var}} [\hat{\tau} (\mathbf{Z}, \mathbf{Y})]$ is positively biased (conservative)

Variance estimation

We showed that the Difference-in-Means estimator's variance is

$$\text{Var} [\hat{\tau} (\mathbf{Z}, \mathbf{Y})] = \frac{1}{N-1} \left(\frac{n_T \sigma_{y_C}^2}{n_C} + \frac{n_C \sigma_{y_T}^2}{n_T} + 2 \sigma_{y_C, y_T} \right)$$

- We have unbiased estimators for $\sigma_{y_C}^2$ and $\sigma_{y_T}^2$, but not σ_{y_C, y_T}
- So what do we do?
- We use **conservative** “plug-in” estimator (Neyman 1923)
- **Conservative** means that

$$\mathbb{E} \left[\widehat{\text{Var}} [\hat{\tau} (\mathbf{Z}, \mathbf{Y})] \right] \geq \text{Var} [\hat{\tau} (\mathbf{Z}, \mathbf{Y})]$$

- When potential outcomes perfectly positively correlated

$$\mathbb{E} \left[\widehat{\text{Var}} [\hat{\tau} (\mathbf{Z}, \mathbf{Y})] \right] = \text{Var} [\hat{\tau} (\mathbf{Z}, \mathbf{Y})]$$

- Otherwise, $\widehat{\text{Var}} [\hat{\tau} (\mathbf{Z}, \mathbf{Y})]$ is positively biased (conservative)
- Potential outcomes will be perfectly positively correlated if and only if τ_i is the same for all $i = 1, \dots, N$ units

Conservative variance estimator

$$\text{Var} [\hat{\tau}(\mathbf{Z}, \mathbf{Y})] = \frac{1}{N-1} \left(\frac{n_T \sigma_{y_C}^2}{n_C} + \frac{n_C \sigma_{y_T}^2}{n_T} + 2 \sigma_{y_C, y_T} \right)$$

Conservative variance estimator

$$\text{Var} [\hat{\tau}(\mathbf{Z}, \mathbf{Y})] = \frac{1}{N-1} \left(\frac{n_T \sigma_{y_C}^2}{n_C} + \frac{n_C \sigma_{y_T}^2}{n_T} + 2\sigma_{y_C, y_T} \right)$$

- The maximum possible value of $2\sigma_{y_C, y_T}$ is $\sigma_{y_C}^2 + \sigma_{y_T}^2$
- So substitute $\sigma_{y_C}^2 + \sigma_{y_T}^2$ for $2\sigma_{y_C, y_T}$:

Conservative variance estimator

$$\text{Var} [\hat{\tau}(\mathbf{Z}, \mathbf{Y})] = \frac{1}{N-1} \left(\frac{n_T \sigma_{y_C}^2}{n_C} + \frac{n_C \sigma_{y_T}^2}{n_T} + 2\sigma_{y_C, y_T} \right)$$

- The maximum possible value of $2\sigma_{y_C, y_T}$ is $\sigma_{y_C}^2 + \sigma_{y_T}^2$
- So substitute $\sigma_{y_C}^2 + \sigma_{y_T}^2$ for $2\sigma_{y_C, y_T}$:

$$\begin{aligned} \text{Var} [\hat{\tau}(\mathbf{Z}, \mathbf{Y})] &= \frac{1}{N-1} \left(\frac{n_T \sigma_{y_C}^2}{n_C} + \frac{n_C \sigma_{y_T}^2}{n_T} + (\sigma_{y_C}^2 + \sigma_{y_T}^2) \right) \\ &= \frac{N}{N-1} \left(\frac{\sigma_{y_C}^2}{n_C} + \frac{\sigma_{y_T}^2}{n_T} \right) \end{aligned}$$

Conservative variance estimator

$$\text{Var} [\hat{\tau}(\mathbf{Z}, \mathbf{Y})] = \frac{1}{N-1} \left(\frac{n_T \sigma_{y_C}^2}{n_C} + \frac{n_C \sigma_{y_T}^2}{n_T} + 2\sigma_{y_C, y_T} \right)$$

- The maximum possible value of $2\sigma_{y_C, y_T}$ is $\sigma_{y_C}^2 + \sigma_{y_T}^2$
- So substitute $\sigma_{y_C}^2 + \sigma_{y_T}^2$ for $2\sigma_{y_C, y_T}$:

$$\begin{aligned} \text{Var} [\hat{\tau}(\mathbf{Z}, \mathbf{Y})] &= \frac{1}{N-1} \left(\frac{n_T \sigma_{y_C}^2}{n_C} + \frac{n_C \sigma_{y_T}^2}{n_T} + (\sigma_{y_C}^2 + \sigma_{y_T}^2) \right) \\ &= \frac{N}{N-1} \left(\frac{\sigma_{y_C}^2}{n_C} + \frac{\sigma_{y_T}^2}{n_T} \right) \end{aligned}$$

- Now all quantities can be estimated!

Conservative variance estimator

$$\text{Var} [\hat{\tau}(\mathbf{Z}, \mathbf{Y})] = \frac{1}{N-1} \left(\frac{n_T \sigma_{y_C}^2}{n_C} + \frac{n_C \sigma_{y_T}^2}{n_T} + 2\sigma_{y_C, y_T} \right)$$

- The maximum possible value of $2\sigma_{y_C, y_T}$ is $\sigma_{y_C}^2 + \sigma_{y_T}^2$
- So substitute $\sigma_{y_C}^2 + \sigma_{y_T}^2$ for $2\sigma_{y_C, y_T}$:

$$\begin{aligned} \text{Var} [\hat{\tau}(\mathbf{Z}, \mathbf{Y})] &= \frac{1}{N-1} \left(\frac{n_T \sigma_{y_C}^2}{n_C} + \frac{n_C \sigma_{y_T}^2}{n_T} + (\sigma_{y_C}^2 + \sigma_{y_T}^2) \right) \\ &= \frac{N}{N-1} \left(\frac{\sigma_{y_C}^2}{n_C} + \frac{\sigma_{y_T}^2}{n_T} \right) \end{aligned}$$


- Now all quantities can be estimated!
- So, just “plug-in” estimators $\hat{\sigma}_{y_C}^2$ and $\hat{\sigma}_{y_T}^2$ for $\sigma_{y_C}^2$ and $\sigma_{y_T}^2$

$$\widehat{\text{Var}} [\hat{\tau}(\mathbf{Z}, \mathbf{Y})] = \frac{N}{N-1} \left(\frac{\hat{\sigma}_{y_C}^2}{n_C} + \frac{\hat{\sigma}_{y_T}^2}{n_T} \right) \quad (1)$$

- For exact expressions of $\hat{\sigma}_{y_C}^2$ and $\hat{\sigma}_{y_T}^2$, see Variance estimators

Variance estimation

- Here is the conservative variance estimator in “village heads” example:



Solid line is true variance of Difference-in-Means estimator, $\text{Var} [\hat{\tau} (\mathbf{Z}, \mathbf{Y})]$

Dashed line is expected value of conservative estimator, $E \left[\widehat{\text{Var}} [\hat{\tau} (\mathbf{Z}, \mathbf{Y})] \right]$

Asymptotic properties

- So far, we have derived
 1. unbiased Difference-in-Means estimator of ATE
 2. variance of Difference-in-Means estimator
 3. conservative estimator of Difference-in-Means estimator's variance

Asymptotics

- So far, we have derived
 1. unbiased Difference-in-Means estimator of ATE
 2. variance of Difference-in-Means estimator
 3. conservative estimator of Difference-in-Means estimator's variance
- All of these derivations were for a fixed N , either small or large
- Now let's see what happens to our estimator when N grows large, $N \rightarrow \infty$
- But first, why do we care?

Asymptotics

- So far, we have derived
 1. unbiased Difference-in-Means estimator of ATE
 2. variance of Difference-in-Means estimator
 3. conservative estimator of Difference-in-Means estimator's variance
- All of these derivations were for a fixed N , either small or large
- Now let's see what happens to our estimator when N grows large, $N \rightarrow \infty$
- But first, why do we care?

N never goes to ∞ in an actual experiment

But properties as $N \rightarrow \infty$ **approximate** properties with fixed, but large N

Consistent estimation

Consistent estimation

- Difference-in-Means estimator is consistent:

- Difference-in-Means estimator is consistent:

$$\lim_{N \rightarrow \infty} \Pr (|\hat{\tau}(\mathbf{Z}, \mathbf{Y}) - \tau| < \epsilon) = 1 \text{ for all } \epsilon > 0$$

Consistent estimation

- Difference-in-Means estimator is consistent:

$$\lim_{N \rightarrow \infty} \Pr(|\hat{\tau}(\mathbf{Z}, \mathbf{Y}) - \tau| < \epsilon) = 1 \text{ for all } \epsilon > 0$$

- In words:

Pick any ϵ you want, no matter how small. There will be some value N^* such that, if size of experiment is greater than N^* , the probability of an estimate within a distance of ϵ from the truth is equal to 1.

Consistent estimation

- Difference-in-Means estimator is consistent:

$$\lim_{N \rightarrow \infty} \Pr (|\hat{\tau}(\mathbf{Z}, \mathbf{Y}) - \tau| < \epsilon) = 1 \text{ for all } \epsilon > 0$$


- In words:

Pick any ϵ you want, no matter how small. There will be some value N^* such that, if size of experiment is greater than N^* , the probability of an estimate within a distance of ϵ from the truth is equal to 1.

- Intuitively, with large experiment, our estimate will be close to true ATE!

Consistent estimation

- “Village heads” example:



Hypothesis testing

Hypothesis tests of the weak null

- The finite population CLT tells us that

$$\frac{\hat{\tau}(\mathbf{Z}, \mathbf{Y}) - \mathbb{E}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]}{\sqrt{\text{Var}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]}} \xrightarrow{d} \mathcal{N}(0, 1)$$

Hypothesis tests of the weak null

- The finite population CLT tells us that

$$\frac{\hat{\tau}(\mathbf{Z}, \mathbf{Y}) - \mathbb{E}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]}{\sqrt{\text{Var}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]}} \xrightarrow{d} \mathcal{N}(0, 1)$$

- Diff-in-Means is unbiased, so write

$$\frac{\hat{\tau}(\mathbf{Z}, \mathbf{Y}) - \tau}{\sqrt{\text{Var}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]}} \xrightarrow{d} \mathcal{N}(0, 1)$$

Hypothesis tests of the weak null

- The finite population CLT tells us that

$$\frac{\hat{\tau}(\mathbf{Z}, \mathbf{Y}) - \mathbb{E}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]}{\sqrt{\text{Var}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]}} \xrightarrow{d} \mathcal{N}(0, 1)$$

- Diff-in-Means is unbiased, so write

$$\frac{\hat{\tau}(\mathbf{Z}, \mathbf{Y}) - \tau}{\sqrt{\text{Var}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]}} \xrightarrow{d} \mathcal{N}(0, 1)$$

- The CLT is an asymptotic results as $N \rightarrow \infty$
- But we can bound error of Normal approximation for fixed N

Hypothesis tests of the weak null

- The finite population CLT tells us that

$$\frac{\hat{\tau}(\mathbf{Z}, \mathbf{Y}) - \mathbb{E}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]}{\sqrt{\text{Var}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]}} \xrightarrow{d} \mathcal{N}(0, 1)$$

- Diff-in-Means is unbiased, so write

$$\frac{\hat{\tau}(\mathbf{Z}, \mathbf{Y}) - \tau}{\sqrt{\text{Var}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]}} \xrightarrow{d} \mathcal{N}(0, 1)$$

- The CLT is an asymptotic results as $N \rightarrow \infty$
- But we can bound error of Normal approximation for fixed N
- Thus, with experiments of at least moderate size and outcomes that aren't too skewed or have extreme outliers,

$$\frac{\hat{\tau}(\mathbf{Z}, \mathbf{Y}) - \tau}{\sqrt{\text{Var}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]}} \stackrel{\text{approx.}}{\sim} \mathcal{N}(0, 1)$$

Hypothesis tests of the weak null

- The finite population CLT tells us that

$$\frac{\hat{\tau}(\mathbf{Z}, \mathbf{Y}) - \mathbb{E}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]}{\sqrt{\text{Var}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]}} \xrightarrow{d} \mathcal{N}(0, 1)$$

- Diff-in-Means is unbiased, so write

$$\frac{\hat{\tau}(\mathbf{Z}, \mathbf{Y}) - \tau}{\sqrt{\text{Var}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]}} \xrightarrow{d} \mathcal{N}(0, 1)$$


- The CLT is an asymptotic results as $N \rightarrow \infty$
- But we can bound error of Normal approximation for fixed N
- Thus, with experiments of at least moderate size and outcomes that aren't too skewed or have extreme outliers,

$$\frac{\hat{\tau}(\mathbf{Z}, \mathbf{Y}) - \tau}{\sqrt{\text{Var}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]}} \stackrel{\text{approx.}}{\sim} \mathcal{N}(0, 1)$$

- This justifies use of standard Normal distribution for hypothesis tests

Hypothesis tests of the weak null

- “Village heads” example:



Hypothesis tests of the weak null

- To test null hypothesis relative to alternative

$H_0 : \tau = \tau_0$ versus either

$H_A : \tau > \tau_0$, $H_A : \tau < \tau_0$ or $H_A : |\tau| > |\tau_0|$

Hypothesis tests of the weak null

- To test null hypothesis relative to alternative

$H_0 : \tau = \tau_0$ versus either

$H_A : \tau > \tau_0$, $H_A : \tau < \tau_0$ or $H_A : |\tau| > |\tau_0|$

- Calculate upper(u), lower(l) or two-sided(t) p-value as

$$p_u = 1 - \Phi \left(\frac{\hat{\tau}(\mathbf{Z}, \mathbf{Y}) - \tau_0}{\sqrt{\widehat{\text{Var}}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]}} \right)$$

$$p_l = \Phi \left(\frac{\hat{\tau}(\mathbf{Z}, \mathbf{Y}) - \tau_0}{\sqrt{\widehat{\text{Var}}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]}} \right)$$

$$p_t = 2 \left(1 - \Phi \left(\frac{|\hat{\tau}(\mathbf{Z}, \mathbf{Y}) - \tau_0|}{\sqrt{\widehat{\text{Var}}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]}} \right) \right)$$

Hypothesis tests of the weak null

- To test null hypothesis relative to alternative

$H_0 : \tau = \tau_0$ versus either

$H_A : \tau > \tau_0$, $H_A : \tau < \tau_0$ or $H_A : |\tau| > |\tau_0|$

- Calculate upper(u), lower(l) or two-sided(t) p-value as

$$p_u = 1 - \Phi \left(\frac{\hat{\tau}(\mathbf{Z}, \mathbf{Y}) - \tau_0}{\sqrt{\widehat{\text{Var}}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]}} \right)$$

$$p_l = \Phi \left(\frac{\hat{\tau}(\mathbf{Z}, \mathbf{Y}) - \tau_0}{\sqrt{\widehat{\text{Var}}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]}} \right)$$

$$p_t = 2 \left(1 - \Phi \left(\frac{|\hat{\tau}(\mathbf{Z}, \mathbf{Y}) - \tau_0|}{\sqrt{\widehat{\text{Var}}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]}} \right) \right)$$

- If p-value is less than size α -level of test, reject. Otherwise, don't
- Note that, since we don't know $\text{Var}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]$, we have used its conservative estimator instead, $\widehat{\text{Var}}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]$

Hypothesis tests of the weak null

Hypothesis tests susceptible to two errors:

- Type I error: Rejecting null hypothesis when it is true
- Type II error: *Failing* to reject null hypothesis when it is false

Hypothesis tests of the weak null

Hypothesis tests susceptible to two errors:

- Type I error: Rejecting null hypothesis when it is true
- Type II error: *Failing* to reject null hypothesis when it is false

A good test controls these errors:

1. Type I error probability is less than or equal to size (α -level) of test
2. Power (1 - type II error probability) is at least as great as α -level
3. Power tends to 1 as $N \rightarrow \infty$

Hypothesis tests of the weak null

- We can prove that tests of weak null satisfy (1) – (3) as $N \rightarrow \infty$
- Thus, when experiments are large, we can often safely use such tests
- But (1) – (3) may not always be satisfied when experiments are small, have skewed outcome distributions or extreme outliers

Confidence intervals

- Equivalence between hypothesis testing and confidence intervals
- Confidence interval is set of null hypotheses we fail to reject

Consider two-sided confidence interval, \mathcal{C}_t :

$$\begin{aligned}\mathcal{C}_t &= \left\{ \tau_0 : \left| \frac{\hat{\tau}(\mathbf{Z}, \mathbf{Y}) - \tau_0}{\sqrt{\widehat{\text{Var}}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]}} \right| \leq z_{1-\alpha/2} \right\} \\&= \left\{ \tau_0 : -z_{1-\alpha/2} \leq \frac{\hat{\tau}(\mathbf{Z}, \mathbf{Y}) - \tau_0}{\sqrt{\widehat{\text{Var}}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]}} \leq z_{1-\alpha/2} \right\} \\&= \left\{ \tau_0 : -z_{1-\alpha/2} \sqrt{\widehat{\text{Var}}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]} \leq \hat{\tau}(\mathbf{Z}, \mathbf{Y}) - \tau_0 \leq z_{1-\alpha/2} \sqrt{\widehat{\text{Var}}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]} \right\} \\&= \left\{ \tau_0 : -\hat{\tau}(\mathbf{Z}, \mathbf{Y}) - z_{1-\alpha/2} \sqrt{\widehat{\text{Var}}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]} \leq -\tau_0 \leq -\hat{\tau}(\mathbf{Z}, \mathbf{Y}) + z_{1-\alpha/2} \sqrt{\widehat{\text{Var}}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]} \right\} \\&= \left\{ \tau_0 : \hat{\tau}(\mathbf{Z}, \mathbf{Y}) + z_{1-\alpha/2} \sqrt{\widehat{\text{Var}}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]} \geq \tau_0 \geq \hat{\tau}(\mathbf{Z}, \mathbf{Y}) - z_{1-\alpha/2} \sqrt{\widehat{\text{Var}}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]} \right\} \\&= \left\{ \tau_0 : \hat{\tau}(\mathbf{Z}, \mathbf{Y}) - z_{1-\alpha/2} \sqrt{\widehat{\text{Var}}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]} \leq \tau_0 \leq \hat{\tau}(\mathbf{Z}, \mathbf{Y}) + z_{1-\alpha/2} \sqrt{\widehat{\text{Var}}[\hat{\tau}(\mathbf{Z}, \mathbf{Y})]} \right\}\end{aligned}$$

Appendix: Estimator of Difference-in-Means estimator's variance

Neyman's conservative estimator of the Difference-in-Means estimator's variance is

$$\widehat{\text{Var}} [\hat{\tau}(\mathbf{Z}, \mathbf{Y})] = \frac{N}{N-1} \left(\frac{\hat{\sigma}_{y_C}^2}{n_C} + \frac{\hat{\sigma}_{y_T}^2}{n_T} \right),$$

where

$$\hat{\sigma}_{y_C}^2 = \left(\frac{N-1}{n(n_C-1)} \right) \sum_{i:Z_i=0}^N (y_{Ci} - \hat{\mu}_{y_C})^2$$

$$\hat{\sigma}_{y_T}^2 = \left(\frac{N-1}{n(n_T-1)} \right) \sum_{i:Z_i=1}^n (y_{Ti} - \hat{\mu}_{y_T})^2$$

$$\hat{\mu}_{y_C} = \left(\frac{1}{n_C} \right) \sum_{i=1}^N (1 - Z_i) y_{Ci}$$

$$\hat{\mu}_{y_T} = \left(\frac{1}{n_T} \right) \sum_{i=1}^N Z_i y_{Ti}$$